

*Experimental Particle Physics Seminar
University of Pennsylvania, October 17, 2007*

Search for Flavor Changing Neutral Currents in Top Quark Decays at CDF



*Ulrich Husemann
Yale University*





Outline of the Talk

What are Flavor Changing Neutral Currents?

The CDF Experiment at the Tevatron

Top Quark Physics at CDF

Search for FCNC in Top Quark Decays

Summary & Conclusions



Outline of the Talk




What are Flavor Changing Neutral Currents?

The CDF Experiment at the Tevatron

Top Quark Physics at CDF


Search for FCNC in Top Quark Decays

Summary & Conclusions





Fermilab 95-759

- **Matter** in the standard model:
12 fermions in three generations
 - Six quarks and their anti-particles
 - Six leptons and their anti-particles
- **Forces** in the standard model:
 - **Strong** force (carrier: gluon)
 - **Electroweak** force (carriers: photon, W^\pm bosons, Z boson)
- Interactions can be described by “**currents**” coupling to gauge bosons, e.g. electromagnetic current




$$\mathcal{L}^{\text{em}} = -q \bar{e} \gamma_\mu e A^\mu \equiv j_\mu^{\text{em}} A^\mu$$

- Flavor changing neutral current (FCNC) interactions:
 - Transition from a quark of **flavor A** and **charge Q** to quark of **flavor B** with the **same charge Q**
 - Examples: $b \rightarrow s\gamma$, $t \rightarrow cH$, ...
- 1960s: only three light quarks (u,d,s) known, **mystery** in neutral kaon system:



10^8 times smaller than...?



- Solution: “**GIM Mechanism**” (Glashow, Iliopoulos, Maiani, 1970)
 - Fourth quark** needed for cancellation in box diagram: prediction of charm quark
 - Cancellation would be **exact** if all quarks had the **same mass**: estimate of charm quark mass

Standard model: **no FCNC at Lagrangian level**

- Massless theory: weak neutral current is **flavor-diagonal**

$$J_\mu^{\text{NC}} = J_\mu^3 - 2 \sin^2 \theta_W j_\mu^{\text{em}} = \bar{u} \left[\frac{1}{2} \gamma_\mu (1 - \gamma_5) - \frac{4}{3} \sin^2 \theta_W \gamma_\mu \right] u - \bar{d} \left[\frac{1}{2} \gamma_\mu (1 - \gamma_5) - \frac{2}{3} \sin^2 \theta_W \gamma_\mu \right] d$$

- Quark masses via **Higgs mechanism**:

- Eigenstates of electroweak interactions are **not mass eigenstates**

$$\mathcal{L}_{\text{Yuk}} = \underbrace{-m_u^{\alpha\beta} \bar{u}'_\alpha u'^\beta - m_d^{\alpha\beta} \bar{d}'_\alpha d'^\beta}_{\text{Mass Terms}} - \underbrace{\frac{1}{\sqrt{2}} f_u^{\alpha\beta} \bar{u}'_\alpha h(x) u'^\beta - \frac{1}{\sqrt{2}} f_d^{\alpha\beta} \bar{d}'_\alpha h(x) d'^\beta}_{\text{Higgs Couplings}} + \text{h.c.}$$

- Unitary transformation of Lagrangian to mass basis, i.e. for physical particles:

$$\begin{aligned} \bar{u}_L &= \bar{u}'_L \mathbf{U}_L^u & u_R &= \mathbf{U}_R^{u\dagger} u'_R & \mathbf{m}_u &= \mathbf{U}_L^{u\dagger} \mathbf{m}'_u \mathbf{U}_R^u \\ \bar{d}_L &= \bar{d}'_L \mathbf{U}_L^d & d_R &= \mathbf{U}_R^{d\dagger} d'_R & \mathbf{m}_d &= \mathbf{U}_L^{d\dagger} \mathbf{m}'_d \mathbf{U}_R^d \end{aligned}$$

- Kinetic terms: **unchanged**
- Higgs couplings proportional to mass terms: **no flavor changing Higgs couplings**
- Neutral currents have **same structure as kinetic terms**: unchanged → **no FCNC**

- Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix (obtained from transformation of charged current to mass basis):

$$J_{\mu}^{\text{CC}} = \bar{u}' \left(\frac{1}{2} \gamma_{\mu} (1 - \gamma_5) \right) d' = \bar{u}'_L \gamma_{\mu} d'_L = \bar{u}_L U_L^{u\dagger} \gamma_{\mu} U_L^d d_L = \bar{u}_L \gamma_{\mu} \mathbf{V}_{\text{CKM}} d_L,$$

- CKM matrix: unitary 3×3 matrix

$$\mathbf{V}_{\text{CKM}} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \quad \text{with}$$


$$\mathbf{V}_{\text{CKM}} \cdot \mathbf{V}_{\text{CKM}}^{\dagger} = \mathbf{V}_{\text{CKM}}^{\dagger} \cdot \mathbf{V}_{\text{CKM}} = \mathbf{1}$$

yields unitarity relations,
e.g. the unitary triangle of
flavor physics (1st vs. 3rd column)

$$V_{ud}^* V_{ub} + V_{cd}^* V_{cb} + V_{td}^* V_{tb} = 0$$

or (used in top FCNC):

$$V_{cd}^* V_{td} + V_{cs}^* V_{ts} + V_{cb}^* V_{tb} = 0$$



FCNC are allowed via higher order mechanisms such as **penguin diagrams**, but heavily suppressed

- Suppression mechanism 1: GIM**

- Penguin matrix element depends on universal functions of **single parameter** $x_i = m_i^2/m_W^2$

$$\mathcal{M} \propto F(x_d) V_{cd}^* V_{td} + F(x_s) V_{cs}^* V_{ts} + F(x_b) V_{cb}^* V_{tb},$$

- Compare to CKM unitarity relation:


$$V_{cd}^* V_{td} + V_{cs}^* V_{ts} + V_{cb}^* V_{tb} = 0$$

Exact cancellation if masses of b, s, and d quarks were the same


- Quark masses more similar for down-type than for up-type: **top FCNC more strongly suppressed than bottom FCNC**, e.g. $\text{BR}(t \rightarrow Zq) \approx 10^{-14}$ vs. $\text{BR}(b \rightarrow sy) \approx 10^{-4}$

- Suppression mechanism 2: smallness of relevant CKM matrix elements**

$$|V_{cd}^* V_{td}| \approx 0.002, \quad |V_{cs}^* V_{ts}| \approx 0.04, \quad |V_{cb}^* V_{tb}| \approx 0.04$$




Universal Function for $t \rightarrow \gamma c$



FCNC are **enhanced** in many models of physics beyond the SM

- Enhancement mechanisms:
 - FCNC interactions at **tree level**
 - Weaker GIM cancellation by **new particles in loop corrections**
- Examples:
 - **New quark singlets**: Z couplings not flavor-diagonal \rightarrow tree level FCNC
 - **Two Higgs doublet** models: modified Higgs mechanism
 - Flavor changing Higgs couplings allowed at tree level
 - Virtual Higgs in loop corrections
 - **Supersymmetry**: gluino/neutralino and squark in loop corrections




Model	$\text{BR}(t \rightarrow Zq)$
Standard Model	$\mathcal{O}(10^{-14})$
$q = 2/3$ Quark Singlet	$\mathcal{O}(10^{-4})$
Two Higgs Doublets	$\mathcal{O}(10^{-7})$
MSSM	$\mathcal{O}(10^{-6})$
R-Parity violating SUSY	$\mathcal{O}(10^{-5})$

[after J.A. Aguilar-Saavedra,
Acta Phys. Polon **B35** (2004) 2695]

FCNC are **enhanced** in many models of physics beyond the SM

- Enhancement mechanisms:
 - FCNC interactions at **tree level**
 - Weaker GIM cancellation by **new particles in loop corrections**
- Examples:
 - **New quark singlets**: Z couplings not flavor-diagonal \rightarrow tree level FCNC
 - **Two Higgs doublet** models: modified Higgs mechanism
 - Flavor changing Higgs couplings allowed at tree level
 - Virtual Higgs in loop corrections
 - **Supersymmetry**: gluino/neutralino and squark in loop corrections




Model	$\text{BR}(t \rightarrow Zq)$
Standard Model	$\mathcal{O}(10^{-14})$
$q = 2/3$ Quark Singlet	$\mathcal{O}(10^{-4})$
Two Higgs Doublets	$\mathcal{O}(10^{-7})$
MSSM	$\mathcal{O}(10^{-6})$
R-Parity violating SUSY	$\mathcal{O}(10^{-5})$

[after J.A. Aguilar-Saavedra,
Acta Phys. Polon **B35** (2004) 2695]

FCNC are **enhanced** in many models of physics beyond the SM

- Enhancement mechanisms:
 - FCNC interactions at **tree level**
 - Weaker GIM cancellation by **new particles in loop corrections**
- Examples:
 - **New quark singlets**: Z couplings not flavor-diagonal \rightarrow tree level FCNC
 - **Two Higgs doublet** models: modified Higgs mechanism
 - Flavor changing Higgs couplings allowed at tree level
 - Virtual Higgs in loop corrections
 - **Supersymmetry**: gluino/neutralino and squark in loop corrections






Model	$\text{BR}(t \rightarrow Zq)$
Standard Model	$\mathcal{O}(10^{-14})$
$q = 2/3$ Quark Singlet	$\mathcal{O}(10^{-4})$
Two Higgs Doublets	$\mathcal{O}(10^{-7})$
MSSM	$\mathcal{O}(10^{-6})$
R-Parity violating SUSY	$\mathcal{O}(10^{-5})$



[after J.A. Aguilar-Saavedra,
Acta Phys. Polon **B35** (2004) 2695]

- Experimental tests of FCNC interactions: **sensitive probes of new physics**
 - Any signal above SM expectations would indicate new physics
 - Measurements constrain allowed phase space for new physics models
- Two types of searches for FCNC in the top sector:
 - Search for **single top production** (LEP, HERA, DØ)
 - Search for **top quark decay via FCNC** (CDF)
- Experiments usually report limits on
 - **Branching fractions** for specific processes, e.g. $\text{BR}(t \rightarrow Zq)$
 - **Coupling parameters** of effective Lagrangian, e.g. for tZq coupling

$$\mathcal{L}_{\text{eff}} = -\frac{g}{2 \cos \theta_W} \cdot \kappa \cdot \left(x_L \cdot \bar{q}_L \gamma_\mu t_L + x_R \cdot \bar{q}_R \gamma_\mu t_R \right) Z^\mu + \dots$$



- **CDF Run I search:**
F. Abe *et al.*, PRL **80** (1998) 2525.
 - Signature: $Z \rightarrow l^+ l^- + 4 \text{ jets}$ (1 b-jet)
 - Limit on $\text{BR}(t \rightarrow Zq)$: **33%**
- **LEP searches:**
P. Achard *et al.* (L3), Phys. Lett. **B549** (2002) 290.
G. Abbiendi *et al.* (Opal), Phys. Lett. **B521** (2001) 181.
J. Abdallah *et al.* (Delphi), Phys. Lett. **B590** (2004) 21.
A. Heister *et al.* (Aleph), Phys. Lett. **B453** (2002) 173.
 - Hadronic top decay (4 jets) or semileptonic top decay (2 jets & lepton)
 - Very similar results among all LEP experiments, best limit on $\text{BR}(t \rightarrow Zq)$: **13.7% (L3)**
- **HERA searches:**
A. Aktas *et al.* (H1), Eur. Phys. J. **C33** (2004) 9.
S. Chekanov *et al.* (ZEUS), Phys. Lett. **B559** (2003) 153.
 - Hadronic top decay (3 jets) or semileptonic top decay (lepton & jet)
 - Most sensitive to $t\gamma q$ vertex, preference for u over c quarks (proton sea)






The H1 result caused some excitement:

Abstract. [...] In the leptonic channel, 5 events are found while 1.31 ± 0.22 events are expected from the Standard Model background. In the hadronic channel, no excess above the expectation for Standard Model processes is found. [...]



- Study **single Top production** via FCNC:



- Artificial neural network** to discriminate signal from background
- World's best limit** on t-c-g and t-u-g couplings $(\kappa/\Lambda)^2 \rightarrow$ previous limits improved by order of magnitude

$$(\kappa_g^c/\Lambda)^2 < 0.023 \text{ TeV}^{-2} \quad (95\% \text{ C.L.})$$

$$(\kappa_g^u/\Lambda)^2 < 0.0014 \text{ TeV}^{-2} \quad (95\% \text{ C.L.})$$

[V. M. Abazov *et al.*, hep-ex/0702005, submitted to PRL]



Outline of the Talk

What are Flavor Changing Neutral Currents?


The CDF Experiment at the Tevatron

Top Quark Physics at CDF

Search for FCNC in Top Quark Decays

Summary & Conclusions


Fermi National Accelerator Laboratory – Aerial View




[Fermilab Visual Media Service]

- Proton-antiproton collider:
 $\sqrt{s} = 1.96 \text{ TeV}$
- 36×36 bunches, collisions every 396 ns
- Record instantaneous peak luminosity:
 $292 \mu\text{b}^{-1} \text{ s}^{-1}$
($1 \mu\text{b}^{-1} \text{ s}^{-1} = 10^{30} \text{ cm}^{-2} \text{ s}^{-1}$)
- Luminosity goal:
5.5–6.5 fb^{-1} of integrated luminosity by 2009,
running in 2010 currently under discussion
- Two multi-purpose detectors: CDF and DØ


- Tevatron continues to perform very well:
 - More than 3 fb^{-1} delivered up to Summer 2007 shutdown
 - More than 2.5 fb^{-1} recorded by CDF




The CDF II Detector




The CDF II Detector



The CDF II Detector



The CDF II Detector




- Cylindrical coordinate system:
 - θ : polar angle w.r.t. to proton direction
 - ϕ : azimuthal angle
 - Pseudorapidity:

$$\eta = -\ln \tan(\theta/2)$$
 - Transverse energy:

$$\vec{E}_T = \sum_{\text{cal towers}} E_i(\sin \theta_i, \phi_i)$$
 - Missing transverse energy (“MET”):

$$\vec{\cancel{E}}_T = -\sum_{\text{jets}} \vec{E}_T - \sum_{\text{leptons}} \vec{p}_T$$





Outline of the Talk



What are Flavor Changing Neutral Currents?

The CDF Experiment at the Tevatron

Top Quark Physics at CDF

Search for FCNC in Top Quark Decays

Summary & Conclusions


Brief history of top quark discovery:

- **1977**: Y discovery – bottom quark
- **1980s**: Searches for “light” top (mass smaller than W boson mass) as isospin partner of bottom at PETRA, Sp \bar{p} S, LEP, CDF Run 0
- **1992/3**: Tevatron Run I starts, first indications for top quark production
- **March 2, 1995**: CDF and DØ announce top quark discovery

The Discovery of the Top Quark

Finding the sixth quark involved the world's most energetic collisions and a cast of thousands

by Tony M. Liss and Paul L. Tipton



[Scientific American, September 1997]

VIOLENT COLLISION between a proton and an antiproton (*center*) creates a top quark (*red*) and an antitop (*blue*). These decay to other particles, typically producing a number of jets and possibly an electron or positron.

- The top is **heavy**: $m_t \approx 170 \text{ GeV}/c^2$ ($40 \times m_b$, approx. mass of gold atom)
- Mass **close to scale of electroweak symmetry breaking** (EWSB), top Yukawa coupling $f \approx 1$:

$$\mathcal{L}_{\text{Yuk},t} = f \frac{v}{\sqrt{2}} \bar{t}_L t_R \equiv m_t \bar{t}_L t_R$$

(vacuum expectation value of Higgs field: $v/\sqrt{2} \approx 178 \text{ GeV}$)


→ Important role in EWSB models

- Top is the **only “free” quark**: lifetime shorter than hadronization time

$$\tau = \frac{1}{\Gamma} \approx \frac{1}{1.5 \text{ GeV}} < \frac{1}{\Lambda_{\text{QCD}}} \approx \frac{1}{0.2 \text{ GeV}}$$


→ No spectroscopy of bound states

→ Spin transferred to decay products




[Fermilab Visual Media Services]


Top production is **rare**: one top quark pair produced every 10 billion collisions



85% $q\bar{q} \rightarrow t\bar{t}$:




15% $gg \rightarrow t\bar{t}$:



$W^- \rightarrow$		hadrons	τ	μ	e
$W^+ \rightarrow$	hadrons	All Hadronic (S/B ≈ 0.04)	Lepton+ τ	Lepton + Jets (S/B ≈ 1)	
	τ	Lepton+ τ			
	μ	Lepton + Jets (S/B ≈ 1)		Dilepton (S/B ≈ 3)	



- Top decay in the Standard Model: $t \rightarrow Wb$ (BR $\approx 100\%$)
- $t\bar{t}$ decay signatures characterized by W decays:
 - All-Hadronic (45% of all decays)
 - Lepton+Jets (30% of all decays)
 - Dilepton (5% of all decays)
- Main background process: production of **W bosons in association with Jets**
- $t\bar{t}$ events contain two b quarks: **b quark identification** (“b-tagging”) crucial

- **SecVtx** b-tagging algorithm: based on significance of 2D impact parameter




- CDF's **single most precise top cross section measurement**: Lepton + Jets channel with SecVtx b-tags
- Results (CDF Public Note 8795)

Single B-Tag	$\sigma_{tt} = 8.2 \pm 0.5 \text{ (stat)} \pm 0.8 \text{ (syst)} \pm 0.5 \text{ (lum)} \text{ pb}$
Double B-Tag	$\sigma_{tt} = 8.8 \pm 0.8 \text{ (stat)} \pm 1.2 \text{ (syst)} \pm 0.5 \text{ (lum)} \text{ pb}$



- From top discovery in 1995 to **precision physics** in 2007:
 - Dataset: 1000s of top events
 - Mass & cross section very precisely measured
 - Evidence for single top production
- Broad program to study **properties** of the top quark: production, decay, quantum numbers, ...
- Measurements of top properties try to answer:

Is the top really the Standard Model top?





Outline of the Talk



What are Flavor Changing Neutral Currents?

The CDF Experiment at the Tevatron

Top Quark Physics at CDF


Search for FCNC in Top Quark Decays

Summary & Conclusions

- Basic question: how often do top quarks decay into Zq ?
→ set limit on **branching fraction**
 $BR(t \rightarrow Zq)$
- Selection of decay channels for $t\bar{t} \rightarrow Zq Wb$:
 - **$Z \rightarrow$ charged leptons**: very clean signature, lepton trigger
 - **$W \rightarrow$ hadrons**: large branching fractions, no neutrinos
→ event can be fully reconstructed
 - Final signature: **$Z + \geq 4$ jets**
- **Analysis Outline:**
 - Baseline Event Selection
 - Initial Background Estimate
 - Optimization of Event Selection
 - Systematic Uncertainties
 - Final Limit Calculation


Z Decay Modes:


- $Z \rightarrow \nu\nu$
- $Z \rightarrow ee/\mu\mu$
- $Z \rightarrow \tau\tau$
- $Z \rightarrow$ hadrons



W Decay Modes:

- $W \rightarrow l\nu$
- $W \rightarrow \tau\nu$
- $W \rightarrow$ hadrons






- Event signature: $Z \rightarrow l^+ l^- + 4 \text{ jets}$
- Motivation for blind analysis: **avoid biases** by looking into the data too early
- Blinding & unblinding strategy:
 - Initial blinded region: $Z + \geq 4 \text{ jets}$
 - Later: add **control region** in $Z + \geq 4 \text{ jets}$ from kinematic constraints
 - Optimization of event selection, prediction of backgrounds, and systematic uncertainties on **data control regions and Monte Carlo (MC) simulation only**
 - Very last step: “**opening the box**”, i.e. look into signal region in data

- Monte Carlo (MC) simulation of FCNC decay $t \rightarrow Zq$ with **PYTHIA**

- $t \rightarrow Zq$ vertex **unknown** to PYTHIA
- Decay generated **flat in $\cos \theta^*$**
(angle between top boost direction and lepton of same charge sign from Z decay, in Z rest frame)





- Solution: **reweight** according to expectation from standard model Higgs mechanism:

$$\frac{d\sigma}{d\cos(\theta^*)} = f^0 \cdot \frac{3}{4} (1 - \cos(\theta^*))^2 + f^- \cdot \frac{3}{8} (1 - \cos(\theta^*))^2 + f^+ \cdot \frac{3}{8} (1 + \cos(\theta^*))^2$$


with $f^0 = 0.65$ (“longitudinal”), $f^- = 0.35$ (“left-handed”), $f^+ = 0$ (“right-handed”)


- Main FCNC signal sample: one top decays $t \rightarrow Zc$, other decays $t \rightarrow Wb$
 - Additional sample required for decay $t \rightarrow Zu$
 - Additional sample for “double FCNC” events, i.e. both tops decay via FCNC $t \rightarrow Zq$






Two leptons
(ee or $\mu\mu$) with
opposite
charge, form
Z boson









- **Simple trigger:** single electron or muon, transverse momentum $>18 \text{ GeV}/c$
- Sharp Z resonance, good lepton momentum resolution
→ cut on lepton pair invariant mass: $76 \text{ GeV}/c^2 < M_{ll} < 106 \text{ GeV}/c^2$
- Enhancing the Z acceptance:
 - Tracking systems have better coverage than calorimeter and muon detectors: allow second lepton to be **isolated track**
→ **doubles acceptance** w.r.t. standard lepton selection
 - Electron tracks lose momentum via bremsstrahlung: **correct track momentum** with calorimeter energy
→ **3% more** dielectron pairs

Bremsstrahlung Recovery: $Z \rightarrow e$ track



- FCNC: **four jet assignments**
 - 1 b-jet from $t \rightarrow Wb$ decay
 - 2 jets from subsequent W decay
 - 1 jet from $t \rightarrow Zq$ decay
- For all **12 possible combinations** of first four jets in the event:
 1. Combine jets #1 and #2 to W, calculate **invariant mass** $m_{W,rec}$
 2. Vary momenta of jets #1 and #2 within their resolutions to match PDG W mass ("**fix W mass**")
 3. Add jet #3 to fixed W, calculate **invariant mass** $m_{t \rightarrow Wb,rec}$
 4. Vary momenta of leptons within their resolutions to match PDG Z mass ("**fix Z mass**")
 5. Add jet #4 to fixed Z, calculate **invariant mass** $m_{t \rightarrow Zq,rec}$



- Pick combination with lowest

$$\chi^2 = \left(\frac{m_{W,rec} - m_{W,PDG}}{\sigma_{W,rec}} \right)^2 + \left(\frac{m_{t \rightarrow Wb,rec} - m_{t,PDG}}{\sigma_{t \rightarrow Wb}} \right)^2 + \left(\frac{m_{t \rightarrow Zq,rec} - m_{t,PDG}}{\sigma_{t \rightarrow Zq}} \right)^2$$

- Widths reflect mass resolutions as measured in MC simulation:
 $\sigma_{W,rec} = 15 \text{ GeV}/c^2$,
 $\sigma_{t \rightarrow Wb,rec} = 24 \text{ GeV}/c^2$
 $\sigma_{t \rightarrow Zq,rec} = 21 \text{ GeV}/c^2$

- How do you search for a signal that is likely not there?
Understand the background!
- Standard model processes that can mimic $Z + \geq 4$ jets signature:
 - **Z+Jets**: Z boson production in association with jets
→ **dominant background** for top FCNC search, most difficult to estimate
 - **Standard model $t\bar{t}$** production
→ **small** background
 - **Dibosons**: WZ and ZZ diboson production → **small** background
 - **W+Jets, WW**: negligible
- Top FCNC background estimate: mixture of data driven techniques and MC predictions

- How do you search for a signal that is likely not there?
Understand the background!
- Standard model processes that can mimic $Z + \geq 4$ jets signature:
 - **Z+Jets**: Z boson production in association with jets
→ **dominant background** for top FCNC search, most difficult to estimate
 - **Standard model $t\bar{t}$** production
→ **small** background
 - **Dibosons**: WZ and ZZ diboson production → **small** background
 - **W+Jets, WW**: negligible
- Top FCNC background estimate: mixture of data driven techniques and MC predictions

Standard Model $t\bar{t}$ Production



- Small background: no real Z, need extra jets from gluon radiation and/or “fake lepton”
- Dilepton channel
($t\bar{t} \rightarrow Wb Wb \rightarrow l\nu b l\nu b$):
dilepton invariant mass can fall into Z mass window
- Lepton+Jets channel
($t\bar{t} \rightarrow Wb Wb \rightarrow l\nu b qq'b$):
misreconstruct one jet as a lepton (“fake”), invariant mass of lepton and fake lepton can fall into Z mass window
- Large fraction of heavy flavor jets: more important in b-tagged samples
- Estimated from MC simulation

- How do you search for a signal that is likely not there?
Understand the background!
- Standard model processes that can mimic $Z + \geq 4$ jets signature:
 - **Z+Jets**: Z boson production in association with jets
→ **dominant background** for top FCNC search, most difficult to estimate
 - **Standard model $t\bar{t}$** production
→ **small** background
 - **Dibosons**: WZ and ZZ diboson production → **small** background
 - **W+Jets, WW**: negligible
- Top FCNC background estimate: mixture of data driven techniques and MC predictions

Diboson Production: WZ, ZZ

- Small background (similar in size to standard model $t\bar{t}$ production)
 - Small cross section but real Z
 - Need extra jets from gluon radiation
- ZZ: Heavy flavor contribution from $Z \rightarrow b\bar{b}$ decay
- Estimated from MC simulation

- MC tool for Z+Jets: **ALPGEN**
 - Modern MC generator for multiparticle final states
 - “MLM matching”** prescription to remove overlap between jets from matrix element and partons showers
- Comparing ALPGEN with data:
 - Leading order generator: **no absolute prediction** for cross section
 - Underestimate** of number of events with large jet multiplicities, **large uncertainties**
- Our strategy: only **shapes** of kinematic distributions **from MC**, **normalization** from **control samples in data**





- **Mass χ^2** : combination of mass constraints – best discriminator

$$\chi^2 = \left(\frac{m_{W,\text{rec}} - m_{W,\text{PDG}}}{\sigma_{W,\text{rec}}} \right)^2 + \left(\frac{m_{t \rightarrow Wb,\text{rec}} - m_{t,\text{PDG}}}{\sigma_{t \rightarrow Wb}} \right)^2 + \left(\frac{m_{t \rightarrow Zq,\text{rec}} - m_{t,\text{PDG}}}{\sigma_{t \rightarrow Zq}} \right)^2$$



- **Transverse mass**: FCNC top decays are more central than Z+jets

$$M_T = \sqrt{(\sum E_T)^2 - (\sum \vec{p}_T)^2}$$

- **Jet transverse energies**: FCNC signal has four “hard” jets, background processes: jets have to come from gluon radiation



- **Advantage** of requiring b-tag:
Better discrimination against main Z+jets background
(heavy flavor backgrounds rather small: SM $t\bar{t}$, $Zb\bar{b}$ + jets)
- **Disadvantage:**
Reduction of data sample size
- **Solution: use both!**
 - **Split sample** in tagged and anti-tagged
 - **Optimize cuts individually** for tagged and anti-tagged samples
 - **Combine samples** in limit calculation
- Main difficulty of this approach:
event migration between samples
 - Systematics may be **correlated or anti-correlated** between samples
 - Taken into account in limit calculation



- Question: **best choice for cut values?**
- Goal: derive **limit** on branching fraction of FCNC process $t \rightarrow Zq$
- No prediction for amount of signal: “signal over background” et al. do not work
- Solution: optimize cuts for **best expected limit** (assuming no signal)


$$\sum_{n_{\text{obs}}} P(n_{\text{obs}} | n_{\text{back}}) \cdot \text{Lim}(n_{\text{obs}} | A, n_{\text{back}})$$

- P: Poisson probability
- L: any limit calculation method
- Our analysis: faster **objective Bayesian** limits for optimization, “better” **Feldman-Cousins** limits for final result (both including systematic uncertainties)
- Correlations among variables: **multi-dimensional** optimization


Final Event Selection

Kinematic Variable	Optimized Cut
Z Mass	$\in [76, 106] \text{ GeV}/c^2$
Leading Jet E_T	$> 40 \text{ GeV}$
Second Jet E_T	$> 30 \text{ GeV}$
Third Jet E_T	$> 20 \text{ GeV}$
Fourth Jet E_T	$> 15 \text{ GeV}$
Transverse Mass	$> 200 \text{ GeV}$
$\sqrt{\chi^2}$	< 1.6 (b -tagged) < 1.35 (anti-tagged)

Expected 95% C.L. Upper Limit
on $\text{BR}(t \rightarrow Zq)$: $7.1\% \pm 3.0\%$




- Total background prediction from control region in data:
 130 ± 28 events
- Tail of mass χ^2 distribution
- Average of cuts at $\sqrt{\chi^2} = 3.0, 3.2$
- Tagging rate: **$15\% \pm 4\%$**
- Tail of mass χ^2 : $16\% \pm 7\%$ (small sample \rightarrow large uncertainties)
- MC prediction of tagging rate: 11% (but: 30% too low for $Z+\leq 3$ Jets)
- Template fit of MC tagging probabilities vs. number of jets: 14%



Source	Without b -tag	Loose SECVTX b -tag
Z+Jets	123.3 ± 28	17.6 ± 6
Standard Model $t\bar{t}$	2.4 ± 0.3	1.7 ± 0.2
Diboson (WZ, ZZ)	4.3 ± 0.2	0.7 ± 0.1
WW, W+Jets	< 0.1	negligible
Total Backgrounds:	130 ± 28	20 ± 6

- **Question:** how to get from event counts to limit on $BR(t \rightarrow Zq)$?
- **Circular dependency #1:** Limit calculation requires knowledge of signal acceptance, but signal acceptance depends on limit
- **Circular dependency #2:** Measure limit on fraction of $t\bar{t}$ production cross section, but cross section changes with changing FCNC contribution
- **Solution:** “running acceptance” – functional form of above dependencies implemented in limit machinery
 - Signal acceptance **dynamically adjusted** as a function of $BR(t \rightarrow Zq)$
 - Signal **normalized to measured $t\bar{t}$ production cross section** measurement
 - $t\bar{t}$ cross section **re-interpreted** as a function of $BR(t \rightarrow Zq)$ to allow for FCNC contribution



- Signal count: probability for one or both tops to decay via FCNC
 $\mathcal{P}(t\bar{t} \rightarrow ZcWb, ZcZc, \dots)$
- Normalization to **double-tagged** $t\bar{t}$ cross section measurement:
 - Double-tagged: **smallest overlap** between acceptances
 - Luminosity uncertainties **cancel**, other uncertainties reduced

$$\begin{aligned}
 \mathcal{B}_Z &\equiv \mathcal{B}(t \rightarrow Zc) = 1 - \mathcal{B}(t \rightarrow Wb) \\
 \mathcal{A}_{WZ} &\equiv \text{FCNC Acceptance} \\
 \mathcal{A}_{ZZ} &\equiv \text{Double FCNC Acceptance} \\
 \mathcal{A}_{LJ_{WW}} &\equiv \text{L+J Acceptance for SM } t\bar{t} \\
 \mathcal{A}_{LJ_{WZ}} &\equiv \text{L+J Acceptance for FCNC} \\
 \mathcal{A}_{LJ_{ZZ}} &\equiv \text{L+J Acceptance for Double FCNC} \\
 K_{ZZ/WZ} &\equiv \mathcal{A}_{ZZ} / \mathcal{A}_{WZ} \\
 \mathcal{R}_{WZ/WW} &\equiv \mathcal{A}_{LJ_{WZ}} / \mathcal{A}_{LJ_{WW}} \\
 \mathcal{R}_{ZZ/WW} &\equiv \mathcal{A}_{LJ_{ZZ}} / \mathcal{A}_{LJ_{WW}}
 \end{aligned}$$

Acceptance Master Formula:

$$N_{\text{signal}} = [(\mathcal{P}(t\bar{t} \rightarrow WbZc) \cdot \mathcal{A}_{WZ}) + (\mathcal{P}(t\bar{t} \rightarrow ZcZc) \cdot \mathcal{A}_{ZZ})] \cdot \sigma_{t\bar{t}} \cdot \int \mathcal{L} dt$$

... 1/2 page of algebra...

$$= \mathcal{B}_Z \cdot \underbrace{(N_{LJ} - B_{LJ})}_{\text{L+J yield}} \cdot \underbrace{\frac{\mathcal{A}_{WZ}}{\mathcal{A}_{LJ_{WW}}}}_{\text{Acc. Ratio}} \cdot \underbrace{\frac{(2 \cdot (1 - \mathcal{B}_Z) + K_{ZZ/WZ} \cdot \mathcal{B}_Z)}{(1 - \mathcal{B}_Z)^2 + 2\mathcal{B}_Z \cdot (1 - \mathcal{B}_Z) \cdot \mathcal{R}_{WZ/WW} + \mathcal{B}_Z^2 \cdot \mathcal{R}_{ZZ/WW}}}_{\text{"Running" Acceptance Correction}}$$

- Signal count: probability for one or both tops to decay via FCNC
 $\mathcal{P}(t\bar{t} \rightarrow ZcWb, ZcZc, \dots)$
- Normalization to **double-tagged** $t\bar{t}$ cross section measurement:
 - Double-tagged: **smallest overlap** between acceptances
 - Luminosity uncertainties **cancel**, other uncertainties reduced

$$\begin{aligned}
 \mathcal{B}_Z &\equiv \mathcal{B}(t \rightarrow Zc) = 1 - \mathcal{B}(t \rightarrow Wb) \\
 \mathcal{A}_{WZ} &\equiv \text{FCNC Acceptance} \\
 \mathcal{A}_{ZZ} &\equiv \text{Double FCNC Acceptance} \\
 \mathcal{A}_{LJ_{WW}} &\equiv \text{L+J Acceptance for SM } t\bar{t} \\
 \mathcal{A}_{LJ_{WZ}} &\equiv \text{L+J Acceptance for FCNC} \\
 \mathcal{A}_{LJ_{ZZ}} &\equiv \text{L+J Acceptance for Double FCNC} \\
 K_{ZZ/WZ} &\equiv \mathcal{A}_{ZZ} / \mathcal{A}_{WZ} \\
 \mathcal{R}_{WZ/WW} &\equiv \mathcal{A}_{LJ_{WZ}} / \mathcal{A}_{LJ_{WW}} \\
 \mathcal{R}_{ZZ/WW} &\equiv \mathcal{A}_{LJ_{ZZ}} / \mathcal{A}_{LJ_{WW}}
 \end{aligned}$$

Acceptance Master Formula:

$$N_{\text{signal}} = [(\mathcal{P}(t\bar{t} \rightarrow WbZc) \cdot \mathcal{A}_{WZ}) + (\mathcal{P}(t\bar{t} \rightarrow ZcZc) \cdot \mathcal{A}_{ZZ})] \cdot \sigma_{t\bar{t}} \cdot \int \mathcal{L} dt$$

... 1/2 page of algebra...

$$= \underbrace{\mathcal{B}_Z \cdot (N_{LJ} - B_{LJ})}_{\text{L+J yield}} \cdot \underbrace{\frac{\mathcal{A}_{WZ}}{\mathcal{A}_{LJ_{WW}}}}_{\text{Acc. Ratio}} \cdot \underbrace{\frac{(2 \cdot (1 - \mathcal{B}_Z) + K_{ZZ/WZ} \cdot \mathcal{B}_Z)}{(1 - \mathcal{B}_Z)^2 + 2\mathcal{B}_Z \cdot (1 - \mathcal{B}_Z) \cdot \mathcal{R}_{WZ/WW} + \mathcal{B}_Z^2 \cdot \mathcal{R}_{ZZ/WW}}}_{\text{"Running" Acceptance Correction}}$$


- Signal systematic evaluated for acceptance ratio A_{WZ}/A_{LJ}
- Distinguish uncertainties: **correlated** or **anti-correlated** between selections
 - **Correlated**: shift anti-tagged & tagged selection into same direction (e.g. lepton SF)
 - **Anti-correlated**: shift anti-tagged & tagged into opposite directions (e.g. b-tagging)

Systematic Uncertainty	Base Selection (%)	Anti-Tagged (%)	Loose Tag (%)
Lepton Scale Factor	0.5	0.5	0.5
Trigger Efficiency	0.2	0.2	0.2
Jet Energy Scale	3.1	2.6	1.9
ISR/FSR	1.3	2.6	6.5
Helicity Re-Weighting	3.5	3.4	3.2
Parton Distribution Functions	0.9	0.9	0.9
Total Correlated	5.0	5.1	7.5
<i>B</i> -Tagging Scale Factor	10.2	16.3	5.5
Mistag $\alpha\beta$ Correction	0.6	1.0	0.4
$\mathcal{B}(t \rightarrow Zc)$ versus $\mathcal{B}(t \rightarrow Zu)$	0.0	4.0	4.0
Total Anti-Correlated	10.2	16.8	6.8

- Signal systematic evaluated for acceptance ratio A_{WZ}/A_{LJ}
- Distinguish uncertainties: **correlated** or **anti-correlated** between selections
 - **Correlated**: shift anti-tagged & tagged selection into same direction (e.g. lepton SF)
 - **Anti-correlated**: shift anti-tagged & tagged into opposite directions (e.g. b-tagging)


Systematic Uncertainty	Base Selection (%)	Anti-Tagged (%)	Loose Tag (%)
Lepton Scale Factor	0.5	0.5	0.5
Trigger Efficiency	0.2	0.2	0.2
Jet Energy Scale	3.1	2.6	1.9
ISR/FSR	1.3	2.6	6.5
Helicity Re-Weighting	3.5	3.4	3.2
Parton Distribution Functions	0.9	0.9	0.9
Total Correlated	5.0	5.1	7.5
<i>B</i> -Tagging Scale Factor	10.2	16.3	5.5
Mistag $\alpha\beta$ Correction	0.6	1.0	0.4
$\mathcal{B}(t \rightarrow Zc)$ versus $\mathcal{B}(t \rightarrow Zu)$	0.0	4.0	4.0
Total Anti-Correlated	10.2	16.8	6.8

- Background systematics **dominated by yield uncertainties**
- Total background yield: 130 ± 28 (21.5% relative uncertainty)
- Tagging rate: $15\% \pm 4\%$ (relative uncertainty: 26.7% tagged, 4.7% anti-tagged)
- Remaining uncertainties: **efficiency of χ^2 cut**
 - Ratio of events with $\sqrt{\chi^2} < 1.6$ (signal region) vs. $\sqrt{\chi^2} > 3.0$ (control region)
 - Dominated by **choice of MC generator and jet energy scale**



Systematic Uncertainty	Anti-Tagged (%)	Loose Tag (%)
Lepton Scale Factor	< 0.1	< 0.1
Trigger Efficiency	< 0.1	< 0.1
Jet Energy Scale	5.1	2.1
B -Tagging Scale Factor	< 0.1	0.3
Mistag $\alpha\beta$ Correction	0.2	0.4
ALPGEN MC Generator	10.0	5.9
Total Uncertainty	11.2	6.3

- Background systematics **dominated by yield uncertainties**
- Total background yield: 130 ± 28 (21.5% relative uncertainty)
- Tagging rate: $15\% \pm 4\%$ (relative uncertainty: 26.7% tagged, 4.7% anti-tagged)
- Remaining uncertainties: **efficiency of χ^2 cut**
 - Ratio of events with $\sqrt{\chi^2} < 1.6$ (signal region) vs. $\sqrt{\chi^2} > 3.0$ (control region)
 - Dominated by **choice of MC generator and jet energy scale**




Systematic Uncertainty	Anti-Tagged (%)	Loose Tag (%)
Lepton Scale Factor	< 0.1	< 0.1
Trigger Efficiency	< 0.1	< 0.1
Jet Energy Scale	5.1	2.1
B-Tagging Scale Factor	< 0.1	0.3
Mistag $\alpha\beta$ Correction	0.2	0.4
ALPGEN MC Generator	10.0	5.9
Total Uncertainty	11.2	6.3

- Opening the box with 1.12 fb^{-1}
 - Event yield consistent with **background only**
 - Fluctuated about 1σ high: slightly unlucky
- Result: The World's Best Limit!**
 $\mathcal{B}(t \rightarrow Zq) < 10.6\% @ 95\% \text{ C.L.}$
 - Expected limit: $7.1\% \pm 3.0\%$
 - 25% better than L3 (13.7%)
 - 3x better than CDF Run I (33%)
 - Above results assumes $m_t = 175 \text{ GeV}/c^2$, limit at $m_t = 170 \text{ GeV}/c^2$: $\text{BR}(t \rightarrow Zq) < 11.2\% @ 95\% \text{ C.L.}$
- Update** with 2 fb^{-1} and improved method in the works



Selection	Observed	Expected
Base Selection	141	130 ± 28
Base Selection (Tagged)	17	20 ± 6
Anti-Tagged Selection	12	7.7 ± 1.8
Tagged Selection	4	3.2 ± 1.1

Mass χ^2 (95% C.L. Upper Limit)




- Large Hadron Collider (LHC):
 - Proton-proton collider at 14 TeV center-of-mass energy (CERN)
 - Two multi-purpose experiments: ATLAS and CMS
 - First data expected in 2008 (2009?)
- Recent ATLAS study on sensitivity for top FCNC
 - Improvement of current limits on BR ($t \rightarrow Zq$) by **2–3 orders of magnitude**
 - Entering interesting regime of 10^{-4} to 10^{-5} : exclusion of first theoretical models
- Caveat: background model
 - Existing MC tools not tuned to new energy regime
 - Tevatron experience: obtain backgrounds from data

ATLAS Detector



[J. Carvalho et al., SN-ATLAS-2007-059]



- Top flavor changing neutral current decays
 - **Extremely rare** in the standard model
 - Enhanced in theories beyond the standard model → any signal would indicate **new physics**
- **First Tevatron Run II search** for FCNC $t \rightarrow Zq$ in top quark decays
 - Event signature: $Z + \geq 4$ jets
 - Main background process: standard model $Z +$ jets production
 - **Mass χ^2** to separate signal from background
- **No evidence for top FCNC found**
 - World's best limit:
 $BR(t \rightarrow Zq) < 10.6\%$ at 95% C.L.
 - Working on improvements, stay tuned!