

QCD Physics Potential of CMS

On behalf of the CMS Collaboration
Klaus Rabbertz
University of Karlsruhe




Outline



- What is QCD?
- Tracks and Hadrons
- Jets
- Photons
- Summary



What is QCD?





What is QCD?



First of all ... it is 98% of us (our mass)
as we have heard yesterday from Raphael!
So we should better know it.




- In Formulae:

- ➔
$$\mathcal{L} = -\frac{1}{4}F_{\alpha\beta}^A F_A^{\alpha\beta} + \sum_{\text{flavours}} \bar{q}_a (i\not{D} - m)_{ab} q_b + \mathcal{L}_{\text{gauge-fixing}}$$

- In Words:


- ➔ QCD is the theory of the strong interaction, one of the four fundamental forces of nature, describing especially
 - ➔ the hard interactions between the coloured quarks and gluons
 - ➔ but also how they bind together to form hadrons.



What is QCD?



- In Pictures:




S. Gieseke



What is QCD?



- Zoomed in:



à la R. Field



QCD in CMS



• CMS QCD Group:

→ **Everything** ... not excluded:

- New particles, exotics, SUSY (Higgs, Exotica, SUSY) **Mostly Background**
- Weak bosons (EWK) **PDFs, Multijets**
- Heaviest quarks (Top, B Physics) **Hadrons, Multijets**
- Very forward topology (Forward Physics) **Jets, PDFs, Evolution**
- Colliding hadrons other than protons (Heavy Ions) **Baseline**


→ **Three subdivisions:**

- Low p_T measurements (tracker, hadrons)
- High p_T measurements (calorimeter, jets)
- Measurements with photons (ECAL, photons)

→ **Plus common efforts (PDFs, ...)**



QCD at Startup




Still enough events/sec left 

• Startup with QCD:

MinBias


- ➔ Not statistically limited
- ➔ First measurements at multi TeV energy scale
- ➔ Re-establishment of Standard Model, i.e. test extrapolations from Tevatron energies
- ➔ **Background** to be understood for **almost everything**
- ➔ Physics commissioning of CMS
- ➔ But be prepared for surprises ...

$$\sigma_{\text{jet}}(E_T^{\text{jet}} > 100 \text{ GeV})$$






Tracks and Hadrons





First Observations



... just a bunch of hadrons!

● Track based analyses:

- ➔ Charged particle rapidity density
- ➔ Charged hadron spectra
- ➔ Underlying event from transverse region

Also at 900 GeV

Phys.Lett.Vol.107B, no. 4


[First UA1 Publication](#)

QCD


17. December 1981

Recall:


**SOME OBSERVATIONS ON THE FIRST EVENTS
SEEN AT THE CERN PROTON-ANTIPROTON COLLIDER**




Hadron Trigger at Startup



At low luminosity want to collect “ideal” data meaning exactly one collision per bunch crossing



CMS PAS QCD-07-002



Need trigger (Pythia):

- ➔ 69% non-diffractive
- ➔ 18% single-diffractive
- ➔ 13% double-diffractive

Possibilities:




- ➔ single or double-sided HF Trigger Tower #
- ➔ others like PixelTracks under examination

Efficiencies single:

- ➔ 81% non-diffractive
- ➔ 15% single-diffractive
- ➔ 15% double-diffractive

Double kills diffr. events!

Charged Particle Rapidity Density from Hits




6 bins in hit multiplicity of equal cross sections

Strategy (used by Phobos at RHIC):

CMS PAS QCD-08-004

- ➔ **No tracking**, just count clusters in the pixel barrel layers (4, 7 and 10 cm radii)
- ➔ Use cluster size to estimate z vertex and to remove hits at high η from non-primary sources
- ➔ Correction for loopers, secondaries; systematic uncertainty expected below 10%



CMS Pixel Triplets



Expect:

- ➔ ~ 2 million events assuming one month with 1 Hz allocated bandwidth

Strategy:

- ➔ Still “no” tracking, Pixel Triplets




CMS pixel detector:

- ➔ 3 barrel layers (4, 7 and 10 cm radii) and 2 end caps on each side


- ➔ $100 \times 150 \mu\text{m}^2$ pixels

Hit triplets:

- ➔ Use pixel hit triplets instead of pairs, loss of acceptance but lower fake rate
- ➔ Reconstructing down to $p_T = 0.075$ GeV




Charged Particle Rapidity Density from Triplets




Simulation result from CMS:

- Charged particle pseudo-rapidity distribution
- Pythia tune DWT





Model expectations for charged particles at $|\eta| = 0$ vs. \sqrt{s} :

- Pythia: $\sim \ln^2(s)$
- Phojet: $\sim \ln(s)$




Charged Hadron Spectra




Empirical fits

CMS PAS QCD-07-001



Limit in p_T at $|y| < 0.1$ for dE/dx differentiation



Technique:

Tracks from pixel triplet seeding
 → Tracking down to p_T of 75 MeV

Klaus Rabbertz

Systematic:

Trigger, feed-down, geom. acceptance, alg. efficiency

Isfahan, Iran, 24.04.2009

Events: ~ 2M

One month with 1 Hz allocated bandwidth


1st IPM Meeting on LHC Physics



The Underlying Event



High P_T Jet Production



The Underlying Event is everything but the hard scatter.


Measurement possibility:

→ Charged particle and p_T sum densities in **transverse region** of leading jet of charged particles


Other "stuff" but the hard scatter

Leading jet

ChgJet #1 Direction



Balancing jet







The Underlying Event



Mix of contributing MinBias and calorimetric jet triggers



Decomposition of trigger contributions to charged particle density in $\Delta\Phi$ plane



CMS PAS QCD-07-003




The Underlying Event




Charged particle density in transverse plane vs. leading charged jet p_T

Extrapolation to LHC from CDF data



Comparison of different Pythia tunes

Statistics as for 100/pb




ATLAS

CMS PAS QCD-07-003





The Underlying Event



Increase sensitivity with tracks from $p_T > 0.5$ GeV instead of > 0.9 GeV

Decrease systematic effects with ratio, but with similar systematic $\rightarrow 0.9 / 1.5$



CMS PAS QCD-07-003



Jets







Theory Picture



Measured (or simulated) Event





Experimental Picture



All jets in the event
satisfying the selection criteria

$$\frac{d^2\sigma}{dp_T dy} = \frac{N_{jets}}{\epsilon \cdot L \cdot \Delta p_T \cdot \Delta y} \times C_{unsm}$$

Master Equation

➔ Jet Efficiency
➔ Event Efficiency

Bins of **corrected** Jet Pt
and Jet rapidity

Unsmearing correction
(due to the finite detector
Pt resolution)

**The JES dominates the
total uncertainty of the
measurement**

K. Kousouris




Generic Jet Analysis



Requires:


- PDFs
- LO & NLO MC
- Det. simulation
- Jet energy scale and resolution
- Calorimeter calibration
- Jet triggers
- Luminosity
- and ...

Data, of course!





Jet Analysis Uncertainties



Theoretical Uncertainties (~ in order of importance):

- ➔ PDF Uncertainty
- ➔ pQCD (Scale) Uncertainty
- ➔ Non-perturbative Corrections
- ➔ PDF Parameterization
- ➔ Electroweak Corrections
- ➔ Knowledge of $\alpha_s(M_Z)$
- ➔ ...

Recall: Jet Algorithms used by CMS:


- ➔ Iterative Cone R = 0.5
- ➔ SISCone R = 0.5, 0.7
- ➔ k_T D = 0.4, 0.6

Experimental Uncertainties (~ in order of importance):

- ➔ Jet Energy Scale (JES)
- ➔ Noise Treatment
- ➔ Pile-Up Treatment
- ➔ Luminosity
- ➔ Jet Energy Resolution (JER)
- ➔ Trigger Efficiencies
- ➔ Resolution in Rapidity
- ➔ Resolution in Azimuth
- ➔ Non-Collision Background
- ➔ ...




QCD Jet Analyses




- **Jet analyses at high transverse momenta:**
 - ➔ **Dijet azimuthal decorrelation**
 - ➔ **Less sensitive to JES, not dependent on luminosity**
 - ➔ **Event shapes**
 - ➔ **Reduced sensitivity to JES & JER, not dependent on luminosity**
 - ➔ **Dijet production ratios & angles**
 - ➔ **Reduced sensitivity to JES, not dependent on luminosity**
 - ➔ **Jet cross section ratios (3-jet / all, $R=0.7$ / $R=0.5$, SISCone / kT)**
 - ➔ **Reduced sensitivity to JES, not dependent on luminosity**
 - ➔ **Jet shapes**
 - ➔ **Multi-jet studies**
 - ➔ **Inclusive jet cross section**
 - ➔ **Most complicated, requires all uncertainties to be under control!**


Inclusive Jets at the Tevatron



CDF Incl. k_T jets, $D=0.7$
Theory: NLO with CTEQ6.1M



D0 Incl. MidPoint cone jets, $R=0.7$
Theory: NLO with CTEQ6.5M




arXiv:0802.2400 [hep-ex]

Phys.Rev.D75:092006,2007


Inclusive Jets at the LHC



$k_T, D=0.6, 10 \text{ TeV}$



SISCone, R=0.7, 10 TeV



LHC reach
3 x 600 GeV
with 100/pb

Log Scale

Bands are PDF uncertainties from CTEQ6.5

NLOJET++, PRL 88 122003 (2002), PR D68 094002 (2003)
fastNLO, hep-ph/0609285
fastjet: PLB641 (2006) [hep-ph/0512210],
SISCone: JHEP 05 (2007) 086 [arXiv:0704.0292 (hep-ph)]




Cross Section Ratios




Cross section ratios in 6 bins in rapidity y


SISCone 0.7 / k_T 0.6 @ 10 TeV



k_T 0.6 10 TeV / 14 TeV



Parton Density Experience



“The data are compared with QCD predictions for various sets of parton distribution functions. The cross section for jets with $E_T > 200$ GeV is significantly higher than current predictions based on $O(\alpha_s^3)$ perturbative QCD calculations. ...”


Explained by change in gluon density which then can be constrained by jets!

Today:


Much better estimates of PDF uncertainties

But beware ...


Phys.Rev.Lett. 77 (1996)




$d^2\sigma/dp_T dy_{PDF} / d^2\sigma/dp_T dy_{CTEQ6.5}$



PDF and Scale Uncertainties




$k_T, D=0.6, 10 \text{ TeV}$



SISCone: Similar result

PDF uncertainties from CTEQ 6.5
At 4 TeV in p_T about -20% to +35%



$k_T, D=0.6, 10 \text{ TeV}$



SISCone: Up to twice as large!

Scale variation of $p_{T,jet}$ to
 $0.5 * p_{T,jet}$ and $2 * p_{T,jet}$

Forward Jets and PDFs



CMS PAS FWD-08-001





Non-perturbative Corrections



To compare with data correct NLO for:

- Multiple Parton Interactions (MPI)
- Hadronization & Decays (Lund, Cluster)

Less compensation of these effects for k_T than for SIScone, not negligible in both cases.
 Need MC tunes for UE with first LHC data!





Compared 3 different tuned MC:

- Pythia Tune D6T
- Herwig++
- Herwig/Jimmy with Tune from ATLAS

Take correction as average of Pythia and Herwig++ and half the spread as uncertainty



JES and New Physics




- ➔ Dominating experimental uncertainty: JES (assumed $\pm 10\%$ at startup)
- ➔ More data and improved JES knowledge needed to start constraining PDFs
- ➔ Sensitive to Contact Interactions beyond Tevatron reach (2.7 TeV) with 10 pb^{-1}

Jet Energy Resolution

- Jet energy resolution from CMS performance study
- JER usually parameterized by:


$$\sigma(p_T) = p_T \cdot \sqrt{C^2 + \frac{S^2}{p_T} + \frac{N^2}{p_T^2}}$$

Derived from MC comparison
Fairly independent of jet algo






Finite detector resolution on a steeply falling jet p_T spectrum leads to strongly asymmetric bin migrations!

Can be derived from dataMC with dijet asymmetry method






Unsmearing Applied



Unsmearing by "Ansatz Method"



- ➔ Artificially smear jets by Gaussian with an arbitrary but reasonable p_T dependent width.
- ➔ Apply ansatz method
- ➔ Method corrects p_T smearing effects on steeply falling spectrum

Unsmearing Steps

Motivation

The **observed** cross section is **higher** than the true one due to the falling shape of the spectrum and the finite p_T resolution. More events migrate into a bin of measured p_T than out of it.


Unsmearing steps:

Analytical expression of the p_T resolution

Ansatz function with free parameters to be determined by the data

Fitting the data with the Ansatz function smeared with p_T resolution.

Unsmearing correction calculated bin by bin.



$$\Rightarrow R(p'_T, p_T) = \frac{1}{\sqrt{2\pi}\sigma(p'_T)} \exp\left[-\frac{(p'_T - p_T)^2}{2\sigma^2(p'_T)}\right]$$

$$\Rightarrow f(p_T) = N \cdot p_T^{-a} \cdot \left(1 - \frac{2 \cosh(y_{min}) p_T}{\sqrt{s}}\right)^b \exp(-\gamma p_T)$$

$$\Rightarrow F(p_T) = \int_0^\infty f(p'_T) R(p'_T, p_T) dp'_T$$

$$\Rightarrow C_{bin} = \frac{\int_{bin} f(p_T) dp_T}{\int_{bin} F(p_T) dp_T}$$



JER Uncertainty



Good knowledge of the resolution required!

A wrong assumption can shift the final spectrum easily by some percent ...


Two scenarios studied:

Very pessimistic:

- Resolution in unsmearing is “real” resolution (in %) - 1 %: 1% better (-)
- Resolution in unsmearing is “real” resolution (in %) + 4%: 4% worse (+)



Optimistic:

- Resolution in unsmearing is 0.95 times “real” resolution (in %): 5 % better (*)
- Resolution in unsmearing is 1.05 times “real” resolution (in %): 5 % worse (*)





Jet Shapes




Jet shape measurements can be used to discriminate between different underlying event models.

Can be used to distinguish gluon originated jets from quark jets.

Measurement of the average integrated (differential) energy flow inside jets.

Jet shape measurements can be used to test the showering models in the MC generators.


Integrated jet shape $\Psi(r)$



CMS PAS QCD-08-005




Event Shapes



Definition:

Transverse global Thrust
 (k_T jets, $E_{T,1} > 80$ GeV, $E_{T,\text{all}} > 60$ GeV)


$$T_{\perp,g} \equiv \max_{\vec{n}_T} \frac{\sum_i |\vec{p}_{\perp,i} \cdot \vec{n}_T|}{\sum_i p_{\perp,i}}$$



Similar as Event Shapes in e^+e^- and ep


- In praxis, need to restrict rapidity range: $|\eta| < 1.3$ → Transverse central Thrust
- Less sensitive to JES & JER uncertainty
- No luminosity uncertainty
- Useful for MC tuning

CMS PAS QCD-08-003



linear

linear




spherical



$$\tau_{\perp,g} \equiv 1 - T_{\perp,g}$$




Event Shapes



- Distributions get more peaked at higher E_T
- Corrected pseudo-data follow behaviour of original Pythia MC
- Alpgen makes different predictions



CMS PAS QCD-08-003




Photons (and Jets)






Photons



Photon processes:


- ➔ Direct photon production
- ➔ Di-photons
- ➔ Photon + n jets



Annihilation Process

LO

Dominant: > 80%



Compton Process

# bunches	β^* (m)	I_b	L ($\text{cm}^{-2}\text{s}^{-1}$)	Pileup	Photons/hour ($p_T > 20 \text{ GeV}$)
1x1	18	10^{10}	10^{27}	low	$3.2 \cdot 10^{-1}$
43x43	18	$3 \cdot 10^{10}$	$3.8 \cdot 10^{29}$	0.05	$1.2 \cdot 10^2$
43x43	4	$3 \cdot 10^{10}$	$1.7 \cdot 10^{30}$	0.21	$5.4 \cdot 10^2$
43x43	2	$4 \cdot 10^{10}$	$6.1 \cdot 10^{30}$	0.76	$2.0 \cdot 10^3$
156x156	4	$4 \cdot 10^{10}$	$1.1 \cdot 10^{31}$	0.38	$3.6 \cdot 10^3$
156x156	4	$9 \cdot 10^{10}$	$5.1 \cdot 10^{31}$	1.9	$1.6 \cdot 10^4$
156x156	2	$9 \cdot 10^{10}$	$1.1 \cdot 10^{32}$	3.9	$3.6 \cdot 10^5$

Photon rate estimations:


- ➔ Photon $p_T > 20 \text{ GeV}$
- ➔ Photon $|\eta| < 2.5$

Not taken into account

S. Ganjour



Photon Isolation



Gauge boson production gives important additional information:


- Luminosity measurement
- Detector calibration
- PDFs
- Background for new physics :-)

Important steps:

- Good efficiency including photon conversions
- Proper photon isolation to suppress background

CMS photon study:

- Photon p_T spectrum for 1/fb
- Background QCD jets in blue
- **After photon isolation cuts**
- Improves S/B > 2 orders of magnitude




Eur. Phys. J. C 53, 49-58 (2008).

P_T^Y



Outlook



- CMS is preparing (again) for first LHC data in autumn
- LHC will explore unknown territory in QCD
- First measurements, even at 900 GeV, will be QCD
- Some tough experimental systematics to deal with, but combining detector parts may help in certain phase space regions (jets+tracks, particle flow)
- Measurements of jets and photons are important tests of QCD:
 - ➔ Angular distributions, inclusive jets, di-jets, photon+jets, di-photons, forward jets
 - ➔ Calibration of the calorimeters
 - ➔ Better understanding of dominant background to many new physics channels
 - ➔ Constraints on PDFs
- New physics might be just ahead!



Thanks to all colleagues helping in preparing this presentation!






Backups






CMS Electromagnetic Calorimeter



Barrel (EB):

- η segments: 2×85
- ϕ segments: 360
- 61200 crystals
(PbWO_4 , $26 X_0$)
- $\Delta\eta \times \Delta\phi \approx$
 0.0174×0.0174



Energy resolution from test beam:


$S = 3.63\%$, $N = 124 \text{ MeV}$, $C = 0.26\%$

$$\left(\frac{\sigma}{E}\right)^2 = \left(\frac{S}{\sqrt{E}}\right)^2 + \left(\frac{N}{E}\right)^2 + C^2$$

End caps (EE):

- (x,y) grid on two halves
- front face $28 \times 28 \text{ mm}^2$
- $2 \times 2 \times 3662$ crystals = 14648
(PbWO_4 , $25 X_0$)

CMS Hadronic Calorimeter




HCAL (tower structure):

- Barrel (HB): $|\eta| < 1.4$, 2304 towers
- End caps (HE): $1.3 < |\eta| < 3.0$, „ towers
- Outside coil (HO): $|\eta| < 1.26$ (tail catcher)
→ 4608 towers (Plastic scintillator tiles, $\approx 10 \lambda_N$)
→ $\Delta\eta \times \Delta\phi \approx 0.087 \times 0.087 \rightarrow 0.350 \times 0.175$

- Forward (HF): $2.9 < |\eta| < 5.0$ (not shown)
→ 2 x 900 towers (Quartz fibers, $\approx 10 \lambda_N$)
→ $\Delta\eta \times \Delta\phi \approx 0.111 \times 0.175 \rightarrow 0.302 \times 0.350$

CASTOR calorimeter (not shown):

- $5.1 < |\eta| < 6.5$, $\approx 22 X_0$, $\approx 10 \lambda_N$





Hadron Spectra Systematics



CMS Pixel triplets


$$\Delta N_{\text{corrected}} = \frac{(1 - \text{fakeRate}) \cdot (1 - \text{feedDown})}{\text{geomAccep} \cdot \text{algoEffic} \cdot (1 - \text{multiCount})} \cdot \Delta N_{\text{measured}}$$

Correction	Dependence on			Corr. [%]	Syst.
	kine	part	mult		
Trigger	no	no	yes	15	5
Geometrical acceptance	yes	yes	no	10-20	2
Algorithmic efficiency	yes	yes	no	10-20	2
Multiple track counting	yes	no	no	small	small
Fake track rate	yes	no	yes	small	small
Feed-down	yes	yes	no	2-15	1-2
η, p_T resolution	no	no	no	1-5	1-5
Total	yes	yes	yes		7-9

CMS PAS QCD-07-001



Tracking Performance



Comparison of tracking performance for:



● **Ideal conditions**

→ **Start-up (misaligned)**

■ **Alignment Position Error application**

Track reconstruction efficiency


Fake rate



CMS PAS QCD-07-003



Dijet Azimuthal Decorrelation



Dijets in pp collisions:

$\Delta\phi$ dijet = $\pi \rightarrow$

Exactly two jets, no further radiation

$\Delta\phi$ dijet small deviations from $\pi \rightarrow$


Additional soft radiation outside the jets

$\Delta\phi$ dijet as small as $2\pi/3 \rightarrow$

One additional high- p_T jet

$\Delta\phi$ dijet small – no limit \rightarrow

Multiple additional hard jets in the event



hep-ex/0409040
PRL 94, 221801 (2005)



Partonic Subprocesses




➤ For $hh \rightarrow$ jets there are **seven** relevant partonic subprocesses:



- 1) $gg \Rightarrow$ jets $\propto H_1(x_1, x_2)$
- 2) $qg, \bar{q}g \Rightarrow$ jets $\propto H_2(x_1, x_2)$
- 3) $gq, g\bar{q} \Rightarrow$ jets $\propto H_3(x_1, x_2)$
- 4) $q_i q_j, \bar{q}_i \bar{q}_j \Rightarrow$ jets $\propto H_4(x_1, x_2)$
- 5) $q_i q_i, \bar{q}_i \bar{q}_i \Rightarrow$ jets $\propto H_5(x_1, x_2)$
- 6) $q_i \bar{q}_i, \bar{q}_i q_i \Rightarrow$ jets $\propto H_6(x_1, x_2)$
- 7) $q_i \bar{q}_j, \bar{q}_i q_j \Rightarrow$ jets $\propto H_7(x_1, x_2)$

➤ Seven linear combinations H_i of PDFs

Subprocess Decomposition




Decomposition of the total ppbar, pp → jets cross section (NLO) into subprocesses
 At central rapidity Subprozesse against the scaling variable $x_T = 2p_T/\sqrt{s}$



Tevatron

LHC, 14 TeV

New Physics from Dijets



New Physics with Jets:


Contact interactions

Resonances

- ★ W' & Z' (Grand Unified Theory)
- ★ E_6 diquarks (D) (Superstrings & GUT)
- ★ Excited quarks (q^*) (Compositeness)
- ★ RS Gravitons (G) (Extra Dimensions)
- ★ Colorons (C) & Axigluons (A) (Extra Color)

Need
 $E_{\text{CMS}} > M$

Di-jet mass distribution



Model	J	Color	Cross Section (pb)					
			M=0.7 TeV		M=2.0 TeV		M=5.0 TeV	
			$ \eta < 1$	$ \eta < 1.3$	$ \eta < 1$	$ \eta < 1.3$	$ \eta < 1$	$ \eta < 1.3$
q^*	1/2	Triplet	7.95×10^2	1.27×10^3	9.01	1.36×10^1	1.82×10^{-2}	2.30×10^{-2}
A,C	1	Octet	3.22×10^2	5.21×10^2	5.79	8.82	1.55×10^{-2}	2.04×10^{-2}
D	0	Triplet	8.11×10^1	1.26×10^2	4.20	5.97	4.65×10^{-2}	5.75×10^{-2}
G	2	Singlet	3.57×10^1	5.47×10^1	1.83×10^{-1}	2.60×10^{-1}	2.64×10^{-4}	3.19×10^{-4}
W'	1	Singlet	1.46×10^1	2.37×10^1	3.49×10^{-1}	5.31×10^{-1}	8.72×10^{-4}	1.17×10^{-3}
Z'	1	Singlet	8.86	1.44×10^1	1.81×10^{-1}	2.77×10^{-1}	5.50×10^{-4}	7.26×10^{-4}

Contact Interactions


- ★ Sensitive to Scale $\Lambda \gg \sqrt{s}$!

$$L_{qq} = \frac{Ag^2}{2\Lambda^2} (\bar{q}_L \gamma^\mu q_L) (\bar{q}_L \gamma_\mu q_L)$$

Contact Interaction



New Physics from Dijets




Search for deviation from expected event rate:

- ➡ QCD from PYTHIA (here) or NLO
- ➡ Contact interaction: PYTHIA or LO


Search for resonances

Possible signals of q^* relative to QCD prediction, visible for < 2 TeV (statistical uncertainty only!)

Cross section ratios



CMS PAS SBM-07-001



One means to avoid systematics is by looking into cross section ratios in η



Recent Limits




Tevatron limit on contact interaction scale (qqqq): **> 2.4 - 2.7 TeV**

Dijet resonance search

CDF Preliminary 03/2008




Resonance	Excluded (GeV)	Resonance	Excluded (GeV)
A or C	260 - 1250	D	290 - 630
ρ_{T8}	260 - 1110	W'	280 - 840
q^*	260 - 870	Z'	320 - 740



Exclusion limits for W' and Z'



Dijet Ratios



- ➔ Sensitivity to new physics from dijet x section ratios in pseudo-rapidity
- ➔ Reduced sensitivity to jet energy scale



Jet Shapes



A possibility to look into details of QCD and jet structure!



Norm. transverse energy distribution:

$$\rho(r) = \frac{\sum p_T(r - \Delta r/2, r + \Delta r/2)}{\Delta r \sum p_T^{Jet}}$$

Good reproduction of general properties (central region $|\eta| < 1$, matched jets)


Jets from generator particles

Jets from calorimeter towers



CMS PAS JME-07-003

Multiple Parton Interactions



Phase space:


As low as possible in pT:

- photon pT > 10 GeV
- jet pT > 20 GeV for calojets
=> could consider jets from tracks

▶ Double-parton-scattering

- four-jet production (→ AFS, UA2, CDF)
- like-sign W production
- **γ + 3-jet production (→ CDF)**

▶ **Need double-parton component to describe the data**





Influence of $\alpha_s(M_Z)$





Cross section ratios at central rapidity
 $\alpha_s(M_Z)$ varies from 0.112 to 0.125

PDG Value $\alpha_s(M_Z) = 0.1176 \pm 0.0020$
Would lead to 2 to 4% variation


With CTEQ6.6 central PDF

With CTEQ6.6A α_s PDF series





Some UA1 Quotations



➔ Quotations from Phys. Lett. Vol. 107B, no. 4:

- ... dipole magnet which produces a field of **0.7 T** over a volume of 7m x 3.5m x 3.5m ...
- ... yields space points at **centimetre** intervals on the detected tracks
- ... two short accelerator development periods in October and November 1981 ...
- The events were **scanned by physicists** on a Megatek display.
- ... was examined independently by all physicists who participated in the scanning. The combined effect of the **scanner variations** ...