

CERN/LHCC 06-???

CMS TDR 8

**** Final Date here ****

CMS Physics

Technical Design Report

Volume II: Physics Performance

Also available at:
<http://cmsdoc.cern.ch/cms/cpt/tdr/index.html>

***** Draft Version 6.0*****

May 26, 2006

CMS Software and Physics, Reconstruction and Selection (PRS) Projects		
CMS Spokesperson	Michel Della Negra, CERN	Michel.Della.Negra@cern.ch
CMS Technical Coordinator	Alain Herve, CERN	Alain.Herve@cern.ch
CMS Collaboration Board Chair	Lorenzo Foa, Pisa	Lorenzo.Foa@cern.ch
CPT Project Manager	Paraskevas Sphicas, CERN and Athens	Paraskevas.Sphicas@cern.ch
CPT PRS Coordinator	Darin Acosta, Florida	Darin.Acosta@cern.ch
CPT PRS Coordinator	Albert De Roeck, CERN	Albert.De.Roeck@cern.ch
CPT Software Coordinator	Lucia Silvestris, Bari	Lucia.Silvestris@cern.ch
CPT Software Coordinator	Avi Yagil, FNAL	Yagil@fnal.gov



Placeholder page for boilerplate, such as:

List of editors...

Acknowledgements...cover design, etc.

ISBN xx-xxxx-xx-xx

Trademark notice: all trademarks appearing in this proposal are acknowledged as such.



CMS Collaboration

Yerevan Physics Institute, Yerevan, ARMENIA

G.L. Bayatian, S. Chatrchyan, G. Hmayakyan, A.M. Sirunyan

Institut für Hochenergiephysik der OeAW, Wien, AUSTRIA

W. Adam, T. Bergauer, M. Dragicevic, J. Erö, M. Friedl, R. Fruehwirth, V. Ghete, P. Glaser, J. Hrubec, M. Jeitler, M. Krammer, I. Magrans, I. Mikulec, W. Mitaroff, T. Noebauer, M. Pernicka, P. Porth, H. Rohringer, J. Strauss, A. Taurok, W. Waltenberger, G. Walzel, E. Widl, C.-E. Wulz

Research Institute for Nuclear Problems, Minsk, BELARUS

A. Fedorov, M. Korzhik, O. Missevitch, R. Zuyevski

National Centre for Particle and High Energy Physics, Minsk, BELARUS

V. Chekhovsky, O. Dvornikov, I. Emeliantchik, A. Litomin, V. Mossolov, N. Shumeiko, A. Solin, R. Stefanovitch, J. Suarez Gonzalez, A. Tikhonov

Byelorussian State University, Minsk, BELARUS

V. Petrov

Vrije Universiteit Brussel, Brussel, BELGIUM

J. D'Hondt, S. De Weirdt, R. Goorens, J. Heyninck, S. Lowette, S. Tavernier, W. Van Doninck^{**1}, L. Van Lancker

Université Libre de Bruxelles, Bruxelles, BELGIUM

O. Bouhali, B. Clerboux, G. De Lentdecker, J.P. Dewulf, T. Mahmoud, P.E. Marage, L. Neukermans, V. Sundararajan, C. Vander Velde, P. Vanlaer, J. Wickens

Université Catholique de Louvain, Louvain-la-Neuve, BELGIUM

S. Assouak, J.L. Bonnet, G. Bruno, J. Caudron, B. De Callatay, J. De Favereau De Jeneret, S. De Visscher, C. Delaere, P. Demin, D. Favart, E. Feltrin, E. Forton, G. Grégoire, S. Kalinin, D. Kcira, T. Keutgen, G. Leibenguth, V. Lemaitre, Y. Liu, D. Michotte, O. Militaru, A. Ninane, S. Ovin, T. Pierzchala, K. Piotrkowski, V. Roberfroid, X. Rouby, D. Teyssier, O. Van der Aa, M. Vander Donckt

Université de Mons-Hainaut, Mons, BELGIUM

E. Daubie, P. Herquet, A. Mollet, A. Romeyer

Universiteit Antwerpen, Wilrijk, BELGIUM

W. Beaumont, M. Cardaci, E. De Langhe, E.A. De Wolf, L. Rurua

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, RJ, BRAZIL

M.H.G. Souza

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, BRAZIL

V. Oguri, A. Santoro, A. Sznajder

Instituto de Fisica - Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ, BRAZIL

M. Vaz

Instituto de Fisica Teorica-Universidade Estadual Paulista, Sao Paulo, SP, BRAZIL

E.M. Gregores, S.F. Novaes

Institute for Nuclear Research and Nuclear Energy, Sofia, BULGARIA

T. Anguelov, G. Antchev, I. Atanasov, J. Damgov, N. Darmanov^{**1}, L. Dimitrov, V. Genchev^{**1}, P. Iaydjiev, B. Panev, S. Piperov, S. Stoykova, G. Sultanov, I. Vankov

University of Sofia, Sofia, BULGARIA

A. Dimitrov, V. Kozhuharov, L. Litov, M. Makariev, A. Marinov, E. Marinova, S. Markov, M. Mateev, B. Pavlov, P. Petkov, C. Sabev, S. Stoynev, Z. Toteva^{**1}, V. Verguilov

Institute of High Energy Physics, Beijing, CHINA

G.M. Chen, H.S. Chen, K.L. He, C.H. Jiang, W.G. Li, H.M. Liu, X. Meng, X.Y. Shen, H.S. Sun, M. Yang, W.R. Zhao, H.L. Zhuang

Peking University, Beijing, CHINA

Y. Ban, J. Cai, S. Liu, S.J. Qian, Z.C. Yang, Y.L. Ye, J. Ying

University for Science and Technology of China, Hefei, Anhui, CHINA

J. Wu, Z.P. Zhang

Technical University of Split, Split, CROATIA

N. Godinovic, I. Puljak, I. Soric

University of Split, Split, CROATIA

Z. Antunovic, M. Dzelalija, K. Marasovic

Institute Rudjer Boskovic, Zagreb, CROATIA

V. Brigljevic, D. Ferencek, K. Kadija, S. Morovic, M. Planinic**2

University of Cyprus, Nicosia, CYPRUS

C. Nicolaou, A. Papadakis, P.A. Razis, D. Tsiakkouri

National Institute of Chemical Physics and Biophysics, Tallinn, ESTONIA

A. Hektor, M. Kadastik, K. Kannike, E. Lippmaa, M. Müntel, M. Raidal

Laboratory of Advanced Energy Systems, Helsinki University of Technology, Espoo, FINLAND

P.A. Aarnio

Helsinki Institute of Physics, Helsinki, FINLAND

S. Czellar, E. Haeggstroem, A. Heikkinen, J. Härkönen, V. Karimäki, R. Kinnunen, T. Lampén, K. Lassila-Perini, S. Lehti, T. Lindén, P.R. Luukka, S. Michal**1, T. Mäenpää, J. Nysten, M. Stettler**1, E. Tuominen, J. Tuominiemi, L. Wendland

Lappeenranta University of Technology, Lappeenranta, FINLAND

T. Tuuva

Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, FRANCE

J.P. Guillaud, P. Nedelec, D. Sillou

DSM/DAPNIA, CEA/Saclay, Gif-sur-Yvette, FRANCE

M. Anfreville, S. Beauceron, E. Bougamont, P. Bredy, R. Chipaux, M. Dejardin, D. Denegri, J. Descamps, B. Fabbro, J.L. Faure, S. Ganjour, F.X. Gentit, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, F. Kircher, M.C. Lemaire**3, B. Levesy**1, E. Locci, J.P. Lottin, I. Mandjavidze, M. Mur, E. Pasquetto, A. Payn, J. Rander, J.M. Reymond, F. Rondeaux, A. Rosowsky, Z.H. Sun, P. Verrecchia

Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, FRANCE

S. Baffioni, F. Beaudette, M. Bercher, U. Berthon, S. Bimbot, J. Bourotte, P. Busson, M. Cerutti, D. Chamont, C. Charlot, C. Collard, D. Decotigny, E. Delmeire, L. Dobrzynski, A.M. Gaillac, Y. Geerebaert, J. Gilly, M. Haguenaer, A. Karar, A. Mathieu, G. Milleret, P. Miné, P. Paganini, T. Romanteau, I. Semeniouk, Y. Sirois

Institut Pluridisciplinaire Hubert Curien, IN2P3-CNRS - ULP, UHA Mulhouse, Strasbourg, FRANCE

J.D. Berst, J.M. Brom, F. Didierjean, F. Drouhin**1, J.C. Fontaine**4, U. Goerlach**5, P. Graehling, L. Gross, L. Houchu, P. Juillot, A. Lounis**5, C. Maazouzi, D. Mangeol, C. Olivetto, T. Todorov**1, P. Van Hove, D. Vintache

Institut de Physique Nucléaire, IN2P3-CNRS, Université Claude Bernard Lyon 1, Villeurbanne, FRANCE

M. Ageron, J.L. Agram, G. Baulieu, M. Bedjidian, J. Blaha, A. Bonnevaux, G. Boudoul**1, E. Chabanat, C. Combaret, D. Contardo**1, R. Della Negra, P. Depasse, T. Dupasquier, H. El Mamouni, N. Estre, J. Fay, S. Gascon, N. Giraud, C. Girerd, R. Haroutounian, J.C. Ianigro, B. Ille, M. Lethuillier, N. Lumb**1, H. Mathez, G. Maurelli, L. Mirabito**1, S. Perries, O. Ravat

Institute of High Energy Physics and Informatization, Tbilisi State University, Tbilisi, GEORGIA
R. Kvatadze

Institute of Physics Academy of Science, Tbilisi, GEORGIA
V. Roinishvili

RWTH, I. Physikalisches Institut, Aachen, GERMANY

R. Adolphi, R. Brauer, W. Braunschweig, H. Esser, L. Feld, A. Heister, W. Karpinski, K. Klein, C. Kukulies, J. Olzem, A. Ostapchuk, D. Pandoulas, G. Pierschel, F. Raupach, S. Schael, G. Schwering, M. Thomas, M. Weber, B. Wittmer, M. Wlochal

RWTH, III. Physikalisches Institut A, Aachen, GERMANY

A. Adolf, P. Biallass, M. Bontenackels, M. Erdmann, H. Fesefeldt, T. Hebbeker, S. Hermann, G. Hilgers, K. Hoepfner**1, C. Hof, S. Kappler, M. Kirsch, D. Lanske, B. Philipps, H. Reithler, T. Rommerskirchen, M. Sowa, H. Szczesny, M. Tonutti, O. Tsigenov

RWTH, III. Physikalisches Institut B, Aachen, GERMANY

F. Beissel, M. Davids, M. Duda, G. Flügge, T. Franke, M. Giffels, T. Hermanns, D. Heydhausen, S. Kasselmann, G. Kaussen, T. Kress, A. Linn, A. Nowack, M. Poettgens, O. Pooth, A. Stahl, D. Tornier, M. Weber

Deutsches Elektronen-Synchrotron, Hamburg, GERMANY

A. Flossdorf, B. Hegner, J. Mnich, C. Rosemann

University of Hamburg, Hamburg, GERMANY

G. Flucke, U. Holm, R. Klanner, U. Pein, N. Schirm, P. Schleper, G. Steinbrück, M. Stoye, R. Van Staa, K. Wick

Institut für Experimentelle Kernphysik, Karlsruhe, GERMANY

P. Blüm, V. Buege, W. De Boer, G. Dirkes**1, M. Fahrner, M. Feindt, U. Felzmann, J. Fernandez Menendez**6, M. Frey, A. Furgeri, F. Hartmann**1, S. Heier, C. Jung, B. Ledermann, Th. Müller, M. Niegel, A. Oehler, T. Ortega Gomez, C. Piasecki, G. Quast, K. Rabbertz, C. Saout, A. Scheurer, D. Schieferdecker, A. Schmidt, H.J. Simonis, A. Theel, A. Vest, T. Weiler, C. Weiser, J. Weng**1, V. Zhukov**7

University of Athens, Athens, GREECE

G. Karapostoli**1, P. Katsas, P. Kreuzer, A. Panagiotou, C. Papadimitropoulos

Institute of Nuclear Physics "Demokritos", Attiki, GREECE

G. Anagnostou, M. Barone, T. Geralis, C. Kalfas, A. Koimas, A. Kyriakis, S. Kyriazopoulou, D. Loukas, A. Markou, C. Markou, C. Mavrommatis, K. Theofilatos, G. Vermisoglou, A. Zachariadou

University of Ioánnina, Ioánnina, GREECE

X. Aslanoglou, I. Evangelou, P. Kokkas, N. Manthos, I. Papadopoulos, G. Sidiropoulos, F.A. Triantis

KFKI Research Institute for Particle and Nuclear Physics, Budapest, HUNGARY

G. Bencze**1, L. Boldizsar, C. Hajdu**1, D. Horvath**8, A. Laszlo, G. Odor, F. Sikler, N. Toth, G. Vesztergombi, P. Zalan

Institute of Nuclear Research ATOMKI, Debrecen, HUNGARY

J. Molnar

University of Debrecen, Debrecen, HUNGARY

N. Beni, A. Kapusi, G. Marian, P. Raics, Z. Szabo, Z. Szillasi, G. Zilizi

Panjab University, Chandigarh, INDIA

H.S. Bawa, S.B. Beri, V. Bhandari, V. Bhatnagar, M. Kaur, R. Kaur, J.M. Kohli, A. Kumar, J.B. Singh

University of Delhi, Delhi, INDIA

A. Bhardwaj, S. Bhattacharya^{**9}, S. Chatterji, S. Chauhan, B.C. Choudhary, P. Gupta, M. Jha, K. Ranjan, R.K. Shivpuri, A.K. Srivastava

Bhabha Atomic Research Centre, Mumbai, INDIA

S. Borkar, M. Dixit, M. Ghodgaonkar, S.K. Kataria, S.K. Lalwani, V. Mishra, A.K. Mohanty, A. Topkar

Tata Institute of Fundamental Research - EHEP, Mumbai, INDIA

T. Aziz, S. Banerjee, S. Bose, N. Cheere, S. Chendvankar, P.V. Deshpande, M. Guchait^{**10}, A. Gurtu, M. Maity^{**11}, G. Majumder, K. Mazumdar, A. Nayak, M.R. Patil, S. Sharma, K. Sudhakar, S.C. Tonwar

Tata Institute of Fundamental Research - HECR, Mumbai, INDIA

B.S. Acharya, S. Banerjee, S. Bheesette, S. Dugad, S.D. Kalmani, V.R. Lakkireddi, N.K. Mondal, N. Panyam, P. Verma

Institute for Studies in Theoretical Physics & Mathematics (IPM), Tehran, IRAN

M. Arabgol, H. Arfaei, M. Hashemi, M. Mohammadi, M. Mohammadi Najafabadi, A. Moshaii, S. Paktinat Mehdiabadi

University College Dublin, Dublin, IRELAND

M. Grunewald

Università di Bari, Politecnico di Bari e Sezione dell' INFN, Bari, ITALY

M. Abbrescia, L. Barbone, A. Colaleo^{**1}, D. Creanza, N. De Filippis, M. De Palma, G. Donvito, L. Fiore, D. Giordano, G. Iaselli, F. Loddo, G. Maggi, M. Maggi, N. Manna, B. Marangelli, M.S. Mennea, S. My, S. Natali, S. Nuzzo, G. Pugliese, V. Radicci, A. Ranieri, F. Romano, G. Selvaggi, L. Silvestris, P. Tempesta, R. Trentadue, G. Zito

Università di Bologna e Sezione dell' INFN, Bologna, ITALY

G. Abbiendi, W. Bacchi, A. Benvenuti, D. Bonacorsi, S. Braibant-Giacomelli, P. Capiluppi, F. Cavallo, C. Ciocca, G. Codispoti, I. D'Antone, G.M. Dallavalle, F. Fabbri, A. Fanfani, P. Giacomelli^{**12}, C. Grandi, M. Guerzoni, L. Guiducci, S. Marcellini, G. Masetti, A. Montanari, F. Navarra, F. Odorici, A. Perrotta, A. Rossi, T. Rovelli, G. Siroli, R. Travaglini

Università di Catania e Sezione dell' INFN, Catania, ITALY

S. Albergo, M. Chiorboli, S. Costa, M. Galanti, G. Gatto Rotondo, F. Noto, R. Potenza, G. Russo, A. Tricomi, C. Tuve

Università di Firenze e Sezione dell' INFN, Firenze, ITALY

A. Bocci, G. Ciralo, V. Ciulli, C. Civinini, R. D'Alessandro, E. Focardi, C. Genta, P. Lenzi, A. Macchiolo, N. Magini, F. Manolescu, C. Marchettini, L. Masetti, S. Mersi, M. Meschini, S. Paoletti, G. Parrini, R. Ranieri, M. Sani

Università di Genova e Sezione dell' INFN, Genova, ITALY

P. Fabbriatore, S. Farinon, M. Greco

Istituto Nazionale di Fisica Nucleare e Università Degli Studi Milano-Bicocca, Milano, ITALY

G. Cattaneo, A. De Min, M. Dominoni, F.M. Farina, F. Ferri, A. Ghezzi, P. Govoni, R. Leporini, S. Magni, M. Malberti, S. Malvezzi, S. Marelli, D. Menasce, L. Moroni, P. Negri, M. Paganoni, D. Pedrini, A. Pullia, S. Ragazzi, N. Redaelli, C. Rovelli, M. Rovere, L. Sala, S. Sala, R. Salerno, T. Tabarelli de Fatis, S. Viganò

Istituto Nazionale di Fisica Nucleare de Napoli (INFN), Napoli, ITALY

G. Comunale, F. Fabozzi, D. Lomidze, S. Mele, P. Paolucci, D. Piccolo, G. Polese, C. Sciacca

Università di Padova e Sezione dell' INFN, Padova, ITALY

P. Azzi, N. Bacchetta^{**1}, M. Bellato, M. Benettoni, D. Bisello, E. Borsato, A. Candelori, P. Checchia, E. Conti, M. De Mattia, T. Dorigo, V. Drollinger, F. Fanzago, F. Gasparini, U. Gasparini, M. Giarin, P. Giubilato, F. Gonella, A. Kaminskiy, S. Karaevskii, V. Khomenkov, S. Lacaprara, I. Lippi, M. Loretì,

O. Lytovchenko, M. Mazzucato, A.T. Meneguzzo, M. Michelotto, F. Montecassiano^{**1}, M. Nigro, M. Passaseo, M. Pegoraro, G. Rampazzo, P. Ronchese, E. Torassa, S. Ventura, M. Zanetti, P. Zotto, G. Zumerle

Università di Pavia e Sezione dell' INFN, Pavia, ITALY

G. Belli, U. Berzano, C. De Vecchi, R. Guida, M.M. Necchi, S.P. Ratti, C. Riccardi, G. Sani, P. Torre, P. Vitulo

Università di Perugia e Sezione dell' INFN, Perugia, ITALY

F. Ambroglini, E. Babucci, D. Benedetti, M. Biasini, G.M. Bilei^{**1}, B. Caponeri, B. Checcucci, L. Fanò, P. Lariccia, G. Mantovani, D. Passeri, M. Pioppi, P. Placidi, V. Postolache, D. Ricci^{**1}, A. Santocchia, L. Servoli, D. Spiga

Università di Pisa, Scuola Normale Superiore e Sezione dell' INFN, Pisa, ITALY

P. Azzurri, G. Bagliesi, A. Basti, L. Benucci, J. Bernardini, T. Boccali, L. Borrello, F. Bosi, F. Calzolari, R. Castaldi, C. Cerri, A.S. Cucoanes, M. D'Alfonso, R. Dell'Orso, S. Dutta, L. Foà, S. Gennai^{**13}, A. Giammanco, A. Giassi, D. Kartashov, F. Ligabue, S. Linari, T. Lomtadze, G.A. Lungu, B. Mangano, G. Martinelli, M. Massa, A. Messineo, A. Moggi, F. Palla, F. Palmonari, G. Petrucciani, F. Raffaelli, A. Rizzi, G. Sanguinetti, G. Segneri, D. Sentenac, A.T. Serban, G. Sguazzoni, A. Slav, P. Spagnolo, R. Tenchini, G. Tonelli, A. Venturi, P.G. Verdini, M. Vos

Università di Roma I e Sezione dell' INFN, Roma, ITALY

S. Baccaro^{**14}, L. Barone, A. Bartoloni, F. Cavallari, S. Costantini, I. Dafinei, D. Del Re^{**9}, M. Diemoz, C. Gargiulo, E. Longo, P. Meridiani, G. Organtini, S. Rahatlou

Università di Torino e Sezione dell' INFN, Torino, ITALY

E. Accomando, M. Arneodo^{**15}, A. Ballestrero, R. Bellan, C. Biino, S. Bolognesi, N. Cartiglia, G. Cerminara, M. Cordero, M. Costa, G. Dellacasa, N. Demaria, E. Maina, C. Mariotti, S. Maselli, P. Mereu, E. Migliore, V. Monaco, M. Nervo, M.M. Obertino, N. Pastrone, G. Petrillo, A. Romero, M. Ruspa^{**15}, R. Sacchi, A. Staiano, P.P. Trapani

Università di Trieste e Sezione dell' INFN, Trieste, ITALY

S. Belforte, F. Cossutti, G. Della Ricca, A. Penzo

Kyungpook National University, Daegu, KOREA

K. Cho, S.W. Ham, D. Han, D.H. Kim, G.N. Kim, J.C. Kim, W.Y. Kim, M.W. Lee, S.K. Oh, W.H. Park, S.R. Ro, D.C. Son, J.S. Suh

Chonnam National University, Kwangju, KOREA

J.Y. Kim

Konkuk University, Seoul, KOREA

S.Y. Jung, J.T. Rhee

Korea University, Seoul, KOREA

B.S. Hong, S.J. Hong, K.S. Lee, I. Park, S.K. Park, K.S. Sim, E. Won

Seoul National University, Seoul, KOREA

S.B. Kim

Universidad Iberoamericana, Mexico City, MEXICO

S. Carrillo Moreno

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, MEXICO

H. Castilla Valdez, A. Sanchez Hernandez

Benemerita Universidad Autonoma de Puebla, Puebla, MEXICO

H.A. Salazar Ibarguen

Universidad Autonoma de San Luis Potosi, San Luis Potosi, MEXICO

A. Morelos Pineda

University of Auckland, Auckland, NEW ZEALAND

R.N.C. Gray, D. Krofcheck

University of Canterbury, Christchurch, NEW ZEALAND

N. Bernardino Rodrigues, P.H. Butler, J.C. Williams

National Centre for Physics, Quaid-I-Azam University, Islamabad, PAKISTAN

Z. Aftab, M. Ahmad, U. Ahmad, I. Ahmed, J. Alam Jan, M.I. Asghar, S. Asghar, M. Hafeez, H.R. Hoorani, M. Ibrahim, M. Iftikhar, M.S. Khan, N. Qaiser, I. Rehman, T. Solaija, S. Toor

Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, POLAND

J. Blocki, A. Cyz, E. Gladysz-Dziadus, S. Mikocki, J. Turnau, Z. Wlodarczyk^{**16}, P. Zychowski

Institute of Experimental Physics, Warsaw, POLAND

K. Bunkowski, H. Czyrkowski, R. Dabrowski, W. Dominik, K. Doroba, A. Kalinowski, M. Konecki, J. Krolkowski, I.M. Kudla, M. Pietrusinski, K. Pozniak^{**17}, W. Zabolotny^{**17}, P. Zych

Soltan Institute for Nuclear Studies, Warsaw, POLAND

M. Bluj, R. Gokieli, L. Gosciolo, M. Górski, K. Nawrocki, P. Traczyk, G. Wrochna, P. Zalewski

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, PORTUGAL

R. Alemany-Fernandez, C. Almeida, N. Almeida, A. Araujo Trindade, P. Bordalo, P. Da Silva Rodrigues, M. Husejko, A. Jain, M. Kazana, P. Musella, S. Ramos, J. Rasteiro Da Silva, P.Q. Ribeiro, M. Santos, J. Semiao, P. Silva, I. Teixeira, J.P. Teixeira, J. Varela^{**1}

Joint Institute for Nuclear Research, Dubna, RUSSIA

S. Afanasiev, K. Babich, I. Belotelov, V. Elsha, Y. Ershov, I. Filozova, A. Golunov, I. Golutvin, N. Gorbounov, I. Gramenitski, V. Kalagin, A. Kamenev, V. Karjavin, S. Khabarov, V. Khabarov, Y. Kiryushin, V. Konoplyanikov, V. Korenkov, G. Kozlov, A. Kurenkov, A. Lanev, V. Lysiakov, A. Malakhov, I. Melnitchenko, V.V. Mitsyn, K. Moisenz, P. Moisenz, S. Movchan, E. Nikonov, D. Oleynik, V. Palichik, V. Perelygin, A. Petrosyan, E. Rogalev, V. Samsonov, M. Savina, R. Semenov, S. Shmatov, S. Shulha, V. Smirnov, D. Smolin, A. Tcheremoukhine, O. Teryaev, E. Tikhonenko, S. Vassiliev, A. Vishnevskiy, A. Volodko, N. Zamiatin, A. Zarubin, P. Zarubin, E. Zubarev

Petersburg Nuclear Physics Institute, Gatchina (St Petersburg), RUSSIA

N. Bondar, V. Golovtsov, A. Golyash, Y. Ivanov, V. Kim, V. Kozlov, V. Lebedev, G. Makarenkov, E. Orishchin, A. Shevel, V. Sknar, I. Smirnov, V. Sulimov, V. Tarakanov, L. Uvarov, G. Velichko, S. Volkov, A. Vorobyev

Institute for Nuclear Research, Moscow, RUSSIA

Yu. Andreev, A. Anisimov, S. Gninenko, N. Golubev, D. Gorbunov, M. Kirsanov, A. Kovzelev, N. Krasnikov, V. Matveev, A. Pashenkov, V.E. Postoev, A. Sadovski, A. Solovey, A. Solovey, D. Soloviev, L. Stepanova, A. Toropin

Institute for Theoretical and Experimental Physics, Moscow, RUSSIA

V. Gavrilov, N. Ilina, V. Kaftanov^{**1}, I. Kiselevich, V. Kolosov, M. Kossov^{**1}, A. Krokhotin, S. Kuleshov, A. Oulianov, G. Safronov, S. Semenov, V. Stolin, E. Vlasov^{**1}, V. Zaytsev

P.N. Lebedev Physical Institute, Moscow, RUSSIA

A.M. Fomenko, N. Konovalova, V. Kozlov, A.I. Lebedev, N. Lvova, S.V. Rusakov, A. Terkulov

Moscow State University, Moscow, RUSSIA

E. Boos, M. Dubinin^{**3}, L. Dudko, A. Ershov, A. Gribushin, V. Ilyin, V. Klyukhin^{**1}, O. Kodolova, I. Lokhtin, S. Petrushanko, L. Sarycheva, V. Savrin, A. Sherstnev, A. Snigirev, K. Teplov, I. Vardanyan

State Research Center of Russian Federation - Institute for High Energy Physics, Protvino, RUSSIA

V. Abramov, I. Azhguirei, S. Bitioukov, K. Datsko, A. Filine, P. Goncharov, V. Grishin, A. Inyakin, V. Kachanov, A. Khmelnikov, D. Konstantinov, A. Korablev, V. Krychkine, A. Levine, I. Lobov, V. Petrov, V. Pikalov, R. Ryutin, S. Slabospitsky, A. Sourkov^{**1}, A. Sytine, L. Tourtchanovitch,

S. Troshin, N. Tyurin, A. Uzunian, A. Volkov, S. Zelepoukine**18

Vinca Institute of Nuclear Sciences, Belgrade, SERBIA

P. Adzic, D. Krpic**19, D. Maletic, P. Milenovic, J. Puzovic**19, N. Smiljkovic**1, M. Zupan

Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, Madrid, SPAIN

M. Aguilar-Benitez, J. Alberdi, J. Alcaraz Maestre, M. Aldaya Martin, P. Arce**1, J.M. Barcala, C. Burgos Lazaro, J. Caballero Bejar, E. Calvo, M. Cardenas Montes, M. Cerrada, M. Chamizo Llatas, N. Colino, M. Daniel, B. De La Cruz, C. Fernandez Bedoya, A. Ferrando, M.C. Fouz, P. Garcia-Abia, J.M. Hernandez, M.I. Josa, J.M. Luque, J. Marin, G. Merino, A. Molinero, J.J. Navarrete, J.C. Oller, E. Perez Calle, L. Romero, J. Salicio, C. Villanueva Munoz, C. Willmott, C. Yuste

Universidad Autónoma de Madrid, Madrid, SPAIN

C. Albajar, J.F. de Trocóniz, M. Fernandez, I. Jimenez, R.F. Teixeira

Universidad de Oviedo, Oviedo, SPAIN

J. Cuevas, J.M. Lopez, H. Naves Sordo, J.M. Vizán García

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, SPAIN

A. Calderon, D. Cano Fernandez, I. Diaz Merino, L.A. Garcia Moral, G. Gomez, I. Gonzalez, J. Gonzalez Sanchez, A. Lopez Virto, J. Marco, R. Marco, C. Martinez Rivero, P. Martinez Ruiz del Arbol, F. Matorras, A. Patino Revuelta**1, T. Rodrigo, D. Rodriguez Gonzalez, A. Ruiz Jimeno, M. Sobron Sanudo, I. Vila, R. Vilar Cortabitarte

CERN, European Organization for Nuclear Research, Geneva, SWITZERLAND

D. Abbaneo, S.M. Abbas, L. Agostino, I. Ahmed, S. Akhtar, N. Amapane, B. Araujo Meleiro, S. Argiro**20, S. Ashby, P. Aspell, E. Auffray, M. Axer, A. Ball, N. Bangert, D. Barney, C. Bernet, W. Bialas, C. Bloch, P. Bloch, S. Bonacini, M. Bosteels, V. Boyer, A. Branson, A.M. Brett, H. Breuker, R. Bruneliere, O. Buchmuller, D. Campi, T. Camporesi, E. Cano, E. Carrone, A. Cattai, R. Chierici, T. Christiansen, S. Cittolin, E. Corrin, M. Corvo, S. Cucciarelli, B. Curé, A. De Roeck, D. Delikaris, M. Della Negra, A. Dierlamm, A. Elliott-Peisert, M. Eppard, H. Foeth, R. Folch, S. Fratianni, W. Funk, A. Gaddi, M. Gastal, J.C. Gayde, H. Gerwig, K. Gill, A.S. Giolo-Nicollerat, F. Glege, R. Gomez-Reino Garrido, R. Goudard, J. Gutleber, M. Hansen, A. Hervé, H.F. Hoffmann, A. Honma, M. Huhtinen, G. Iles, V. Innocente, W. Jank, P. Janot, K. Kloukinas, C. Lasseur, M. Lebeau, P. Lecoq, C. Leonidopoulos, M. Letheren, C. Ljuslin, R. Loos, G. Magazzu, L. Malgeri, M. Mannelli, A. Marchioro, F. Meijers, E. Meschi, R. Moser, M. Mulders, J. Nash, R.A. Ofierzynski, A. Oh, P. Olbrechts, A. Onnela, L. Orsini, I. Pal, G. Papotti, R. Paramatti, G. Passardi, B. Perea Solano, G. Perinic, P. Petagna, A. Petrilli, A. Pfeiffer, M. Pimiä, R. Pintus, H. Postema, R. Principe, J. Puerta Pelayo, A. Racz, J. Rehn, S. Reynaud, M. Risoldi, P. Rodrigues Simoes Moreira, G. Rolandi, P. Rosinsky, P. Rumerio, H. Sakulin, D. Samyn, F.P. Schilling, C. Schwick, C. Schäfer, I. Segoni, A. Sharma, P. Siegrist, N. Sinanis, P. Sphicas**21, M. Spiropulu, F. Szoncsó, O. Teller, J. Troska, E. Tsemelis, D. Tsirigkas, A. Tsirou, D. Ungaro, F. Vasey, M. Vazquez Acosta, L. Veillet, P. Vichoudis, P. Wertelaers, A. Wijnant, M. Wilhelmsson, I.M. Willers

Paul Scherrer Institut, Villigen, SWITZERLAND

W. Bertl, K. Deiters, W. Erdmann, K. Gabathuler, S. Heising, R. Horisberger, Q. Ingram, H.C. Kaestli, D. Kotlinski, S. König, D. Renker, T. Rohe, M. Spira

Institute for Particle Physics, ETH Zurich, Zurich, SWITZERLAND

B. Betev, G. Davatz, G. Dissertori, M. Dittmar, L. Djambazov, J. Ehlers, R. Eichler, G. Faber, K. Freudenreich, J.F. Fuchs**1, C. Grab, A. Holzner, P. Ingenito, U. Langenegger, P. Lecomte, G. Leshev, A. Lister**22, P.D. Luckey, W. Lustermann, J.D. Maillefaud**1, F. Moortgat, A. Nardulli, F. Nessi-Tedaldi, L. Pape, F. Pauss, H. Rykaczewski**23, U. Röser, D. Schinzel, A. Starodumov**24, F. Stöckli, H. Suter, L. Tauscher, P. Trüb**25, H.P. von Gunten, M. Wensveen**1

Universität Zürich, Zürich, SWITZERLAND

E. Alagoz, C. AMSLER, V. Chiochia, C. Hoermann, K. Prokofiev, C. Regenfus, P. Robmann, T. Speer, S. Steiner, L. Wilke

National Central University, Chung-Li, TAIWAN

S. Blyth, Y.H. Chang, E.A. Chen, A. Go, C.C. Hung, C.M. Kuo, W. Lin

National Taiwan University (NTU), Taipei, TAIWAN

P. Chang, Y. Chao, K.F. Chen, Z. Gao^{**1}, Y. Hsiung, Y.J. Lei, J. Schümann, J.G. Shiu, K. Ueno, Y. Velikzhanin, P. Yeh

Cukurova University, Adana, TURKEY

S. Aydin, M.N. Bakirci, S. Cerci, I. Dumanoglu, S. Erturk, S. Esen, E. Eskut, A. Kayis Topaksu, P. Kurt, H. Ozkurt, A. Polatöz, K. Sogut, H. Topakli, M. Vergili, T. Yetkin, G. Önengüt

Middle East Technical University, Physics Department, Ankara, TURKEY

H. Gamsizkan, C. Ozkan, S. Sekmen, M. Serin-Zeyrek, R. Sever, E. Yazgan, M. Zeyrek

Bogaziçi University, Department of Physics, Istanbul, TURKEY

A. Cakir^{**26}, K. Cankocak^{**27}, M. Deliomeroglu, D. Demir^{**26}, K. Dindar, E. Gülmez, E. Isiksal^{**28}, M. Kaya^{**29}, O. Kaya, S. Ozkorucuklu^{**30}, N. Sonmez^{**31}

Institute of Single Crystals of National Academy of Science, Kharkov, UKRAINE

B. Grinev, V. Lyubynskiy, V. Senchyshyn

National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, UKRAINE

L. Levchuk, P. Sorokin

University of Bristol, Bristol, UNITED KINGDOM

D.S. Bailey, T. Barrass, J.J. Brooke, R. Croft, D. Cussans, D. Evans, R. Frazier, N. Grant, M. Hansen, G.P. Heath, H.F. Heath, B. Huckvale, C. Lynch, C.K. Mackay, S. Metson, D.M. Newbold^{**32}, V.J. Smith, R.J. Tapper

Rutherford Appleton Laboratory, Didcot, UNITED KINGDOM

S.A. Baird, K.W. Bell, R.M. Brown, D.J.A. Cockerill, J.A. Coughlan, P.S. Flower, V.B. Francis, M. French, J. Greenhalgh, R. Halsall, J. Hill, L. Jones, B.W. Kennedy, L. Lintern, A.B. Lodge, J. Maddox, Q. Morrissey, P. Murray, M. Pearson, S. Quinton, J. Salisbury, A. Shah, C. Shepherd-Themistocleous, B. Smith, M. Sproston, R. Stephenson, S. Taghavirad, I.R. Tomalin, J.H. Williams

Imperial College, University of London, London, UNITED KINGDOM

F. Arteche^{**1}, R. Bainbridge, G. Barber, P. Barrillon, R. Beuselinck, F. Blekman, D. Britton, D. Colling, G. Daskalakis, G. Dewhurst, S. Dris^{**1}, C. Foudas, J. Fulcher, S. Greder, G. Hall, J. Jones, J. Leaver, B.C. MacEvoy, O. Maroney, A. Nikitenko^{**24}, A. Papageorgiou, D.M. Raymond, M.J. Ryan, C. Seez, P. Sharp^{**1}, M. Takahashi, C. Timlin, T. Virdee^{**1}, S. Wakefield, M. Wingham, A. Zabi, Y. Zhang, O. Zorba

Brunel University, Uxbridge, UNITED KINGDOM

C. Da Via, I. Goitom, P.R. Hobson, P. Kyberd, C. Munro, J. Nebrensky, I. Reid, O. Sharif, R. Taylor, L. Teodorescu, S.J. Watts, I. Yaselli

Boston University, Boston, Massachusetts, USA

E. Hazen, A.H. Heering, D. Lazic, E. Machado, D. Osborne, J. Rohlf, L. Sulak, F. Varela Rodriguez, S. Wu

Brown University, Providence, Rhode Island, USA

D. Cutts, R. Hooper, G. Landsberg, R. Partridge, S. Vanini^{**33}

University of California, Davis, Davis, California, USA

R. Breedon, M. Case, M. Chertok, J. Conway, P.T. Cox, R. Erbacher, J. Gunion, B. Holbrook, W. Ko, R. Lander, D. Pellett, J. Smith, A. Soha, M. Tripathi, R. Vogt

University of California, Los Angeles, Los Angeles, California, USA

V. Andreev, K. Arisaka, D. Cline, R. Cousins, S. Erhan^{**1}, M. Felcini^{**1}, J. Hauser, M. Ignatenko, B. Lisowski, D. Matlock, C. Matthey, B. Mohr, J. Mumford, S. Otwinowski, G. Rakness, P. Schlein,

Y. Shi, J. Tucker, V. Valuev, R. Wallny, H.G. Wang, X. Yang, Y. Zheng

University of California, Riverside, Riverside, California, USA

R. Clare, D. Fortin, D. Futyan^{**1}, J.W. Gary, M. Giunta^{**1}, G. Hanson, G.Y. Jeng, S.C. Kao, H. Liu, G. Pasztor^{**34}, A. Satpathy, B.C. Shen, R. Stringer, V. Sytnik, R. Wilken, D. Zer-Zion

University of California, San Diego, La Jolla, California, USA

J.G. Branson, E. Dusinberre, J. Letts, T. Martin, M. Mojaver, H.P. Paar, H. Pi, M. Pieri, A. Rana, V. Sharma, A. White, F. Würthwein

University of California, Santa Barbara, Santa Barbara, California, USA

A. Affolder, C. Campagnari, C. Hill, J. Incandela, S. Kyre, J. Lamb, J. Richman, D. Stuart, D. White

California Institute of Technology, Pasadena, California, USA

J. Albert, A. Bornheim, J. Bunn, J. Chen, G. Denis, P. Galvez, M. Gataullin, I. Legrand, V. Litvine, Y. Ma, D. Nae, H.B. Newman, S. Ravot, S. Shevchenko, S. Singh, C. Steenberg, X. Su, M. Thomas, V. Timciuc, F. van Lingen, J. Veverka, B.R. Voicu^{**1}, A. Weinstein, R. Wilkinson, X. Yang, Y. Yang, L.Y. Zhang, K. Zhu, R.Y. Zhu

Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

T. Ferguson, M. Paulini, J. Russ, N. Terentyev, H. Vogel, I. Vorobiev

University of Colorado at Boulder, Boulder, Colorado, USA

J.P. Cumalat, W.T. Ford, D. Johnson, U. Nauenberg, K. Stenson, S.R. Wagner

Cornell University, Ithaca, NY, USA

J. Alexander, D. Cassel, K. Ecklund, B. Heltsley, C.D. Jones, V. Kuznetsov, J.R. Patterson, A. Ryd, J. Thom, P. Wittich

Fairfield University, Fairfield, Connecticut, USA

C.P. Beetz, G. Cirino, V. Podrasky, C. Sanzeni, D. Winn

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

S. Abdullin^{**24}, M.A. Afaq^{**1}, M. Albrow, J. Amundson, G. Apollinari, M. Atac, W. Badgett, J.A. Bakken, B. Baldin, L.A.T. Bauerdick, A. Baumbaugh, U. Baur, P.C. Bhat, F. Borchering, K. Burkett, J.N. Butler, H. Cheung, I. Churin, S. Cihangir, M. Demarteau, D.P. Eartly, J.E. Elias, V.D. Elvira, D. Evans, I. Fisk, J. Freeman, P. Gartung, F.J.M. Geurts, D.A. Glenzinski, E. Gottschalk, G. Graham, D. Green, G.M. Guglielmo, Y. Guo, O. Gutsche, A. Hahn, J. Hanlon, S. Hansen, R.M. Harris, T. Hesselroth, S.L. Holm, B. Holzman, S. Iqbal, E. James, M. Johnson, U. Joshi, B. Klima, J. Kowalkowski, T. Kramer, S. Kwan, E. La Vallie, M. Larwill, S. Los, L. Lueking, G. Lukhanin, S. Lusin^{**1}, K. Maeshima, P. McBride, S.J. Murray, V. O'Dell, M. Paterno, J. Patrick, D. Petravick, R. Pordes, O. Prokofyev, V. Rasmislovich, N. Ratnikova, A. Ronzhin, V. Sekhri, E. Sexton-Kennedy, T. Shaw, D. Skow, R.P. Smith, W.J. Spalding, L. Spiegel, M. Stavrianakou, G. Stiehr, I. Suzuki, P. Tan, W. Tanenbaum, S. Tkaczyk, S. Veseli, R. Vidal, H. Wenzel, J. Whitmore, W.J. Womersley, W.M. Wu, Y. Wu, A. Yagil, J. Yarba, J.C. Yun

University of Florida, Gainesville, Florida, USA

D. Acosta, P. Avery, V. Barashko, P. Bartalini, D. Bourilkov, R. Cavanaugh, A. Drozdetskiy, R.D. Field, Y. Fu, L. Gray, D. Holmes, B.J. Kim, S. Klimenko, J. Konigsberg, A. Korytov, K. Kotov, P. Levchenko, A. Madorsky, K. Matchev, G. Mitselmakher, Y. Pakhotin, C. Prescott, P. Ramond, J.L. Rodriguez, M. Schmitt, B. Scurlock, H. Stoeck, J. Yelton

Florida International University, Miami, Florida, USA

W. Boeglin, V. Gaultney, L. Kramer, S. Linn, P. Markowitz, G. Martinez, B. Raue, J. Reinhold

Florida State University, Tallahassee, Florida, USA

A. Askew, M. Bertoldi, W.G.D. Dharmaratna, Y. Gershtein, S. Hagopian, V. Hagopian, M. Jenkins, K.F. Johnson, H. Prosper, H. Wahl

Florida Institute of Technology, Melbourne, Florida, USA

M. Baarmand, L. Baksay**35, S. Guragain, M. Hohlmann, H. Mermerkaya, R. Ralich, I. Vodopyanov

University of Illinois at Chicago (UIC), Chicago, Illinois, USA

M.R. Adams, R.R. Betts, C.E. Gerber, E. Shabalina, C. Smith, T. Ten

The University of Iowa, Iowa City, Iowa, USA

U. Akgun, A.S. Ayan, A. Cooper, P. Debbins, F. Duru, M. Fountain, N. George, E. McCliment, J.P. Merlo, A. Mestvirishvili, M.J. Miller, C.R. Newsom, E. Norbeck, Y. Onel, I. Schmidt, S. Wang

Iowa State University, Ames, Iowa, USA

E.W. Anderson, O. Atramentov, J.M. Hauptman, J. Lamsa

Johns Hopkins University, Baltimore, Maryland, USA

B.A. Barnett, B. Blumenfeld, C.Y. Chien, D.W. Kim, P. Maksimovic, S. Spangler, M. Swartz

The University of Kansas, Lawrence, Kansas, USA

P. Baringer, A. Bean, D. Coppage, O. Grachov, E.J. Kim, M. Murray

Kansas State University, Manhattan, Kansas, USA

D. Bandurin, T. Bolton, A. Khanov**24, Y. Maravin, D. Onoprienko, F. Rizatdinova, R. Sidwell, N. Stanton, E. Von Toerne

University of Maryland, College Park, Maryland, USA

D. Baden, R. Bard, S.C. Eno, T. Grassi, N.J. Hadley, R.G. Kellogg, S. Kunori, F. Ratnikov, A. Skuja

Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

R. Arcidiacono, M. Ballintijn, G. Bauer, P. Harris, I. Kravchenko, C. Loizides, S. Nahn, C. Paus, S. Pavlon, C. Roland, G. Roland, K. Sumorok, S. Vaurynovich, G. Veres, B. Wyslouch

University of Minnesota, Minneapolis, Minnesota, USA

D. Bailleux, S. Corum, P. Cushman, A. De Benedetti, A. Dolgoplov, R. Egeland, G. Franzoni, W.J. Gilbert, J. Grahl, J. Haupt, Y. Kubota, J. Mans, N. Pearson, R. Rusack, A. Singovsky

University of Mississippi, University, Mississippi, USA

L.M. Cremaldi, R. Godang, R. Kroeger, D.A. Sanders, D. Summers

University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom, D.R. Claes, A. Dominguez, M. Eads, C. Lundstedt, S. Malik, G.R. Snow, A. Sobol

State University of New York at Buffalo, Buffalo, New York, USA

I. Iashvili, A. Kharchilava

Northeastern University, Boston, Massachusetts, USA

G. Alverson, E. Barberis, O. Boeriu, G. Eulisse, Y. Musienko**36, S. Muzaffar, I. Osborne, S. Reucroft, J. Swain, L. Taylor, L. Tuura, D. Wood

Northwestern University, Evanston, Illinois, USA

B. Gobbi, M. Kubantsev, H. Schellman, M. Schmitt, E. Spencer, M. Velasco

University of Notre Dame, Notre Dame, Indiana, USA

B. Baumbaugh, N.M. Cason, M. Hildreth, D.J. Karmgard, N. Marinelli**21, R. Ruchti, J. Warchol, M. Wayne

The Ohio State University, Columbus, Ohio, USA

B. Bylsma, L.S. Durkin, J. Gilmore, J. Gu, D. Herman, P. Killewald, K. Knobbe, T.Y. Ling

Princeton University, Princeton, New Jersey, USA

P. Elmer, D. Marlow, P. Piroué, D. Stickland, C. Tully, T. Wildish, S. Wynhoff, Z. Xie

Purdue University, West Lafayette, Indiana, USA

A. Apresyan, K. Arndt, K. Banicz, V.E. Barnes, G. Bolla, D. Bortoletto, A. Bujak, A.F. Garfinkel, O. Gonzalez Lopez, L. Gutay, N. Ippolito, Y. Kozhevnikov**1, A.T. Laasanen, C. Liu, V. Maroussov, P. Merkel, D.H. Miller, J. Miyamoto, N. Neumeister, C. Rott, A. Roy, A. Sedov, I. Shipsey

Purdue University Calumet, Hammond, Indiana, USA

N. Parashar

Rice University, Houston, Texas, USA

G. Eppley, S.J. Lee, J. Liu, M. Matveev, T. Nussbaum, B.P. Padley, J. Roberts, A. Tumanov, P. Yepes

University of Rochester, Rochester, New York, USA

A. Bodek, H. Budd, Y.S. Chung, P. De Barbaro^{**1}, R. Demina, R. Eusebi, G. Ginther, Y. Gotra, A. Hocker, U. Husemann, S. Korjenevski, W. Sakumoto, P. Slattery, P. Tipton, M. Zielinski

Rutgers, the State University of New Jersey, Piscataway, New Jersey, USA

E. Bartz, J. Doroshenko, E. Halkiadakis, P.F. Jacques, M.S. Kalelkar, D. Khits, A. Lath, A. Macpherson^{**1}, L. Perera, R. Plano, K. Rose, S. Schnetzer, S. Somalwar, R. Stone, G. Thomson, T.L. Watts

Texas Tech University, Lubbock, Texas, USA

N. Akchurin, K.W. Carrell, K. Gumus, C. Jeong, H. Kim, V. Papadimitriou, A. Sill, M. Spezziga, E. Washington, R. Wigmans, L. Zhang

Vanderbilt University, Nashville, Tennessee, USA

T. Bapty, D. Engh, W. Johns, T. Keskinpala, E. Luiggi Lopez, S. Neema, S. Nordstrom, S. Pathak, P. Sheldon, E.W. Vaandering, M. Webster

University of Virginia, Charlottesville, Virginia, USA

M.W. Arenton, S. Conetti, B. Cox, R. Hirosky, R. Imlay, A. Ledovskoy, D. Phillips II, H. Powell, M. Ronquest, D. Smith

University of Wisconsin, Madison, Wisconsin, USA

Y.W. Baek, J.N. Bellinger, D. Bradley, D. Carlsmith, I. Crotty^{**1}, S. Dasu, F. Feyzi, T. Gorski, M. Grothe^{**37}, W. Hogg, M. Jaworski, P. Klabbers, A. Lanaro, R. Loveless, M. Magrans de Abril, D. Reeder, W.H. Smith, D. Wenman

Yale University, New Haven, Connecticut, USA

G.S. Atoyan^{**36}, S. Dhawan, V. Issakov, H. Neal, A. Poblaguev, M.E. Zeller

Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Ulugbek, Tashkent, UZBEKISTAN

B.S. Yuldashev

^{**1}: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

^{**2}: Also at University of Zagreb, Zagreb, Croatia

^{**3}: Also at California Institute of Technology, Pasadena, USA

^{**4}: Also at Université de Haute-Alsace, Mulhouse, France

^{**5}: Also at Université Louis Pasteur, Strasbourg, France

^{**6}: Now at Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

^{**7}: Also at Moscow State University, Moscow, Russia

^{**8}: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary

^{**9}: Also at University of California, San Diego, La Jolla, USA

^{**10}: Also at Tata Institute of Fundamental Research - HECR, Mumbai, India

^{**11}: Also at University of Visva-Bharati, Santiniketan, India

^{**12}: Also at University of California, Riverside, Riverside, USA

^{**13}: Also at Centro Studi Enrico Fermi, Roma, Italy

^{**14}: Also at ENEA - Casaccia Research Center, S. Maria di Galeria, Italy

- **15: Now at Università del Piemonte Orientale, Novara, Italy
- **16: Also at Institute of Physics, Swietokrzyska Academy, Kielce, Poland
- **17: Also at Warsaw University of Technology, Institute of Electronic Systems, Warsaw, Poland
- **18: Also at Institute for Particle Physics, ETH Zurich, Zurich, Switzerland
- **19: Also at Faculty of Physics of University of Belgrade, Belgrade, Serbia
- **20: Also at INFN-CNAF, Bologna, Italy
- **21: Also at University of Athens, Athens, Greece
- **22: Now at University of California, Davis, Davis, USA
- **23: Now at ESO, Munich-Garching, Germany
- **24: Also at Institute for Theoretical and Experimental Physics, Moscow, Russia
- **25: Also at Paul Scherrer Institut, Villigen, Switzerland
- **26: Also at Izmir Institute of Technology (IYTE), Izmir, Turkey
- **27: Also at Mugla University, Mugla, Turkey
- **28: Also at Marmara University, Istanbul, Turkey
- **29: Also at Kafkas University, Kars, Turkey
- **30: Also at Suleyman Demirel University, Isparta, Turkey
- **31: Also at Ege University, Izmir, Turkey
- **32: Also at Rutherford Appleton Laboratory, Didcot, United Kingdom
- **33: Also at Università di Padova e Sezione dell' INFN, Padova, Italy
- **34: Also at KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary
- **35: Also at University of Debrecen, Debrecen, Hungary
- **36: Also at Institute for Nuclear Research, Moscow, Russia
- **37: Also at Università di Torino e Sezione dell' INFN, Torino, Italy



Executive Summary

The Large Hadron Collider will provide extraordinary opportunities in particle physics based on its unprecedented collision energy and luminosity when it begins operation in 2007. The principal aim of this Technical Design Report is to present the strategy of CMS to explore the rich physics programme offered by the LHC: Volume 1 covering the operational procedures and reconstruction tools necessary to perform physics at the LHC, and Volume 2 demonstrating the physics capability of the CMS experiment based on this foundation. A description of the procedures and reconstruction tools specifically for LHC start-up, including the performance of the High-Level Trigger algorithms and the early physics opportunities, will be published in an addendum to this Report.

In the first volume we highlight the final detector configuration as it will appear shortly after LHC start-up, including new detectors in the forward regions and for determining the beam luminosity. Results on the performance of the CMS detectors as obtained from detailed simulations are presented for realistic operating conditions, and validated where possible against test-beam or cosmic-ray data. Schemes to synchronize, calibrate, align, and monitor the detectors before, during and after LHC start-up are also given. Reconstruction algorithms developed to perform measurements of muons, electrons, photons, jets, taus, heavy-flavour quarks and the missing transverse energy using these detector subsystems are described. The performance of the reconstruction algorithms is determined from detailed simulations for realistic operating conditions, but techniques to measure the performance from LHC data are described as well. Parameterizations of the performance have been derived to facilitate faster simulations for some of the physics studies described in Volume 2. Included in this first volume are descriptions of the software components needed to implement all of the above, covering simulation, reconstruction, calibration and alignment, monitoring, and visualization.

The second volume covers the capability of the CMS experiment to address physics at the LHC. The prime goals of CMS are to explore physics at the TeV scale and to study the mechanism of electroweak symmetry breaking—through the discovery of the Higgs particle or otherwise. To carry out this task, CMS must be prepared to search for new particles, such as the Higgs boson or supersymmetric partners of the Standard Model particles, from the start-up of the LHC since new physics at the TeV scale may manifest itself with modest data samples of the order of a few fb^{-1} or less. The experience of the Magnet Test and Cosmic Challenge, scheduled for second quarter 2006, plays a crucial role in the preparation of CMS experiment, whereby calibration, alignment and reconstruction procedures are tested and made ready in advance of the LHC pilot and first physics runs. Lessons drawn from this test, as well as the plans for the first physics runs, will be described in an addendum to this Report.

The tools that have been prepared in Volume 1 are applied in Volume 2 to study in great detail and with all the methodology of performing an analysis on CMS data specific benchmark processes upon which to gauge the performance of CMS. These processes cover several

Higgs boson decay channels, the production and decay of new particles such as Z' and supersymmetric particles, B_s production and processes in heavy ion collisions. The simulation of these benchmark processes includes subtle effects such as possible detector miscalibration and misalignment. Besides these benchmark processes, the physics reach of CMS is studied for a large number of signatures arising in the Standard Model and also in theories beyond the Standard Model for integrated luminosities of 10 fb^{-1} , and in a few cases up to the asymptotic value of 300 fb^{-1} . The Standard Model processes include QCD, B -physics, diffraction, detailed studies of the top quark properties, and electroweak physics topics such as the W and Z^0 boson properties. The production and decay of the Higgs particle is studied for many observable decays, and the precision with which the Higgs boson properties can be derived is determined. About ten different supersymmetry benchmark points are analysed, and methods to extract e.g. the sparticle masses are tested. Furthermore, the discovery reach for a plethora of alternative models for new physics is explored, notably extra dimensions, new vector boson high mass states, little Higgs models, technicolour and others.

In summary, the content of these two volumes is meant to serve as a comprehensive reference for new CMS collaborators. It provides an entry point to the documentation of the standard simulation, reconstruction, and analysis tools and provides a measure of the expected detector performance and physics reach as we head into the LHC era.

Structure of Volume 2

Chapter 1, the Introduction, describes the context of this document.

Chapters 2-6 describe examples of full analyses, with photons, electrons, muons, jets, missing E_T , B -mesons and τ 's, and for quarkonia in heavy ion collisions.

Chapters 7-15 describe the physics reach for Standard Model processes, Higgs discovery and searches for new physics beyond the Standard Model.



Contents

1	Introduction	1
I	Complete analyses	2
2	Physics Studies with Photons and Electrons	3
2.1	Introduction	3
2.2	Benchmark Channel: $H \rightarrow \gamma\gamma$	3
2.2.1	Higgs Boson Production and Decay	4
2.2.2	Backgrounds	4
2.2.3	Reconstruction, selection, and signal significance calculation	6
2.2.4	Cut-based Analysis	8
2.2.5	Optimized Analysis Estimating s/b for Each Event	13
2.2.6	Summary	21
2.3	Benchmark Channel: $H \rightarrow ZZ^{(*)} \rightarrow 4$ electrons	21
2.3.1	Datasets for Signal and Background Processes	22
2.3.2	Data Reduction	24
2.3.3	Event Selection and Kinematic Reconstruction	25
2.3.4	Systematics	30
2.3.5	$H \rightarrow 4e$ Observability, Mass and Cross-Section Measurements	35
3	Physics Studies with Muons	39
3.1	Benchmark Channel: $H \rightarrow ZZ^{(*)} \rightarrow 4$ muons	39
3.1.1	Physics Processes and Their Simulation	39
3.1.2	Event Selection	40
3.1.3	Higgs Boson Search Analysis	42
3.1.4	Measurement of the Higgs Boson Properties at $\mathcal{L} = 30 \text{ fb}^{-1}$	49
3.1.5	Conclusions	52
3.2	Benchmark Channel: $H \rightarrow WW^{(*)} \rightarrow 2$ muons	53
3.2.1	Introduction	53
3.2.2	Physics processes	54
3.2.3	Event selection	55
3.2.4	The trigger selection	56
3.2.5	JET reconstruction and the JET veto	58
3.2.6	Missing energy reconstruction and the MET cut	58

3.2.7	The selection results	61
3.2.8	Background estimation and systematics	62
3.2.9	$t\bar{t}$ background normalization	63
3.2.10	WW background normalization	65
3.2.11	Other backgrounds normalization	67
3.2.12	Detector misalignment systematics	68
3.2.13	Signal significance	68
3.2.14	Conclusions	69
3.3	Benchmark Channel: $Z' \rightarrow \mu\mu$	69
3.3.1	Introduction	69
3.3.2	Signal and background processes	70
3.3.3	Event selection	72
3.3.4	Signal observability	73
3.3.5	Distinguishing among Z' models	80
3.3.6	Discriminating between different spin hypotheses	81
4	Physics Studies with Jets and E_T^{miss}	85
4.1	Benchmark Channel: new physics from dijets	85
4.1.1	Dijet Analysis	85
4.1.2	Rates and Efficiencies from Jet Triggers	85
4.1.3	Dijet Mass Distribution from QCD	87
4.1.4	Searches using Dijet Mass	87
4.1.5	Searches using Dijet Mass and Angle	89
4.1.6	Systematic Uncertainties	89
4.2	Benchmark Channel: Low Mass Supersymmetry	91
4.2.1	Introduction	91
4.2.2	Jets and missing transverse energy at CMS	92
4.2.3	Clean-up requirements	93
4.2.4	Analysis Path	93
4.2.5	Missing transverse energy in QCD production	93
4.2.6	Indirect Lepton Veto	95
4.2.7	The Standard Z boson candle calibration	96
4.2.8	Analysis Results	97
4.2.9	Systematic Uncertainties	98
4.2.10	Discussion	100
5	Physics Studies with tracks, B mesons, and taus	104
5.1	Benchmark Channels: Study of the decay $B_s \rightarrow J/\psi\phi$	104
5.1.1	Introduction	104
5.1.2	Event generation	105
5.1.3	Trigger Selection	106
5.1.4	Offline Selection and Reconstruction	108
5.1.5	The maximum likelihood analysis	110

5.1.6	Result	113
5.1.7	Systematics and detector effects	114
5.1.8	Conclusion	117
5.2	Associated production of MSSM heavy neutral Higgs bosons $b\bar{b}H(A)$ with $H(A) \rightarrow \tau\tau$	118
5.2.1	Introduction	118
5.2.2	Event generation	118
5.2.3	Level-1 and High Level trigger selections	119
5.2.4	Off-line event selection strategy	119
5.2.5	Method of the Higgs boson mass reconstruction	120
5.2.6	$H \rightarrow \tau\tau \rightarrow$ two τ jet analysis	120
5.2.7	$H \rightarrow \tau\tau \rightarrow \mu$ +jet analysis	127
5.2.8	$H \rightarrow \tau\tau \rightarrow e$ +jet analysis	132
5.3	Benchmark Channels: $t\bar{t}H, H \rightarrow b\bar{b}$	138
5.3.1	Introduction	138
5.3.2	Event Generation and Simulation	139
5.3.3	L1 and High Level Trigger Selections	141
5.3.4	Reconstruction	142
5.3.5	Event Selection	145
5.3.6	Discussion of Systematic Uncertainties	151
5.3.7	Combined Significance	153
6	Physics Studies with Heavy Ions	156
6.1	Benchmark Channel: $PbPb \rightarrow Q\bar{Q} + X \rightarrow \mu^+\mu^- + X$	156
6.1.1	Simulation of Physics and Background Processes	156
6.1.2	Reconstruction and Analysis	157
6.1.3	Results	159
6.1.4	Conclusions	159
II	CMS Physics Reach	162
7	Physics of Strong Interactions	163
7.1	The structure of the proton	163
7.2	QCD and jet physics	163
7.2.1	Introduction	163
7.2.2	Jet algorithms	163
7.2.3	Trigger scheme, event selection and phase space	165
7.2.4	Input data	166
7.2.5	Jet energy calibration	166
7.2.6	NLO calculation	166
7.2.7	Experimental and theoretical uncertainties	166
7.2.8	Summary and Outlook	167

7.3	Underlying Event Studies	168
7.3.1	Definition of the physics process and status of the art	168
7.3.2	Underlying event observables discussed for charged jet events	169
7.3.3	Feasibility studies	171
7.3.4	Conclusions	173
7.4	Physics of b-quarks and hadrons	173
7.4.1	Inclusive b-quark production	173
7.4.2	Study of B_c hadrons	179
7.5	Diffraction and forward physics	183
7.5.1	Introduction	183
7.5.2	The interest of diffractive interactions	184
7.5.3	A survey of the accessible diffractive/forward processes	184
7.6	Physics with heavy ions	190
7.6.1	High-density QCD: heavy-ion physics	190
7.6.2	Hard probes of QCD matter at LHC	191
7.6.3	Gluon saturation and QGP colour screening via Quarkonia	191
8	Physics of Top Quarks	193
8.1	Selection of $t\bar{t}$ events and measurement of the cross sections	193
8.1.1	Introduction	193
8.1.2	Di-leptonic channel	193
8.1.3	Semi-leptonic channel	197
8.1.4	Fully hadronic channel	200
8.2	Measurement of the top quark mass	203
8.2.1	Di-leptonic events	203
8.2.2	Semi-leptonic events	204
8.2.3	Fully hadronic events	207
8.2.4	Top quark mass from J/ψ final states	210
8.2.5	Summary of top mass determinations	215
8.3	Spin Correlation in Top-Quark Pair Production	216
8.3.1	Introduction	216
8.3.2	Simulation of $t\bar{t}$ With Spin Correlation	217
8.3.3	Online and Offline Event Selection	217
8.3.4	Estimation of Correlation Coefficient	218
8.4	Single top quark production	220
8.4.1	Introduction	220
8.4.2	Selection and cross section - t -channel	223
8.4.3	Selection and cross section - tW -channel	225
8.4.4	Selection and cross section - s -channel	228
8.4.5	Conclusion	231
8.5	Search for flavor changing neutral currents in top decays	232
8.5.1	Introduction	232

8.5.2	Signal and background generation	233
8.5.3	Selections	233
8.5.4	Sensitivity estimation	234
9	Electroweak Physics	237
9.1	Production of W and Z bosons	237
9.1.1	Introduction	237
9.1.2	W/Z into Electrons	237
9.1.3	W/Z into Muons	241
9.1.4	Parton distribution functions and parton luminosities	245
9.2	Muon pairs from the Drell-Yan process	245
9.2.1	Introduction	245
9.2.2	Cross section measurements	246
9.2.3	Prospects on the measurement of the forward-backward asymmetry	249
9.3	Determination of the W mass	250
9.3.1	Introduction	250
9.3.2	Event selections	251
9.3.3	$W \rightarrow e\nu$	251
9.3.4	$W \rightarrow \mu\nu$	253
9.3.5	Expected precision and systematic uncertainties	254
9.4	Multi-boson production	255
9.4.1	Introduction	255
9.4.2	Signal definition and modeling	256
9.4.3	Background processes	257
9.4.4	$W^\pm Z^0$ selection	257
9.4.5	$Z^0 Z^0$ selection	258
9.4.6	Systematic uncertainties	259
9.4.7	Results	259
10	Standard Model Higgs Bosons	261
10.1	Introduction	261
10.2	Higgs boson channels	264
10.2.1	Inclusive Higgs boson production with $H \rightarrow ZZ^{(*)} \rightarrow e^+e^-\mu^+\mu^-$	264
10.2.2	Inclusive Higgs boson production with $H \rightarrow WW^{(*)} \rightarrow 2\ell 2\nu$	274
10.2.3	The vector boson fusion production with $H \rightarrow \tau\tau$	280
10.2.4	Searching for Standard Model Higgs via Vector Boson Fusion in $H \rightarrow W^+W^- \rightarrow \ell^\pm\nu jj$ with m_H from 120 to 250 GeV/ c^2	283
10.2.5	Vector boson fusion production with $H \rightarrow \gamma\gamma$	290
10.2.6	Associated WH production with $H \rightarrow WW^{(*)} \rightarrow 2\ell 2\nu$	298
10.2.7	Associated $t\bar{t}H$ production with $H \rightarrow \gamma\gamma$	307
10.2.8	Associated WH,ZH production with $H \rightarrow \gamma\gamma$	317
10.2.9	Associated WH production with $H \rightarrow WW^{(*)} \rightarrow 2\ell 2\nu$	324
10.3	Discovery reach	331

10.3.1	Accuracy of the Higgs boson mass measurement	332
10.3.2	Discovery reach for the Standard Model Higgs boson	332
10.3.3	Study of CP properties of the Higgs boson using angle correlation in the $\Phi \rightarrow ZZ \rightarrow e^+e^-\mu^+\mu^-$ process	335
11	MSSM Higgs bosons	340
11.1	Introduction	340
11.2	Higgs boson channels	349
11.2.1	Associated $b\bar{b}H$ production with $H \rightarrow \tau\tau \rightarrow e^\pm\mu^\mp + E_T^{\text{miss}}$	350
11.2.2	Associated $b\bar{b}H$ production with $H \rightarrow \mu^+\mu^-$	353
11.2.3	Associated $b\bar{b}H$ production with $H \rightarrow b\bar{b}$	359
11.2.4	Charged Higgs boson of $M_H < m_t$ in $t\bar{t} \rightarrow H^\pm W^\mp b\bar{b}$ production with $H^\pm \rightarrow \tau^\pm\nu$, $\tau \rightarrow \text{hadrons } \nu$ and $W^\mp \rightarrow \ell^\mp\nu$	364
11.2.5	Charged Higgs boson of $M_H > m_t$ in $gg \rightarrow tbH^\pm$ production with $H^\pm \rightarrow \tau^\pm\nu$, $\tau \rightarrow \text{hadrons } \nu$ and $W^\mp \rightarrow jj$	369
11.2.6	Charged Higgs boson of $M_H > m_t$ in $gg \rightarrow tbH^\pm$ production with $H^\pm \rightarrow tb$	376
11.2.7	Search for the $A \rightarrow Zh$ decay with $Z \rightarrow \ell^+\ell^-$, $h \rightarrow b\bar{b}$	380
11.2.8	Search for $A^0/H^0 \rightarrow \chi_2^0\chi_2^0 \rightarrow 4l + E_T^{\text{miss}}$ channel in mSUGRA	385
11.3	Discovery reach and measurement of MSSM parameters	387
11.3.1	Benchmark scenarios for MSSM Higgs boson searches	387
11.3.2	Discovery reach in the M_A - $\tan\beta$ plane	393
12	Search for Higgs boson in non SUSY models	397
12.1	Scalar sector of 5D Randal-Sundrum model	397
12.1.1	The $\phi \rightarrow hh$ analysis with the $\gamma\gamma b\bar{b}$ and $\tau\tau b\bar{b}$ final states.	397
12.2	Search for pair production of doubly charged Higgs boson in the Littlest Higgs model	400
12.2.1	Search for the final state with four muons	401
12.2.2	Search for the final states with τ leptons	406
13	Supersymmetry	416
13.1	Introduction	416
13.2	Summary of Supersymmetry	416
13.2.1	The MSSM	416
13.2.2	MSUGRA parameters and spectrum	417
13.3	Scope of present searches	418
13.3.1	Sparticle production and cascade decays	418
13.3.2	Test points for MSUGRA	420
13.4	Hemisphere algorithm for separation of decay chains	423
13.4.1	Basic idea and goal	423
13.4.2	Seeding methods	424
13.4.3	Association methods	425
13.4.4	Results	425

13.5	Inclusive analysis with missing transverse energy and jets	426
13.5.1	Analysis path and results	427
13.6	Inclusive analysis with missing transverse energy, jets, and muons	428
13.6.1	Signal Selection and Backgrounds Considered	429
13.6.2	Results for 10 fb^{-1} using Full Detector Simulation and Reconstruction	429
13.6.3	CMS Reach using Inclusive Muons with Jets and Missing Energy	430
13.7	Inclusive analyses with same sign dileptons	431
13.7.1	Signal selection and backgrounds	432
13.7.2	Results for full detector simulated mSUGRA samples	432
13.7.3	CMS Inclusive Reach	433
13.8	Inclusive analyses with opposite sign dileptons	434
13.8.1	Signal selection and backgrounds	434
13.8.2	Results for point LM1	435
13.8.3	CMS inclusive reach	437
13.9	Inclusive analyses with di-taus	438
13.9.1	Event selection and background studies	439
13.9.2	Discovery potential of mSUGRA with di-taus final states	440
13.10	Inclusive analyses with Higgs	442
13.10.1	Signal selection and backgrounds	442
13.10.2	Results at LM5 and systematics	443
13.10.3	CMS reach for inclusive Higgs production	444
13.11	Inclusive SUSY search with Z^0	444
13.11.1	Topology of the signal	445
13.11.2	Event Selection	446
13.11.3	Results and Systematic Uncertainties	446
13.11.4	CMS reach for inclusive Z^0 search	448
13.12	Inclusive analyses with top	449
13.12.1	Top quark and lepton reconstruction and identification	449
13.12.2	Signal selection and backgrounds	449
13.12.3	Results at point LM1	451
13.12.4	CMS reach for inclusive top search	451
13.13	Final states with di-taus	452
13.13.1	Extraction of mSUGRA mass spectra from the measurement of the end points of invariant mass distributions.	452
13.14	Direct $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ production in tri-leptons	454
13.14.1	Signal selection and backgrounds	455
13.14.2	Results at LM9 and systematics	456
13.14.3	CMS reach for the tri-lepton final state	457
13.15	Production of $\tilde{l} \tilde{l}$	458
13.15.1	Simulation details	458
13.15.2	Sleptons production and decays	458
13.15.3	Signature and backgrounds	459

13.15.4	Results	459
13.16	Lepton flavour violation in neutralino decay	460
13.16.1	Signal selection and backgrounds	460
13.16.2	Results at CMS test points and reach	461
13.17	Look beyond MSUGRA	463
13.17.1	Non Universal Higgs Masses	463
13.18	Summary of the reach with inclusive analyses	469
13.18.1	Summary of the MSUGRA studies	469
14	Extra Dimensions and new vector boson high mass states	470
14.1	Introduction	470
14.1.1	Models with heavy vector bosons	470
14.1.2	Arkani-Hamed Dimopoulos Dvali (ADD) models	470
14.1.3	Inverse TeV sized extra dimensions	473
14.1.4	Randall Sundrum (RS) models	474
14.2	High mass dielectron final states	475
14.2.1	Event selection and correction	476
14.2.2	Mass peak distributions	477
14.2.3	Discovery potential of CMS	477
14.2.4	Systematic uncertainties	480
14.2.5	Identification of new particles	480
14.3	High mass dimuon final states	482
14.3.1	The Randall-Sundrum Model in the dimuon channel	482
14.3.2	The ADD model in the dimuon channel	484
14.4	High energy single lepton final states	486
14.4.1	Introduction	486
14.4.2	Data Samples	487
14.4.3	Event selection and analysis	488
14.4.4	Discovery and Exclusion Potential	489
14.4.5	Systematic Uncertainties	489
14.4.6	Summary	490
14.5	High mass di-jet final states	491
14.5.1	Dijet Resonances and Contact Interactions	491
14.5.2	Dijet Resonance Search	491
14.6	High mass di-photon final states	494
14.6.1	Introduction	494
14.6.2	Event generation and kinematics preselection	494
14.6.3	Offline selection and analysis	495
14.6.4	K-factors	495
14.6.5	Results	495
14.6.6	Limits uncertainties for 30 fb^{-1}	498
14.7	Single γ final state with E_T^{miss} from Extra Dimensions	499

14.7.1	Topology of single-photon final states	499
14.7.2	Backgrounds from the Standard Model	499
14.7.3	Event Selection	500
14.7.4	Systematic uncertainties and discovery potential	501
14.8	Black Holes (BHs)	501
14.8.1	Introduction to Higher-dimensional BHs	501
14.8.2	Analysis selection path and results	502
15	Alternative BSM signatures	505
15.1	Technicolour	505
15.1.1	The $\rho_{TC} \rightarrow W + Z$ channel	505
15.2	Search for Contact Interactions with dimuons	508
15.2.1	Analysis Path	509
15.3	Search for Contact Interactions with dijets	513
15.4	Heavy Majorana neutrinos and right-handed bosons	515
15.4.1	Introduction	515
15.4.2	Heavy Majorana neutrino production and decay	515
15.4.3	Selection of candidate events and the analysis variables	516
15.4.4	The background	517
15.4.5	Results	519
15.5	Little Higgs models	520
15.5.1	Introduction	520
15.6	Same Sign top	522
A	95% CL limits and 5σ discoveries	526
A.1	On the True Significance of a Local Excess of Events	526
B	Systematic errors	529
B.1	Hard process description and parametric uncertainties	529
B.2	Hard process scale	531
B.3	PDF description	531
B.4	QCD Radiation: the Parton Shower Monte Carlo	532
B.5	Fragmentation	533
B.6	Minimum bias and Underlying event	535
B.7	Pile up and LHC cross sections	536
B.8	Decays	537
B.9	LHAPDF and PDF uncertainties	538
C	Monte Carlo models and generators	541
C.1	Introduction	541
C.2	General scheme of generator's usage in CMS	542
C.3	<u>CMKIN</u>	543
C.4	Full event simulation generators	544

C.4.1	<u>PYTHIA</u>	544
C.4.2	<u>HERWIG</u>	546
C.4.3	<u>ISAJET</u>	547
C.4.4	<u>HIJING</u>	547
C.5	Tree level matrix element generators	547
C.5.1	<u>ALPGEN</u>	547
C.5.2	<u>CompHEP</u>	547
C.5.3	<u>MadGraph</u> and <u>MadEvent</u>	548
C.5.4	<u>TopReX</u>	548
C.6	Supplementary packages	548
C.6.1	<u>PHOTOS</u>	548
C.6.2	<u>TAUOLA</u>	549
C.6.3	<u>PYQUEN</u>	549
C.6.4	<u>HYDJET</u>	549
D	GARCON: Genetic Algorithm for Rectangular Cuts Optimization	550
E	Online Selection	552
E.1	Introduction	552
E.2	Description of trigger tools	552
E.2.1	L1 reconstruction	553
E.2.2	HLT reconstruction	553
E.3	Triggering with Forward Detectors	554
E.3.1	Objective	554
E.3.2	Level-1 Trigger Rates for Forward Detectors Trigger Stream	555
E.3.3	Level-1 Signal Efficiencies	557
E.3.4	Effect of pile-up, beam-halo and beam-gas backgrounds	559
E.3.5	HLT strategies	559
E.4	High-Level Trigger paths	560
E.4.1	L1 conditions	560
E.4.2	Evolution of DAQ-TDR triggers	561
E.4.3	New triggers	562
E.5	Performance	567
E.5.1	L1 Rates	568
E.5.2	L1 Trigger Object Corrections	569
E.5.3	HLT Rates	569
E.5.4	Trigger Tables	569
F	References	577



Chapter 1

Introduction

ADR to add

Part I

Complete analyses

Chapter 2

Physics Studies with Photons and Electrons



2.1 Introduction



Probably remove this chapter introduction...

2.2 Benchmark Channel: $H \rightarrow \gamma\gamma$

The $H \rightarrow \gamma\gamma$ channel has been studied since the initial planning of the LHC and SSC as an important channel for the discovery of Higgs particles at masses beyond the upper reach of LEP and below about 150 GeV [1, 2, 3]. The signature sought in the inclusive analysis is two high E_T isolated photons. The challenge for discovery of a Higgs in this mode is the small branching fraction of about 0.002, since in this mass range the dominant decay mode of the Higgs is $b\bar{b}$. The $\gamma\gamma$ decay mode can be well identified experimentally but the signal rate is small compared to the backgrounds coming both from two prompt photons (irreducible), and from those in which one or more of the photons are due to decay products or misidentified particles in jets (reducible). It has long been understood that $H \rightarrow \gamma\gamma$ can be detected as a narrow mass peak above a large background. The background magnitude can be determined from the region outside the peak. After event selection, for an integrated luminosity of 20 fb^{-1} and for a Higgs boson mass of 120 GeV, we expect approximately 350 signal events in a mass window of 2 GeV over 7000 background events.

In this study we present two complementary inclusive analyses for the $H \rightarrow \gamma\gamma$ channel: a standard cut based analysis and a high performance, discovery-oriented analysis, based on the method described in [4, 5]. Both are carried out with our present knowledge of the expected background, estimated with full detector simulation. Further details can be found in [6]. The study concentrates on the first years of LHC operation and uses simulated events with pileup corresponding to a luminosity of $2 \times 10^{33} \text{ cm}^{-2}\text{sec}^{-1}$.

The idea of measuring the rate of background by using the mass regions adjoining the Higgs peak is extended to also measure the characteristics of the background, and using this information to help separate background from signal. The $H \rightarrow \gamma\gamma$ channel is particularly well suited to this technique because the signal is relatively small and can be confined to a narrow mass region thanks to the excellent photon energy and position resolution of the CMS detector [7].

By using photon isolation and photon kinematic information, significant additional discrimination between signal and background can be achieved. The optimized analysis uses this

information to discriminate between signal and background by comparing data in mass side-bands with signal Monte Carlo. Use is made of a neural network, but likelihood variables or other techniques may prove to be better in the future. The expected purity in terms of signal/background, corresponding to each event, can be estimated based on this information and each event then can be used optimally to evaluate the likelihood of a signal plus background hypothesis compared to a background-only hypothesis.

In the optimized analysis the expected signal to background ratio is calculated for each event. By dividing the cut-based analysis in various categories with different s/b ratios results improve toward those that are obtained with the optimized analysis. If the maximum s/b ratio in the optimized analysis is limited to the best category used in the cut-based analysis, the performances of the two analyses are nearly identical.

The optimized, discovery-oriented analysis is particularly appropriate to the $H \rightarrow \gamma\gamma$ channel because the Higgs signal appears in a narrow mass peak allowing analysis of the large background in the mass side-bands. The analysis will not be limited by the poor simulation of the background once data will be available.

The study described requires a comprehensive understanding and simulation of the CMS detector. The electromagnetic calorimeter is used to make the primary measurements of photon energy and position. The tracker is used to measure the position of the interaction vertex. The tracker, ECAL and HCAL are used to determine if the photon candidate is well isolated. While background characteristics will be measured from data, the signal must be well simulated to perform the analysis described below. This requires a detailed understanding of the detector performance as well as its calibration.

2.2.1 Higgs Boson Production and Decay

For this inclusive study the Higgs boson production mechanisms with the largest cross-sections in the Standard Model have been simulated: gluon fusion, qqH production through Weak Vector Boson Fusion (WBF), associated Higgs production with W or Z bosons, and Higgs production associated with a $t\bar{t}$ pair. The cross sections for the different production processes [8] and the $H \rightarrow \gamma\gamma$ branching ratios [9] are summarized in Table 2.1. The analysis described in this chapter has been limited to careful measurement of the inclusive di-photon channel, to address the main detector issues, and no use has been made of tagging leptons or jets. In the future, channel identification, based on additional leptons and jets, will improve the sensitivity. For the moment these 'tagged' channels are investigated individually in other studies [10, 11, 12].

2.2.2 Backgrounds

Backgrounds with two real prompt high E_T photons are called "irreducible", although they can be somewhat reduced due to kinematic differences from signal processes in which high mass particles are produced. Two photons can be produced from two gluons in the initial state through a "box diagram" or from initial quark and anti-quark annihilation.

Backgrounds in which at least one final state jet is interpreted as a photon are called "reducible" and are much harder to simulate since jets are copiously produced at the LHC and Monte Carlo samples that correspond to 10 fb^{-1} are much too large to fully simulate. Selections at generator level have been devised in order to be able to select multi-jet and γ plus

Table 2.1: Next to Leading Order cross sections for the different Higgs boson production processes and $H \rightarrow \gamma\gamma$ branching ratios.

M_H	115 GeV	120 GeV	130 GeV	140 GeV	150 GeV
σ (gg fusion)	39.2 pb	36.4 pb	31.6 pb	27.7 pb	24.5
σ (WVB fusion)	4.7 pb	4.5 pb	4.1 pb	3.8 pb	3.6
σ (WH, ZH, $t\bar{t}H$)	3.8 pb	3.3 pb	2.6 pb	2.1 pb	1.7
Total σ	47.6 pb	44.2 pb	38.3 pb	33.6 pb	29.7
$H \rightarrow \gamma\gamma$ Branching Ratio	0.00208	0.00220	0.00224	0.00195	0.00140
Inclusive $\sigma \times B.R.$	99.3 fb	97.5 fb	86.0 fb	65.5 fb	41.5 fb

jets events that contribute to the background of the $H \rightarrow \gamma\gamma$ channel and reject events that have negligible chance of producing background to the final analysis.

The γ + jet can be viewed, for the selection point of view, as the combination of two different processes: one where another photon is radiated during the fragmentation of the jet (two prompt photons), the other where there is only one prompt photon in the final state and the other photon candidate corresponds to a mis-identified jet or isolated π^0 (one prompt plus one fake photon). These two processes have been separated using generator level information, and are listed separately in the tables below. Also, different K-factors are applied.

The generator level preselection of γ + jet events that contribute to the $H \rightarrow \gamma\gamma$ background is straightforward. For $pp \rightarrow$ jets, a much tighter set of cuts at the particle kinematics level was carefully developed and studied. Groups of particles, protocandidates, which might form a photon candidate after event simulation are identified. Cuts are applied on the transverse energy of two protocandidates and on their invariant mass, and this involves an estimate on the lower and upper limits to the energy of the photon candidates that might be reconstructed from the protocandidates after the simulation. An estimate is also made on likely level of isolation of the resulting photon candidate.

With such selection a rejection of a factor of about 41000 can be obtained, with an estimated inefficiency of 14% for $pp \rightarrow$ jets events generated with PYTHIA with $\hat{p}_\perp > 30$ GeV (transverse momentum of the products of the hard interaction). The inefficiency after the final analysis selection was estimated by using a looser preselection similar to that used for the $pp \rightarrow \gamma$ + jet simulation. Further details can be found in [6]. Events rejected by the preselection have rather low E_T photons and are not very important for the final analysis.

The Monte Carlo samples used are summarized in Table 2.2. All events were generated with PYTHIA, (missing reference cite PYTHIA) simulated with the Geant based (missing reference cite GEANT) CMSIM [13] or OSCAR [14], and reconstructed with ORCA version 8.7.3 [15]. Pile-up events from minimum bias interactions were added to the hard interaction, assuming a luminosity of $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$.

K-factors are applied to take into account the expected differences between the lowest order cross sections given by PYTHIA and the NLO cross sections of the different background processes [16, 17, 18, 19]. ~~NAME: BINOTH3, hep-ph/0200364, doesn't exist~~ The K-factors used for each background are summarized in Table 2.3 and are estimated to have an uncertainty of 20-30%.

Table 2.2: Monte Carlo samples used in the $H \rightarrow \gamma\gamma$ analysis with LO cross section from PYTHIA and total corresponding integrated luminosities of the analyzed samples.

Process	\hat{p}_\perp (GeV)	M_H (GeV)	σ (pb)	Presel. σ (pb)	Events Analyzed	Int Lum. (fb^{-1})
$H \rightarrow \gamma\gamma$ (gg fusion)	-	120	-	-	181K	-
$H \rightarrow \gamma\gamma$ (WB fusion)	-	120	-	-	193K	-
$H \rightarrow \gamma\gamma$ (gg fusion)	-	115–150	-	-	20K	-
$H \rightarrow \gamma\gamma$ (WB fusion)	-	115–150	-	-	20K	-
$H \rightarrow \gamma\gamma$ (WH,ZH,ttH)	-	115–150	-	-	20K	-
$pp \rightarrow \gamma\gamma$ (born)	> 25	-	82	44	920K	30
$pp \rightarrow \gamma\gamma$ (box)	> 25	-	82	31	668K	20
$pp \rightarrow \gamma + \text{jet}$	> 30	-	5×10^4	2.5×10^3	5.5M	2.2
$pp \rightarrow \text{jets}$	> 50	-	2.8×10^7	4.7×10^3	4.5M	1.0
Drell Yan ee	-	-	4×10^3	4×10^3	460K	0.1

Table 2.3: Background K-factors applied to PYTHIA cross sections.

$pp \rightarrow \gamma\gamma$ (born)	1.5
$pp \rightarrow \gamma\gamma$ (box)	1.2
$pp \rightarrow \gamma + \text{jet}$ (2 prompt)	1.72
$pp \rightarrow \gamma + \text{jet}$ (1 prompt+ 1 fake)	1
$pp \rightarrow \text{jets}$	1

2.2.3 Reconstruction, selection, and signal significance calculation

2.2.3.1 Trigger

$H \rightarrow \gamma\gamma$ events are selected with extremely high efficiency both by the Level-1 and High Level triggers that are described in details in reference [20]. Since in the analysis selection tighter E_T and isolation cuts are applied, the inefficiency due to the trigger is negligible.

2.2.3.2 Photon Reconstruction

Photons are reconstructed with the standard ECAL algorithms [21, 7]. At this level the photon reconstruction efficiency is over 99.5% for photons in the region covered by the ECAL.

The energy resolution of reconstructed photons is excellent for photons that do not convert or that convert late in the tracker. Energy resolution deteriorates somewhat for photons that convert early in the tracker. Nevertheless, photon energy resolution is substantially less affected by tracker material than is electron energy resolution and the Higgs reconstruction in the calorimeter is quite reliable even for converted photons.

For signal events, where this effect is relevant, the energy response of the individual crystals of the ECAL has been smeared using a miscalibration file randomly generated to correspond to the intercalibration precision expected after calibration with $W \rightarrow e\nu$ events obtained with

an integrated luminosity of 10 fb^{-1} , as described in [7]. The precision is 0.3% in the central part on the barrel, growing up to 1.0% at the edge of the barrel and in the endcaps.

The tools that have been developed to identify and reconstruct photon conversions in the tracker [22], and π^0 rejection tools developed for the endcap silicon preshower detector and the barrel crystals, have not yet been included in the analysis.

2.2.3.3 Primary Vertex Identification

The bunch length at LHC has an rms width of 75 mm resulting in a longitudinal spread of interaction vertices of 53 mm. If the mean longitudinal position is used (nominal vertex), the invariant mass of a two-photon state, such as the $H \rightarrow \gamma\gamma$, is smeared by about 1.5 GeV, due to the mis-measurement of the angle between the two photons related to the uncertainty of the photon directions.

The two high E_T photons coming from the Higgs boson decay are produced in association with other tracks that may come from the underlying event and initial state gluon radiation or from the other particles produced with the Higgs boson in the case of WBF fusion, WH or ZH production and $t\bar{t}H$ production.

The charged tracks associated to the Higgs production vertex are typically harder than those coming from minimum bias interactions. Therefore the vertex can be identified by reconstructing the primary vertices in the event and selecting the one that most likely corresponds to the Higgs boson production, based on charged tracks.

At low luminosity ($2 \times 10^{33} \text{ cm}^{-2}\text{sec}^{-1}$) we are able to identify the correct vertex, defined as being within 5 mm of the actual vertex, in about 83% of the signal events passing the selection described in section 2.2.4.1. Clearly these results will be affected by any significant variation of the characteristics of the pileup events from what is simulated in our pileup events.

2.2.3.4 Photon Isolation

Detailed studies have been made of photon isolation and its optimization [23, 24]. Fake photon signals due to jets can be rejected by looking for additional energetic particles accompanying the photon candidate. Charged pions and kaons can be detected in the tracker or in the calorimeters. Neutral pions and other particles decaying to photons can be detected in the ECAL. The hadron calorimeter may be important for detecting charged particles not efficiently reconstructed in the tracker, particularly at high η , or other particles like neutrons or K_{long}^0 .

2.2.3.5 Separation into Categories based on Lateral Shower Shape and Pseudo-rapidity

The shower shape variable R_9 , defined as the fraction of the supercluster energy found inside the 3×3 array of crystals centred around the highest energy crystal, is effective in distinguishing photon conversions in the material of the tracker. Photon candidates with large values of R_9 either did not convert or converted late in the tracker and have good energy resolution. Photons converting early have lower values of R_9 and worse energy resolution.

The variable R_9 has been shown to be very useful also in discriminating between photons and jets. This occurs both because of the conversion discrimination – either of the photons

from a π^0 can convert – and because, looking in a small 3×3 crystal area inside the super-cluster, the R_9 variable can provide very local isolation information about narrow jets.

In the multi-category analysis the events are separated into categories based on R_9 so as to take advantage of better mass resolution where it is expected (the unconverted photons), and yet still use all the events (since the mass resolution varies by at most a factor of 2). This separation also tends to put background events involving jets into categories with lower R_9 .

We also find that photons detected in the endcaps have worse energy resolution and more background than photons detected in the barrel so that it is useful to separate events with one or more photons in the endcaps from those with both photons in the barrel.

2.2.3.6 Calculation of Confidence Levels

Confidence levels are computed by using the Log Likelihood Ratio frequentistic method, as described in [25]. Given the expected signal and background distributions in the final variable (the mass distribution for the cut-based analysis), we simulate many possible outcomes of the experiment by means of Monte Carlo. This is done both in the hypothesis that the signal exists and that it does not exist. To compute a confidence level, we order our trials according to an estimator. This is a single number that is useful to order random trials from most background-only-like to most signal-plus-background-like. The simplest and probably best estimator is the Log Likelihood Ratio (LLR) which compares the likelihood of the data to come from a background-only distribution to the likelihood to come from a signal-plus-background distribution. Each likelihood is the product of probabilities from all the bins. The median confidence level is computed both for discovery and for exclusion.

2.2.3.7 Effect of Systematic Errors

To include systematic errors the background and signal expectation are randomized by the systematic error during the generation of the random trials, while keeping their expectations at the nominal value. If necessary the correlations between the errors on the different analysis bins is included. It is observed that the signal systematic error has no effect on the median LLR of signal-plus-background experiments, nor on that of background-only experiments. Of course the distribution corresponding to the signal-plus-background experiments is enlarged by the systematic error on the signal and this makes exclusion more difficult. On the other hand the effect of the systematic error on the background is very large, because of the small signal over background ratio. The mean of the distributions is still unchanged but the widths are enlarged both for background-only experiments and for signal-plus-background experiments. This decreases both the discovery and exclusion sensitivities.

2.2.4 Cut-based Analysis

2.2.4.1 Selection

Two photon candidates are required with pseudo-rapidity $|\eta| < 2.5$, with transverse energies larger than 40 GeV and 35 GeV respectively, and satisfying the following isolation requirements:

- no tracks with p_T larger than 1.5 GeV/c must be present inside a cone with $\Delta R < 0.3$ around the photon candidate. We only consider tracks with hits in at least two layers of the silicon pixel detector, therefore converted photons are likely to be rejected only if they convert before the second pixel layer;
- the total E_T of all ECAL island basic clusters with $0.06 < \Delta R < 0.35$ around the direction of the photon candidate, regardless of whether they belong to the super-cluster or not must be less than 6 in the barrel and 3 GeV in the endcaps;
- the total transverse energies of HCAL towers within $\Delta R < 0.3$ around the photon candidate must be less than 6 GeV in the barrel and 5 GeV in the endcaps.

In order to further reduce the background that is higher when at least one of the photons is detected in the electromagnetic calorimeter endcaps and to increase the performance of the analysis in the forward region additional isolation requirements are applied for events where one, or more, of the candidates has $|\eta| > 1.4442$. For these events the candidate in the barrel is required to satisfy the tighter isolation selection that is applied to photons in the endcaps: ECAL isolation less than 3 GeV and HCAL isolation less than 5 GeV.




Figure 2.1: Di-photon invariant mass spectrum after the selection for the cut-based analysis. Events are normalized to an integrated luminosity of 1 fb^{-1} and the Higgs signal, shown for different masses, is scaled by a factor 10.

Table 2.4: Expected background after the selection for Higgs boson masses between 115 and 150 GeV, expressed in fb/GeV

Process	M_H	115 GeV	120 GeV	130 GeV	140 GeV	150 GeV
$pp \rightarrow \gamma\gamma$ (born)		48	44	36	29	24
$pp \rightarrow \gamma\gamma$ (box)		36	31	23	16	12
$pp \rightarrow \gamma + \text{jet}$ (2 prompt)		43	40	32	26	22
$pp \rightarrow \gamma + \text{jet}$ (1 prompt + 1 fake)		40	34	22	19	14
$pp \rightarrow \text{jets}$		29	27	20	18	14
Drell-Yan ee		2	2	1	1	1
Total background		203	178	134	109	86

Table 2.5: Selection efficiency for the Higgs signal in different mass windows.

$M_H(\text{GeV})$	Window $\pm 1 \text{ GeV}$	Window $\pm 1.5 \text{ GeV}$	Window $\pm 2.5 \text{ GeV}$	Window $\pm 5 \text{ GeV}$	Window Total
115	17%	21%	25%	28%	29%
120	18%	22%	26%	29%	30%
130	18%	22%	27%	31%	32%
140	18%	23%	28%	32%	34%
150	28%	24%	29%	33%	36%

Figure 2.1 shows the mass distribution after the selection. The efficiency for a 120 GeV Higgs boson is 30% and the total expected background is 178 fb/GeV. The number of expected background events for the different types of background is shown in Table 2.4 while the Higgs efficiency in different mass windows is shown in Table 2.5, the efficiency is computed using all generated signal events. The signal contribution to the total number of events is very small, particularly outside the mass region under study. The background can be estimated by a fit to the data mass distribution.

The error on the background estimation comes from two sources:

- the statistical precision which decreases with the size of the mass range that is used to perform the fit;
- the systematic error related to the shape of the function that is used to fit the distribution.

It is not possible to know the exact functional form of the background shape and the error must be estimated by assuming a function, simulating a distribution and then using a different function to fit the data. Clearly this error grows with the size of the mass range used. For a reasonable mass range of $\pm 10 - 20 \text{ GeV}$ excluding $+3$ and -5 GeV from the Higgs boson mass under study and for an integrated luminosity of 20 fb^{-1} the statistical and systematic errors are estimated to be 0.4% and 0.5% respectively. The statistical error decreases with the integrated luminosity while the systematic error is constant.

2.2.4.2 Splitting into Categories

Changing the cuts or adding new discriminating variables to this analysis does not give large improvements in the sensitivity. This can be seen, for example, from the fact that it is not possible to use the very powerful variable, R_9 , to reject events without losing performance. This is because the increase in s/b ratio does not compensate the loss in efficiency.

The way to improve the sensitivity of the analysis is to keep all selected events but to split the sample into categories with different s/b ratios.

The following 3 possibilities are considered:

- 1 single category;
- 4 categories from 2 R_9^{\min} ranges (R_9^{\min} larger or smaller than 0.93) times 2 pseudo-rapidity regions $|\eta|^{\max}$ in barrel or endcaps;
- 12 categories from 3 R_9^{\min} ranges ($R_9^{\min} > 0.948$, $0.9 < R_9^{\min} < 0.948$ and $R_9^{\min} < 0.9$) times 4 pseudo-rapidity regions ($|\eta|^{\max} < 0.9$, $0.9 < |\eta|^{\max} < 1.4442$, $1.4442 <$

$$|\eta|^{\max} < 2.1 \text{ and } |\eta|^{\max} > 2.1).$$




Figure 2.2: Invariant mass spectrum after the selection relative to the cut-based analysis with four categories defined in the text: barrel with large R_9 (a), barrel with small R_9 (b), endcaps with large R_9 (c) and endcaps with small R_9 (d). Events are normalized to an integrated luminosity of 1 fb^{-1} and the Higgs signal, shown for different masses, is scaled by a factor 10.

Figure 2.2 shows the mass spectrum after splitting into four categories. The signal over background ratio is much larger in the best category and the composition of the background varies between the different samples: irreducible backgrounds dominate for large R_9 and reducible backgrounds are larger for small R_9 .

Table 2.6 shows, for the 12 category analysis, the fraction of events along with the maximum s/b ratio in each category.

Table 2.6: Fractions of events in each of the 12 categories and maximum s/b in the mass region of 120 GeV.

	$ \eta ^{\max} > 0.9$		$0.9 < \eta ^{\max} < 1.4442$		$1.4442 < \eta ^{\max} < 2.1$		$ \eta ^{\max} > 2.1$	
	frac.	s/b	frac.	s/b	frac.	s/b	frac.	s/b
$R_9^{\min} > 0.948$	15.5%	14.7%	13.1%	9.0%	10.8%	6.1%	8.5%	4.5%
$0.9 < R_9^{\min} < 0.948$	9.4%	12.2%	6.8%	6.8%	6.7%	4.8%	2.7%	2.8%
$R_9^{\min} < 0.9$	8.3%	7.6%	11.1%	4.3%	5.4%	3.2%	1.7%	2.2%

2.2.4.3 Systematic errors

It has been seen that the total error on the background is approximately 0.65% and is due to the uncertainty of the function fit to the side-bands of the mass distribution, estimated to be 0.5%, plus the statistical error on the fit that is approximately 0.4% for an integrated luminosity of 20 fb^{-1} .

An error of 0.65% has a very large effect on the discovery CL when only one category is used. The reason is that a large fraction of signal events corresponds to a very low s/b , of the order of a percent. The effect can be reduced by applying a cut on the signal over background s/b . This corresponds to using events in a mass window around the analyzed mass, until s/b becomes smaller than the chosen cut. The optimal cut for this analysis is 0.02.

When the events are split into categories the number of background events in each category is reduced on average by $1/N_{\text{cat}}$ and this increases the statistical error on the background estimation by approximately a factor $\sqrt{N_{\text{cat}}}$, but this error is completely uncorrelated between the different categories. The error related to the uncertainty of the fit function remains constant and it is also uncorrelated between the different categories because, due to the different cuts the background shapes are different and described by different functions. The total error is then less than the total error reduced by $1/N_{\text{cat}}$. This reduces the effect of the systematic error on the discovery.

The effect of the systematic error on the background estimation is also related to the signal over background of the analysis. A more sensitive analysis, for which a larger part of the signal has a higher s/b ratio, is less affected by the same relative uncertainty on the background.

Clearly the current understanding of the background is affected by much larger uncertainties such as: cross section, di-photon kinematic distributions and efficiency of the selection (mainly affected by jet fragmentation, pile-up and by the structure of the underlying events).

The systematic error on the signal, that as has been mentioned has no effect on the discovery CL, has contributions from the theoretical uncertainty of the cross section (+15 -12% from the scale variation and +4 -5%), from the measurement of the integrated luminosity ($\sim 5\%$), from the trigger ($\sim 1\%$), from the analysis selection (that will be measured for example with $Z \rightarrow \mu\mu\gamma$) and from the uncertainties on the photon energy resolution. Other effects that could modify the ability to discover the Higgs boson are: uncertainties on the structure of the underlying events, that could change the efficiency of the primary vertex determination and the amount of material in the tracker before the electromagnetic calorimeter.

The effect on the performances of the analysis of an increase of 20% of the tracker material has been evaluated. The main effects on such change on the analysis would be:

- increase of the inefficiency of the track isolation requirements for early photon conversions, before or inside the second layer of the pixel detector.
- increase of the inefficiency of ECAL isolation cut;
- decrease of the value of R_9 for all photons that would cause a migration of events from more sensitive categories to less sensitive categories.

It was estimated that such change would increase the luminosity needed to achieve a given discovery CL of approximately 6%. Given that the amount of tracker material will be known with a precision of $\sim 2\%$ the related systematic error is less than 1%.

In what follows a conservative 20% systematic error on the signal is assumed. It affects

exclusion of a signal, not discovery, since the signal rate is directly measured from data in case of discovery.

2.2.4.4 Results of the Cut-Based Analysis

Table 2.7: Integrated luminosity needed to discover or exclude the Higgs boson with mass 120 GeV with or without taking into account the systematic errors (fb^{-1}).

Analysis	5σ	5σ	3σ	3σ	95%	95%
	discovery no syst	discovery syst	evidence no syst	evidence syst	exclusion no syst	exclusion syst
counting exp.	27.4	48.7	10.0	13.2	4.5	6.5
1 category	24.5	39.5	8.9	11.5	4.1	5.8
4 categories	21.3	26.0	7.5	9.1	3.5	4.8
12 categories	19.3	22.8	7.0	8.1	3.2	4.4

Table 2.7 shows the integrated luminosity needed to obtain 5σ discovery or 95% CL exclusion for a 120 GeV mass Higgs boson with the different splittings. The effect of the systematic errors is also shown. We can observe how the performance increases and the effect of the error on the background estimation decreases with the number of categories. In the three cases (1, 4 and 12 categories) the event selection is the same and that the differences in performance come from the splitting of the total sample in different sub-samples with different sensitivities (s/b). In the split category analyses the computation of the log-likelihood ratio estimator is made separately for each 1 GeV bin in mass, whereas in the “counting experiment” only a single (optimum) mass window is evaluated.

The integrated luminosity needed for discovery and exclusion, using the 12-category analysis, for the mass range studied between 115 and 150 GeV are shown in the plots at the end of the section (Figure 2.8). The Higgs boson can be discovered with mass between 115 and 140 GeV with less than 30 fb^{-1} and excluded in the same mass range, at 95% CL, with less than 5 fb^{-1} .

As mentioned before, all these results have been obtained assuming an intercalibration of the ECAL, after having collected an integrated luminosity of 10 fb^{-1} . With the whole ECAL intercalibrated to a precision better than 0.5% over all the solid angle, the results improve such that approximately 10% less integrated luminosity is needed for discovery.

2.2.5 Optimized Analysis Estimating s/b for Each Event

In the optimized analysis 6 categories are used, 3 in which both photons are in the barrel and 3 in which at least 1 photon is in an endcap. The 3 categories are defined, as for the cut-based analysis, to have the lowest R_9 photon candidate with $R_9 > 0.948$, $0.948 > R_9 > 0.90$ and $R_9 < 0.90$ respectively. The categories are labelled with numbers from 0 to 5 starting with the highest R_9 category in the barrel and ending with the lowest R_9 category in the endcaps.

2.2.5.1 Mass Distributions in Categories

The di-photon mass distributions enable the separation of signal from background. Signal peaks sharply at the Higgs mass while the backgrounds are quite smooth. This allows good estimation of the magnitude of the background under the peak.

The best mass resolution and the best s/b ratio in the peak is found in category 0, with high R_9 in the barrel.

2.2.5.2 Loose Selection of Events for Optimized Analysis

Isolation requirements are applied to photon candidates prior to the computation of the neural network isolation variables NN_{isol} :

- the transverse E_T of the photon candidates must be larger than 40 GeV and the absolute value of their pseudo-rapidity less than 2.5;
- no tracks with p_T larger than 1.5 GeV/c must be present inside a cone with $\Delta R < 0.1$ around the photon candidate;
- the total E_T of all ECAL island basic clusters with $\Delta R < 0.3$ around the photon candidate, excluding those belonging to the super-cluster itself must be less than 5 GeV;
- the total transverse energies of HCAL towers within $\Delta R < 0.35$ around the photon candidate must be less than 35 GeV;
- the sum of the transverse momenta of charged tracks within $\Delta R < 0.2$ around the photon candidate must be less than 100 GeV.

Before optimizing the final analysis, some additional cuts are applied. These both simplify the neural network training and slightly improve the performance. It is required that:

- the events pass the double photon High Level Trigger;
- the isolation neural net output is greater than 0.25 for both photons.

2.2.5.3 Optimized use of Kinematic Variables to Separate Signal and Background

In addition to the mass, there are kinematic differences between signal and background. In particular the signal has a harder photon E_T distribution than the background – the background can have a high mass by having a large η difference between the photon candidates. Weak Boson Fusion and associated production of a Higgs with other massive particles enhance these differences between signal and background. The large, reducible backgrounds often have photon candidates that are not well isolated.

As with the Higgs searches performed at LEP, higher performance can be achieved if the expected signal over background, s/b , is estimated for each event. This is particularly effective if the s/b varies significantly from event to event. This is the case here due to wide variations in photon isolation and photon E_T . There is also significant dependence of the s/b on photon conversion and on location in the detector.

One photon isolation variable NN_{isol} for each photon, is combined with kinematic variables to help separate signal and background. A neural net is trained to distinguish background events, taken from the mass side-bands, from signal Monte Carlo events. There is no danger

of over-training since background events from the signal mass region are not used and independent samples are used for the signal Monte Carlo. The input variables are devised to be insensitive to the di-photon mass so that the background rejection due to the kinematics and isolation is independent of the background rejection from the mass distribution. A neural net has been used because it is easy to re-train when changes are made to the input variables or data samples. There are many other methods that could be used to replace the neural net.




Figure 2.3: Distribution of the inputs to the neural network for signal and background sources. The signal is multiplied by 50 for visibility.

Six variables are used as inputs to a neural net. They are the isolation NN outputs NN_{isol} for the 2 photons, the transverse energies of the 2 photons, the absolute value of the rapidity difference between the 2 photons, and the longitudinal momentum of the photon pair.

The distributions of the input variables are shown for signal and background in Figure 2.3. Kinematic information that are likely to be highly sensitive to higher order corrections to the background simulation has not been used. Such information, like the E_T of the Higgs boson candidate, the E_T transverse to the photon direction, and information about additional jets will ultimately be useful but may not be reliable until better simulations or actual data are

available to train on.

The neural net is trained in each of the 6 categories independently. The net has 6 input nodes, 12 intermediate nodes in a single layer, and 1 output node. The error function has been modified from the standard to improve training toward a high signal over background region. A minimum neural net output cut is applied that eliminates 1% of the signal in each category and a function is fit to the distribution above that cut. These functions are used to bin the data and to smooth the background in a limited region.

It is useful to examine the neural net output distribution for events from different sources (Figure 2.4). Low NN outputs are dominated by photon candidates from jets which are not well isolated. The large peak at 0.85 represents both signal and background where the photon is relatively well isolated and the photon E_T is normal. Higher photon E_T events are found in the peak near 1. There is an enhancement of the signal, particularly for the WBF and associated production processes. The background there is dominated by events with at least one jet interpreted as a photon.





Figure 2.4: The neural net output for events in the barrel for each signal ($M_H=120$ GeV) and background source. The signal is multiplied by 10 for visibility.

2.2.5.4 Estimation of Signal to Background Ratio for Each Event

In order to get the most information out of each event, the signal over background is estimated for each event. In the simplest analyses, cuts are applied to select only high signal over background events and those are counted. Such a simple analysis loses information because some of the events that are cut could contribute to the measurement and because some of the events that are accepted are not used optimally.

Events in the mass peak for the Higgs mass hypothesis under consideration have high signal over background expectation while events outside the peak have lower expected s/b . Similarly, events at high NN_{kin} output have higher s/b expectation. The kinematics and isolation information in NN_{kin} has been made independent of mass information so the two s/b ratios



c

Figure 2.5: The di-photon mass distribution for each source for barrel events with kinematic neural net output greater than 0.85 (top) and 0.97 (bottom). The signal ($M_H=120$ GeV) is multiplied by 10 for visibility.

can be multiplied to get a good estimate of the s/b expectation for the event:

$$\left(\frac{s}{b}\right)_{est.} = \left(\frac{s}{b}\right)_{mass} \times \left(\frac{s}{b}\right)_{kin}$$

This is an estimate that is to bin signal and background events. If the estimate is bad, the performance of the analysis suffers because good s/b events are not well separated from bad ones. It is not possible for a bad estimate to make the analysis appear to perform too well. The s/b estimate need not be normalized correctly, since it is a relative number used to bin events.

The events are binned according to the s/b estimate. Histograms are made in each of the six categories. The actual signal to background ratio is computed for the binned events and used to calculate confidence levels that data are consistent with a background-only hypothesis or with a signal-plus-background hypothesis.

2.2.5.5 Smoothing the Background

The $H \rightarrow \gamma\gamma$ channel has the good feature that the mass is essentially independent of isolation and suitably chosen kinematic variables. With this factorization assumption, background can be smoothed well even in regions with low statistics.

The background expectation in a bin must be reliably estimated in order to correctly calculate confidence levels. Downward fluctuations in the background estimation can have a significant impact on the CL. The number of simulated events for the irreducible (jet) backgrounds is about one seventh of the number that will be available in the data at the time it would be expected to discover the Higgs. Therefore problems with background estimation are even more difficult now than they will be when we have data.

The background distributions are very smooth in the mass variable, so the distribution in mass can be reliably smoothed. This is done by spreading each event over a ± 5 GeV region according to the functions fit to the mass distribution. A wider mass region could be used but this would interfere with the training of the analysis on an independent sample in the mass side-bands.

The background distribution in the neural net output is also smoothed over a region of ± 0.05 using the fit functions. It is therefore quite important that the background fit functions accurately represent the neural net distribution. In the smoothing process, the normalization of the background is carefully maintained to high accuracy.




Figure 2.6: The Mass distribution for each source for barrel events with kinematic neural net output greater than 0.97 and $R_9 > 0.948$. The signal for $M_H=120$ GeV is multiplied by a factor 10.

With this two-dimensional smoothing accurate background expectations are obtained except in the regions with extremely small amounts of background. In such regions, bins must be combined until sufficient background events are available. If a s/b bin has too few MC background events contributing to it, it is combined with the nearest (lower s/b) bin. This is continued until there are sufficient events. This combination clearly reduces the sensitivity of the analysis but cannot be avoided without a more detailed understanding of the background, which is a goal for the future. At present, at least 20 Monte Carlo background events

are required in a bin. Since the current MC samples contain about seven times less events than expected in the data, significant improvements are possible, allowing higher s/b bins to be used, resulting in better performance.

Figure 2.5 shows the mass distributions for barrel events with two different cuts on the neural net output. The looser cut simply excludes most of the obviously non-isolated candidates. It can be seen that all of the backgrounds are important at this level. The tighter cut highly enhances the s/b ratio and emphasises the importance of smoothing, which has not been applied to the background in this distribution.

Figure 2.6 shows the mass distribution for neural net output greater than 0.97 in category 0. Again it is clear that smoothing in two dimensions is needed to get a reasonable estimate of the background. It is useful to note that even in this very high s/b region, the largest contribution to the signal is from gluon fusion, although the relative contributions of the other production processes has increased.

2.2.5.6 Combination of Categories into Final s/b Distribution

At this point the signal and background is binned in s/b in six categories. These could be used to calculate the confidence level, however, it seems most useful in the light of future plans to analyze separate channels, to combine the categories into one s/b plot in a similar way as may be used to re-combine channels. The six histograms are combined into one which can be used calculate confidence levels. The combination is based on the actual signal to background in each bin. In principle, this is the same as combining results from different channels or even from different experiments in a way that makes optimal use of all channels and does not pollute high quality channels with data of lesser purity.

The final binning of data into s/b bins is shown in figure 2.7. The plot extends from very low signal to background to a small number of events with $s/b > 1$.

The relative contribution of barrel and endcap categories can be estimated from the total LLR computed and LLRs computed excluding each category. The six categories have rather widely varying contributions to the Log Likelihood Ratio and hence to the performance of the analysis. Table 2.8 shows the fraction of signal and the fraction of the LLR for each category.

Table 2.8: Performance in the six categories for $M_H = 120$ GeV.

Category	Signal %	LLR %
0	27.8	48.0
1	16.1	24.8
2	21.7	11.9
3	16.6	9.7
4	9.0	4.1
5	8.8	1.5

Some of the categories have a fairly small effect on the final result. This remains true after the application of systematic normalization uncertainties described below. It is clear that photon conversions result in a significant deterioration of the performance. It is hoped to mitigate this somewhat by using the conversion track reconstruction in the future, but the




Figure 2.7: The final distribution of binned signal ($M_H=120$ GeV) and background in $\log(s/b)$ for an integrated luminosity of 7.7 fb^{-1} .

poorer mass resolution cannot be recovered and a big effect is not expected.

2.2.5.7 Results of the Optimized Analysis

The same estimates of systematic error are used to obtain the results in the optimized analysis as are used in the cut-based analysis. Most of the development and studies have been made for a Higgs mass of 120 GeV. For this mass, a 5σ discovery can be made with about 7 fb^{-1} luminosity. A 1% background normalization uncertainty corresponds to an increase of the luminosity needed for a 5σ discovery from 7 fb^{-1} to 7.7 fb^{-1} .




Figure 2.8: Integrated luminosity needed for a 5σ discovery (left) and discovery sensitivity with an integrated luminosity of 30 fb^{-1} (right) with the optimized analysis. The results from the cut-based analysis in 12 categories are also shown for comparison.

There is a great deal of uncertainty in this benchmark estimate of luminosity due to our poor understanding of the backgrounds we will contend with when the LHC starts running, however, this is not considered here as a systematic error on a discovery since it is proposed to measure the background from the data. Figure 2.8 shows the luminosity needed for a 5σ discovery and the discovery sensitivity with an integrated luminosity of 30 fb^{-1} for several Higgs masses, both for the fully optimized analysis and for the cut-based analysis using 12 categories described in Section 2.2.4.4. It seems possible to discover, or at least have strong evidence for a low mass Higgs in the first good year of running.

2.2.6 Summary

A standard cut-based analysis can discover the Higgs boson with 5σ significance between the LEP lower limit and 140 GeV with less than 30 fb^{-1} of integrated luminosity. Approximately 5 fb^{-1} are needed to exclude its existence in the same mass range.

It has been shown that the $H \rightarrow \gamma\gamma$ channel can be used to discover a low mass Higgs with an integrated luminosity not too different from that needed for higher mass Higgs, 7.7 fb^{-1} at 120 GeV with an analysis using an event by event estimation of the s/b ratio. Because of the excellent mass resolution expected in the di-photon channel, the background rate and characteristics from the data can be determined from di-photon events at masses away from the Higgs mass hypothesis.

An inclusive analysis has been presented. In future the various signal channels will be identified by looking for additional jets, leptons, or missing energy. This will clearly improve the sensitivity of the analysis.



Benchmark Channel: $H \rightarrow ZZ^{(*)} \rightarrow 4$ electrons

One of the most promising road towards a discovery at the LHC of the Higgs boson postulated in the SM is via single H production followed by a cascade decay into charged leptons, $H \rightarrow ZZ^{(*)} \rightarrow l^+l^-l^+l^-$.

The single Higgs boson production benefits from a high production cross-section, with values of about $40 \times 10^3 \text{ fb}$ at $m_H = 130 \text{ GeV}/c^2$ and decreasing monotonically to about $10 \times 10^3 \text{ fb}$ around $m_H = 300 \text{ GeV}/c^2$. The production cross-section is dominated ($\gtrsim 80\%$) over this mass range by gluon-gluon fusion processes via triangular loops involving heavy quark (mostly the top quark) flavours. The branching ratio for the $H \rightarrow ZZ^{(*)}$ decay in the SM is sizeable for any m_H value above $130 \text{ GeV}/c^2$. It remains above 2% for $m_H \leq 2 \times M_W$ with a peak above 8% around $m_H \simeq 150 \text{ GeV}/c^2$, and rises to values of 20 to 30% for $m_H \geq 2 \times m_Z$. The Z bosons have a 10% probability to yield a pair of charged leptons. Thus, the decay chain $H \rightarrow ZZ^{(*)} \rightarrow l^+l^-l^+l^-$ (in short $H \rightarrow 4l$) offers a possibly significant and very clean and simple multi-lepton final state signature for the SM Higgs boson at the LHC.

Ultimately, the channel can provide a precision determination of the Higgs boson mass and production cross-section. The anti-correlation of the Z spin projections in the $H \rightarrow ZZ$ decay and the polarization of each Z boson can be used to constrain, and eventually determine, the spin and CP quantum numbers of the Higgs resonance. Furthermore, the $ZZ^{(*)}$ and $WW^{(*)}$ decay modes are related via $SU(2)$ and the combination of channels could allow for cancellation of some systematic uncertainties in a determination of the Higgs coupling. But

first and foremost is the necessity to be best prepared for a discovery at the LHC. The $H \rightarrow 4l$ channel is on the critical path of such a discovery over an extended range of possible m_H values.

In this section, the discovery potential of the CMS experiment for the SM Higgs boson is discussed in the mass range of $120 \leq m_H \leq 300 \text{ GeV}/c^2$, focusing on the $4e$ channel. The analysis [26] relies on a detailed simulation of the detector response in the experimental conditions of the first years of low luminosity LHC running. The signal and background Monte Carlo datasets used for this prospective are described in Section 2.3.1. The detailed High Level Trigger (HLT) and reconstruction algorithms used at each step of this analysis have been presented in [7]. Basic, and in part compulsory, triggering and pre-selection steps for data reduction are described in Section 2.3.2. Simple observables from the electron reconstruction are used to characterize the event signature for this pre-selection step. The final event selection relies on more involved requirements for primary electrons coupled with basic event kinematics and is presented in Section 2.3.3. The selection is optimized to preserve a best signal detection efficiency and highest significance for a discovery, while allowing for a control of experimental systematics and of systematics on physics background rates. Realistic strategies for the taming and measurements of systematics are described in Section 2.3.4. Results on the expected discovery reach of the SM Higgs boson in CMS in the $H \rightarrow 4e$ channel and for the measurement of its mass, width and cross-section are finally presented in Section 2.3.5.

2.3.1 Datasets for Signal and Background Processes

Monte Carlo data samples for the signal from single SM Higgs boson production as well as for SM background from $ZZ^{(*)}$ pair production, $t\bar{t}$ pair production and $Zb\bar{b}$ associated production are used. The signal and background processes are generated for pp collisions at the LHC at a centre-of-mass energy $\sqrt{pp} = 14 \text{ TeV}$, with pile-up conditions from multiple collisions as expected in a collider machine configuration providing an instantaneous luminosity of $2 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$ (of $\mathcal{O}(10) \text{ fb}^{-1}/\text{year}$). All cross-sections are normalized within acceptance to Next-to-Leading-Order (NLO) calculations. The event generators are interfaced with PHOTOS [27, 28] for the simulation of QED final state radiations. The non-perturbative parton density functions in the proton are taken from the so-called CTEQ6 distributions [29] which were obtained in 2002 from a global QCD analysis combining all existing relevant deep inelastic and jet cross-section measurement results.

The Higgs boson is produced via either gluon fusion and weak boson fusion processes. The $4e$ signal samples are generated at various m_H with PYTHIA [30]. The Higgs boson is forced to decay into a Z boson pair. The Z bosons are subsequently forced to undergo a decay in electron-positron pair. The signal is normalized to the value of total cross-section at NLO calculated including all Higgs boson production processes via HIGLU [31], with branching ratios $BR(H \rightarrow ZZ^{(*)})$ calculated via HDECAY [32].

In the $4e$ channel (and similarly for the 4μ channel), an additional enhancement of the signal is considered which is due to the constructive final state interference between like-sign electrons originating from different $Z^{(*)}$ bosons [33]. This enhancement has been re-evaluated with CompHEP [34] and amounts to a factor 1.130 ± 0.006 at $m_H = 115 \text{ GeV}/c^2$, slowly decreasing to a negligible value when approaching $m_H \approx 2m_Z$.

The $ZZ^{(*)}$ SM background continuum is generated using PYTHIA [30]. This includes only

the t -channel contribution with $q\bar{q}$ in the initial state. The missing s -channel might contribute up to 10% for low Higgs boson masses and can be neglected for higher masses. The differential cross-section is re-weighted using m_{4e} dependent NLO K -factors obtained with MCFM 4.1, with an average K -factor of $\langle K_{NLO} \rangle = 1.35$. Both Z bosons are constrained within the mass range $5 - 150 \text{ GeV}/c^2$ and are forced to decay into charged lepton pairs, with the τ leptons subsequently forced to undergo leptonic decays via $\tau \rightarrow \mu\nu$ or $\tau \rightarrow e\nu$. The missing gg contribution is estimated to be of order 20% at LO [33], with $\pm 8\%$ uncertainties for masses below $200 \text{ GeV}/c^2$ and with unknown NLO K -factors. Recent calculations with TopREX [35] of the gluon fusion production process of two real Z confirm above assumptions, and this contribution has been shown to remain stable after kinematic cuts for a $H \rightarrow 4l$ analysis. The cross-section here is simply increased by the mean expected contribution.

The $t\bar{t}$ background sample is also generated with PYTHIA [30], with W bosons and τ leptons forced to leptonic decays, but with b quarks left to decay freely. Both gluon fusion and quark annihilation initial states are simulated and the cross-section is normalized to the NLO value of $840 \pm 5\%(\text{scale}) \pm 3\%(\text{pdf}) \text{ pb}$ [36].

The $Zb\bar{b}$ background is generated using all lowest order $gg \rightarrow e^+e^-b\bar{b}$ and $qq' \rightarrow e^+e^-b\bar{b}$ diagrams (excluding diagrams involving the SM Higgs boson) calculated with CompHEP [34] and interfaced with PYTHIA [30] for showering and hadronization. All possible combinations of quarks are considered in the initial state. The total LO cross-section for $m_{ee} > 5 \text{ GeV}/c^2$ is 115 pb of which about 89% originates from gg processes, 7.7% involve u-like quarks and 3.2% involve d-like quarks in the initial state. The hadronization and decay of the b quarks are left free. A NLO K -factor of 2.4 ± 0.3 is applied. Signal and background

$m_H \text{ (GeV}/c^2)$	$\sigma_{NLO} \text{ (pb)}$	$\sigma_{NLO} \times BR \times Acc. \text{ (fb)}$	$N_{simul.}$
115	47.73	0.27	10000
120	44.30	0.48	10000
130	38.44	1.11	10000
140	33.69	1.78	10000
150	29.81	1.94	10000
160	26.56	0.92	10000
170	23.89	0.43	10000
180	21.59	0.98	10000
190	19.67	3.58	10000
200	17.96	3.94	10000
250	12.37	3.07	10000
300	9.58	2.60	10000
$ZZ^{(*)}$	29.0	20.2	150 000
$Zb\bar{b}$	276.3	120.4	87 000
$t\bar{t}$	840	194.0	500 000

Table 2.9: Total cross-sections at NLO (pb), cross-section in the $4e$ channel within acceptance (fb), and number of accepted events in data samples available for analysis.

events are filtered at generator level for further analysis if satisfying the following acceptance requirements: $\geq 2e^+$ and $\geq 2e^-$ with $p_T^e > 5 \text{ GeV}/c$ in $|\eta| < 2.7$. In addition for the $Zb\bar{b}$ background, at least two e^+e^- pairs with invariant mass in the range $5 - 400 \text{ GeV}/c^2$ are required. In Table 2.9 cross-sections at NLO and after pre-selection, as well as number of events in data samples available for analysis after pre-selection are given.

Detailed simulation of the CMS detector is performed using the official CMS simulation OSCAR. Reconstruction of physics objects is performed in ORCA.

2.3.2 Data Reduction

The events of interest for the Higgs boson search in the $H \rightarrow 4e$ channel must satisfy a minimal set of requirements.

A first and compulsory condition for the events is to satisfy the CMS Level 1 (hardware) trigger conditions and the filtering of the (software) HLT. This triggering step is described in Section 2.3.2.1. The basic electron triggers are expected to be saturated by SM processes such as the single Z and W production. Further filtering is obtained with a minimal set of additional electron requirements as described in Section 2.3.2.2.

The pre-selection must preserve the signal acceptance, and especially the electron reconstruction efficiency, until later stages where the analysis can best profit from more involved algorithms applied to reduced event samples.

2.3.2.1 Triggering

The events must have satisfied the *single e*, *double e* or *double relaxed e* requirements at L1/HLT level. The *single e* trigger requires one isolated (charged) “electromagnetic” object with a threshold set at a reconstructed transverse energy in the electromagnetic calorimeter (ECAL) of $E_T = 26$ GeV. The *double e* trigger requires two isolated (charged) “electromagnetic” objects, each above a threshold of $E_T = 12.0$ GeV. In contrast, the *double relaxed e* trigger does not impose isolation for the (charged) “electromagnetic” objects and the increased rate is compensated by a higher threshold of $E_T = 19.0$ GeV.

The trigger efficiency for the Higgs boson signal, normalized to the cross-section within acceptance as defined in Section 2.3.1, is above 95% for masses $m_H > 130$ GeV/ c^2 .

2.3.2.2 Pre-selection of four electron candidates

Following the L1 and HLT filtering steps, the event candidates must further satisfy basic electron pre-selection requirements. These requirements are designed to tame possible background sources involving “fake” electron contamination from QCD jets.

For Higgs bosons with a mass m_H below 300 GeV/ c^2 , the $4e$ final state always involves at least one (or few) low p_T^e electron(s). In the range of m_H values below the Z pair production threshold, where the Z and Z* bosons themselves receive in general only small transverse momentum, the mean p_T^e of the softest electron falls in a range where a full combination of tracking and calorimetry information becomes important. The p_T^e spectra for signal events at $m_H = 150$ GeV/ c^2 is shown for example in Fig. 2.9a. The softest electron, which generally couples to the off-shell Z*, has a most probable p_T^e value below 10 GeV/ c for masses $m_H \lesssim 140$ GeV/ c^2 . Hence, an excellent electron reconstruction is essential down to very low p_T^e values, well below the range of $p_T^e \simeq 40 - 45$ GeV/ c for which the reconstruction will be best constrained in CMS via measurements with SM single Z and single W production. The control of systematic uncertainties from experimental data is a major issue for such low p_T^e electrons and this will be discussed in detail in Section 2.3.4.

This analysis makes use of the elaborate reconstruction procedures which have been introduced very recently in CMS and have been described in detail in Ref. [37]. The electron identification and momentum measurements are hampered by the amount of tracker material which is distributed in front of the ECAL, and by the presence of a strong magnetic field aligned with the collider beam z axis. The procedures introduced in Ref. [37] provide new useful observables that allow to better deal with these detector effects, combining information from the pixel detector, the silicon strip tracker and the ECAL.

The pre-selection of the signal event candidates relies on the presence of at least 2 e^+ and 2 e^- candidates within the acceptance $|\eta| < 2.5$ and each with $p_T > 5 \text{ GeV}/c$, verifying the following characteristics:

- $E_{sc}/p_{in} < 3$, where E_{sc} is the supercluster energy and p_{in} the track momentum at the interaction vertex,
- $|\Delta\phi_{in}| = |\phi_{sc} - \phi_{in}^{extrap}| < 0.1$, where ϕ_{sc} is the energy weighted ϕ position of the supercluster and ϕ_{in}^{extrap} is the ϕ of the track at vertex, extrapolated to the ECAL assuming a perfect helix,
- $|\Delta\eta_{in}| = |\eta_{sc} - \eta_{in}^{extrap}| < 0.02$, with notations as in previous,
- $H/E < 0.2$, where H is the energy deposited in the HCAL tower just behind the electromagnetic seed cluster and E the energy of the electromagnetic seed cluster,
- and $\sum_{cone} p_T^{tracks}/p_T^e < 0.5$, a loose track isolation requirement, whose calculation will be described in Section 2.3.3.1.

The electron pre-selection efficiency is shown in Fig. 2.9b and Fig. 2.9c as a function of p_T^e and η^e for the electrons from Higgs boson events at $m_H = 150 \text{ GeV}/c^2$. The efficiency steeply rises and reaches a plateau around 86% for $p_T^e \gtrsim 20 \text{ GeV}/c$. The efficiency is above 90% for $|\eta| \lesssim 1.1$ and decreases towards the edge of the tracker acceptance when approaching $|\eta| \simeq 2.5$. The pre-selection efficiency for electrons from the same sample is represented in Fig. 2.9d as a two-dimensional map in the p_T versus η plane.

The absolute efficiencies for the Higgs boson signal at different m_H values and for the backgrounds are shown in Fig. 2.10a after triggering and the multi-electron pre-selection step. The acceptance for the Higgs boson signal is maintained above 50% in the full relevant mass range.

The signal and background events fulfilling the triggering and pre-selection steps are represented in the reconstructed invariant mass m_{4e} spectrum in Fig. 2.10b. The Higgs boson signal is seen to emerge above the background for masses around $150 \text{ GeV}/c^2$ and above $\simeq 2m_Z$. More background suppression is required elsewhere.

2.3.3 Event Selection and Kinematic Reconstruction

The further steps of the event selection rely on a more detailed characterization of the electron candidates and simple kinematic expectations. The electrons from the Higgs boson, in contrast to at least one e^+e^- pair from the $t\bar{t}$ and $Zb\bar{b}$ backgrounds, are isolated and originate from a common primary vertex. The corresponding analysis requirements are discussed in Section 2.3.3.1. Profiting from the expectation of a narrow resonance in the m_{4e} spectrum, and of the likely presence of a real Z boson in the final state, the kinematics and its simple evolution with m_H can be further exploited. The electrons of the e^+e^- pair at lowest m_{ee}







Figure 2.9: Electrons in SM Higgs boson $4e$ decay channel for $m_H = 150 \text{ GeV}/c^2$; a) transverse momentum of each of the four final state electrons; b) efficiency at pre-selection as a function of p_T^e ; c) efficiency at pre-selection as a function of η_e ; d) efficiency in the p_T^e versus η_e plane.





Figure 2.10: Higgs boson signal and dominant background sources after pre-selection step; a) overall pre-selection efficiency for m_H in the range from 115 to 300 GeV/c^2 and the background from $ZZ^{(*)}$ continuum, from $Zb\bar{b}$ and $t\bar{t}$; b) separate signal and background contributions to the spectrum of reconstructed invariant mass m_{4e} .

have on average a much harder p_T^e spectrum for the Higgs boson signal than for the $t\bar{t}$ and $Zb\bar{b}$ backgrounds. Moreover, the combination of the Z and $Z^{(*)}$ mass spectra distinguishes the Higgs boson signal from the $ZZ^{(*)}$ SM background continuum. These kinematic requirements are discussed in Section 2.3.3.2.

2.3.3.1 Isolated primary electrons

A very loose vertex constraint is first imposed on the longitudinal impact parameter for the four electron candidates in each event. All electrons should verify $IP_L/\sigma_L < 13$, where σ_L is the error on the longitudinal impact parameter IP_L . The main vertex constraint is imposed on the transverse impact parameter of the electrons to suppress secondary vertices. Secondary electrons appear for instance in semi-leptonic decays in the hadronization of the b quark jets in $Zb\bar{b}$ and $t\bar{t}$ background events. The sum of the transverse impact parameter significance (IP_T/σ_T), i.e. the ratio of the transverse impact parameter IP_T over its error σ_T , is shown separately in Fig. 2.11 for the e^+e^- pairs with the highest and lowest reconstructed mass. For both of these background sources, the displaced vertices are most likely to appear




Figure 2.11: Sum of the transverse impact parameter significance (IP_T/σ_T) of e^+e^- pairs for a Higgs boson at $m_H = 150 \text{ GeV}/c^2$, for the $ZZ^{(*)}$ continuum, and for $Zb\bar{b}$ and $t\bar{t}$; a) $\sum IP_T/\sigma_T$ from the electrons of the e^+e^- pair with a reconstructed mass m_{ee} best matching the Z boson mass; b) $\sum IP_T/\sigma_T$ from the second (softest) e^+e^- pair.

in the softest pair of reconstructed electrons. A best rejection power is obtained by imposing $\sum IP_T/\sigma_T < 30$ for the highest reconstructed m_{ee} and a more stringent cut of $\sum IP_T/\sigma_T < 15$ for the lowest reconstructed m_{ee} .

Another powerful discriminant against secondary electrons in b jets or in general against fake electrons in QCD jets, is provided by isolation requirements. The electrons coupled to the Z or $Z^{(*)}$ in the $H \rightarrow 4e$ channel are expected to be on average well isolated from hadronic activity. Hadronic activity in single Higgs boson production appears in NLO processes, in the recoil against the Higgs boson. The Higgs boson itself generally receives a significant longitudinal boost in the laboratory reference frame but, as a scalar, decays uniformly in its centre-of-mass reference frame. In contrast, the electrons in the b jets from $t\bar{t}$ or $Zb\bar{b}$ are accompanied by significant hadronic activity.

Two partly complementary observables can be best used for the isolation of low p_T^e electrons. These rely either on measurements of primary tracks or on the energy flow in the hadronic

calorimeter (HCAL). Both observables are insensitive to the eventual electron-induced electromagnetic showering in the tracker material. For the “track isolation”, an isolation cone




Figure 2.12: Electron isolation observables for the signal at $m_H = 150 \text{ GeV}/c^2$ and the SM backgrounds; a) track isolation, $\sum_{\text{cone}} p_T^{\text{tracks}}/p_T^e$; b) hadronic isolation, $\sum_{\text{cone}} E_T^{\text{HCAL}}/p_T^e$, for the second least isolated electrons.

of size $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.2$ is defined around the electron direction, and tracks with $p_T > 1.5 \text{ GeV}/c$ originating from the same primary vertex within $|\Delta IP_L| < 0.1 \text{ cm}$ are considered. To avoid suppressing signal events, tracks attached to an electron candidate of opposite charge, and giving $m_{e^+e^-} > 10 \text{ GeV}/c^2$, are discarded. All the 4 electrons from the Higgs boson candidate events must satisfy $\sum_{\text{cone}} p_T^{\text{tracks}}/p_T^e < 0.1$. Distributions of this track isolation observable are shown in Fig. 2.12a. For the “hadronic isolation”, all HCAL towers in an isolation cone size as above, and contributing with $E_T > 0.5 \text{ GeV}$ are considered in the ratio $\sum_{\text{cone}} E_T^{\text{HCAL}}/p_T^e$. This ratio is required to be below 0.05 for at least three electrons. The cut is relaxed to 0.2 for the fourth electron. Distributions of this hadronic isolation observable are shown in Fig. 2.12b.

Further electron identification requirements must be imposed to suppress the possible background, involving “fake” electrons, from Drell-Yan processes at NLO where a Z^* recoils against jet(s). Different electron identification cuts are used depending on the distinct classes of track-supercluster electron patterns [37] in order to preserve the electron detection efficiency at all η^e . More details can be found in Ref. [26]. This tightening of the electron identification entails an absolute efficiency loss for the Higgs boson signal below 5%.

2.3.3.2 Kinematics

The cascade $H \rightarrow ZZ^{(*)} \rightarrow 4e$ for a Higgs boson, mostly produced at small transverse momentum, leads to very distinctly ordered p_T^e spectra for the four final state electrons. Moreover, the p_T^e spectra of the softest electrons for the Higgs boson signal is on average harder than the one expected from secondary electrons from the $Zb\bar{b}$ or $t\bar{t}$ backgrounds. Thus, it is advantageous to profit from the knowledge of the expected p_T^e distributions for the Higgs boson signal. A best set of p_T^e cuts as a function of m_H is given in Table 2.10. The cut on the

softest electron is maintained to a lowest value for simplicity and to preserve the signal efficiency at low m_H . Otherwise the p_T^e cuts are seen to slowly evolve for as long as $m_H < 2m_Z$ and then rise faster above the Z pair production threshold. The p_T^e cuts lead for example [26]

m_H (GeV/ c^2)	p_T^1	p_T^2	p_T^3	p_T^4	$m_{Z_1}^{min}$	$m_{Z_1}^{max}$	$m_{Z_2}^{min}$	$m_{Z_2}^{max}$
		(GeV/ c)			(GeV/ c^2)			
115	7	10	10	15	51	101	10	50
120	7	12	15	15	51	101	10	50
130	7	12	15	15	61	101	10	60
140	7	12	15	15	71	101	10	65
150	7	12	15	15	71	101	15	65
160	7	15	15	15	71	101	15	70
170	7	15	15	20	81	101	20	80
180	7	15	15	20	81	101	30	90
190	7	15	20	30	81	101	40	100
200	7	15	25	30	81	101	40	110
250	7	20	40	50	51	131	20	200
300	7	30	40	60	51	131	15	300

Table 2.10: Electron p_T cuts, from the lowest to the highest p_T electron and reconstructed Z_1 and Z_2 invariant mass cuts.

to a reduction by a factor of 5 to 10 of the $Zb\bar{b}$ background, and a factor of 3 to 5 of the $t\bar{t}$ background for $m_{4e} < 2m_Z$. Both backgrounds are also heavily suppressed above $2m_Z$.

Labeling Z_1 the boson reconstructed with an m_{ee} closest to the nominal Z mass and Z_2 the one reconstructed from the second e^+e^- pair, one expects for $m_{4e} < 2m_Z$ in the case of the the Higgs boson signal that $m_{4e} \simeq m_{Z_1} + m_{Z_2}$ with most often the presence of a Z boson on its mass shell, $m_{Z_1} \simeq m_Z$. The Z boson masses saturate the phase space and are dominantly produced with small velocity in the Higgs boson rest frame. The requirement of one real Z boson suppresses further the $t\bar{t}$ backgrounds for low m_{4e} . The cut on Z_2 is powerful against the $ZZ^{(*)}$ continuum and further suppresses the $Zb\bar{b}$ and $t\bar{t}$ backgrounds. A set of optimal Z_1 and Z_2 cuts is given in Table 2.10 as a function of m_H . The cuts lead for example [26] for $m_{4e} \simeq 150 \text{ GeV}/c^2$ to a reduction of the $ZZ^{(*)}$ continuum by a factor of about 6.5 and a reduction of the $t\bar{t}$ background by a factor of about 2.5.

m_H	115	120	130	140	150	160	170	180	190	200	250	300
	(GeV/ c^2)											
Signal	24.3	26.0	31.2	35.2	36.0	37.4	38.0	39.9	40.9	42.5	41.2	38.6
$ZZ^{(*)}$	5.24	4.94	5.68	5.95	5.14	5.23	6.87	17.8	25.1	26.2	22.3	13.9
$Zb\bar{b}$	0.22	0.16	0.17	0.17	0.16	0.10	0.097	0.068	0.037	0.031	0.013	0.001
$t\bar{t}$	0.054	0.044	0.043	0.033	0.032	0.022	0.021	0.011	0.008	0.008	0.013	0.006

Table 2.11: Summary of selection efficiencies normalized to the generation pre-selection efficiency.

The Fig. 2.13a shows as an illustration the expected m_{4e} invariant mass distributions for the signal at $m_H = 150 \text{ GeV}/c^2$ and for backgrounds after triggering and pre-selection. The further background suppression from the isolated primary electron requirement, the p_T^e and Z mass cuts is seen by comparison in Fig. 2.13b. The global selection efficiency (normalized to the acceptance defined at the generation level) is given in Table 2.11 for the signal and backgrounds. The Fig. 2.13c and Fig. 2.13d show for illustration the possible outcome of two random Monte Carlo experiments corresponding to favorable and less favorable fluctuations of the Higgs boson signal for an integrated luminosity of 30 fb^{-1} . The Poissonian probability




Figure 2.13: Distributions of the reconstructed invariant mass m_{4e} for the SM Higgs bosons signal at $m_H = 150 \text{ GeV}/c^2$ and for the SM backgrounds after (a) pre-selection step and (b) after all cuts. The number of events are normalized in cross-section. Single Monte Carlo experiments corresponding to an integrated luminosity of 30 fb^{-1} for (c) a favorable case and (d) a less favorable case.

to have equal or more favorable (respectively equal or less favorable) fluctuations is of about 5% for the example cases shown.

2.3.4 Systematics

In this section the systematic errors are discussed in the context of a discovery via a simple event counting method. The “theoretical” and “experimental” sources of errors are distinguished. The theoretical uncertainties concern the estimation of the background rates within the cuts defining the acceptance of the Higgs boson signal and are discussed in Section 2.3.4.1. The experimental uncertainties take into account the limited knowledge of the detector responses and efficiencies, and of the corresponding Monte Carlo modelling. These are discussed in Section 2.3.4.2. A comparison of different methods for the control of background systematics is presented in Section 2.3.4.3.

2.3.4.1 Theoretical errors

The theoretical uncertainty on the number of background events in the signal region from PDFs and QCD scales variations has been estimated by the MCFM program [38]. CTEQ6M

PDF are used and 20 eigenvector parameters have been varied by $\pm 1\sigma$. Both QCD normalization and factorization scales have been varied independently up and down for a factor two from their nominal values of $2m_Z$. The resulting uncertainties from PDF and QCD scale are of the order of 6% for direct estimation of ZZ background, from 2 to 8% for normalization to single $Z \rightarrow 2e$, and from 0.5 to 4% for the normalization to sidebands discussed further in Section 2.3.4.3. The gluon fusion cross-section uncertainties in the ZZ background of 8% is also considered as a part of theoretical uncertainties.

The uncertainty on the normalization of the measurements to the pp luminosity of the LHC collider is estimated to be of the order of 3% for an integrated luminosity above 10 fb^{-1} .

2.3.4.2 Experimental errors

The main remaining sources of experimental systematics expected in the CMS experiment after having collected of $\mathcal{O}(10) \text{ fb}^{-1}$, and relevant for the $H \rightarrow 4e$ channel, originate from uncertainties on knowledge of the amount of tracker material in front of the ECAL, from the precision of the (pattern dependent) energy calibration of electron objects, and from the control of electron efficiencies. The strategy adopted consists of relying on experimental data, and in particular on single Z and W production, to minimize these systematic errors. The electrons from $W \rightarrow e\nu$ and $Z \rightarrow ee$ decays are used to control the energy measurements and reconstruction efficiencies.

A change of the integral amount of tracker material traversed by electrons before reaching the ECAL is susceptible of affecting the electron selection and identification efficiencies, as well as energy measurement scales and resolution. The uncertainty on the material budget will limit the precision of the acceptance calculations, when using the Monte Carlo model to extrapolate away from the kinematic domain best constrained via single Z and W measurements.

There are many observables that are directly or indirectly sensitive to the amount of tracker material, and that have been used in collider experiments. Examples are the distribution of converted photon vertices, or the shape of the E/p comparing tracker momentum measurement p to the energy E measured in the calorimeter in finite cluster volume, or a comparison of data and Monte Carlo for the Z mass resolution, etc. A new technique is used which is based on the electron GSF tracking introduced recently in Ref. [37]. The difference between the momentum magnitude at vertex and at the last hit, $p_{\text{in}} - p_{\text{out}}$, is a measure of the integral amount of bremsstrahlung. The mean fraction f_{brem} of the energy radiated along the complete trajectory is roughly proportional to the integral amount of material traversed. Hence, one can relate f_{brem} to the material thickness X/X_0 where X_0 is the characteristic radiation length via the formula $\langle X \rangle / X_0 \simeq -\ln(1 - f_{\text{brem}})$, where $f_{\text{brem}} = (p_{\text{in}} - p_{\text{out}})/p_{\text{in}}$.

The amount of tracker material measured in this way for single electron data is shown in Fig. 2.14a. The results obtained in the configuration corresponding to the nominal tracker material coincide very well with the known material distribution as given in Ref. [7]. The Fig. 2.14b shows the ratio of the measured material thickness obtained in configurations where the amount of material was changed by $\pm 10\%$, normalized to the measurement results in the nominal case. The ratio is found to be remarkably stable as a function of η , despite the fact that the integral amount of material has a strong η dependence. Thus, single electrons can be used in CMS to tune the Monte Carlo model of the tracker material per η slice. The Fig. 2.14c shows that in a given η slice the measured material thickness is linearly

correlated to a change (at least within a range of $\pm 10\%$) of the true material thickness. Similar results are obtained when considering various restricted range of p_T^e within a sample of uniformly distributed electrons in the p_T^e range from 5 to 100 GeV/ c . With the electron statistics expected from single Z production for an integrated LHC luminosity of $\mathcal{O}(10) \text{ fb}^{-1}$, it should be possible to determine the tracker material thickness to a precision better than 2% over the full acceptance in η . The Fig. 2.14d shows that such a 2% uncertainty on the material budget have almost no effect on electron reconstruction efficiency.




Figure 2.14: Sensitivity to variations of the tracker material budget from electron measurements based on GSF tracks; a) measured amount of material as a function of $|\eta|$ for the nominal tracker configuration and for an integral material budget changed by $\pm 10\%$; b) ratio of the measured amount of material as a function of $|\eta|$; c) measured versus true thickness in X_0 of the tracker material; d) effect of a change of 2% of the material budget on the electron reconstruction efficiency.

Electron reconstruction efficiencies and energy scales will be controlled by electrons from $W \rightarrow e\nu$ and $Z \rightarrow ee$ decay. Huge cross-sections of these two processes will allow for a significant reduction of reconstruction uncertainties already after few fb^{-1} . Electrons from $Z \rightarrow ee$ are produced centrally with a characteristic Jacobian p_T distributions around 45 GeV/ c . It is therefore expected that the best control of experimental systematics is obtained in the central part of the detector and for electrons around the Jacobian peak.

Electron reconstruction uncertainties as a function of η and p_T are given in Fig. 2.15a and Fig. 2.15b respectively, for an integrated luminosity of 0.2 fb^{-1} . The expected behaviour of an increased uncertainties when moving away from the Jacobian peak or from the central η region can be clearly seen. From the expected reconstruction errors evolution with the

luminosity, all reconstruction efficiency uncertainties can be safely absorbed in a single factor of 1% per electron, for integrated luminosities larger than 10 fb^{-1} .

The second important systematic effect is the uncertainty on the energy scale determination. Using single Z production, it has been shown in Ref. [39] that the absolute energy scale for electrons can in principle be controlled with great precision with average uncertainties reaching values below 0.1%. The systematic uncertainties has to be studied as a function of p_T^e and η^e given the different electron spectrum in $H \rightarrow ZZ^{(*)} \rightarrow 4e$ and $Z \rightarrow ee$ decays. The reachable precision depends on the amount of integrated LHC luminosity. In this analysis, the second leg of a Z boson decay, tagged as an electron by imposing stringent electron identification requirements on the first leg combined with a kinematic constraint to the Z boson mass, is used as a probe to estimate systematics on the energy scale. Uncertainties versus η and p_T for golden and showering electrons are shown in Fig 2.15c and Fig 2.15d, for the integrated luminosity of 0.2 fb^{-1} . With expected evolution of these uncertainties with the luminosity, it is found that an uncertainty in energy scale of 0.5% in the barrel region, and 1% in the endcaps, for integrated luminosities larger than 10 fb^{-1} , can be safely considered.




Figure 2.15: Control of experimental uncertainties using SM data; uncertainties on measurements of electron reconstruction, isolation and identification as a function of (a) η and (b) p_T ; uncertainties on measuring the energy scale for golden and showering electrons as a function of (c) η and (d) p_T .

2.3.4.3 Control of background rates

Following the primary and isolated electron selection and the application of basic kinematic requirements, only the $ZZ^{(*)}$ continuum remains as the dominant or sole background over the full mass range in consideration for the SM Higgs boson search. Thus, the determination of the mean expected number of SM $ZZ^{(*)}$ background events in the signal region, defined e.g. by a simple sliding window in the m_{4e} spectrum, remains as a key issue.

The three main methods for the estimation of $ZZ^{(*)}$ continuum contribution to the background in the signal region are:

- direct simulation of the $ZZ^{(*)} \rightarrow 4e$ process,
- normalization to the $Z \rightarrow 2e$ data,
- normalization to the sidebands.

The first method entirely relies on existing SM constraints and the theoretical knowledge, with uncertainties coming from the PDFs used to describe the colliding protons and from QCD scale variations. It furthermore is reliant on the LHC luminosity uncertainties, and on the Monte Carlo modeling of the acceptance and detector response for the uncertainties arising from electron reconstruction and selection. Otherwise, the method potentially benefits from the fact that the statistical precision on the mean background expectation is only limited by the Monte Carlo statistics, and can therefore be assumed negligible in the context of a prospective for an analysis to be performed in a future CMS experiment.

The second method aims at profiting from the fact that the SM single Z production cross-sections is measured with great precision in an experiment which will have integrated a luminosity of $\mathcal{O}(10) \text{ fb}^{-1}$ at the LHC. Using the ratio of $ZZ \rightarrow 4e$ to $Z \rightarrow 2e$ rates allows to profit from a full cancellation of pp luminosity uncertainties, while providing a partial cancellation of PDF and QCD scale variations uncertainties (due to their correlations in a part of the initial state phase space) and a partial cancellation of experimental uncertainties.

In the method of the normalization from sidebands, the number of background events *inside* the acceptance of the signal region is determined from the number of background events measured *outside* the signal region, by multiplying the latter with the ratio α_{MC} between *inside* and *outside* expectations as determined using Monte Carlo simulation. Using the sidebands one also expects to fully cancel luminosity uncertainties, to reduce PDF and QCD scale variation uncertainties and substantially reduce experimental uncertainties too. Statistical errors with sidebands normalization come from the statistics of the background rate outside the signal region and can be a limiting factor for the method. By relaxing some of late analysis cuts, such as invariant Z mass, the background events rate outside the signal region increases, reducing therefore statistical errors for this method. The price to pay is an increased background rate in the signal region too and, therefore, some balancing is needed.

Using results from previous sections, both theoretical and experimental uncertainties are evaluated for two methods: normalization to the $Z \rightarrow 2e$ measurements and normalization to the sidebands. For the normalization to single $Z \rightarrow 2e$ measurements results are shown in Figure 2.16a. The overall systematic uncertainty with this method is of about 5%. Experimental uncertainties are seen to dominate for $m_H \simeq 2m_Z$ while theoretical errors take over above the pair production threshold. Uncertainties for the sidebands normalization are shown in Figure 2.16b. Statistical uncertainties scale as the square root of the number of background events outside the signal region and are shown for an integrated luminosity of




Figure 2.16: Theoretical and experimental uncertainty estimations for both methods for evaluation of background from data: (a) normalization to the single $Z \rightarrow 2e$ measurements and (b) normalization to the sidebands. Expected statistical errors for sidebands are also shown, for integrated luminosity of 30 fb^{-1} .

30 fb^{-1} and for two analysis scenarios: after all analysis cuts and without cuts on the mass of both Z bosons. A trade-off in the second method is in a somewhat lower nominal significance (for about 8%) while statistical errors decrease by a factor of about 2.5. Full significance calculations with and without systematics and statistical uncertainties are presented in the following section.

2.3.5 $H \rightarrow 4e$ Observability, Mass and Cross-Section Measurements

2.3.5.1 Discovery reach

A simple counting experiment is used here to quantify the sensitivity of the experiment to the presence of a Higgs boson signal. The expected number of signal (N_S) and background (N_B) events are evaluated in a sliding window whose central position m_{4e} varies between 100 and 320 GeV/c^2 . The size of the optimal window increases progressively from 6 GeV/c^2 at $m_{4e} = 115 \text{ GeV}/c^2$ to 24 GeV/c^2 at $m_{4e} = 300 \text{ GeV}/c^2$. The Table 2.12 presents for each Higgs boson mass hypothesis the mean expected number of signal and background events, and associated uncertainties.

The significance of the $H \rightarrow 4e$ signal observation is shown as a function of m_H in Fig. 2.17 (left) as expected for an integrated luminosity of 30 fb^{-1} . The results are given for both the S_{cP} and the S_{cL} significance estimators. The S_{cP} is defined as the probability for a Poisson distribution with mean N_B to observe a number of events equal or greater than $N_S + N_B$, converted in the equivalent number of standard deviations of a Gaussian distribution. The S_{cL} corresponds to the widely used log-likelihood ratio significance [40] and is given for comparison. The effect of including experimental and theoretical systematics, described in section 2.3.4 and listed in Table 2.12, on the significance S_{cP} [41] is also shown, for two different methods of controlling the background uncertainties. A signal observation with a significance above 3 standard deviations is expected in the $H \rightarrow 4e$ channel alone for m_H in the range from 130 to 160 GeV/c^2 , and above 180 GeV/c^2 . The integrated luminosity needed for a 5 standard deviations discovery of the SM Higgs boson in the $H \rightarrow 4e$ channel alone is

m_H	115	120	130	140	150	160	170	180	190	200	250	300
	(GeV/c^2)											
N_S	1.52	2.97	8.18	15.80	17.19	8.38	3.76	9.95	34.05	38.20	27.68	21.69
N_B	2.26	1.94	3.71	4.31	3.68	3.10	3.37	6.42	14.62	17.29	13.40	7.63
δN_B												
exp. :	0.063	0.089	0.126	0.167	0.105	0.148	0.145	0.187	0.551	0.505	0.466	0.187
theo. :	0.039	0.049	0.079	0.098	0.095	0.084	0.100	0.191	0.440	0.549	0.602	0.417

Table 2.12: Expected number of Higgs boson signal (N_S) and SM background (N_B) events for an integrated luminosity of 30 fb^{-1} , in the optimized window for the reconstructed invariant mass m_{4e} . The uncertainties (δN_B) are given for systematics from experimental (exp.) and theoretical (theo.) sources, for an analysis where the $ZZ^{(*)}$ continuum has been normalized to the measurement of single Z production.




Figure 2.17: (a) Significance S_{cP} for an integrated luminosity of 30 fb^{-1} as a function of the Higgs boson mass without and with systematics included in both options of $ZZ^{(*)}$ normalization to the measured sidebands or the measured single Z production cross-section. The significance S_{cL} is also shown. (b) luminosity needed for a 3σ observation and 5σ discovery with the systematics included using $ZZ^{(*)}$ normalization to the Z cross-section.

also shown as a function of m_H in Fig. 2.17b. Systematic errors from normalisation to the Z cross-section have been included.

2.3.5.2 Mass, width and cross-section measurements

At an early stage of the Higgs boson search and discovery in the $H \rightarrow 4e$ channel, given very low statistics, a robust and simple estimation of m_H can be obtained by a simple mean (or weighted mean) of the m_{4e} values measured for individual events. The events falling in the pre-defined optimal mass window introduced in the above Section 2.3.5.1 and used to establish the signal significance, can be used for such purposes. For higher statistics, a fit of the m_{4e} mass distribution to a signal plus background shape can be used to extract simultaneously the mass and the cross-section \times branching ratio of a Higgs boson signal. Detector effects dominate the Higgs boson mass resolution below the Z pair production threshold and a sensitivity to the Higgs boson intrinsic width is expected only for masses well above $2m_Z$.

The precision on the parameter measurements for the Higgs boson depend on the quality of the reconstructed electrons and can, in general, be improved using event-by-event errors

on the electron momentum estimation [37]. Example cases for two different sub-samples of Higgs boson events differing by the pattern of the four reconstructed electrons are presented in Fig. 2.18. Clearly, event candidates built from four *non-showering* electrons in the barrel




Figure 2.18: Mass measurements: (a) example case for two different event sub-samples differing by the pattern of the four reconstructed electrons; (b) relative errors as a function of the Higgs boson mass using the mean mass and the fitted mass as obtained for an integrated luminosity of 30 fb^{-1} .

part of the ECAL, a subset representing only about 1.76% of all signal events, allow for a much better m_H measurement (smallest errors on average and least dispersion of the mass measurement errors) than candidates built mainly from e.g. *showering* electrons in the end-caps part of the ECAL. About 36.7% of the signal event candidates contain three or more *showering* electrons. A weighted mean of the events of the m_{4e} distribution falling in the signal window has been considered for the estimation of the Higgs boson mass in Ref. [26]. A simple mean can be also used for simplicity.

The reconstructed Higgs boson mass and its error obtained from the mean value for events falling in the expected signal window is presented in Fig. 2.18b. The error is obtained from the dispersion of the mean values obtained from large number of Monte Carlo experiments at an integrated luminosity of 30 fb^{-1} . The results are shown as a function of the Higgs boson mass. The systematic bias on the mass estimate for the low m_H cases for this simple mean approach is due to the asymmetric shape of the reconstructed signal and can be modelled. In the mass ranges where the Higgs boson signal significance exceeds 3 standard deviations, the uncertainty on the mass determination is found to be everywhere below 0.4%. It reaches values below 0.2% for $m_H \simeq 200 \text{ GeV}/c^2$. For comparison, results obtained by fitting the m_{4e} distribution are also shown. The fit method requires a significant number of events (typically $\gtrsim \mathcal{O}(10)$) to converge and provide reasonably stable results. The m_{4e} distribution is fitted by a signal plus background shape. The signal contribution is modeled with two Gaussians, describing respectively the core and the low m_{4e} tail of the signal distribution. The tail parameters (fraction, mean and dispersion) are fixed by fitting the “signal only” expectation. The background is modeled using a flat distribution up to about $m_{4e} \approx 2m_Z$ and a linear function (non-zero slope) for higher Higgs boson masses. This has been found to provide a sufficiently good model of the observation in a restricted mass range around the signal region. A likelihood fit is then performed on each Monte Carlo experiments and the reconstructed mass and precision are extracted from the distribution of the fitted values of the peak of the Gaussian core. Where the fit can be performed, the Fig. 2.18b shows that an unbiased estimation of m_H is obtained within errors.

The fitted number of signal events is used to estimate the production cross-section by correcting for the global acceptance efficiency. The statistical precision on this measurement is here also obtained from the width of the distribution of the fitted parameters in Monte Carlo experiments. An unbiased measurement of the cross-section is obtained over the full mass range considered here, with a precision of the cross-section measurement between 20 and 30%. With such a precision, the influence of the detector systematics (about 5%) and of the uncertainty on the luminosity measurement (less than 3% for 30 fb^{-1}) is marginal. For an integrated luminosity of 60 fb^{-1} , the precision on the cross-section measurement improves to about 15%.

A measurement of the width is possible only for Higgs boson masses above $\gtrsim 2m_Z$ where at the same time the Higgs natural width is becoming large and the detector resolution is improving. A Gaussian width with central values of about $2.3 \text{ GeV}/c^2$ for $m_H = 200 \text{ GeV}/c^2$ and $4.2 \text{ GeV}/c^2$ for $m_H = 300 \text{ GeV}/c^2$ is obtained from the fit, but with a rather large uncertainty of about 50%.

Chapter 3

Physics Studies with Muons



Benchmark Channel: $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$

The $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ process is one of the cleanest channels for discovering the Standard Model Higgs boson at LHC. This section presents the CMS potential for discovering the Higgs boson in this decay mode and measuring its mass, width, and production cross section, in the range of Higgs boson masses from 115 GeV to 600 GeV. Both signal and background event samples are generated at the Leading Order (LO) approximation, and Next to Leading Order (NLO) production cross sections, computed using different methods, are used for their normalization. To simulate the detector response and reconstruct physics objects, the full CMS detector simulation and reconstruction software was used. A full treatment of the most important theoretical and instrumental systematic uncertainties are presented, together with their effect on the evaluation of the significance of the Higgs boson observation and on the measurement of its parameters. To minimize systematic uncertainties, new methods of reconstructing the most important corrections directly from data were developed.

3.1.1 Physics Processes and Their Simulation

The Higgs boson event samples for 18 Higgs boson mass points and the three main background processes, $t\bar{t}$, $(Z^{(*)}/\gamma^*)b\bar{b}$ and $(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$ were simulated using the CMS simulation [14] and reconstruction [15] software. These three backgrounds will be hereafter referred to as $t\bar{t}$, $Zb\bar{b}$ and ZZ , respectively. Details on the generator-level simulation conditions, cross sections and K-factors can be found in [42]. Many other plausible background candidates, $b\bar{b}b\bar{b}$, $b\bar{b}c\bar{c}$, $c\bar{c}c\bar{c}$, single-top, $Zc\bar{c}$, $Wb\bar{b}$, $Wc\bar{c}$, fake and π/K decay muons in QCD, were considered and found to be negligible.

Only events with at least $2\mu^+$ and $2\mu^-$ in pseudorapidity range $|\eta| < 2.4$ and with $p_T > 3\text{ GeV}$ were retained for further analysis. Muons outside these kinematic limits could not be reconstructed in the CMS detector. Additional cuts were applied on dimuon invariant masses for the Higgs boson samples ($m_Z > 5\text{ GeV}$) and for ZZ and $Zb\bar{b}$ samples ($m_{\mu^+\mu^-} > 5\text{ GeV}$). The first $\mu^+\mu^-$ pair in the ZZ and $Zb\bar{b}$ samples was defined as the one with its invariant mass closest to m_Z , while the second $\mu^+\mu^-$ pair was made out of the two remaining highest p_T muons of opposite charge. These cuts do not bias the Monte Carlo samples since all the analysis cuts, described below, are tighter.

The Higgs boson samples were generated with PYTHIA 6.225 [30] (LO gluon and weak boson fusion, $gg \rightarrow H$ and $q\bar{q} \rightarrow q\bar{q}H$) interfaced via CMKIN [43]. Events were re-weighted to correspond to the total NLO cross section $\sigma(\text{pp} \rightarrow H) \cdot BR(H \rightarrow ZZ) \cdot BR(Z \rightarrow 2\ell)^2$ (Fig. 3.1).




Figure 3.1: Standard Model NLO cross section for the process $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ vs. Higgs boson mass.




Figure 3.2: Distributions of $m_{4\mu}$ after preselection cuts for $t\bar{t}$, $Zb\bar{b}$, ZZ and a Higgs boson signal of $m_H = 140$ GeV.

The cross section $\sigma(pp \rightarrow H)$ and the branching ratio $BR(H \rightarrow ZZ)$ were taken from [44]; $BR(Z \rightarrow 2\ell) = 0.101$ [45]. Interference of permutations of identical leptons originating from different Z bosons results in an enhancement to the cross section for $H \rightarrow ZZ^{(*)} \rightarrow 4\ell$ processes with identical leptons [42], which is about 15% for $m_H = 115$ GeV and steadily goes to zero for $m_H = 180$ GeV. This correction was calculated with CompHEP.

The $t\bar{t}$ sample was generated with PYTHIA 6.225 (LO $gg \rightarrow t\bar{t}$ and $q\bar{q} \rightarrow t\bar{t}$). Events were reweighted to correspond to the total NLO cross section $\sigma(pp \rightarrow t\bar{t}) \cdot BR(W \rightarrow \ell\nu)^2$. The NLO cross section $\sigma(pp \rightarrow t\bar{t}) = 840$ pb was taken from [46] and the branching ratio $BR(W \rightarrow \ell\nu) = 0.320$ from [45].

The $Zb\bar{b} \rightarrow \mu^+\mu^-b\bar{b}$ sample was generated with the CompHEP 4.2p1 [47] matrix element generator, interfaced to PYTHIA 6.225 for showering and hadronization. Included subprocesses were $q\bar{q}/gg \rightarrow (Z/\gamma^*)b\bar{b} \rightarrow \mu^+\mu^-b\bar{b}$. The corresponding CompHEP LO cross section was found to be 116 pb. To obtain the NLO cross section a NLO K-factor $K_{NLO} = 2.4 \pm 0.3$, computed with MCFM [48], was used.

The $q\bar{q} \rightarrow ZZ \rightarrow 4\mu$ and $q\bar{q} \rightarrow ZZ \rightarrow 2\mu 2\tau$ event samples were generated with CompHEP, including both the t - and s -channel diagrams [49]. The CompHEP events were further interfaced to PYTHIA 6.225 for showering and hadronization. The CompHEP LO cross sections for the two sub-processes were 113 fb and 157 fb. To account for contributions due to all the NLO diagrams and due to the NNLO gluon fusion ($gg \rightarrow ZZ$, known to contribute $\sim 20\%$ with respect to the LO [33] cross section), events are reweighted with the $m_{4\mu}$ -dependent K-factor $K(m_{4\mu}) = K_{NLO}(m_{4\mu}) + 0.2$. The NLO K-factor $K_{NLO}(m_{4\mu})$ was obtained with MCFM. The details on the dynamic differences between NLO and LO are summarized elsewhere [50].

The $m_{4\mu}$ distributions for a Higgs boson signal of $m_H = 140$ GeV and the main backgrounds are shown in Fig. 3.2 after the preselection cuts described above.

3.1.2 Event Selection

3.1.2.1 Trigger and offline muon selection

CMS has been designed and optimized to detect and reconstruct muons. These particles provide a very clean signature and thus a very high trigger efficiency, with an average over 98% for the Level-1 Global Muon Trigger [7]. The inclusive muon triggers based on the selection of a single muon with $p_T > 19\text{ GeV}$ or dimuons with $p_T > 7\text{ GeV}$ assures an efficiency of practically 100% for collecting events with four high- p_T muons.

In order to minimize muon reconstruction systematic uncertainties, we select only those reconstructed muons that have transverse momentum $p_T > 7\text{ GeV}$, if they are in the central pseudo-rapidity region ($|\eta| < 1.1$), or with momentum $p > 13\text{ GeV}$, if they are in the endcaps ($|\eta| > 1.1$) [51]. These cuts do not affect the number of accepted signal events significantly.

Also, we require that all four possible combinations of the reconstructed dimuon masses be above 12 GeV , $m_{\mu^+\mu^-} > 12\text{ GeV}$. As in the previous case, this cut has a very little effect on the Higgs boson events and is primarily intended to suppress poorly simulated hadron background contributions originating from charmonium and bottomium dimuon decays.

3.1.2.2 Discriminating variables

The $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ signal presents a characteristic topology, which consists of two opposite charge muon-pairs in the final state. All four muons are isolated, have a high transverse momentum and point to the same Z-boson mass, depending on the restrictions in the phase space introduced by the Higgs boson mass itself. The four-muon invariant mass peaks at the Higgs boson mass, within the detector resolution. The width of the resonant peak accounts for the natural Higgs boson width and the detector resolution.

In $Zb\bar{b}$ and $t\bar{t}$ background events, two of the muons come from b-quark decays and are usually found within a jet (i.e., non-isolated), have lower transverse momenta and often exhibit detectable displaced vertices. The isolation is defined as the amount of transverse energy in the calorimeter (calorimeter isolation), or the sum of the transverse momentum of the tracks reconstructed in the tracker (tracker isolation), inside a cone in η - ϕ space with a radius $R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ around each muon. Figure 3.3 (left) shows the distribution of the calorimeter isolation variable for the least isolated muon, for two Higgs boson signals, 150 GeV and 300 GeV , and for the background. Requiring a maximum isolation in all four muons drastically suppresses $t\bar{t}$ and $Zb\bar{b}$ contamination.

Further restrictions on the p_T spectrum of the 2 lowest p_T muons in the event (see Figure 3.3 (right), for the 2nd lowest p_T muon) reduces even more the $t\bar{t}$ and $Zb\bar{b}$ contamination. In this way, the ZZ background, which presents a topology very similar to that of the signal, becomes a dominant and irreducible background. Only the four-muon mass distribution, the main discriminant variable, allows the resonant Higgs signal to be identified over the continuum ZZ production.

Distinction on the basis of dimuon invariant mass or displaced vertices does not increase the Higgs boson signal over the ZZ background. However, they may play an important role in eliminating other possible unaccounted for backgrounds, arising from the primary interactions, accelerator beam halo, detector mis-performance, etc.




Figure 3.3: Examples of discriminating variables: (left) muon calorimeter-based isolation $\sum E_T$ for the least isolated muon and (right) transverse momentum of the 2nd lowest p_T muon. The hatched histograms represent the Higgs boson signals of masses 150 GeV and 300 GeV, while the solid, dashed and dash-dotted lines indicate the contribution from the $t\bar{t}$, ZZ and $Zb\bar{b}$ backgrounds, respectively. The arrows indicate the positions of the cuts.

Additional variables that may help discriminating H from the dominant ZZ background have been studied: $p_T(4\mu)$, number of jets and their E_T , etc. However, these variables are driven by the NLO production processes, while our samples were generated at the Leading Order by PYTHIA and CompHEP. Therefore, any conclusions that we might derive from these samples would not be reliable. Some angular distributions built from muons also have some differences originating from the underlying spin structures, but they are not sufficiently discriminating to be used for cuts and may be strongly affected by the NLO diagrams.

3.1.3 Higgs Boson Search Analysis

3.1.3.1 Search using $m_{4\mu}$ -independent cuts

Given the clear signature of the Higgs boson events, the signal extraction has been performed with a unique set of cuts, independent the Higgs boson mass, the details can be found in [52]. A Higgs mass-independent analysis is expected to minimize the dependence on the simulation of the discriminating variables in the Monte Carlo and the sensitivity to systematic errors. It is also readily applicable to real data and robust under variations of the detector conditions (calibrations, resolutions, efficiencies). Moreover, in our case, a mass-dependent selection does not significantly increase the significance of observing a signal.

The unique set of selection cuts has been designed to make the analysis robust when applied to real data. As explained below, some of the cuts (dimuon invariant mass, Pt cuts on the two hardest muons and isolation cuts on the two most isolated muons) slightly decrease the signal significance but make the selection more robust under imperfect conditions in the detector.

A loose requirement on the invariant mass of the pair of unlike-sign muons in the event

which is closer to the nominal Z-boson mass, namely, $70 \text{ GeV} < m_{\mu^+\mu^-} < 100 \text{ GeV}$, leaves more than 90% of the signal, while eliminating around 50% of the $t\bar{t}$ contamination. The loss in the signal is due to the internal bremsstrahlung and $Z \rightarrow 2\tau \rightarrow 2\mu 4\nu$ decays.

Cuts of 12 GeV and 8 GeV are set on the p_T of the two lowest- p_T muons. The p_T of the two highest- p_T muons must be larger than 15 GeV. The latter cut almost does not affect neither the signal nor the background, but is considered useful for eliminating unexpected background in real data. The efficiency of the p_T cuts in the signal is close to 90% while it suppresses around 50% of the remaining $Zb\bar{b}$ events, 40% of the $t\bar{t}$ events and about 20% of the ZZ background.

For the purposes of the isolation cut optimization, different cone radii and several energy and transverse momentum thresholds have been studied. Those yielding the maximum signal significance are, for calorimeter isolation, a cone radius of 0.24 and energy thresholds of 5 GeV and 9 GeV, while for tracker isolation a cone radius of 0.20 and p_T thresholds of 2.5 GeV and 4 GeV. The numbers are given for the two least isolated muons. Although a requirement on the isolation of the two most isolated muons does not increase the signal significance, following the same argument as in the case of the p_T cuts, a cut of 3.5 GeV and 5 GeV for the calorimeter isolation and 2 GeV and 2.5 GeV for the tracker isolation is set for the two most isolated muons.

After these cuts, $Zb\bar{b}$ and $t\bar{t}$ events are suppressed to a negligible level in comparison to the remaining ZZ background. The efficiencies of each selection cut over the signal, for the 18 Higgs mass points studied, are shown in Figure 3.4 (left). The four-muon mass distributions for signal and background events that survive the selection cuts are displayed in Figure 3.4 (right).




Figure 3.4: (Left) $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ efficiency vs. m_H after different cuts are applied. (Right) Reconstructed four-muon invariant mass distribution, for an integrated luminosity of 30 fb^{-1} , for background (shaded histograms) and several Higgs signals (hatched), after the selection criteria are applied.

In order to estimate the statistical significance of the signal, the log-likelihood ratio (LLR) statistical method [53, 54] is used. The distribution to discriminate signal and background is the

four-muon invariant mass (Fig. 3.4 (right)). This distribution, for each Higgs boson mass hypothesis and for the background, is used to calculate the log likelihood ratio, $-2 \ln Q$, which is then used to evaluate the compatibility of the data with either the *signal plus background* or the *background-only* hypothesis [?]. The $-2 \ln Q$ estimator is sensitive both to the normalization and the shape of the discriminant. Each event in the sum has a weight $\ln(1 + s/b)$ which depends on the signal-to-background ratio, s/b , in the bin where it is found, which in turn depends on the m_H hypothesis. The whole spectrum of the discriminant variable enters the LLR calculation. This avoids any ambiguity in the definition of a signal region for determining the signal significance, present in counting methods.

Figure 3.5 (left) shows the statistical significance, $S_L \equiv \sqrt{\langle 2 \ln Q \rangle}$, for an integrated luminosity 30 fb^{-1} at different $m_{4\mu}$ invariant masses, should the Higgs boson exit at one of these masses. Based on this distribution, the plot on the right depicts the integrated luminosity required to reach a statistical significance of the signal of 3σ and 5σ , as function of m_H . The expected integrated luminosity required to exclude the signal at the 95% confidence level in a background-only experiment is also shown as function of m_H . The effect of including systematic uncertainties (subsection 3.1.3.3) in the calculation of S_L is at the level of 15%-20% of the statistical accuracy of the expected significance, supporting that this analysis is not




Figure 3.5: (Left) Statistical significance of the signal, S_L , as function of the Higgs boson mass for an integrated luminosity of 30 fb^{-1} , for mass-independent cuts (red squares, solid line) and mass-dependent cuts (black dots, dashed line). The shaded band represents the statistical uncertainty on S_L . (Right) Integrated luminosity, for mass-independent (solid lines) and mass-dependent cuts (dashed), required to achieve a statistical significance of three (green) and five (red) standard deviations, as a function of the Higgs mass. The integrated luminosity required for excluding a Higgs boson signal at the 95% C.L. in a background-only experiment is also displayed (gray).

In order to more accurately quantify the degree of compatibility of the observed data with any of the two hypotheses, the confidence levels CL_b and CL_s are defined using the $-2 \ln Q$ probability density functions, *p.d.f.*, for both the background-only and the signal-plus-background hypotheses (details can be found in Refs. [53, 52]).

The presence of a signal can be inferred from the behavior of $1 - CL_b$ for the background-only hypothesis, which is the probability of observing in a sample of simulated background-only experiments a more signal-like value of $-2 \ln Q$. The observation of the value $1 - CL_b = 2.85 \times 10^{-7}$ indicates a 5σ excess in the data with respect to the background expectation. While CL_b quantifies the lack of compatibility of an excess of observed events with the background-only hypothesis, CL_s gives information about how compatible it is with an actual signal (Fig. 3.6).




Figure 3.6: Mean values for $1 - CL_b$ (left) and $1 - CL_s$ (right) as a function of the Higgs boson mass hypothesis, assuming existence of Higgs boson at 250 GeV mass and for an integrated luminosity of 10 fb^{-1} . The observation of the Higgs is just a little shy of the 5σ discovery (left). The mass points for which the curve $1 - CL_s$ is above 0.95 are excluded at 95% CL (right). The 1σ and 2σ bands on $1 - CL_b$ and $1 - CL_s$, originating from the Poisson statistical fluctuations of the number of background events in each bin of the discriminant distribution, are also shown.

3.1.3.2 Search using $m_{4\mu}$ -dependent cuts

One can take advantage of the fact that the Higgs boson resonance $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ is relatively narrow and use $m_{4\mu}$ -dependent cuts for its search. All details of such search strategy can be found in [42]. The analysis steps in this case would be as follows:

- First, events with 4 muons ($2\mu^+2\mu^-$) satisfying p_T , p , and $m_{\mu^+\mu^-}$ quality cuts as described in section 3.1.2.1 are selected. This ensures that muons are reliably reconstructed and removes a “contamination” originating from heavy quarkonia decays.
- Second, after reconstructing a four-muon invariant mass, the $m_{4\mu}$ -dependent cuts are applied. The cuts, being smooth functions of $m_{4\mu}$, are optimized in such a way that they maximize the significance of the Higgs signal excess *at all Higgs boson mass points*.
- And finally, the resulting $m_{4\mu}$ distribution is analyzed for the presence of a Higgs boson resonance. The search can be done using either the LLR significance S_L estimator built for the whole spectrum or the LLR S_{cL} estimator built for a single-

bin, or signal window (counting experiment). The direct comparison of the results can be found in [42].

To perform the desired $m_{4\mu}$ -dependent cut optimization, we used a recently developed program GARCON [55]. The counting experiment significance estimator S_{cL} is the natural tool for such optimization. The first half of the available Monte Carlo statistics was used for the cut optimization. The results for the 18 Higgs mass points were then fit to obtain smooth $m_{4\mu}$ -dependent cuts. It was found that, given the level of the expected dominant backgrounds ($t\bar{t}$, $Zb\bar{b}$, ZZ), there are only three critical discriminating cuts (details are given in Ref. [42]):

- Muon isolation cut, both tracker- and calorimeter-based, on the worst isolated muon, or equivalently one common cut on all four muons. This cut strongly suppresses $t\bar{t}$ and $Zb\bar{b}$ backgrounds. The cuts gets tighter and tighter as $m_{4\mu}$ gets smaller as the role of $Zb\bar{b}$ and $t\bar{t}$ increases (Fig. 3.2).
- p_T on the second lowest p_T muon, or equivalently one common cut on the three highest p_T muons. This cut helps to further suppress $Zb\bar{b}$ background to the level well below ZZ and reduces the ZZ background at high four-muon invariant masses. This cut becomes more stringent with increasing $m_{4\mu}$.
- And, of course, $m_{4\mu}$ window being used for scanning over the background. It roughly corresponds to the $\pm 2\sigma$ width, where σ is the Higgs boson peak width that includes the detector resolution and the Standard Model Higgs boson width.

The final results are obtained by applying these cuts to the second half of the available Monte Carlo statistics. The observed stability of the results ensures that the cut optimization did not pick peculiar phase space corners corresponding to statistical flukes. After applying the cuts, the $t\bar{t}$ and $Zb\bar{b}$ backgrounds are now suppressed well below the irreducible ZZ background.

Figure 3.5 shows, for different Higgs boson masses, the expected significance S_L at $\mathcal{L} = 30 \text{ fb}^{-1}$ (left) and the average integrated luminosities at which a “ 5σ -discovery”, “ 3σ -evidence”, and exclusion at 95%CL are expected (right). The gain in significance with respect to the flat, $m_{4\mu}$ -independent, cuts can be easily translated into probabilistic terms. For example, the Higgs boson with $m_H = 130 \text{ GeV}$ is right at the “ 5σ -discovery” threshold for an integrated luminosity $\mathcal{L} = 30 \text{ fb}^{-1}$. The difference in the average expected significance, 5.1 and 6.0, means in this case that the chances of observing significance in excess of 5 for $m_H = 130 \text{ GeV}$ at $\mathcal{L} = 30 \text{ fb}^{-1}$ are 55% for the flat cuts and 80% for the $m_{4\mu}$ -dependent cuts.

3.1.3.3 Systematic Errors

The analysis of the systematic errors can be sub-divided into two distinct stages. First, one needs to understand the level of uncertainties in predicting the level of background in vicinity of a particular $m_{4\mu}$ point being investigated for a possible event excess. Second, these uncertainties in the background need to be included in the evaluation of the significance of an excess of events, should it be observed.

Uncertainties in the signal are not very important for establishing an excess of events over the background. It is the uncertainties in the background that are of main concern. After applying the analysis cuts as described earlier, the ZZ production is the dominant irreducible background with all other processes giving much smaller contributions. This reduces the analysis of systematic errors to those of the $ZZ \rightarrow 4\mu$ process.

One can try to evaluate/guess the theoretical and detector performance related uncertain-

ties starting from the first principles. However, the credibility of the detector performance systematic errors estimated this way is always shaky, especially during the earlier stages of the detector operation when the changes in the system are frequent and hard to monitor and timely incorporate into the detector Monte Carlo simulation. Therefore, we developed methods evaluating various corrections, such as muon reconstruction efficiency, muon isolation cut efficiency, directly from data in order to minimize our relying on the Monte Carlo simulation, and, thus, significantly reducing the associated systematic errors. Also, throughout this analysis, we estimate the background around a particular $m_{4\mu}$ area in a reference to a *measured* control sample. Note that this completely eliminates uncertainties associated with measuring the luminosity and reduces the sensitivity to PDF and QCD-scales. For the control sample, we use either the inclusive $Z \rightarrow 2\mu$ process or sidebands of the $m_{4\mu}$ spectrum itself.

The main uncertainties can be grouped as follows:

1. Uncertainties associated with the background production rates, *i.e.* not directly related to CMS Detector performance itself:
 - ZZ: PDF and QCD scale uncertainties described in details in Ref. [38].
 - ZZ: NLO and NNLO contributions vs LO described in details in Ref. [50] plus some related issues are discussed in Ref. [33]. These possible uncertainties are not taken into account in the results shown below, for details see Ref. [42].
 - LHC luminosity: when we estimate the ZZ background events in the signal region via the measured number of events in the control samples, the luminosity uncertainties largely cancel out.
2. Uncertainties associated with the CMS detector performance (hardware/software) and our analysis-specific cuts:
 - ZZ: Trigger efficiency, being very close to 100% due to presence of four muons, does not have substantial systematic errors.
 - ZZ: Muon reconstruction efficiency is reconstructed directly from data [51]. The associated systematic error is less than 1% per muon. Using normalization to the measured $Z \rightarrow 2\mu$ process, this leaves us with 2% uncertainty per event for the $ZZ \rightarrow 4\mu$ background production.
 - ZZ: Muon isolation cut efficiency is also reconstructed directly from data [56] with about 2% uncertainty per event.
 - Higgs: $m_{4\mu}$ resolution is affected by muon p_T resolution. This almost does not affect the background distribution. In [42], we show that even making a mistake in the $m_{4\mu}$ distribution width by as much as 25% has only a tiny effect on evaluating a significance of an excess of events. The muon p_T resolution is fairly easy to measure from data using the measured J/ψ and Z peak widths with the precision much better than needed.
 - ZZ: $m_{4\mu}$ scale. The effect of these uncertainties on the number of background events in a signal window appears only on steep slopes of the $m_{4\mu}$ distribution. For the steepest part of the $m_{4\mu}$ distribution in the 180 GeV-200 GeV range, we obtain $\delta b/b \sim 0.1 \delta m_{4\mu}$, where $m_{4\mu}$ is in GeV and b is the number of background events. This implies that to be able to neglect this effect, one needs to know the momentum scale with precision of 0.1 GeV at

$p_T \sim 50$ GeV. This can be easily achieved with just a few hundreds of $Z \rightarrow 2\mu$ events.

Fig. 3.7 summarizes all systematic errors on the expected number of events in the $Z \rightarrow 4\mu$ background for the two methods: via referencing to the total measured $Z \rightarrow 2\mu$ cross section and via referencing to the event count in the sidebands of the $m_{4\mu}$ spectrum itself.

Significance with the background uncertainties included

For the Gaussian-like signal over relatively flat background, the S_L and S_{cL} estimators are strongly correlated, with the typical difference of 5%-10% [42]. This stems from the fact that the signal peak is very localized and the background is relatively flat. This allows us to study the effect of systematic errors on the evaluation of significance *at the time of measurements* using the counting experiment approach, for which everything can be done analytically. All details on the method we use can be found in Ref. [42]. The method allows to account for the theoretical and instrumental systematic errors as well as for statistical errors when a control sample with a limited event count is used.

The final result of these studies is presented in Figure 3.8. Starting from an integrated luminosity at which the statistical significance of a Higgs boson observation would be equal to 5 (if the level of background without any errors was known), the figure shows how this significance must be de-rated due to the systematic errors at the time of the measurements as described in the previous sub-section. The effect of systematic errors at lower or higher luminosities is not as important: at lower luminosity the significance is not sufficient to make serious claims, anyway; while after surpassing the significance of 5, the existence of the Higgs boson can be considered established and the gears must be switched to measuring its parameters.

The two curves with filled and empty circles show the difference of the two methods for evaluating the background in the signal region: via normalization to the measured $Z \rightarrow 2\mu$ cross section, and via normalization to the event count in sidebands (100 GeV to 700 GeV, excluding the signal region). The effect of systematic errors at lower luminosities becomes smaller for the former method and quickly diverges for the latter. As the luminosity increases, the trends obviously reverse. Around the threshold of $S = 5$, the difference between the two methods is not very dramatic; the true benefit of using two approaches to estimating background from data is in their complementarity.

Local significance and overall statistical fluctuation probability

In a search for a relatively narrow 4μ invariant mass peak over a broad background spectrum, one must take into account that the probability of observing a background fluctuation giving an excess of events consistent with a Higgs hypothesis of *some particular mass* might be considerably higher than the local significance calculated for a given mass might imply. This over-estimation of significance strongly depends on how the analysis is set and what constraints/priors on the “phase space” of parameters are used. *E.g.*, in a search specifically tailored for the Standard Model Higgs, the only free parameter is the Higgs boson mass; its width, production cross section, and decay branching ratios are being dependent on the mass. To make the search even more constrained, one can use a prior on the Higgs mass as it comes out from the precision electroweak measurements. And so on. A specific case study showing the potential scope of the effect, which may be comparable or even larger than the effect of the systematic errors discussed above, is given in Appendix A.1.




Figure 3.7: Uncertainties in the count of the $ZZ \rightarrow 4\mu$ background events in the signal region window at different $m_{4\mu}$. The window size is $\pm 2\sigma$ of the expected experimental Higgs resonance width. (Left) The background event count in the signal region is derived from the measured number of $Z \rightarrow 2\mu$ events. (Right) The background event count in the signal region, b , is calculated from the number of $ZZ \rightarrow 4\mu$ events B in the range 100 GeV-700 GeV (excluding the signal region window), *i.e.* $b = \rho \cdot B$.

3.1.4 Measurement of the Higgs Boson Properties at $\mathcal{L} = 30 \text{ fb}^{-1}$

The capabilities of the CMS detector to measure the mass, cross section and width of the Higgs boson are determined for an integrated luminosity of 30 fb^{-1} [57, ?]. These parameters are measured using a binned maximum likelihood fit to the reconstructed four-muon invariant mass, which includes the signal and background contributions after all the selection cuts have been applied (Figure 3.4 (right)). The ‘observed’ distribution, f_{sb} , is expressed in terms of the signal, p_s , and background, p_b , probability density functions (*p.d.f.*) as:

$$f_{sb}(m_{4\mu}; m_{fit}, \Gamma, N_s, N_b) = N_s \cdot p_s(m_{4\mu}; m_{fit}, \Gamma) + N_b \cdot p_b(m_{4\mu})$$

N_s is the number of signal events, N_b the number of background events, m_{fit} the position of the mass peak and Γ the intrinsic width of the Higgs boson. The signal *p.d.f.* is the sum of two contributions: a convolution of a Breit-Wigner signal shape with a Gaussian distribution that accounts for detector resolution, p_{core} , and a function that reproduces the radiative tail due to internal bremsstrahlung, p_{tail} :

$$p_s = \beta \cdot p_{core}(m_{4\mu}; m_{fit}, \Gamma, \sigma) + (1 - \beta) \cdot p_{tail}(m_{4\mu}; m_{fit}, \tau)$$

where $1 - \beta$ is the fraction of signal events in the radiative tail. The tail shape is parametrized *ad hoc* as

$$p_{tail} = \frac{(m_{4\mu} - m_{fit})^2}{2\tau^3} \exp\left(\frac{m_{4\mu} - m_{fit}}{\tau}\right)$$

if $m_{4\mu} < m_{fit}$ and is zero otherwise [58]. Figure 3.9 (left) illustrates the different contributions to f_{sb} . The p_s function is fitted to the signal-only distributions to obtain the parameters of the radiative tail, which remain fixed in the fit to the signal plus background spectra.




Figure 3.8: Effect of including systematic errors into evaluation of significance at the time of measurements. The reference luminosities, dependent on the Higgs boson mass, are chosen to correspond to an observation of significance $S = 5$, if not for the presence of systematic errors. Solid circles show de-rating of significance for the case of systematic errors when the background is evaluated from the measured $Z \rightarrow 2\mu$ cross section. Open circles show the effect for the case when the background in signal region is normalized to the sidebands.

For Higgs boson masses below 190 GeV, the intrinsic width is negligible compared to the mass spread introduced by the experimental resolution and the signal is thus approximated by a Gaussian shape. For masses above 400 GeV, the natural width of the Higgs is much larger than the experimental resolution, hence the description using a pure Breit-Wigner function yields similar parameters as those obtained from the convolution.

The detector resolution is extracted from the $m_{4\mu}$ distribution of ZZ events with a four-muon mass above $2m_Z$, for which the kinematics is similar to that of the signal. For masses below $2m_Z$, the intrinsic Higgs boson width is negligible, therefore the resolution is measured directly from the width of the $m_{4\mu}$ distribution. This width has been found to be consistent with the extrapolation of the resolution determined using ZZ events.

The background *p.d.f.*, p_b , is approximated by either a polynomial or an exponential function, depending on the mass region under study. The parameters are determined performing a binned maximum likelihood fit to the background sample. The parameters defining the shape of the background are fixed in the global fit to signal plus background, but not its normalization.

The values of the parameters, together with their errors, are obtained directly from the fit. The result of the fit to the signal plus background distribution is shown in Figure 3.9 (right) for a Higgs boson signal of $m_H = 250$ GeV. Figure 3.10 (left) depicts the relative shift of the fitted Higgs boson mass with respect to the true mass, together with its statistical error. These values are compatible with zero in the full range of masses, which means that the true mass is accurately recovered after applying the fitting method to the reconstructed sample. The evolution of the relative error as a function of the true mass is displayed in Figure 3.10 (right), showing that the mass can be measured with precisions from 0.1% to 5.4%. The increase in




Figure 3.9: (Left) Example of the shapes of the different contributions to f_{sb} . (Right) Data-like distribution expected for a Higgs boson signals of $m_H = 250 \text{ GeV}$, for an integrated luminosity of 30 fb^{-1} , together with the result of the fit (solid line) and the expected background (shaded area). This pseudo-experiment is selected randomly.

this error around 170 GeV is due to the smaller signal statistics caused by the suppression of the $H \rightarrow ZZ^{(*)}$ decay at this mass. The increasing uncertainty at higher masses is due to the smaller production cross sections, the larger intrinsic width of the Higgs boson and, to a lesser extent, the worse resolution for high p_T muons.

The number of signal and background events is obtained from the fit. The relative error in the cross-section measurement is determined from the number of signal events (N_s) and its statistical uncertainty (ΔN_s) as $\Delta N_s / N_s$, shown in Figure 3.11 (left) as function of the Higgs boson mass. The contribution of the background is properly taken into account, as its normalization is a free parameter in the fit. The cross section can be determined with a precision between 20% and 45%, except for masses below 130 GeV , where the statistics is low.

The measured width, together with its statistical error, is presented in Figure 3.11 (right) as function of the true mass. The width can be determined with an error between 35% and 45% above 190 GeV . Below this mass there is no sensitivity to the Higgs boson width and upper limits at 95% confidence level (C.L.) are set. For the sake of comparison, the width obtained by fitting only a Gaussian for masses below 200 GeV and only a Breit-Wigner for masses above 200 GeV is also shown, together with the statistical uncertainty. The Breit-Wigner-only fits do not take into account the detector resolution, and therefore the intrinsic theoretical values are not recovered.

The measurement of the parameters is affected by systematic uncertainties in the muon momentum resolution (determined from data), in the muon reconstruction efficiency (around 2%) and those associated to the selection cuts (close to 1%) [52]. These systematic uncertainties are mostly uncorrelated. The impact in the measured mass and width is small. The cross-section measurement is also affected by the uncertainty in the luminosity determination, which is around 3% (Figure 3.11 (left)).




Figure 3.10: (Left) Relative shift of the fitted value of the Higgs boson mass with respect to the input m_H value, as function of m_H . The shaded area is the error in the determination of the peak value from the fit, also shown as function of the Higgs boson mass (Right). The dots correspond to the result of the convolution and the triangles to the Gaussian approximation.

The results obtained for Higgs boson masses around 170 GeV and above 500 GeV, for which the expected number of events is somewhat low for $\mathcal{L} = 30 \text{ fb}^{-1}$, have to be taken as representative results for the typical expected distributions. The higher errors of the parameters for those m_H values are consistent with statistics. For extending the measurement of the Higgs boson parameters to smaller masses or to lower luminosities, it should be more appropriate to extract the parameters from a large set of randomly chosen four-muon mass distributions with the correct statistics.

3.1.5 Conclusions

Discovery of the Standard Model Higgs boson and measurement of its mass, production cross section and width in the “golden” decay mode $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ were analyzed in the context of the CMS Detector. The explored range of Higgs boson masses was 115 GeV–600 GeV. The Monte Carlo samples were normalized to represent the NLO cross sections, including $m_{4\mu}$ -dependent K-factors. To simulate the detector response and reconstruct physics objects, the full CMS Detector simulation and reconstruction software was used. The Higgs boson discovery potential was explored for different analysis variations, including the use of $m_{4\mu}$ -dependent and flat cuts, log-likelihood ratio based on the full $m_{4\mu}$ spectrum and a straightforward counting experiment approach. A full treatment of the most important theoretical and instrumental systematic errors and their effect on evaluation of significance of the Higgs boson observation as well as measuring its parameters were presented. To minimize systematic errors, a number of methods of reconstructing the necessary corrections directly from data were developed.

It was shown that at $\sim 2 \text{ fb}^{-1}$ of integrated luminosity, CMS would be able to start excluding the Standard Model Higgs boson at 95% CL for m_H in vicinity of 200 GeV. By the time CMS reaches $\sim 30 \text{ fb}^{-1}$, it would exclude the Standard Model Higgs boson in its four-muon decay




Figure 3.11: (Left) Relative error in the cross-section measurement, $\Delta N_s/N_s$, as a function of the m_H . ΔN_s is the statistical error of N_s obtained from the fit. The dots correspond to the result of the convolution and the triangles to the Gaussian approximation. The dashed line indicates the impact of the systematic uncertainties. (Right) Measured Higgs boson width (squares), its statistical error (green band) and the theoretical calculation of Γ_H (dashed line). Upper limits to the width at 95% C. L. are shown (red line) for $m_H < 190 \text{ GeV}$. The result of Gaussian (triangles) and Breit-Wigner (dots) fits are also shown for comparison.

mode in the mass range $m_H = 120 \text{ GeV}-600 \text{ GeV}$, if indeed it does not exist.

The discoveries at the level of “ 5σ ” local significance could be already possible at $\sim 10 \text{ fb}^{-1}$ for m_H in the range $140 \text{ GeV}-150 \text{ GeV}$ and $190 \text{ GeV}-400 \text{ GeV}$. By the time $\sim 30 \text{ fb}^{-1}$ are collected, the discovery range would open up to $130 \text{ GeV}-160 \text{ GeV}$ and $180 \text{ GeV}-500 \text{ GeV}$. An observation of the Higgs boson with the mass $m_H \sim 170 \text{ GeV}$ or $\sim 600 \text{ GeV}$ in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ decay channel would require an integrated luminosity of the order of 100 fb^{-1} .

At the integrated luminosity of $\sim 30 \text{ fb}^{-1}$, the Higgs boson mass could be measured with a precision between 0.1 % and 5.4 %, depending on its mass. The intrinsic width could be measured only for the Higgs boson heavier than 190 GeV , with a precision $\sim 35\%$. For lower masses, the Higgs boson width becomes much smaller than the detector resolution and only upper limits of the order of a few GeV could be set. The production cross section would be determined with a precision $\sim 30\%$.

Benchmark Channel: $H \rightarrow WW^{(*)} \rightarrow 2 \text{ muons}$

3.2.1 Introduction

Previous studies [59] [60] demonstrated the relevance of the $H \rightarrow WW^{(*)} \rightarrow 2l2\nu$ channel for the Higgs discovery with an integrated luminosity of less than 5 fb^{-1} . The physics study was performed on the data produced at the end of the full simulation, trigger and off-line detector

reconstruction chain, including realistic assumptions for the sub-detectors misalignments. The goal of this study is to provide the discovery significance as a function of the Higgs mass using the most up-to-date detector simulation and reconstruction code, considering all the relevant background contributions and providing an as much as possible complete estimation of the systematic errors. The muon reconstruction has an average efficiency in the detector geometrical acceptance ($\eta < 2.4$) of 95 – 99% for the transverse momentum ranging from 5 GeV up to $P_T = 1$ TeV, as extensively discussed in [61], while the percentage of mis-assigned charge for muons with $P_T = 100$ GeV is less than 0.1%.

In Section 2 the signal and the most important background processes are described, reporting the relevant production cross sections predicted by the theoretical computations at the leading and next-to-leading orders. The event selection and cut optimization is discussed in Section 3. Section 4 addresses the problem of evaluating the systematic uncertainties and the background normalization, reporting the final results in terms of expected significance of the signal as a function of the mass of the Higgs boson.

3.2.2 Physics processes

3.2.2.1 Signal processes

The signal was studied in the range between 130 to 180 GeV using 7 samples of datasets (Tab. 3.1). The generation was done using the PYTHIA program [62], considering the most relevant signal sources:

$$g g \rightarrow H \rightarrow WW^{(*)} \rightarrow 2\mu 2\nu \quad (3.1)$$

$$q \bar{q} \rightarrow VV q' \bar{q}' \rightarrow H q' \bar{q}' ; H \rightarrow WW^{(*)} \rightarrow 2\mu 2\nu \quad (3.2)$$

The detector simulation was performed using the CMSIM 133 simulation program [13], based on the GEANT3 package. In the “digitization” process, i.e. the simulation of the detector response producing the raw data, performed using the version ORCA 871 of the CMS reconstruction program[15], the effect of the event pile up expected at the machine luminosity $2 \times 10^{33} \text{cm}^{-2} \text{sec}^{-1}$ was included.

3.2.2.2 Background processes

The dominant background giving the largest contribution at the end of the complete selection chain, is the irreducible one from the continuum production of W pairs decaying into muons and neutrinos. Other significant or critical sources of backgrounds are the production of top quarks and the Drell-Yan muon pairs. The most important backgrounds are thus the processes:

$$q \bar{q} \rightarrow W^+ W^- \rightarrow 2\mu 2\nu \quad (3.3)$$

$$g g \rightarrow t \bar{t} \rightarrow 2\mu 2\nu \quad (3.4)$$

$$q \bar{q} \rightarrow \gamma^*, Z \rightarrow 2\mu \quad (3.5)$$

Further contributions from $b\bar{b}$, $ggWW$, WZ , ZZ , and Wt production processes were also considered. A part from Wt and $gg \rightarrow WW$, all the processes but the latter have been generated with PYTHIA. For the former process, TOPREX Monte Carlo [35] has been used which

correctly takes into account the top mass and the spin correlations throughout the decay chain. The latter dataset has been simulated starting from a Monte Carlo sample produced by N. Kauer et al. [63]. The full list of dataset samples used for the background study is given in Tab. 3.2

3.2.2.3 Cross sections at NLO

All the processes considered in this study have been simulated with Born-level accuracy. In order to approximate the NLO predictions for the signal and the W -pair background, phase space depended reweighting (k -)factors has been applied [64]. These factors have been obtained by matching respectively the p_T distribution of the Higgs and of the W^+W^- system provided by PYTHIA to the one predicted by MC@NLO [65]¹. The $k(p_T)$ factors used for each p_T intervals are given in Appendix of [66]. The absolute cross sections for Higgs production through gluon-gluon fusion and vector boson fusion have been calculated by M.Spira [8] and are listed in Tab. 3.1.

Higgs mass (GeV)	$\sigma^{NLO} \times BR(2l)$ Gluon Fusion (pb)	$\sigma^{NLO} \times BR(2l)$ VBF (pb)	$\sigma^{NLO} \times BR(2l)$ num. of events
130	0.94	0.12	20000
140	1.39	0.19	20000
150	1.73	0.25	17000
160	2.03	0.31	44000
165	2.04	0.32	49000
170	1.95	0.31	40000
180	1.71	0.28	20000

Table 3.1: The cross section at the next-to-leading order for Higgs production through gluon fusion and vector boson fusion (VBF) processes and the number of generated events are reported.

No reweighting has been applied to the other processes, whose total cross sections have been simply rescaled accordingly the NLO calculation performed J.Campbell, F.Tramontano, K.Ellis and F.Maltoni using the MCFM Monte Carlo program [?] [67] [46]. These cross sections are reported in Tab. 3.2.

3.2.3 Event selection

The signal selection requires the identification of two high p_T isolated muons. The background reduction is obtained applying suitable kinematic cuts to the reconstructed muons, a veto on the presence of central jets and a high missing E_T (MET) in the event. As discussed in the following sections, separate optimizations were performed independently on the muon isolation variables, jet and missing energy thresholds and on the muons kinematical variables.

¹For the signal, only the Higgs production through the gluon-gluon mechanism has been reweighted with $k(p_T)$ factors accordingly to NLO description

Channel	$\sigma^{NLO} \times BR(pb)$	num. of ev.
$qq \rightarrow WW \rightarrow 2l$	11.7	164000
$t\bar{t}$	840	548000
$gg \rightarrow WW \rightarrow 2l$	0.54 (LO)	50000
γ^*, Z	145000	2700000

Channel	$\sigma^{NLO} \times BR(pb)$	num. of ev.
$b\bar{b} \rightarrow 2\mu$	710 (LO)(*)	640000
$ZW \rightarrow 3l$	1.63	72000
$tWb \rightarrow 2l$ (toprex)	3.4	191000
$ZZ \rightarrow 2l$	1.52	99000

Table 3.2: The cross section at the next-to-leading order for the background processes. The $gg \rightarrow WW$ process is generated using a matrix element program linked to PYTHIA for the showering [63]. This process is only known at LO. (*) For $b\bar{b} \rightarrow 2\mu$ the preselection $p_T > 20, 10 \text{ GeV}/c$ was applied.

3.2.4 The trigger selection

Events passing the global L1 trigger must be reduced with a more restricted trigger requirement to limit the recorded event rate. Two trigger streams were considered in this analysis:

1. the HLT double muon stream;
2. the OR of the HLT single muon and double muon stream.

Before any selection the single or double muon HLT trigger efficiency is 92 %, while the double muon HLT trigger efficiency is 80 % [68]. After the off-line cuts for the Higgs selection, which will be described in detail in the following section, the overall efficiency of the first stream relative to the second one is found to be $(97 \pm 1)\%$, for $m_H = 165 \text{ GeV}$. In the following, the trigger selection used was the HLT double muon stream, for which the trigger rate is predicted to be a factor 7 smaller than the single muon one [68].

3.2.4.1 The muon identification and isolation

A first event selection based on the identification of two prompt muons required :

- L1 and HLT di-muon trigger bits found;
- two oppositely charged muons reconstructed by the Global Muon reconstructor algorithm developed in ORCA, as described in [61].

The first requirement assures the events to be found in the CMS di-muon data stream, which currently foresees a symmetric threshold of $7 \text{ GeV}/c$ on the p_T of both muons as reconstructed by the High Level Trigger algorithm, for operations at a machine luminosity of $2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$; in addition, at least one of the muons must fulfill the HLT isolation criteria [68]. As discussed in ref.[68], the trigger rate for this datastream is predicted to be about 4 Hz.

At the off-line reconstruction and selection stage, two cones were considered for the isolation around each reconstructed muon tracks. The $\sum P_T$ summed over all the charged track candidates found in the Tracker detector was accounted inside the first cone. The $\sum Et$ over

the energy deposits in the ECAL and HCAL towers was accounted in the second cone. The size of a cone around a muon track is defined as $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2}$. A muon is considered to be isolated if the $\sum Pt$ ($\sum Et$) inside the considered cones of size $\Delta R_{Tracker}$ (ΔR_{Calo}) is below the threshold P_T (max) (E_T (max)). An optimization study was performed to find the four parameters:

$$(1) \Delta R_{Tracker} \quad (2) P_T \text{ (max)} \quad (3) \Delta R_{Calo} \quad (4) E_T \text{ (max)}$$

searching for the highest signal over background ratio. The optimization was performed using the signal dataset with $m_H = 165 \text{ GeV}$ (Tab. ??) and the $b\bar{b}$ background dataset, which is the most sensitive to the isolation cut. At this first stage of the selection, the background reduction was not requested to be very large, thus keeping the signal reduction relatively small; for each combination of the cones:

$$\Delta R_{Tracker} = 0.25, 0.3, 0.35, 0.4 \quad \Delta R_{Calo} = 0.25, 0.3, 0.35, 0.4 \quad (3.6)$$

the cut efficiency of 85% for the signal was requested. With two free parameters, E_T (max) and P_T (max), several solutions are possible. A reasonable choice is to give the same weight to the Tracker and Calorimeter isolation cuts. The mean and the r.m.s. values of the p_T and energy deposition for the signal dataset within different cones are reported in [69]. For each set of isolation cones ($\Delta R_{Tracker}, \Delta R_{Calo}$) the E_T and P_T thresholds were chosen as follows:

$$E_T^{thresh} = \langle E_T \rangle + x \cdot \sigma(E_T) \quad (3.7)$$

$$P_T^{thresh} = \langle P_T \rangle + x \cdot \sigma(P_T) \quad (3.8)$$

where the parameter x was set to the value giving the required 85% efficiency for the signal.

Figure 3.12 shows the resulting background selection efficiency.




Figure 3.12: $b\bar{b}$ background efficiencies for the 16 combinations of cones considered for the muon isolation selection cut.

The best selection is obtained with:

$$\Delta R_{Tracker} = 0.25 \quad P_T < 2.0 \text{ GeV} \quad \Delta R_{Calo} = 0.3 \quad E_T < 4.7 \text{ GeV} \quad (3.9)$$

corresponding to $x = 1.8$ for the energy deposition and P_T cut. The isolation cuts used in the analysis were:

$$\Delta R_{Tracker} = 0.25 \quad P_T < 2.0 \text{ GeV} \quad \Delta R_{Calo} = 0.3 \quad E_T < 5.0 \text{ GeV} \quad (3.10)$$

3.2.5 JET reconstruction and the JET veto

The reconstruction of jets is needed to obtain a strong $t\bar{t}$ background reduction by applying a JET veto. The JET reconstruction algorithms can use the raw energy sum of the ECAL and HCAL towers, either with a fixed energy threshold or with eta dependent thresholds. The eta dependent threshold did not show to improve the $t\bar{t}$ background rejection with respect to a fixed combined E_T and E thresholds [66]. The " γ energy calibrated Jets" were also considered, but the calibration tool available up to end of 2005 showed a distortion for the E_T correlation between "generated" and reconstructed jets in the low energy region ("generated" jets are those built using the tracks at generator level, i.e. without the detector simulation). The jets reconstructed from raw energies with fixed E_T and E thresholds were finally chosen to be used for the JET veto. A strong E_T cut helps in the background reduction. However, below $E_T = 25 \text{ GeV}$ the fraction of jets matching with a generated jet starts to decrease, because of ghost jet candidates mainly due to pileup effects. The matching was defined within a cone around the reconstructed jet candidate $\Delta R_{rec-gen jet} < 0.3$. In order to reduce the number of fake jets, a quality parameter was introduced:

$$\alpha = \sum_{\text{selected tracks}} P_T/E_T(\text{jet}) \quad (3.11)$$

where the selected tracks are those inside the jet ($\Delta R_{trk-jet} < 0.5$) with more than 5 associated hits, pointing to the primary interaction vertex ($|z_{trk} - z_{vtx}| < 0.4 \text{ cm}$). The mean value of α is 0.66 (two third of the jet energy on average is due to charged particles). A reconstructed jet candidate with E_T in the low energy region ($< 20 \text{ GeV}$) was considered only if $\alpha > 0.2$. It has been shown [66] that this selection significantly reduces the number of fake jets (the fraction of matched jets being greater than 90% for $E_T > 15 \text{ GeV}$) with negligible loss of reconstruction efficiency for true jets. Different JET reconstruction algorithms were tested. The best signal ($m_H = 165 \text{ GeV}$) / background ($t\bar{t}$) ratio was obtained using an iterative cone algorithm [70] with a cone size $R=0.5$ and calorimeter towers having raw energies $E_T^{tower} > 0.5 \text{ GeV}$ and $E^{tower} > 0.8$. To summarize, the JET veto is applied if:

$$E_T > 15 \text{ GeV} \quad |\eta_{jet}| < 2.5 \quad (3.12)$$

and the α cut is required in the jet energy range $15 \text{ GeV} < E_T < 20 \text{ GeV}$.

3.2.6 Missing energy reconstruction and the MET cut

The transverse missing energy is reconstructed with the sum of the ECAL and HCAL tower raw energies, corrected for the muons energy contribution. The most sensible background to the MET cut is the di-muon production from Drell-Yan (DY) process. The right plot in Fig. 3.13 shows the MET distributions for DY events having a reconstructed di-muon invariant mass inside the Z mass region (shown by the black area in the left plot), and for signal events with $m_{Higgs} = 165 \text{ GeV}$. The signal and background distribution were normalized to an integrated luminosity $L = 10 \text{ fb}^{-1}$.




Figure 3.13: Reconstructed di-muon invariant mass for Drell-Yan events selected inside the Z mass region(left, black area); MET distributions for the selected Drell-Yan events and for signal events scaled at the integrated luminosity $L = 10fb^{-1}$ (right).

A MET threshold of $47 GeV$ is 4σ over the mean value for the background and 1.5σ under the mean value for the signal. Drell-Yan events are thus strongly suppressed by applying a MET threshold. The cut used in this analysis was $MET > 50 GeV$.

3.2.6.1 The kinematic cuts

The kinematic of the two muons is different for signal and background:

- signal events from gluon-gluon scattering are more central than the W^+W^- background from $q\bar{q}$ scattering, thus resulting in a slightly more central rapidity distribution for the decay muons;
- due to the scalar nature of the Higgs boson and of the V-A structure of the weak interaction, for Higgs masses close to $2 \cdot M_W$, the W^+W^- spin correlation plays in favor of small opening angles between the two muons;
- signal events have a lepton P_T spectra peak close to $M_W/2$;
- DY background has a two muons invariant mass peak at M_Z .

In addition, the muons from b quarks (as in the case of the $b\bar{b}$ background and eventually from $t\bar{t}$) have large impact parameters. The following cuts were applied before the optimization of the kinematical cuts:

1. $|\eta(\mu_1)|, |\eta(\mu_2)| < 2.0$ (pseudorapidity of the two muons);
2. $IP(\mu_1), IP(\mu_2) < 3\sigma$ (impact parameter of the two muons);
3. $P_T(\mu_{max}) < 55 GeV$ (transverse momentum of the two muons);
4. $m_{\mu_1\mu_2} > 12 GeV$ (invariant mass of the two muons);
5. $\Delta\phi_{\mu_1\mu_2} < 0.8$ (opening angle between the two muons).

The cut 1. is useful for the WW background reduction, as well as cuts 3 and 5. The cut 2. reduces the $b\bar{b}$ events, while cut 4. rejects potential background from b-resonances. After the requirement of the muon isolation described in Section 2, the overall signal efficiency for cuts 1 to 4 is about 90%. The distribution of the variable $\Delta\phi_{\mu_1\mu_2}$ will be used to search for the Higgs signal.

The optimization study was performed by varying the following cuts:

$$P_T(\mu_{max}) > 25, 30, 35, 40 \text{ GeV} \quad P_T(\mu_{min}) > 15, 20, 25, 30 \text{ GeV} \quad m_{\mu_1\mu_2} < 35, 40, 45, 50, 55, 60 \text{ GeV} \quad (3.13)$$

to find the set of cuts giving the best significance. The estimator S_{cP} was used, which gives the significance using the Poisson distribution [71]. The input of the estimator are the number of signal and background events, the statistical uncertainties and the theoretical systematics in the background. The optimization was performed using as before the signal dataset with $M_H = 165 \text{ GeV}$, and using all the background contributions, properly normalized considering their production cross sections.

The optimization result could depend on the statistics of the event data samples and on the estimated systematic errors. We searched for the maximum significance in four different conditions:

$$L = 1 \text{ fb}^{-1} \quad L = 2 \text{ fb}^{-1} \quad \text{syst. err.} = 10\% \quad \text{syst. err.} = 15\% \quad (3.14)$$

Fig. 3.14 shows, as an example, the significance result expected as a function of $p_T(\mu_{max})$ and $p_T(\mu_{min})$ cuts for two different values of the di-muon invariant mass cut, for the case of an integrated luminosity $L = 1 \text{ fb}^{-1}$ and an overall 10% systematic error.




Figure 3.14: Significance as a function of P_T cuts for $m_{\mu_1\mu_2} < 40 \text{ GeV}$ (left) and for $m_{\mu_1\mu_2} < 50 \text{ GeV}$ (right) with $L = 1 \text{ fb}^{-1}$ and $\text{syst. err.} = 10\%$

The following cuts:

$$P_T(\mu_{max}) > 35 \text{ GeV} \quad P_T(\mu_{min}) > 25 \text{ GeV} \quad m_{\mu_1\mu_2} < 50 \text{ GeV} \quad (3.15)$$

give the maximum significance (about 3.0 for $L = 1 \text{ fb}^{-1}$ and an assumed $\text{syst. err.} = 10\%$) in all the four conditions.

3.2.7 The selection results

The optimized selection cuts discussed above were applied to the background and signal samples. The list of cuts is described in Tab. 3.3. The expected number of events for a luminosity of 1 fb^{-1} are given in Tab. 3.4 for the signals and the backgrounds.

1	L1+HLT di-muon	6	$MET > 50 \text{ GeV}$
2	2μ opposite charge	7	$35 \text{ GeV} < P_T(\mu_{max}) < 55 \text{ GeV}$
3	Isolation	8	$25 \text{ GeV} < P_T(\mu_{min})$
4	$\eta < 2.0$ $IP < 3\sigma$	9	$m_{\mu_1\mu_2} < 50 \text{ GeV}$
5	Jet Veto	10	$\Delta\phi_{\mu_1\mu_2} < 0.8$

Table 3.3: The list of cuts applied to the signal and background samples

	L1+HLT di-muon	All cuts	ϵ_{tot}
$m_H = 130 \text{ GeV}$	112	0.68 ± 0.19	$(0.07 \pm 0.02)\%$
$m_H = 140 \text{ GeV}$	162	1.7 ± 0.4	$(0.12 \pm 0.03)\%$
$m_H = 150 \text{ GeV}$	228	5.3 ± 0.8	$(0.26 \pm 0.04)\%$
$m_H = 160 \text{ GeV}$	256	12.6 ± 0.7	$(0.58 \pm 0.04)\%$
$m_H = 165 \text{ GeV}$	264	14.3 ± 0.8	$(0.64 \pm 0.04)\%$
$m_H = 170 \text{ GeV}$	259	11.0 ± 0.7	$(0.53 \pm 0.03)\%$
$m_H = 180 \text{ GeV}$	233	5.9 ± 0.8	$(0.30 \pm 0.04)\%$
$qq \rightarrow WW$	1040	4.1 ± 0.5	$(0.036 \pm 0.005)\%$
$t\bar{t} \rightarrow 2\mu 2\nu$	17007	2.6 ± 0.3	$(0.012 \pm 0.001)\%$
$gg \rightarrow WW$	58	1.0 ± 0.1	$(0.18 \pm 0.02)\%$
$\gamma^*, Z \rightarrow 2\mu$	720653	0.3 ± 0.3	$(4 \pm 4)10^{-5}\%$
$b\bar{b} \rightarrow 2\mu 2\nu$	69374	0	0%
Wt	615	0.57 ± 0.10	$(0.017 \pm 0.003)\%$
ZZ	218	0.18 ± 0.05	$(0.012 \pm 0.003)\%$
ZW	384	0.13 ± 0.05	$(0.008 \pm 0.003)\%$

Table 3.4: The expected number of events for a luminosity of 1 fb^{-1} for the signal with Higgs masses between 130 and 180 GeV and for the backgrounds.

Figure 3.15 shows the distributions of the MET, $P_T(\mu_{max})$, $P_T(\mu_{min})$ and $m_{\mu_1\mu_2}$ variables for the signal and the three most important backgrounds after the jet-veto and the following selection cuts applied in the order reported in the Tab. 3.3.

Figure 3.16 shows the final distribution obtained for the azimuth angle difference between the muons, expected for an integrated luminosity $L = 10 \text{ fb}^{-1}$ and for the Higgs signal of mass $m_H = 165 \text{ GeV}/c$.

As stated above, all the numbers at the various selection steps refer to the analysis applied to the HLT di-muon stream. For comparison the event numbers after all the selection cuts were also studied for the case in which the analysis were performed on the data including the single muon trigger data stream. The inclusion of this datastream, which is foreseen to have a rate about 7 times larger than the di-muon stream [68], would result in a $(3 \pm 1)\%$ increase of the overall signal selection efficiency. The Higgs search with mass appreciably different than 165 GeV can take advantage from a dedicated cut optimization like reported




Figure 3.15: Distributions of the missing energy, transverse momentum and invariant mass for a luminosity of 10 fb^{-1} following the cut list order.

in [69].

3.2.8 Background estimation and systematics

The precise understanding of the backgrounds is the most critical issue concerning this Higgs discovery channel. The direct use of the Monte Carlo predictions, i.e. $N_{bkg,MC} = \sigma_{bkg,MC} \cdot \epsilon_{ff}$, leads to introduce high systematic uncertainties due either to theoretical calculation and to experimental systematics. The most reliable approach to address this problem is to measure the different sources of background directly from the data. The commonly used method to extrapolate the background contribution directly from the data consists of selecting a signal-free phase space region (control region) where a given background process is enhanced. The normalization from data for the two most relevant background, i.e. $t\bar{t}$ and WW has been addressed. For both backgrounds, a dedicated control region was defined. The number of background events in the signal region can then be estimated through:

$$N_{signal.reg} = \frac{N_{MonteCarlo}^{signal.reg}}{N_{MonteCarlo}^{control.reg}} N_{control.reg} \quad (3.16)$$




Figure 3.16: Distribution of the angle between the two muons for a luminosity of 10 fb^{-1} at the end of the selection.

where $N_{\text{signal.reg}}^{\text{MonteCarlo}}$ and $N_{\text{control.reg}}^{\text{MonteCarlo}}$ are the numbers of events predicted by the Monte Carlo simulation in the signal and control region. The error on the ratio $N_{\text{signal.reg}}^{\text{MonteCarlo}}/N_{\text{control.reg}}^{\text{MonteCarlo}}$ accounts for a theoretical contribution (scale variation, PDF uncertainty) and detector systematics effects. The precision with which the number of $N_{\text{signal.reg}}$ can be predicted depends also on the statistical error on $N_{\text{control.reg}}$.

3.2.9 $t\bar{t}$ background normalization

Since the presence of two b-tagged jets is a striking evidence for $t\bar{t}$ events, the most natural control region for this process is then defined by applying the same selection cuts as for the signal region but the jet veto, with the additional request of two b-tagged jets in the detector acceptance². The $t\bar{t}$ evaluation from the data for the $H \rightarrow WW^{(*)}$ channel has been studied in Ref. [72] to which we refer for further details. In this study a jet is tagged as a b-jet if its measured E_T is greater than 20 GeV and if there are at least two tracks belonging to the jet (i.e. within a cone of 0.5 around the jet axis) whose σ_{IP} is higher than 2. With such settings the double b-tagging efficiency for $t\bar{t}$ events is $\mathcal{O}(30\%)$. The mistagging rate has been calculated from the ratio between the number of b-tagged jets and the total number of jet with $E_T > 20 \text{ GeV}$ in the fully simulated DY sample and it resulted to be $\mathcal{O}(3\%)$.

In the following we consider the background processes in the $t\bar{t}$ control region. For 1 fb^{-1} the number of $t\bar{t}$ events in the control region just defined is foreseen to be 17, whereas the contribution from the signal and Wt is completely negligible (in both cases smaller than 0.1 events).

Not all the processes with $2\mu + 2b + E_t^{\text{miss}}$ as final state have been fully simulated for this analysis, nevertheless general considerations and fast Monte Carlo generator level cross checks lead to exclude other relevant sources of backgrounds, as briefly outlined in the following. The more natural concurrent process is the not resonant $W^+W^- \rightarrow 2\mu + b\bar{b}$ which is anyway suppressed with respect to $t\bar{t}$. Its cross section is indeed expected to be smaller than 0.3 pb .

² in Ref. [72] an additional control region for $t\bar{t}$ events defined by requiring two high E_T jets instead of two b-tagged jets has been proposed. However, it has been shown there that due to the high contamination from Drell-Yan events, this control region is less indicate for same flavour lepton final states

Assuming the same efficiency for the kinematic selections as for the $W^+W^- \rightarrow 2\mu$ ($\sim 0.07\%$) and including the double-b tagging efficiency, less than 0.1 events are expected for 1 fb^{-1} in the control region.

In the fully simulated Drell-Yan sample used in this analysis, the eventual additional $b\bar{b}$ pair comes only from a gluon splitting, the main mechanism of $\gamma^*/Z^* + 2b$ not being included. For an estimation of the contamination of the $t\bar{t}$ control region due to this process we thus used a parton level sample generated with a matrix element Monte Carlo (MadGraph [73]). Applying the signal kinematic selections, but the E_T cut on the latter sample, ~ 10 events are expected for 1 fb^{-1} . The rejection due to E_T cut has been calculated from the fully simulated sample where actually two b-quarks were present in the final state and it turned to be smaller than 1%. Considering also the efficiency for the double b-tagging, we can safely exclude this as a dangerous background.

In the following the various contribution of uncertainty in the $t\bar{t}$ normalization procedure are listed and described. The results are summarized in Table 3.5 for 1, 5 and 10 fb^{-1} .

- **Theoretical uncertainty.**

The theoretical uncertainty of the $t\bar{t}$ cross section ratio $\sigma_{\text{signal.reg}}/\sigma_{\text{control.reg}}$ has been studied in [74] at parton level with LO precision by varying the renormalization and factorization scale. The error has been estimated to range between 3% to 10% mostly due to the choice of PDF. Some studies were done also at NLO: E_T spectra and multiplicity of jets resulted not to be affected by higher order contributions but the estimation of the theoretical error at NLO was not provided. In the following we will assume the theoretical uncertainty on the $t\bar{t}$ normalization procedure to be 10%.

- **Jet Energy Scale (JES) uncertainty.**

In the background normalization procedures we proposed, the JES uncertainty is particularly important since it affects in an opposite sense the signal region, defined by vetoing the jets, and the control region where the presence of two jets is required. To take into account this sort of anticorrelation of $\epsilon_{\text{signal.reg}}$ and $\epsilon_{\text{control.reg}}$, we estimate the effect of the JES uncertainty directly on their ratio by rescaling the measured jet four momentum by a fractional uncertainty (i.e. $P_{\text{jet}}^\mu = (1 + \lambda)P_{\text{jet}}^\mu$).

The relative variation of $\frac{N_{\text{signal.reg}}^{\text{MonteCarlo}}}{N_{\text{control.reg}}^{\text{MonteCarlo}}}$ for various values of λ is reported in [69]. The JES uncertainty foreseen at CMS is $\mathcal{O}(5\%)$ for 1 fb^{-1} and it is expected to decrease down to $\sim 3\%$ for 5 fb^{-1} (thanks to the calibration on the W mass) [61]. The effect of the JES uncertainty is 10% for 1 fb^{-1} and 6% for 5 fb^{-1} .

- **α criterion uncertainty.**

To estimate the systematic uncertainty due to α criterion, the value of the cut has been varied from 0.15 to 0.25. Moreover different values of the minimum p_t for a track to be included in the sum have been tried, from 2 to 3 GeV. The consequent variation of the jet veto efficiency (affecting only $N_{\text{signal.reg}}^{\text{MonteCarlo}}$) is relatively small, i.e. of the order of 4%.

- **b-Tagging uncertainty.**

The uncertainty on the b-tagging efficiency will be estimated exploiting $t\bar{t}$ events as calibration samples. The precision with which the b-tagging efficiency will be known is expected to be $\pm 11\%$ for 1 fb^{-1} integrated luminosity and it is foreseen to improve to $\pm 7\%$ with 10 fb^{-1} [75].

- **Uncertainties on the composition of the control region.**

As it has been shown in the previous section, $t\bar{t}$ is the dominant process in the chosen control region, other processes contributing for less than 1%. It is then safe to simply neglect this source of systematic error.

- **Statistical uncertainty on $N_{control_reg}$**

Assuming a poissonian behavior, the statistical uncertainty scales with the integrated luminosity as the square root of the number of $t\bar{t}$ events in the control region.

Luminosity (fb^{-1})	Theoretical error	Detector systematics			Statistical error	Total error
		JES	α criterion	b-tagging		
1	10%	10%	4%	11%	24%	30%
5	10%	6%	4%	9%	11%	19%
10	10%	6%	4%	7%	8%	16%

Table 3.5: Sources of uncertainty for the $t\bar{t}$ background normalization procedure. Results are shown for 1, 5 and 10 fb^{-1} .

3.2.10 WW background normalization

In contrast to the $t\bar{t}$ background normalization, which can be performed using an almost completely pure $t\bar{t}$ control sample, it is impossible to isolate the WW background in a clean way, which means that contributions of other processes have to be subtracted and their systematic uncertainties have to be taken into account during the normalization procedure of the WW background, including $gg \rightarrow W^+W^-$ events. In Fig. 3.17 the overall background normalization strategy is illustrated. There are four phase space regions involved in the WW background normalization. Each region is defined with a certain set of cuts:

- **signal region:** the selection of events in the signal region as described above.
- **WW region:** same as in the signal region, but $\Delta\phi_{\mu_1\mu_2} > 0.8$ and $50 \text{ GeV} < m_{\mu_1\mu_2} < 80 \text{ GeV}$.
- **DY (WW) region:** same as in the WW region, but $80 \text{ GeV} < m_{\mu_1\mu_2} < 100 \text{ GeV}$.
- **$t\bar{t}$ (WW) region:** same as in the WW region, but the jet veto is replaced with the requirement of two b-tagged jets ($E_t > 20 \text{ GeV}$ and two tracks with $\sigma_{IP} > 2$).

In all cases, the selection is independent of the Higgs mass hypothesis. The total number of events in each region is given in Tab. 3.6, and the contributions of individual processes are represented in form of pie charts in Fig. 3.17. The main contamination of the WW region is due to Drell-Yan, $t\bar{t}$ and the Higgs signal. The number of Drell-Yan and $t\bar{t}$ is determined by extrapolating the corresponding numbers from relatively clean control regions and are subtracted from the WW region. Additional small contributions from other backgrounds in the WW region are determined from Monte Carlos and then subtracted. So far, no concrete method has been established to subtract Higgs events from the WW control region. Therefore, we choose the conservative approach to treat these Higgs events as an additional background in the WW region.

- **Theoretical uncertainties**

The theoretical uncertainties of W pair production with subsequent decay to lep-

tons have been studied in detail in Ref. [76], and the main sources of potential uncertainties of the shapes of kinematic variables turn out to be spin correlations, underlying event, and scale dependence. The effect of spin correlations can be taken into account properly with the correct choice of an event generator, and the underlying event is expected to be measured from the data with sufficient precision. The shape dependence on the choice of the renormalization and factorization scales is sizable in case of the contribution from the $gg \rightarrow W^+W^-$ subprocess, because the higher order corrections are unknown in this case. For the cuts, described below, this uncertainty is about 9% and is taken into account in the following.

- **Statistical error and uncertainties on the composition of the control region.**

All background normalization uncertainties are calculated in the following way:

$$\delta_{\text{extrapolation}} = \sum_i \sqrt{n_{\text{total}} + (n_i \times \delta_i)^2} \times \varepsilon_{\text{control} \rightarrow \text{target}} \quad (3.17)$$

where n_{total} ³ is the total number of events in the corresponding control region, $n_i \times \delta_i$ is the product of the number of events and the systematic uncertainty of an individual process in the control region, and $\varepsilon_{\text{control} \rightarrow \text{target}}$ is the extrapolation efficiency from the control region to the target region, e.g. the signal region.

The WW background normalization requires three extrapolations from control regions to target regions:

- DY (WW) region \Rightarrow WW region: with an extrapolation uncertainty of 5% [77] the extrapolated number of events and the uncertainty from Eq. 3.17 is 15.86 ± 1.23 events (79.29 ± 4.49 events) for 1 fb^{-1} (5 fb^{-1}) of integrated luminosity.
- tt (WW) region \Rightarrow WW region: with an extrapolation uncertainty of 20% (15%) [72] the extrapolated number of events and the uncertainty from Eq. 3.17 is 6.19 ± 1.75 events (30.93 ± 5.41 events) for 1 fb^{-1} (5 fb^{-1}) of integrated luminosity.
- WW region \Rightarrow signal region: as illustrated in Fig. 3.17, the first two items are inputs to this extrapolation, which means that the obtained numbers of Drell-Yan and $t\bar{t}$ events are subtracted in the WW region and the corresponding uncertainties are propagated. The extrapolation uncertainty of WW events, which is mainly due to the unknown higher order correction of the $gg \rightarrow W^+W^-$ contribution [76], amounts to 9% for the cuts used in this analysis. In addition, the remaining backgrounds are estimated and subtracted with the following uncertainties: $\delta_{Wt} = 40\%$, $\delta_{ZW} = 20\%$ and $\delta_{ZZ} = 20\%$. According to Eq. 3.17 we obtain 7.35 ± 3.04 events (36.77 ± 7.85 events) for 1 fb^{-1} (5 fb^{-1}) of integrated luminosity.

The results of the last item are used for the calculation of the Higgs discovery potential with $m_h = 165 \text{ GeV}$, and an integrated luminosity of either 1 fb^{-1} or 5 fb^{-1} .

Furthermore, it should be pointed out that the entire background normalization procedure is performed using only the di-muon data set and therefore no additional data sets are needed. In this way, potential uncertainties due to different trigger efficiencies and different integrated luminosities of other data sets do not play a role.

³This term takes into account the statistical fluctuations of the control sample.




Figure 3.17: Scheme for background normalization from the data in different phase space regions: the signal region, the $t\bar{t}$ region, the WW region, the DY (WW) region, and the $t\bar{t}$ (WW) region. The arrows indicate the extrapolation of the number of events determined in the corresponding “control region” into the corresponding “target region”. Each region is represented by a pie chart that shows the fractions of certain types of events: h_{165} is the Higgs signal with $m_h = 165 \text{ GeV}$, WW is the sum of WW backgrounds, $t\bar{t}$ is the $t\bar{t}$ background, DY is the Drell-Yan background, and *other* is the sum of the Wt , ZW and ZZ backgrounds. The number of expected events in each region is reported in Tab. 3.6.

Channel	Signal region	$t\bar{t}$ region	WW region	$t\bar{t}$ (WW) region	DY (WW) region
Signal	14.3	0.0	6.0	0.0	0.1
$t\bar{t}$	2.6	17.0	6.2	24.7	3.2
WW	5.1	0.0	11.5	0.0	4.4
DY	0.3	0.0	15.0	0.0	267
Wt, ZZ, WZ	0.8	0.1	1.9	0.1	7.3
all	23.1	17.1	40.6	24.8	282

Table 3.6: Number of expected events in all the regions with an integrated luminosity of 1 fb^{-1} . The signal region numbers are referred to $m_H = 165 \text{ GeV}$.

3.2.11 Other backgrounds normalization

The Drell-Yan background has been normalized to estimate the contamination in the WW region. The same results can be achieved in the signal region. Fig. 3.15 demonstrates that the invariant mass cut 80 GeV to 100 GeV defines a clean control region. ZW background can be normalized by requiring one additional lepton in the final state and removing the $\Delta\phi$ and the invariant mass cuts. ZZ background can be normalized by requiring two additional leptons in the final state and removing the $\Delta\phi$ and the invariant mass cuts. They are expected to contribute to the total background by only 3% (DY), 1% (ZW) and 1% (ZZ). For the Wt

background it is not easy to define a normalization region. As this process is expected not to represent a sizable fraction of the total background ($\sim 6\%$), the Monte Carlo prediction will be then directly used, the cross section theoretical uncertainty is estimated to be about 30% at LO and 10% at NLO [67].

3.2.12 Detector misalignment systematics

A study for the misalignment impact on the track reconstruction has been done [78]. In the first data scenario ($100 \text{ pb}^{-1} - 1 \text{ fb}^{-1}$) the muon chamber position uncertainty is expected to be 1 mm and the orientation uncertainty about 0.2 mrad. The tracker position uncertainty is expected to be about 5 μm for TPE, 10 μm for TPB, 50 μm for TEC and TOB, 100 μm for TIB and 400 μm for TID. The results from simulation show the muon reconstruction efficiency will be unaffected, while the momentum resolution (for 100 GeV tracks) will be reduced from 1-2 % to 4-5%. Under these circumstances, the systematic contribution to the signal and background selection is expected to be negligible with respect to the background normalization systematics.

3.2.13 Signal significance

The signal significance can be obtained using counting or Likelihood methods. Here the counting S_{cP} method (with the "scp" program [79]) was used. S_{cP} is the probability, converted in equivalent number of sigmas, to observe at least $N_s + N_b$ events from Poisson distribution with mean N_b . The presence of systematic errors influences the significance calculations. The hypothesis is to find the same number of signal and background events predicted by the Monte Carlo. The systematic errors due to the $t\bar{t}$ and WW background normalization methods was included. Two options was considered:

1. the signal contamination in the WW control region can be subtracted;
2. the signal contamination in the WW control region must be considered as additional background.

The option 1 was considered to have a comparison with the $H \rightarrow WW \rightarrow 2l2\nu$ analysis [66]. Table 3.7 summarizes the total backgrounds and errors for different integrated luminosities. The systematics and statistical errors due to the limited Monte Carlo statistics are included.

Option	Luminosity	Total background	Total error
1.	1 fb^{-1}	8.8	3.2 (36%)
	5 fb^{-1}	44.0	8.3 (19%)
2.	1 fb^{-1}	11.0	3.2 (29%)
	5 fb^{-1}	55.3	8.3 (15%)

Table 3.7: Total background and error for integrated luminosity of 1 and 5 fb^{-1} . The two options for the signal contamination in the WW control region were considered.

The signal to background ratio as a function of different Higgs masses and the signal significance are shown in Fig. 3.18.




Figure 3.18: Signal to background ratio for the option 1. as a function of different Higgs masses. Error bars are the statistical contribution due to the limited Monte Carlo statistics (left). Significance as a function of different Higgs masses with a luminosity of 1 and 5 fb^{-1} , solid line for the option 1., dashed line for the option 2 (right).

3.2.14 Conclusions

The possibility to discover the Higgs boson particle through its decay channel into $WW^{(*)} \rightarrow 2\mu 2\nu$ was studied in detail using the most up to date CMS simulation and reconstruction programs, including the L1 trigger and HLT selections. Particular care has been put in the event selection optimization, in the determination of the number of background events from the data and in the evaluation of the experimental and theoretical systematical uncertainties. Taking all these effects into account, it was shown that in the Higgs mass range $155\text{--}175 \text{ GeV}$ a signal significance bigger than 3 standard deviations can be achieved with 5 fb^{-1} integrated luminosity. On the other hand, with 1 fb^{-1} luminosity only a 2 sigma significance can be achieved even in the most favourable case $m_H \sim 2m_W$, when this final state topology alone is used for the Higgs search.



Benchmark Channel: $Z' \rightarrow \mu\mu$



3.3.1 Introduction

FIXME: Where will the general Z' and R-S graviton introductions go? If it is somewhere else, then this intro can be deleted and just reference elsewhere.

Additional heavy neutral gauge bosons (Z') are predicted in many superstring-inspired [80, 81] and grand unified theories (GUTs) [82], as well as in dynamical symmetry breaking [83] and “little Higgs” [84] models. There are no reliable theoretical predictions, however, of the Z' mass scale. Current lower limits on the Z' mass are (depending on the model) of the order of $600\text{--}900 \text{ GeV}$ [45]. The mass region up to about 1 TeV is expected to be explored at Run II at the Tevatron [85, 86]. The LHC offers the opportunity to search for Z' bosons in a mass range significantly larger than 1 TeV .

Observability of the $Z' \rightarrow \mu^+ \mu^-$ channel in CMS is discussed in Sections 3.3.2–3.3.4. Since narrow graviton resonances such as those in Randall-Sundrum models [87] can also decay to lepton pairs (Section 14.3.1), much of the discussion in these sections is also applicable to them. If a new resonance is discovered, the characterization of its spin and couplings will proceed via the traditional methods of measuring production and decay probabilities and distributions. For example, the two-photon decay should be observable for a graviton and not for a Z' , as discussed in Section 14.6. The measurement of forward-backward asymmetries of leptonic decay products, both at the resonance peak and off the peak, yields information on parity-violating couplings and hence can help distinguish among different Z' models (Section 3.3.5). Angles of the decay products can also be used for spin discrimination (Section 3.3.6).

3.3.2 Signal and background processes

3.3.2.1 Signal $Z' \rightarrow \mu^+ \mu^-$

Signal and background samples were generated with PYTHIA [62] version 6.227 (with photon emission off incoming or outgoing quarks and leptons switched on) and the CTEQ6L set of parton distribution functions [29] from LHAPDF [88] version 4.1.1.

From a large variety of Z' bosons described in the literature, we consider six which are frequently discussed, and whose properties are representative of a broad class of extra gauge bosons:

- Z_{SSM} within the Sequential Standard Model (SSM), which has the same couplings as the Standard Model Z^0 ; it is available in PYTHIA [30].
- Z_ψ , Z_η and Z_χ , arising in E_6 and $SO(10)$ GUT groups. Couplings to quarks and leptons were obtained from Refs. [89, 90].
- Z_{LRM} and Z_{ALRM} , arising in the framework of the so-called “left-right” [91] and “alternative left-right” [85, 86] models. Their couplings were obtained from Ref. [85, 86], with the choice of $g_R = g_L$.

The generation of signal events with PYTHIA includes the full $\gamma^*/Z^0/Z'$ interference structure. We assume that Z' bosons decay only to three ordinary families of quarks and leptons and that no exotic decay channels are open. Properties for these models are in Table 3.8. The cross sections are shown at leading order (LO), as predicted by PYTHIA. We scale them by a constant K factor of 1.35 [92] in order to take into account the next-to-next-to-leading order (NNLO) QCD corrections. Electroweak higher-order corrections are not yet accounted for (see discussion in Section 3.3.4.4.1).

3.3.2.2 Background from Drell-Yan production and other processes

The dominant (and irreducible) background to $pp \rightarrow Z' \rightarrow \mu^+ \mu^-$ is the Drell-Yan production of muon pairs, $pp \rightarrow \gamma/Z^0 \rightarrow \mu^+ \mu^-$. The Drell-Yan cross section in PYTHIA was scaled by the same K factor of 1.35 [92] to get an agreement with the NNLO QCD calculations.

The overall contribution from ZZ , ZW , WW , and $t\bar{t}$ was found to be at the level of only a few percent of the Drell-Yan background and can be further suppressed by signal-selection criteria with almost no reduction in signal efficiency; we neglect this contribution. A few

Table 3.8: Summary of expected properties of Z' bosons for six studied models. For each model, the first column shows the ratio of the total Z' decay width Γ to its mass M , the second column shows the dimuon branching ratio Br. The three middle columns, labeled $\sigma^{\text{LO}} \cdot \text{Br}$, give the product of the pure- Z' leading-order production cross section and the branching ratio for three studied Z' masses; the last three columns give $\sigma^{\text{LO}} \cdot \text{Br}$ obtained when the full $\gamma^*/Z^0/Z'$ interference structure is included. The numbers quoted are for the mass intervals above 400 GeV for $M = 1$ TeV, above 1.5 TeV for $M = 3$ TeV, and above 3 TeV for $M = 5$ TeV. The values of $\sigma \cdot \text{Br}$ in the three middle columns correspond to Z' -only samples not used in our study; the values in the last three columns refer to the full-interference samples that we did use.

Model	Γ/M , %	Br ($Z' \rightarrow \mu^+\mu^-$), %	$\sigma^{\text{LO}} \cdot \text{Br}$, fb (PYTHIA)			$\sigma^{\text{LO}} \cdot \text{Br}$, full interference, fb (PYTHIA)		
			1 TeV	3 TeV	5 TeV	1 TeV	3 TeV	5 TeV
Z_{SSM}	3.1	3.0	480	1.9	0.034	610	2.8	0.050
Z_{ψ}	0.6	4.0	130	0.5	0.009	340	1.7	0.032
Z_{η}	0.7	3.4	150	0.6	0.011	370	1.8	0.035
Z_{χ}	1.3	5.7	280	1.0	0.014	500	2.2	0.038
Z_{LRM}	2.2	2.3	310	1.2	0.020	500	2.3	0.040
Z_{ALRM}	1.6	8.6	580	2.6	0.051	740	3.7	0.077

other potential background sources (like cosmics, jet-jet, W-jet, $b\bar{b}$, hadron punchthroughs, and poorly measured $Z^0 \rightarrow \mu^+\mu^-$ events) have not been studied yet, but their contribution is expected to be small.

3.3.2.3 Simulation and reconstruction

The detector response was simulated with the GEANT4-based OSCAR package [14], version 3.9.8. The digitization (simulation of the electronic response), the emulation of the Level-1 and High-Level Triggers (HLT), and the offline reconstruction were performed with the CMS full-reconstruction ORCA package [15], version 8.13.2. Both in-time and out-of-time pile-up of inelastic and diffractive collisions has been included, with an average of 5.0 and 25.0 minimum-bias collisions per bunch crossing for low luminosity ($\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$) and high luminosity ($\mathcal{L} = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$) phases of LHC operation, respectively. Misalignments of the tracker and of the muon system expected at the initial and at the well-advanced stages of the data taking have been taken into account by using two misalignment scenarios developed in the framework of the CMS reconstruction, referred to as the “first data” and the “long term” scenarios [78]:

- The “first data” scenario gives an estimate of the alignment achieved with an integrated luminosity of about 0.1 fb^{-1} and corresponds to the situation when the pixel detector is aligned with tracks and the first information from the Laser Alignment System (LAS) is available for the muon detectors.
- The “long term” scenario describes the expected residual alignment uncertainties once the performance of the LAS reaches its design level and the alignment with tracks is done in all tracking detectors. The current estimate is that this can be achieved with an integrated luminosity of about 1 fb^{-1} .

As a result, for each of the Z' models above, several sets of simulated samples corresponding to different possible combinations of luminosities and misalignment scenarios were produced at each of three mass values of 1, 3, and 5 TeV. Since the Drell-Yan cross section falls rapidly with the mass of the muon pair, Drell-Yan background was generated in six mass intervals (with lower mass bounds of 0.2, 0.4, 1, 1.5, 2, and 3 TeV), again for different combinations of luminosities and misalignment scenarios.

3.3.3 Event selection

For $\mu^+\mu^-$ invariant mass between 1 TeV and 5 TeV, the fraction of Drell-Yan events with both muons within the full geometrical acceptance of the muon system ($|\eta| < 2.4$) increases from about 80% at 1 TeV to almost 95% at very high masses. The acceptance of $Z' \rightarrow \mu^+\mu^-$ events is very similar.

We require that the event pass the logical OR of single-muon and dimuon triggers, both Level-1 and HLT. We use the default ORCA implementations of low-luminosity and high-luminosity muon trigger algorithms described in Refs. [68, 7], with the exception of the HLT calorimeter isolation criterion requiring that the weighted sum of energy deposits in ECAL and HCAL in a cone around the muon direction be below a pre-defined threshold. Its current implementation leads to significant efficiency losses for *isolated* high- p_T muons (since they are often accompanied by electromagnetic showers) and is currently being worked on; we do not apply HLT calorimeter isolation in this study (tracker isolation is applied). An increase in the trigger rate in the absence of calorimeter isolation should be mitigated by higher p_T thresholds; we have checked that raising the p_T thresholds of the single-muon HLT by 10–20 GeV with respect to their nominal values changes trigger efficiency for our signals by a negligible amount. For the Z' models that we study (as well as for the Drell-Yan background), the combined Level-1/HLT trigger efficiency is about 98% at 1 TeV and decreases with the Z' mass down to about 95% at 5 TeV. At high luminosity, the trigger efficiency is 95% at 1 TeV and 93% at 5 TeV. These efficiencies are relative to having at least one muon inside the geometrical acceptance of the muon trigger ($|\eta| < 2.1$) and both muons from the Z' decay inside the full acceptance of the muon system. No dependence of trigger efficiency on tracker and muon misalignment has been observed, in agreement with the results reported in Ref. [93].

We require that at least two muons of opposite sign charge be reconstructed offline. Detailed description of offline muon reconstruction can be found in Ref. [7]. For each muon candidate, we examine the results of fits to two subsets of hits associated to this candidate: 1) excluding all muon hits except for those in the innermost muon station, and 2) excluding hits in muon chambers appearing to contain electromagnetic showers. Optimal performance for high- p_T muons is achieved by choosing the best fit on a track-by-track basis using goodness-of-fit variables. The fraction of Z' events with an opposite-sign dimuon reconstructed offline is about 97% at 1 TeV for both the “first data” and the “long term” misalignment scenarios, and decreases slightly with the Z' mass, to about 95% at 5 TeV for the “long term” misalignment scenario. The efficiencies quoted are calculated relative to the number of events accepted by the trigger and with both muons from the Z' decay within the full geometrical acceptance of the muon system.

The overall efficiency – including acceptance, trigger and offline reconstruction – for $Z' \rightarrow \mu^+\mu^-$ events with a mass between 1 and 5 TeV lies in the range of 77–85% at low luminosity, and of 75–83% at high luminosity.

3.3.4 Signal observability

The search for a new resonance is performed with an unbinned maximum likelihood fit to the $\mu^+\mu^-$ invariant mass spectrum over a range which includes Drell-Yan continuum as well as a possible peak. The fit takes as input the presumed signal and background shapes, and determines the best-fit background normalization. More details are in Refs. [94, 95].

3.3.4.1 Mass spectra and fitting procedure

Prior to the calculation of the invariant mass of an opposite-sign muon pair, $M_{\mu\mu}$, a search for photon candidates in a cone with a radius of $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} < 0.1$ around the trajectory of each muon is performed, and the 4-momentum of the photon candidate with the smallest ΔR in the cone is added to the 4-momentum of the muon. This procedure recovers some of the energy lost by the muon via final state radiation and radiative processes in the detector, thus improving the invariant mass resolution.

The resolution for $M_{\mu\mu}$ depends strongly on the misalignment scenario, and weakly on the amount of pile-up. If the ‘‘long term’’ misalignment scenario for the tracker and the muon chambers is considered, the sigma of the Gaussian fit to the mass resolution curves varies from 4.2% at 1 TeV to 9.0% at 5 TeV; the RMS truncated at $\pm 30\%$ is $\sim 6\%$ at 1 TeV and $\sim 10\%$ at 5 TeV. The corresponding numbers for the ‘‘first data’’ misalignment scenario at 1 TeV are $\sigma = 12.5\%$ and RMS $\sim 12\%$. The bias in the mass resolution does not exceed 1% for the ‘‘long term’’ scenario at all masses considered and for the ‘‘first data’’ scenario at 1 TeV.

An example of the $M_{\mu\mu}$ spectra showing 1 TeV Z_η signal and Drell-Yan background is in Figure 3.19. The left-hand plot shows generated mass spectra (100% efficiency with no detector- and reconstruction-related effects); it can be compared to the right-hand plot for fully-reconstructed events using the ‘‘first data’’ misalignment scenario. Signal peak is clearly visible in spite of the poor mass resolution.

The mass spectra in Figure 3.19 are obtained by re-scaling the simulated spectra with large statistics down to a modest number of events characteristic for the regime close to the discovery limit; the statistical fluctuations are thus not to scale. In what follows, we use ensembles of Monte Carlo pseudo-experiments selected from available large-statistics samples. The number of events in each experiment, N_{evt} , fluctuates according to a Poisson distribution with a mean of $\sigma \cdot \text{Br} \cdot \int \mathcal{L} dt \cdot \varepsilon$, where $\int \mathcal{L} dt$ is the integrated luminosity and ε is the combined trigger and reconstruction efficiency.

In order to test for the existence of a resonance and to measure its parameters if it is found to exist, an unbinned maximum likelihood fit of the $M_{\mu\mu}$ values in each MC experiment is appropriate. One can imagine that, in the initial data analysis, one is confident about the background shape but not the absolute normalization. In this case, data can be fit with a sum of signal and background shapes, presumed known, with the signal fraction as a free parameter. In the presence of a signal, one can fix or let vary the mass and the width as well. Thus, as a model of the probability density function (pdf), p , of the parent population of the observed mass spectra, we use

$$p(M_{\mu\mu}; f_s, m_0, \Gamma) = f_s \cdot p_s(M_{\mu\mu}; m_0, \Gamma) + (1 - f_s) \cdot p_b(M_{\mu\mu}). \quad (3.18)$$

Here




Figure 3.19: Histograms of the $\mu^+\mu^-$ invariant mass for 1 TeV Z_η plus background (open histogram) and for background only (shaded histogram), at the event-generator level (left) and for events selected by the Level-1/HLT triggers and reconstructed assuming the “first data” misalignment scenario (right). The number of events per bin is normalized to an integrated luminosity of 0.1 fb^{-1} .

- p_s , the pdf of the signal, is a convolution of a Breit-Wigner signal shape with a Gaussian accounting for mass resolution smearing. The convolution includes the dependence of the mass resolution on $M_{\mu\mu}$, but the radiative tail of the signal is not yet accounted for.
- p_b , the pdf of the background, is modeled as an exponential, $\exp(-k \cdot M_{\mu\mu}^{0.3})$, with the parameter k determined from fits to Drell-Yan events. This pdf, with the value of k of 2.0, gives a good description of the background shape in the whole mass region between 400 and 5000 GeV.

There are three free parameters in the fit: the signal fraction $f_s = N_s/(N_s + N_b)$, the position of the mass peak m_0 , and the full width at half maximum (FWHM), Γ , of the signal. The shape of the background distribution is fixed, while its level is determined by the fit: f_s is a free parameter. Therefore, the fit explores the difference *in shape* between the signal and the background, and is not sensitive to uncertainties in the expected signal and background levels.

The background shape is currently determined from fits to large-statistics background-only simulated distributions in the full mass region of interest, including the region under the signal peak. In the real experiment, the shape will likely have to be extracted from the data in signal-free regions. The accuracy of predicting the background shape is an important contribution to the systematic uncertainty of the analysis and is discussed in Section 3.3.4.4.3.

Ref. [94] contains examples of results of fits to M.C. small-event samples. With even the small number of events needed to give evidence of a resonance, the mass is determined fairly well, with a precision of 4–8% depending on the resonance mass and alignment uncertainties. However, for the narrow resonances under study, typically little information can be obtained about the width.

Table 3.9: Average values of the likelihood-ratio significance estimator $S_{\mathcal{L}}$ for six different Z' models, at three signal mass points and for a few representative values of an integrated luminosity. The uncertainties shown are statistical only.

Mass	1 TeV	3 TeV	5 TeV
$\int \mathcal{L} dt$	0.1 fb^{-1}	10 fb^{-1}	300 fb^{-1}
Z_{SSM}	12.4 ± 0.2	10.1 ± 0.2	5.8 ± 0.1
Z_{ψ}	5.1 ± 0.2	4.4 ± 0.1	2.4 ± 0.2
Z_{η}	5.5 ± 0.2	5.1 ± 0.1	2.9 ± 0.1
Z_{χ}	9.1 ± 0.2	6.7 ± 0.2	3.2 ± 0.1
Z_{LRM}	9.0 ± 0.2	7.4 ± 0.2	4.1 ± 0.1
Z_{ALRM}	13.3 ± 0.3	11.8 ± 0.2	7.7 ± 0.2

3.3.4.2 Significance estimator

We follow closely the approach of Ref. [96], which is based on the theorem of S.S. Wilks [97]. The test statistic is the likelihood-ratio estimator $S_{\mathcal{L}}$:

$$S_{\mathcal{L}} = \sqrt{2 \ln (\mathcal{L}_{s+b} / \mathcal{L}_b)}, \quad (3.19)$$

where \mathcal{L}_{s+b} is the maximum likelihood value obtained in the full signal-plus-background fit, and \mathcal{L}_b is the maximum likelihood from the background-only fit. Studies show [94] that in the small-statistics low-background regime characteristic of a Z' search, the asymptotic conditions of Wilks's theorem [97] are satisfied well enough and $S_{\mathcal{L}}$ is the number of Gaussian-equivalent standard deviations a measurement lies from the value predicted by a background-only (null) hypothesis. This requires fixing both m_0 and Γ in the fits using the pdf of Eq. (3.18).

We follow a common convention in using the (arbitrary, but useful for comparison) specification that $S > 5$ is necessary to establish a discovery. This S refers to the local excess without accounting for the degree of freedom due to the unknown mass; how one might de-rate S in a time-dependent way in this context as data comes in will be the subject of a future study.

3.3.4.3 Discovery potential in $Z' \rightarrow \mu^+ \mu^-$ channel

Table 3.9 gives a summary of the signal significance expected for different Z' models, masses and integrated luminosities. The numbers shown are for the ‘‘first data’’ misalignment scenario and low luminosity parameters for $\int \mathcal{L} dt = 0.1 \text{ fb}^{-1}$, the ‘‘long term’’ misalignment scenario and low luminosity parameters for 10 fb^{-1} , and the ‘‘long term’’ misalignment scenario and high luminosity parameters for 300 fb^{-1} . $S_{\mathcal{L}}$ scales as expected with the square root of $\int \mathcal{L} dt$.

We use the same combinations of luminosities and misalignment scenarios to calculate the integrated luminosity needed to reach 5σ significance. The results for various Z' models are shown in Figure 3.20 as a function of Z' mass. One can see that

- A very low integrated luminosity, less than 0.1 fb^{-1} , and non-optimal alignment of the tracker and the muon detectors should be sufficient to discover Z' bosons at 1 TeV, a mass value which will likely be above the Tevatron reach. One would need about 50% less data to reach the same signal significance if the optimal alignment is achieved.




Figure 3.20: Integrated luminosity needed to reach 5σ significance ($S_{\mathcal{L}} = 5$) as a function of Z' mass for (top to bottom) Z_{ψ} , Z_{η} , Z_{χ} , Z_{LRM} , Z_{SSM} and Z_{ALRM} . Symbols indicate fully-simulated mass-luminosity points, lines are the results of interpolations between the points.

- An integrated luminosity of 10 fb^{-1} is sufficient to reach 5σ significance at 3 TeV for most (but not all) of the Z' models considered if the optimal alignment is available: depending on the model, the mass reach is in the range between 2.9 and 3.8 TeV.
- An integrated luminosity of 100 fb^{-1} does not allow one to obtain 5σ significance at 5 TeV with only the $Z' \rightarrow \mu^+ \mu^-$ channel for any of the models considered: the corresponding mass reach lies in the region between 3.9 and 4.9 TeV.

These estimates of signal significance do not incorporate systematic uncertainties, which we discuss in the next section.

3.3.4.4 Systematic uncertainties

The main sources of systematic uncertainties are expected to be a) theoretical uncertainties (parton distributions, higher-order corrections, etc.), b) uncertainties arising from an imperfect knowledge of the detector (alignment, calibration, magnetic field), and c) uncertainties in the fitting procedure (background shape, functional forms of pdf's, mass resolution, etc.).

3.3.4.4.1 Theoretical uncertainties Our current estimates of the Z' mass reach depend on the accuracy of the modeling of the Standard Model processes and of the Z' boson production. The following sources of theoretical uncertainties have been studied:

- **Higher-order QCD corrections.** We use a constant $K_{\text{QCD}}^{\text{NNLO}}$ factor of 1.35 to rescale PYTHIA cross sections for Drell-Yan and Z' bosons to NNLO QCD predictions. This is an approximation, since such a reweight does not take into account variations of the ratio of NNLO and LO cross sections with the invariant mass and other observables, such as rapidity and p_T . It was shown [92] that the variations of the $K_{\text{QCD}}^{\text{NNLO}}$ factor with the mass in the mass interval between 500 GeV and 5 TeV is in the range of $\Delta K_{\text{QCD}} = \pm 0.05$; the dependence on other observables and the ensuing impact on acceptance, efficiency, etc. remains to be studied. Since K is expected to be nearly identical for the signal and dominant background, the effect of changes in K from the nominal value $K_0 = 1.35$ is to scale the expected significance by $\sqrt{K/K_0}$.
- **Higher-order electroweak corrections.** Only preliminary estimates of electroweak next-to-leading order corrections exist for the LHC and $M_{\mu\mu} > 1$ TeV [98, 99]. Currently, we use $K_{\text{EW}} = 1$ for the central values of signal and background cross-sections, and assign an uncertainty of $\Delta K_{\text{EW}} = \pm 0.10$ based on discussions in Refs. [98, 99].
- **Parton distribution functions (PDFs).** We use the CTEQ6.1M eigenvector PDF sets [29] and the “master” equations in Ref. [100] to evaluate the uncertainties characterizing current knowledge of the parton distributions. The effect on the total cross section σ was found to be similar for the Drell-Yan background and for the studied Z' models at any given mass, with prediction uncertainties lying in the range of $\frac{\Delta\sigma}{\sigma} = \begin{smallmatrix} -7\% \\ +4\% \end{smallmatrix}$ at $M_{\mu\mu} = 1$ TeV, raising to $\begin{smallmatrix} -10\% \\ +12\% \end{smallmatrix}$ at $M_{\mu\mu} = 3$ TeV, and reaching as much as $\begin{smallmatrix} -20\% \\ +30\% \end{smallmatrix}$ at $M_{\mu\mu} = 5$ TeV. The effect on other observables and on the acceptance has not been studied yet, but is expected to be small.
- **Hard process scale.** The dependence of the observables on the choice for renormalization and factorization Q^2 scales, μ_R and μ_F , is unphysical and is commonly taken as a rough estimate of the uncertainty due to unaccounted higher orders in QCD calculations. The study of the sensitivity of the Drell-Yan cross section to the choice for the QCD scale is described in Ref. [92]. Both μ_F and μ_R were varied in the range of $M_{\mu\mu}/2 < \mu < 2M_{\mu\mu}$ around the default choice of $\mu = M_{\mu\mu}$, and the mass-dependent variations of the cross section obtained. At NNLO, they are smaller than $\pm 1\%$ at 1 TeV, but as large as -25% (for $\mu = 2M_{\mu\mu}$) and $+5\%$ (for $\mu = M_{\mu\mu}/2$) at 5 TeV. We use the NNLO estimates given in Ref. [92] for both the Drell-Yan and the Z' bosons.

Since our analysis relies only on the background shape and not on any assumptions about background normalization, the uncertainties in signal and background cross sections described in this section will not have any direct impact on the calculation of significance once a data set is in hand. They do effect, however, estimates of the Z' mass reach based on Monte Carlo predictions for the signal and the background. We combine them in quadrature, and use the obtained mass-dependent band as 1σ uncertainty in the expected number of signal and background events. This band is then translated into 1σ uncertainty in the prediction of the mean integrated luminosity needed to reach 5σ significance for any given Z' model. This uncertainty, and the best estimates of the luminosity, is shown in Figure 3.21 for the models




Figure 3.21: Integrated luminosity needed to reach 5σ significance ($S_{\mathcal{L}} = 5$) as a function of Z' mass for Z_{ψ} and Z_{ALRM} models. Solid lines show the best estimates, dashed lines indicate boundaries of the band corresponding to the predictions with $\pm 1\sigma$ theoretical uncertainty.

with the smallest and the largest values of $\sigma \cdot \text{Br}$ among the models studied, Z_{ψ} and Z_{ALRM} .

3.3.4.4.2 Uncertainties in the detector performance The key element in the performance of high- p_{T} muon reconstruction and, therefore, for the Z' mass reach is the alignment of the tracker and the muon system. Unlike the muons in the region of low and moderate p_{T} values, where the influence of the tracker alignment is predominant, both the tracker alignment and the muon system alignment play an important role for the muons at TeV scale. We take them into account by using the two realistic misalignment scenarios developed in the CMS reconstruction, the “first data” and the “long term”. These scenarios, however, are only based on the current best estimates (and sometimes guesses) of expected alignment uncertainties and will be refined as better estimates from alignment studies become available. Therefore, they have intrinsic uncertainties, which at the moment cannot be evaluated. As discussed above and in Ref. [93], neither the trigger efficiency nor the offline reconstruction efficiency for high- p_{T} muons is affected by the misalignment even in the worst-case scenario once the alignment position uncertainties are used in reconstruction algorithms [78]. So uncertainties in alignment translate mainly into uncertainties in the invariant mass resolution. We show below that even sizable variations in the width of the mass resolution have only a

small impact on the Z' mass reach.

Another potentially important source of systematic uncertainties is the uncertainty in the calibration precision of the muon chambers. The impact of uncertainties in the calibration of the Drift Tube chambers on the Z' mass reach has been studied by 1) changing the t_0 offsets for all chambers by ± 2 ns, and 2) scaling drift velocity (changing time-to-distance relationship) by $\pm 3\%$. These variations represent conservative upper bounds on corresponding effects [101]. The effect of changing t_0 offset was found to be negligible for Z' samples at all studied mass values and for both misalignment scenarios considered. The scaling of drift velocity has a negligible impact for the “first data” misalignment scenario with its rather poor mass resolution, but results in an increase of 5–10% in the width of the mass resolution for the “long term” scenario (no change in trigger and dimuon reconstruction efficiencies). This translates into a negligible effect in the Z' mass reach. Uncertainties in the calibration of the Cathode Strip Chambers are less critical and hence are expected to have a negligible impact on the Z' detection as well.

The effect of uncertainties in the knowledge of the magnetic field remains to be studied.

3.3.4.4.3 Uncertainties in background shape and mass resolution Many experimental uncertainties have a negligible or small impact on the results of our studies because the proposed analysis method is not sensitive to uncertainties in the predicted levels of signal and background processes. For example, only the mass dependence of the uncertainty in the muon reconstruction efficiency needs to be taken into account, not the absolute uncertainty. The same is true for the trigger efficiency and for the uncertainty in the $M_{\mu\mu}$ scale. Among those uncertainties that do not cancel out, two seem to be particularly important: the uncertainty in the background shape, and the uncertainty in the mass resolution.

As described above, the background shape is currently determined from fits to background distributions predicted by the Monte Carlo simulation. In the analysis of real data, this MC-based shape will be compared with (and perhaps tuned to) the background shape in the region of low masses where one has high statistics of background events. The issue is then the reliability of the extrapolation from the steeply falling spectrum into the candidate signal region. This will have to be studied in detail once the real data starts to be available. What is interesting to explore at this stage of analysis is how rapidly the significance deteriorates as the ratio of background events in the high-statistics normalization region to background events in the candidate signal region is wrongly predicted by the MC-motivated background shape. To study this, we multiply our background pdf (p_b in Eq. (3.18)) by a function which is unity in the high-statistics background-only region and smoothly transitions to a tunable value, f , under the candidate mass peak. Values of integrated luminosity were chosen to correspond to 5σ significance for each model at $f = 1$. For $f = 2$ (assuming twice as much background in the signal region as there really is), 5σ becomes 4.2σ for Z_{ALRM} and is about 3.7σ for Z_ψ . For f around 1.1 or 1.2, the change in S is of the order of a few per cent.

Sensitivity of the Z' mass reach to uncertainties in the invariant mass resolution has been studied by applying extra Gaussian smearing to the reconstructed values of $M_{\mu\mu}$ of both the signal and background events and comparing the signal significance obtained with modified $M_{\mu\mu}$ values to that calculated with the nominal $M_{\mu\mu}$ values. We found that an increase of 10% in the mass resolution width, σ_M , reduces the signal significance by less than 2% at the values of $S_{\mathcal{L}}$ close to 5; 20% worse resolution gives 5% or less smaller $S_{\mathcal{L}}$. The effect is not very big, indicating that an approximate knowledge of σ_M should suffice. (This exercise does not

check, however, the effect of extreme tails of the mass resolution being bigger than expected, which could lead to a background shape (and amount) different from that obtained from the simulation.) The knowledge of σ_M as a function of $M_{\mu\mu}$ is also used in the pdf of the signal in Eq. (3.18), where it defines the width of a Gaussian accounting for resolution smearing of the signal shape. This does not need to be very precise either: assuming resolution 20% better than it really is reduces $S_{\mathcal{L}}$ by less than 1%.

3.3.5 Distinguishing among Z' models

The forward-backward asymmetry, A_{FB} , of the leptonic decay products provides information on parity-violating couplings, on and off resonance, as discussed for example in Refs. [102, 103].

FIXME: If the following is somewhere else in the PTDR, delete it. In any case, refer to and coordinate text with DY Standard Model part of the TDR.

The forward-backward asymmetry for $q\bar{q} \rightarrow \mu^+\mu^-$ interactions is defined as (e.g., Refs. [103, 104])

$$A_{\text{FB}} = \frac{\sigma_F - \sigma_B}{\sigma_F + \sigma_B}, \quad (3.20)$$

where

$$\sigma_F \equiv \int_0^1 \frac{d\sigma(q\bar{q} \rightarrow \mu^+\mu^-)}{d\cos\theta^*} d\cos\theta^*, \quad \sigma_B \equiv \int_{-1}^0 \frac{d\sigma(q\bar{q} \rightarrow \mu^+\mu^-)}{d\cos\theta^*} d\cos\theta^*, \quad (3.21)$$

and where θ^* is the angle in the dimuon center-of-mass (CM) reference frame between the negative muon and the incident quark. For spin-1 $\gamma^*/Z^0/Z'$ propagators, the probability density function $P(\cos\theta^*)$ is most generally of the form

$$P(\cos\theta^*; A_{\text{FB}}, b) = \frac{3}{2(3+b)}(1 + b\cos^2\theta^*) + A_{\text{FB}}\cos\theta^*. \quad (3.22)$$

Although $b = 1$ from general considerations, in the fits described here b is typically left as a free parameter. In Ref. [90], Rosner expresses A_{FB} for $f\bar{f} \rightarrow \gamma^*/Z^0/Z' \rightarrow \mu^+\mu^-$ events in terms of the left- and right-handed couplings of the photon, Z^0 , and Z' to u quarks, d quarks, and charged leptons. More details, including the couplings for the models studied, are given in Ref. [105].

For CM energies well above the Z^0 peak, the Drell-Yan background has a characteristic A_{FB} of about 0.6 [103], and provides a useful starting point.

3.3.5.1 Uncertainty in the sign of $\cos\theta^*$ in pp collisions

In proton-proton interactions, the quark direction is ambiguous experimentally since a quark can originate with equal probability from either proton, and the sign of $\cos\theta^*$ is not directly measurable. We follow Ref. [106] and infer the sign of $\cos\theta^*$ by assuming that the longitudinal motion of the dimuon system is in the direction of the proton contributing the annihilating quark, since a quark in a proton typically carries a larger momentum fraction x than does an antiquark. We refer to the inference of the wrong sign of $\cos\theta^*$ as “mistagging” the sign. If not accounted for, the mistagged events, particularly at low y , reduce (“dilute”) the apparent value of A_{FB} . Some authors deal with this problem by removing events below a chosen y threshold [106], or by examining A_{FB} in bins of y [107]; in Ref. [105], an approached

is described which assigns the probability of a mistag on an event-by-event basis, thus using all events in a given sample. As knowledge of the mistagging probability depends on the Parton Distribution Functions, the effect of uncertainties in PDFs must be evaluated, and will be the subject of future work.

3.3.5.2 Other uncertainties

The transverse momentum p_T of the annihilating quark and/or antiquark provides another source of uncertainty in the measurement of $\cos \theta^*$, since the observable quantity is the vector sum of these transverse momenta. We use the Collins-Soper reference frame [108], in which angles are measured with respect to the axis that bisects the target and beam axes in the dimuon CM frame, to minimize the effect of p_T on the measurement of $\cos \theta^*$, and let θ_{CS}^* denote the polar angle of the μ^- in this frame.

As described in Ref. [105], the effect of detector acceptance, combined with high mistag probability for events near $y = 0$, means that events lying near the edges of acceptance carry the largest information for the A_{FB} measurement. Hence, in addition to trying to obtain maximum acceptance, it is particularly important to understand the effect of any asymmetries in the acceptance which may arise as a result of the real detector efficiencies not being perfectly symmetric or of the beam crossing not being perfectly centered.

3.3.5.3 Likelihood function and fitting procedure

Since a Z' can be discovered with a small number of events (Section 3.3.4), and since the search for anomalous A_{FB} in the highest mass continuum Drell-Yan events at any given luminosity will use a restricted sample of events, we consider an unbinned likelihood fit. The procedure and results with statistical errors only are described in Ref. [105]. The results of numerous fits can be summarized simply with a nominal statistical uncertainty in A_{FB} of 0.09 in a fit with 400 events for 1 TeV Z' samples, and of 0.08 with 400 events for 3 TeV samples. Ref. [105] also reviews an appropriate hypothesis-testing methodology for distinguishing between Z' models. Results incorporating misalignment and systematic errors including those due to Parton Distribution Functions are in progress.

3.3.6 Discriminating between different spin hypotheses

In order to distinguish the spins of a spin-1 Z' bosons and a spin-2 gravitons in a dilepton decay mode, Ref. [109] considers an unbinned likelihood ratio statistic incorporating the angles of the decay products. The statistical interpretation of this statistic is discussed in detail in Ref. [110], also considering the possibility of spin 0.

To leading order, the subdiagram for Z' formation is quark-antiquark ($q\bar{q}$) annihilation, while for a graviton there exist both $q\bar{q}$ annihilation and gluon-gluon (gg) fusion. One defines θ^* as the angle in the dilepton center-of-mass reference frame between the negative lepton ℓ^- and the incident quark or gluon. In this section we consider only the parity-conserving terms; inference from these terms can be combined with that of the parity-violating terms giving rise to A_{FB} .

For light lepton decay products, the angular probability density functions in the absence of interference are in Table 3.10. These are determined from angular momentum considerations

and do not depend on the couplings. For the spin-2 graviton, only the relative fractions of $q\bar{q}$ annihilation, gluon fusion, and background (predominantly from the Drell-Yan continuum) events are needed to arrive at a parameter-free form for the expected distribution. (For spin 1, the resonance and the Drell-Yan background have the same form.)

Table 3.10: Angular distributions for the decay products of spin-1 and spin-2 resonances, considering only even terms in $\cos\theta^*$.

Channel	d -functions	Normalized density for $\cos\theta^*$
$q\bar{q} \rightarrow G^* \rightarrow f\bar{f}$	$ d_{1,1}^2 ^2 + d_{1,-1}^2 ^2$	$P_q = \frac{5}{8} (1 - 3\cos^2\theta^* + 4\cos^4\theta^*)$
$gg \rightarrow G^* \rightarrow f\bar{f}$	$ d_{2,1}^2 ^2 + d_{2,-1}^2 ^2$	$P_g = \frac{5}{8} (1 - \cos^4\theta^*)$
$q\bar{q} \rightarrow \gamma^*/Z^0/Z' \rightarrow f\bar{f}$	$ d_{1,1}^1 ^2 + d_{1,-1}^1 ^2$	$P_1 = \frac{3}{8} (1 + \cos^2\theta^*)$

The fractions of generated events arising from these processes are denoted by ϵ_q , ϵ_g , and ϵ_1 , respectively, with $\epsilon_q + \epsilon_g + \epsilon_1 = 1$. Then the form of the probability density $P(\cos\theta^*)$ is

$$P(\cos\theta^*) = \epsilon_q P_q + \epsilon_g P_g + \epsilon_1 P_1. \quad (3.23)$$

As in the A_{FB} measurements, we let θ_{CS}^* denote the polar angle of the ℓ^- in the Collins-Soper frame. Experimentally one will obtain a set of events with θ_{CS}^* measured along with other quantities such as dilepton transverse momentum p_T^{dil} and rapidity y^{dil} . From these, one can construct the probability density $P_{\text{acc}}(\cos\theta_{CS}^*)$ for events accepted (observed) in an experiment for each hypothesis H_i , where i labels the model such as Z' or G^* . In this study, we consider only the angular information and integrate over p_T^{dil} , y^{dil} , and any other relevant quantities; if one has confidence that these quantities are well described by the event generators, more variables can be added to P_{acc} . Since we do not add this information, P_{acc} for accepted events approximately factorizes:

$$P_{\text{acc}}(\cos\theta_{CS}^*|H_i) = P(\cos\theta_{CS}^*|H_i) \bar{\Omega}(\cos\theta_{CS}^*), \quad (3.24)$$

where $P(\cos\theta_{CS}^*|H_i)$ is from Eq. (3.23) with the ϵ_j set appropriately for the model considered (e.g. for the spin-1 hypothesis, we set $\epsilon_1 = 1$ and $\epsilon_q = \epsilon_g = 0$), and $\bar{\Omega}$ is the acceptance averaged over p_T , y , etc.

Eq. (3.24) has no free parameters if the fractions ϵ_q , ϵ_g , and ϵ_1 are considered to be fixed. For each observed event, one evaluates $P_{\text{acc}}(\cos\theta_{CS}^*|H_i)$ at the observed $\cos\theta_{CS}^*$ to obtain the *likelihood* $\mathcal{L}(H_i)$ of that event under the given hypothesis. The combined likelihood of the data set under a hypothesis is then the product of the events' likelihoods; henceforth in this paper, $\mathcal{L}(H_i)$ refers to this product unless otherwise stated. As Ref. [?] discusses, the absence of free parameters means that the Neyman-Pearson hypothesis testing for *simple* hypothesis testing is applicable.

For testing a simple null hypothesis H_A of one spin against another simple alternative spin hypothesis H_B , we use the likelihood ratio $\lambda = \mathcal{L}(H_A)/\mathcal{L}(H_B)$, with critical region again chosen such that $\alpha = \beta$. For investigating and summarizing which values of λ^{cut} correspond to which values of α and β , the quantity $-2\ln\lambda = 2\ln\mathcal{L}(H_B) - 2\ln\mathcal{L}(H_A)$ is particularly useful. For simple hypotheses H_A and H_B , the central limit theorem implies that $-2\ln\lambda$ tends to a Gaussian.

3.3.6.1 Testing spin 1 versus spin 2

A detailed discussion of the intermediate steps in applying the above method for discriminating spin 1 from spin 2 is in Ref. [?], using large samples of Z' and G^* events (from the Randall-Sundrum (RS) model [87]) generated with HERWIG. (Generator-level results using PYTHIA are completely compatible.) The ratio λ of the likelihoods of the hypotheses is calculated for each event, assigning spin 1 as the null hypothesis H_A and spin 2 as the alternative hypothesis H_B . In taking the ratio, the average acceptance cancels to good approximation and one essentially recovers the ratios of the angular forms. Histograms of $-2 \ln \lambda$ for these events are highly asymmetric and strongly peaked at one side [?]. In view of the asymmetries in the underlying event histograms, the convergence of the sums of $-2 \ln \lambda$ values for N selected events toward Gaussians is quite striking. The means and rms deviations of the sums are in excellent agreement with the means and rms deviations of the respective event histograms scaled by N and \sqrt{N} , respectively, as expected from the central limit theorem.

The statistical technique of Ref. [?] has been applied to fully-reconstructed Z' and G^* events [111]. Details of simulation, trigger and reconstruction are described in Sections 3.3.2, 3.3.3 and 14.3.1. From ensembles of pseudo-experiments, we determine the number N of events per experiment corresponding to various values of $\alpha = \beta$, expressed in equivalent number of Gaussian standard deviations “ σ ” for *one-tailed* tests, e.g., for $\alpha = 0.159$ we report $\alpha = 1\sigma$, and so on. The values of α so obtained scale as expected as \sqrt{N} .

Table 3.11 contains, for different studied masses and values of the Randall-Sundrum ratio $c = k/\bar{M}_{\text{Pl}}$, the integrated luminosity needed for a 2σ significance, and the corresponding numbers of signal and background events. All numbers are for the “long term” misalignment scenario; the cross section for Z' production is assumed to be equal to that of G^* with the given c value. Of course, because the production cross section falls rather steeply with mass, the integrated luminosity needed for spin discrimination increases with mass. For RS gravitons, the production cross section scales as c^2 ; therefore, the integrated luminosity required for spin discrimination quickly increases as c gets smaller, and so does the number of signal events, because of a larger background contamination. The region in the plane of $M_{G^*}-c$ in which Randall-Sundrum G^* can be distinguished from Z' with 2σ significance if one treats two spin hypotheses symmetrically is shown in Figure 3.22 for a few representative values of the integrated luminosity.

Table 3.11: Integrated luminosity and numbers of signal and background events N_s and N_b required to discriminate spin-1 and spin-2 hypotheses with $\alpha = \beta$ corresponding to 2σ (one-tailed). The first column indicates the mass of the resonance; the second column shows the values of the RS ratio $c = k/\bar{M}_{\text{Pl}}$; the third column specifies the integrated luminosity needed for 2σ discrimination; the last two columns show the corresponding numbers of signal and background events.

$M_{\mu\mu}$, TeV	c	$\int \mathcal{L} dt$, fb $^{-1}$	N_s	N_b
1.0	0.01	50	200	87
1.0	0.02	10	146	16
1.5	0.02	90	174	41
3.0	0.05	1200	154	22
3.0	0.10	290	148	6

Alternatives to the $\alpha = \beta$ criterion, in particular tests in which α is minimized for one hy-




Figure 3.22: Region in the plane of $M_{G^*}-c$ in which Randall-Sundrum G^* can be distinguished from Z' having an equal cross section with 2σ significance if one treats two spin hypotheses symmetrically, for a few representative values of the integrated luminosity. The region which can be probed lies to the left of the lines.

pothesis at the cost of increase in β , are discussed in Ref. [?].

3.3.6.2 Discrimination from spin 0

While the motivation of discriminating Z' from G^* has focused studies on discriminating spin 1 from spin 2, another possibility to be considered is spin 0 resonance (which is uniform in $\cos\theta^*$). For accepted spin-0 events, the probability density for $\cos\theta_{CS}^*$ is somewhat in between the mostly concave-upward function for spin 1 and the predominantly concave-downward function for spin 2.

As discussed in Ref. [?], discriminating either spin 1 or spin 2 from spin 0 requires significantly more events than discriminating spin 2 from spin 1.

Chapter 4

Physics Studies with Jets and E_T^{miss}



4.1 Benchmark Channel: new physics from dijets

Inclusive dijet production ($pp \rightarrow 2 \text{ jets} + X$) is the dominant LHC process. Simple to observe, and rich in potential signals of new physics, dijets are expected to be one of the earliest CMS measurements. In this section we discuss the measured distributions and their systematic uncertainties [112]. In section 14.5.2 and 15.3 we use these distributions to estimate our sensitivity to specific models of new physics.

4.1.1 Dijet Analysis

We use samples produced during Data Challenge 2004, generated using Pythia dijet processes mixed with pileup of minimum bias interactions for an assumed luminosity of $2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$, simulated with OSCAR 2.4.5, and later reconstructed with ORCA 8.7.1. Jets are reconstructed as localized energy depositions in the CMS calorimeters arranged in a projective tower geometry. The jet energy E is defined as the scalar sum of the calorimeter tower energies inside a cone of radius $R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.5$, centered on the jet direction. The jet momentum \vec{P} is the corresponding vector sum of energies, with the vector pointing in the tower direction. Both the jet energy and momentum are corrected back to the particles in the jet cone originating from the hard interaction excluding pileup [113]. We define the dijet system as the two jets with the highest p_T in an event (leading jets) and define the dijet mass $m = \sqrt{(E_1 + E_2)^2 - (\vec{P}_1 + \vec{P}_2)^2}$. We select events in which the leading jets each have $|\eta| < 1$. This cut enhances our sensitivity to new physics, produced at low $|\eta|$, compared to the predominantly t -channel processes from the QCD background. In all plots that are a function of dijet mass, we plot in bins of width equal to the Gaussian resolution measured in section 4.1.4.1.

4.1.2 Rates and Efficiencies from Jet Triggers

We use simulated data from the single jet triggers discussed in section E.4.3.2. From the three trigger tables for luminosities of $L = 10^{32}, 10^{33}, 10^{34} \text{ cm}^{-2}\text{s}^{-1}$ we expect initial samples of size at least $100 \text{ pb}^{-1}, 1 \text{ fb}^{-1},$ and 10 fb^{-1} respectively. This is from 10^6 seconds of collisions, equivalent to one month of continuous operation at 40% efficiency. In Fig. 4.1 we show the rate expected from these triggers as a function of dijet mass. By construction there are comparable events in each trigger, and a high statistics overlap between triggers for a given table. We see that the highest mass dijet is expected to be 5, 6 and 7 TeV for samples of size




Figure 4.1: Rate of jet trigger as a function of dijet mass. The 3 plots correspond to 3 trigger tables, and each plot shows multiple triggers with various p_T thresholds and prescales.

100 pb^{-1} , 1 fb^{-1} , and 10 fb^{-1} respectively. In Fig. 4.2 we show the trigger efficiency vs. dijet mass, measured for each trigger using the neighboring trigger with a lower p_T threshold, explicitly show the mass cuts that are fully efficient. In Fig. 4.3 we show the data we




Figure 4.2: Jet trigger efficiency (points) and fully efficient dijet mass cuts (lines).



use to measure the cross section. We use each trigger where it is fully efficient and stop





Figure 4.3: Rate of jet trigger for cross section measurement. Same triggers as Fig. 4.1.

using the trigger where the next trigger is fully efficient. Fig. 4.3 shows there are adequate numbers of fully efficient events for analysis.

4.1.3 Dijet Mass Distribution from QCD

In Fig. 4.4 we combine the triggers to produce a cross section across the full mass spectrum.  prescaled triggers allow us to measure mass down to 300 GeV, or even smaller if we





Figure 4.4: Cross section vs. dijet mass and the contributing jet triggers.

can understand the efficiency of the lowest threshold trigger. The mass measured with the prescaled triggers will allow us to connect to dijet masses measured at the Tevatron.

In Fig. 4.5 we show the fractional statistical error on the cross section, the simplest measure  our sensitivity to new physics. Fig. 4.5 shows that our prescaled triggers will allow a




Figure 4.5: Fractional statistical error on the jet cross section for the samples in Fig. 4.5.

measurement of QCD with 1-3% statistical accuracy. The unprescaled triggers will have 1% error at threshold and the first unprescaled sample begins at a mass of 670 GeV, giving us full sensitivity to new physics in a region that overlaps with previous dijet mass measurements at the Tevatron.

4.1.4 Searches using Dijet Mass

Here we will discuss the signal and background distributions that are needed for a dijet resonance search using the mass distribution. In section 14.5.2 we use these techniques to estimate our sensitivity to seven models of narrow dijet resonances.

4.1.4.1 Narrow Dijet Resonance Shapes

The simulated shape of a narrow dijet resonance in CMS is shown in Figure 4.6. The shape is composed of a Gaussian distribution from jet energy resolution and a long tail to low mass. The measured RMS of the Gaussian component is $\sigma/M = 0.045 + 1.3/\sqrt{M}$. The long tail to low mass comes predominately from final state QCD radiation (extra jets) which reduce the reconstructed mass. All resonances with a natural width significantly less than our resolution should look similar to this in the CMS detector. The model used in Figure 4.6 was a Z' from Pythia.

4.1.4.2 QCD Background to Dijet Resonances

Figure 4.6 compares a Z' signal cross section to the QCD background found in section 4.1.3. The differential cross section for the QCD background is well fit by a simple parameterization




Figure 4.6: Left) The dijet mass distribution from a 2 TeV/c^2 Z' (histogram) is fit with a Gaussian (solid curve) from the peak region to high mass and the Gaussian is extended to lower mass (dashed curve). Right) The differential cross section as a function of dijet mass for the QCD background and three Z' signals with a mass of 0.7, 2, and 5 TeV/c^2 .

of the form

$$\frac{d\sigma}{dm} = \frac{p_0(1 - m/\sqrt{s})^{p_1}}{m^{p_2}} \quad (4.1)$$

where m is the dijet mass, $\sqrt{s} = 14000 \text{ GeV}/c^2$ is the collision energy, and p_0, p_1, p_2 are arbitrary parameters. The resonance sensitivity estimates in section 14.5.2 use this parameterization to smooth away background fluctuations in our simulation sample. In a search with real data, a similar parameterization could be used to simply model the measured background, as was done by CDF [114], or a full NLO QCD calculation smeared with the jet resolution could be used to model the background, as was done by D0 [115].

4.1.5 Searches using Dijet Mass and Angle

Here we will discuss the signal and background distributions that are used for searches for new physics in the dijet mass and angular distribution simultaneously. This technique can be used to confirm resonances observed in the dijet mass distribution, and measure their spin, or to discover other new physics that could affect the dijet angular distribution. In section 15.3 we use these techniques to estimate our sensitivity to a model of quark contact interactions.

4.1.5.1 Dijet Ratio: $N(|\eta| < 0.5)/N(0.5 < |\eta| < 1.0)$

The ratio of the number of dijets in which both jets have $|\eta| < 0.5$ to the number of dijets in which both jets have $0.5 < |\eta| < 1.0$ was first introduced by D0 to search for contact interactions as a function of dijet mass [116]. It is the simplest measure of the most sensitive part of the angular distribution, providing a single number we can measure as a function of dijet mass. In Figure 4.7 we show our lowest order calculation of the dijet ratio from QCD compared with a left-handed contact interaction among quarks [117, 118] at three different values of the contact interaction scale. For this calculation we used the same code as [117] with modern parton distributions [29]. Lowest order QCD gives a fairly flat dijet ratio




Figure 4.7: Left) A lowest order calculation of the dijet ratio from QCD (solid curve) is compared with QCD plus a quark contact interaction at a scale Λ^+ of 15 TeV (dashed), 10 TeV (dotted) and 5 TeV (dot-dashed). Right) The dijet ratio in the CMS simulation from QCD (points) is compared to the value 0.6 (line).

around 0.6 while the contact interactions produce an increase in the dijet ratio at high mass. Figure 4.7 also shows that a full CMS detector simulation of the dijet ratio from QCD, using the samples discussed in section 4.1.3, is indistinguishable from a flat ratio of 0.6 within the simulation statistical uncertainty.

4.1.6 Systematic Uncertainties

In figure 4.8 we present estimates of systematic uncertainties on both the dijet cross section and the dijet ratio. The systematics discussed below have a large affect on the cross section and little affect on the dijet ratio.




Figure 4.8: Left) Systematic uncertainty on the dijet cross section due to jet energy scale (solid curve), parton distributions (dashed curve), and calorimeter energy and η resolution (dotted curve) are compared to the statistical uncertainties for 10fb^{-1} (error bars). Right) Systematic bounds on the dijet ratio from uncertainties in the relative jet energy scale (dashed curve), parton distributions (dotted curve), and calorimeter energy and eta resolution (dot dash curve), are compared to the expectations of QCD and three contact interaction scales (solid line and curves).

4.1.6.1 Absolute Jet Energy Scale

We have concluded that an overall uncertainty on the jet energy scale in the barrel of $\pm 5\%$ is achievable [120]. We have propagated this energy scale error to the dijet mass cross section by measuring the effect of a $\pm 5\%$ change in mass on a smooth fit to the dijet mass cross section. As shown in figure 4.8, the resulting upper uncertainty on the cross section varies from 30% at a dijet mass of $0.3\text{ TeV}/c^2$ to 80% at a dijet mass of 6.5 TeV . This large systematic uncertainty, increasing with dijet mass, is the primary reason we do not use the dijet mass distribution to search for quark contact interactions. For the dijet ratio the absolute jet energy scale uncertainty has no effect, because the dijet ratio is flat versus dijet mass. The uncertainty cancels out in the ratio.

4.1.6.2 Relative Jet Energy Scale

We have shown that by using dijet balance an uncertainty of $\pm 0.5\%$ is achievable [121] for the relative jet energy scale as a function of η within the barrel, in 0.1 steps in η . Here we assume that the relative jet energy scale, defined in this analysis as the uniformity in energy scale in the region $0.5 < |\eta| < 1.0$ compared to $|\eta| < 0.5$, can be determined to $\pm 0.5\%$. For the cross section as a function of mass this uncertainty is negligible compare to the $\pm 5\%$ error in the absolute energy scale. We have propagated this error to the dijet ratio by measuring the effect of a $\pm 0.5\%$ change in dijet mass for the measurement of $N(0.5 < |\eta| < 1)$ while keeping $N(|\eta| < 0.5)$ unchanged. As shown in figure 4.8, the resulting upper uncertainty in the ratio varies from 0.013 (2%) at a mass of $0.3\text{ TeV}/c^2$ to 0.032 (5%) at a mass of 6.5 TeV .

4.1.6.3 Resolution

The effect of calorimeter resolution is the difference between the measurement with jets constructed from MC particles (Gen Jets) and the measurement with jets constructed from calorimeter depositions and corrected (Rec Jets). This difference, often called the smearing due

to calorimeter resolution, is taken as a bound on the size of the systematic uncertainty due to resolution. For the cross section, the difference between Rec Jets and Gen Jets is small. This smearing varies from 15% at 0.3 TeV to 3% at 6.5 TeV, as shown in Figure 4.8. For the ratio, there is no change between Gen Jets and corrected Rec Jets within the Monte Carlo statistics presented in Fig. 4.7, and the statistical error on the simulation gives a bound on the systematic of 0.02 (3%) in the ratio, which is shown in Figure 4.8.

4.1.6.4 Parton Distributions

CTEQ 6.1 provides parton distributions with systematic uncertainties coming from the world data on parton distribution functions (PDFs) and the fit to world data [29]. CTEQ 6.1 has 40 different error PDFs, 20 PDFs at positive error, and 20 PDFs at negative error. We have used these 40 PDFs and the recommended procedure [29] to calculate the PDF uncertainties on both the cross section and the dijet ratio using our lowest order QCD calculation. As shown in figure 4.8, the resulting upper uncertainty in the cross section varies from 5% at a dijet mass of 0.3 TeV to 32% at a dijet mass of 6.5 TeV. As shown in figure 4.8, the resulting uncertainty in the dijet ratio peaks at a value of 0.02 (3%) in the ratio at a mass of around 3.5 TeV, and declines at both lower and higher masses.

4.1.6.5 Luminosity, Efficiency and Acceptance

The luminosity uncertainty on the cross section is around 10%, small compared to other uncertainties, and has no affect on the dijet ratio. For the masses we consider in this analysis there is full efficiency for finding a dijet in the event with negligible uncertainty. The acceptance for jets is defined by the cut in η , and any measured jet distributions must be compared to calculations using the same cuts, with negligible uncertainty in the comparison of measured and calculated jet η .



Benchmark Channel: Law Mass Supersymmetry



ME: fix for references, one wide table

ME: need to incorporate Gigi comments

4.2.1 Introduction

R -parity conserving SUSY leads to characteristic signatures with missing transverse energy in the final state due to the stable lightest supersymmetric particle (LSP). In the search described below for the bosonic partners of quarks (squarks) and the fermionic partners of gluons (gluinos) it is assumed that the LSP is weakly interacting, as is the case for most of the MSSM parameter space.

This analysis focuses on gluino and squark production within the minimal supergravity model (mSUGRA). In this model the entire SUSY mass spectrum is essentially determined by only five unknown parameters: the common scalar mass at the GUT scale, M_0 ; the common gaugino mass at the GUT scale, $M_{1/2}$; the common trilinear coupling at the GUT scale, A_0 ; the sign of the Higgsino mixing parameter, $sign(\mu)$; and the ratio of the Higgs vacuum expectation values, $\tan \beta$.

We investigate whether the production and decay of gluinos and scalar quarks is observable in the rate of ≥ 3 -jet events with large missing transverse energy. The large missing energy originates from the two LSPs in the final states of the squark and gluino decays. The three or more hadronic jets result from the hadronic decays of the squarks and/or gluinos. We use the ISAJET Monte Carlo (MC) program interfaced with PYTHIA which provides the coherent parton shower and underlying event model to generate squark and gluino production with parameters $M_0 = 60 \text{ GeV}/c^2$, $M_{1/2} = 250 \text{ GeV}/c^2$, $A_0 = 0$, $\mu > 0$ and $\tan \beta = 10$ (LM1 test point). For this set of parameters $m(\tilde{g}) > m(\tilde{q})$ and production of $\tilde{g}\tilde{q}$ is 53%, $\tilde{q}\tilde{q}$ 28% and $\tilde{g}\tilde{g}$ 12%. The decay $\tilde{g} \rightarrow q\tilde{L},R + q$ is dominant. Specifically the gluino and squark decays proceed as follows:

$$\tilde{g} \rightarrow q q\tilde{L},R, \quad \text{or} \quad \tilde{g} \rightarrow \tilde{q}\tilde{q}_{L,R} \quad (4.2)$$

$$\tilde{q}_R \rightarrow q\tilde{\chi}_1^0, (100\%) \quad (4.3)$$

$$\tilde{q}_L \rightarrow q + \tilde{\chi}_2^0, (30\%) \quad (4.4)$$

$$\tilde{q}_L \rightarrow q + \tilde{\chi}_1^+, (70\%) \quad (4.5)$$

$$(4.6)$$

while the charginos and neutralinos decay as follows:

$$\tilde{\chi}_2^0 \rightarrow \tilde{\ell}_R \ell, (11.2\%) \quad (4.7)$$

$$\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1 \tau, (46\%) \quad (4.8)$$

$$\tilde{\chi}_1^+ \rightarrow \tilde{\nu}_L \ell, (36\%) \quad (4.9)$$

The total LO production cross section for squarks and gluinos at this point of the mSUGRA parameter space is 49 pb. The major Standard Model background components for a multi-jet plus large missing transverse energy search include production of Z +jets with the Z decaying invisibly, W +jets, top-antitop pairs, dibosons, single top and QCD jets.

4.2.2 Jets and missing transverse energy at CMS

The CMS detector is described in detail in Ref. [7] The momenta of charged particles are measured in the silicon and pixel tracking devices which are positioned inside a 4 T superconducting solenoidal magnet. The central and end-cap electromagnetic and hadronic calorimeters (also within the magnet) are arranged in a semi-projective tower geometry, cover the pseudorapidity region $|\eta| < 3$ and are used to identify jets. Jets are defined as localized energy depositions in the calorimeters and are reconstructed using an iterative clustering algorithm with a fixed cone of radius $\Delta R \equiv \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.5$ in $\eta - \phi$ space [7]. Jets are ordered in transverse energy, $E_T = E \sin \theta$, where E is the scalar sum of energy deposited in the calorimeter towers within the cone, and θ is the angle formed by the beam-line, the event vertex, and the cone center. Jets with uncorrected $E_T > 30 \text{ GeV}$ and with $|\eta| < 3$ are used throughout this analysis.

The offline missing transverse energy is defined as the negative vector sum of the transverse energy in the electromagnetic and hadronic calorimeter towers, $E_T^{\text{miss}} = -\sum_i (E_i \sin \theta_i) \hat{n}_i$, where E_i is the energy of the i -th tower, \hat{n}_i is a transverse unit vector pointing to the center of each tower, and θ_i is the polar angle of the tower; the sum extends to $|\eta| < 5$. The data sample is selected with a hardware trigger which requires $E_T^{\text{miss},L1} > 46 \text{ GeV}$ ($|\eta| < 5$ coverage) and a central jet of $E_T > 88 \text{ GeV}$. A parameterization of the L1 trigger efficiency as measured in

Table 4.1: Cleanup Preselection efficiency.

Sample/Requirement	$F_{em} > 0.1$ (%)	$F_{ch} > 0.175$ (%)	Both(%)
LM1	99.88	91.32	91.24

a dijet sample is applied to all data analyzed. For the confirmation of the High Level Trigger (HLT) the E_T^{miss} is required to be above 200 GeV where the HLT trigger is fully efficient. In the following sections we detail the methodology and analysis strategies towards a search for SUSY using a dataset of events collected according to the missing transverse energy plus jet Level-1 and HLT trigger path.

4.2.3 Clean-up requirements

In anticipation of real data a preselection is used to reject accelerator- and detector-related backgrounds (such as beam halo and noise), and cosmic ray events. At least one primary vertex is required in the event and the preselection uses the event electromagnetic fraction, F_{em} (defined as the E_T -weighted jet ElectroMagnetic Fraction sum over the electromagnetic calorimeter acceptance, $|\eta_d| \leq 3.0$) and event charged fraction, F_{ch} (defined as the average ratio of the jets calorimetric transverse energy over the sum of the E_T of the associated to the jet tracks for jets within $|\eta| < 1.7$) to distinguish between real and fake jet events. The preselection requirements and their efficiency on the signal are shown in table 4.1. The values of the requirements are chosen based on the Tevatron data where similar requirements have been used to clean the high p_T multijet plus large missing transverse energy datasets from a number of spurious and instrumental backgrounds that tend to appear as spikes in the low end of the event electromagnetic and charge fraction distributions.

4.2.4 Analysis Path

Events that are accepted by the preselection requirements, proceed through the analysis path if they have missing transverse energy $E_T^{miss} > 200$ GeV and at least three jets with $E_T \geq 30$ GeV within $|\eta| < 3$. In addition the leading jet is required to be within the central tracker fiducial volume *i.e.* $|\eta| < 1.7$. These requirements directly define the searched for signal signature. The rest of the analysis path is designed based on elimination of the major classes of backgrounds: the QCD production, top-antitop pairs and the W/Z -QCD associated production. In table 13.5 the path is shown with a remark indicating the reason and aim of each selection step.


In the following sections the motivation and details of the analysis path are discussed.

4.2.5 Missing transverse energy in QCD production

Due the very high QCD production cross section the Standard Model background to a large missing transverse energy plus jets data-sample is dominated by QCD events. The observed missing transverse energy in QCD jet production is largely a result of jet mismeasurements and detector resolution. In figure 4.9 the missing transverse energy full spectrum is shown for QCD 3-jet events in the P_T region between 120 GeV and 1.8 TeV.

Table 4.2: The E_T^{miss} + multijet SUSY search analysis path

Requirement	Remark
Level 1	L1 trigger efficiency parameterization
HLT, $E_T^{\text{miss}} > 200$ GeV	trigger/signal signature
primary vertex ≥ 1	primary cleanup
$F_{em} \geq 0.175, F_{ch} \geq 0.1$	primary cleanup
$N_j \geq 3, \eta_d^{1j} < 1.7$	signal signature
$\delta\phi_{\min}(E_T^{\text{miss}} - jet) \geq 0.3$ rad, $R_1, R_2 > 0.5$ rad, $\delta\phi(E_T^{\text{miss}} - j(2)) > 20^\circ$	QCD rejection
$I_{SO}^{\text{lead trk}} = 0$	ILV (I) $W/Z/t\bar{t}$ rejection
$f_{em(j(1))}, f_{em(j(2))} < 0.9$	ILV (II), $W/Z/t\bar{t}$ rejection
$E_{T,j(1)} > 180$ GeV, $E_{T,j(2)} > 110$ GeV	signal/background optimization
$H_T > 500$ GeV	signal/background optimization
SUSY LM1 signal efficiency 13%	

Figure 4.9: E_T^{miss} distribution in QCD 3-jet events.

It is to be noted that due to finite computing resources and the large production cross section it is unrealistic to fully simulate and reconstruct samples with adequate Monte Carlo statistics. It is also unrealistic due to the trigger and data acquisition bandwidth constraints to collect QCD datasets with low E_T thresholds during data-taking. However the CMS trigger table includes a large number of prescaled QCD trigger-paths that will be used to extract the shape of the missing transverse energy and the direct normalization for the QCD background component in all-hadronic events with large missing energy. In addition, topological requirements are designed to eliminate as possible the QCD contribution. In a QCD multijet event with large missing energy, the highest E_T jet is typically the most accurately measured. When the second or third jet is mismeasured, the E_T^{miss} direction is pulled close in ϕ to the mismeasured jet direction. We eliminate the residual QCD component by using the correlation in the $\delta\phi_1 = |\phi_{j(1)} - \phi(E_T^{\text{miss}})|$ versus $\delta\phi_2 = |\phi_{j(2)} - \phi(E_T^{\text{miss}})|$ plane, as shown in figure 4.10. Events with $R_1 > 0.5$ rad and $R_2 > 0.5$ rad, where $R_1 = \sqrt{\delta\phi_2^2 + (\pi - \delta\phi_1)^2}$ and $R_2 = \sqrt{\delta\phi_1^2 + (\pi - \delta\phi_2)^2}$, are accepted. In addition we require that no jet in the event be closer than 0.3 rad to the missing energy direction and that the second jet be further than 20° from

Figure 4.11).




Figure 4.10: $\delta\phi_1$ versus $\delta\phi_2$ for (left) SUSY signal and (right) QCD dijet events





Figure 4.11: $\delta\phi_2 = |\phi_{j(2)} - \phi(E_T^{miss})|$ for (left) SUSY signal and (right) QCD dijet events

After a baseline selection of $N_j \geq 2$ and $E_T^{miss} > 93$ GeV the cumulative efficiency of the angular requirements is $\sim 90\%$ for the SUSY signal. They reject $\sim 85\%$ of all QCD dijet events.

4.2.6 Indirect Lepton Veto

In this analysis there is no explicit lepton identification. Leptons in the signal SUSY events result from cascade decays of squarks and gluinos through charginos and neutralinos. To reduce the large background contribution mainly from $W(\rightarrow \ell\nu) + jets$ and $t\bar{t}$ production and decays, an *indirect lepton veto* scheme is designed. The aim of the indirect lepton veto is twofold: a) to retain large signal efficiency b) to achieve large rejection of the $W, Z, t\bar{t}$ backgrounds (independent of the MC used, namely parton shower only versus complete matrix element in particular for the higher jet multiplicity bins).

Given that electrons are also clustered as jets, the jet electromagnetic fraction, f_{em} , which is close to 1 for electrons, is efficient in rejecting background events containing electrons while retaining good efficiency in the LM1 SUSY inclusive signal. Events are selected if the two highest E_T jets are not be purely electromagnetic, *i.e.* $f_{em,j(1)} < 0.9$ and $f_{em,j(2)} < 0.9$. The leading and second jet electromagnetic fraction distributions for $W \rightarrow e\nu + \geq 2$ jets are shown in figure 4.12. The corresponding distributions for the SUSY LM1 signal are shown

in figure 4.13. The signal efficiency is $\sim 87\%$ while 90% of the $W \rightarrow e\nu + \geq 2$ jets are rejected. A systematic uncertainty of 5% on the background rejection efficiency is assigned due to a ation between PYTHIA and ALPGEN +PYTHIA samples.





Figure 4.12: Electromagnetic fraction of (left) leading and (right) second jet in $W \rightarrow e\nu + \geq 2$ events.




Figure 4.13: Electromagnetic fraction of (left) leading and (right) second jet in SUSY LM1 events

To further reject electrons, muons and taus from W and Z decays while retaining the SUSY signal efficiency a tracking isolation strategy is employed as follows: if the leading track in the event has $p_T \geq 15$ GeV and the ratio of the sum of the p_T of all tracks around it in a cone of $\Delta R=0.35$ over the p_T of the track is less than 10% the event is dropped. The requirement of accepting events with a non-isolated leading track is noted as $\text{Iso}^{\text{trk}}=0$.

The leading isolated track veto has $\sim 92\%$ signal efficiency while it rejects $\sim 50\%$ of the W/Z +jets events (in PYTHIA as well as ALPGEN generated samples). The cumulative W/Z +jets rejection efficiency when both requirements of the indirect lepton veto are applied is between 50% and 90% depending on the lepton flavor with lower rejection as expected when the boson decay product includes a τ lepton. When applied in the full analyses path it rejects 40% of $t\bar{t}$ inclusive events. The cumulative SUSY signal efficiency is $\sim 80\%$.

4.2.7 The Standard Z boson candle calibration

Events with large missing transverse energy and ≥ 3 jets in the final state are expected from $Z(\rightarrow \nu\bar{\nu}) + \geq 3$ jets and $W(\rightarrow \tau\nu) + \geq 2$ jets (the third jet originating from the hadronic τ

decay) processes. Additional residual contribution is expected also from $W(\rightarrow \mu\nu)$, $e\nu + \geq 3$ jets. The Monte Carlo predictions for events with ≥ 3 jets can be normalized to the observed $Z(\rightarrow \mu\mu) + 2$ jets data sample (which has a larger cross section) via the measured $\frac{N_{jet}}{N_{jet+1}}$ ratio, where N_{jet} is the number of jets with $\sim 1 \text{ fb}^{-1}$ of data.

The ratio $\rho \equiv \frac{\sigma(pp \rightarrow W(\rightarrow \mu\nu) + jets)}{\sigma(pp \rightarrow Z(\rightarrow \mu^+\mu^-) + jets)}$ can be used to normalize the W +jets MC predictions. Assuming lepton universality, the predictions for the number of events with ≥ 2 - and ≥ 3 -jets from W and Z production and decays to all flavors can be normalized to the $Z(\rightarrow \mu^+\mu^-) + \geq 2$ jets data. By normalizing the MC predictions to data we will avoid large systematic effects due to the renormalization scale, the choice of parton density functions, initial- and final-state radiation, and the jet energy scale. The total uncertainty ($\sim 5\%$) is then dominated by the uncertainty on the luminosity measurement, the uncertainty on the measured ratio $\frac{N_{jet}}{N_{jet+1}}$ (now unknown), and the uncertainty on the ratio ρ as a function of N_{jet} . It is to be noted that the method can be used to absolutely normalize the Monte Carlo predictions for $Z(\rightarrow \nu\bar{\nu}) + \geq 3$ jets assuming that after detector simulation they can be tuned to reproduce the ratios mentioned above as well as the kinematic distributions observed in the ‘‘candle’’ data sample, or the actual data ‘‘candle’’ sample can be used stand-alone to predict the rate and event kinematics of the $Z(\rightarrow \nu\bar{\nu}) + \geq 3$ jets process.

In this study the $Z \rightarrow \mu\mu + \geq 2$ jets with $Z_{PT} > 200 \text{ GeV}$ is the ‘‘candle’’ data sample. Both the muon and electron decays of the Z can and will be used as the standardizable candle, but for the purposes of demonstrating the method, the Z muon decays are chosen. The additional advantage of the muon channel is the supreme CMS muon detection due to the tracking and muon systems. Since the completely raw missing transverse energy is used (as is expected to be the case at the start-up of the experiment), the shape of the E_T^{miss} distribution in the measured using a the $Z \rightarrow \mu\mu + \geq 2$ jet events will be very close to the shape of the invisible $Z \rightarrow \nu\nu + \geq 2$ jet events as shown in figure 4.14. The muon decays of the Z are selected from an inclusive sample using the following requirements as baseline selection: a) at least one primary vertex, b) at least 2 jets with $E_T \geq 30 \text{ GeV}$, and $|\eta_d| \leq 3$ c) $E_T^{miss} > 200 \text{ GeV}$ and d) for the Z boson identification two reconstructed muons with invariant mass closest to the measured Z boson mass ($91.2 \text{ GeV}/c^2$) and within $20 \text{ GeV}/c^2$. The ‘‘Z-mass’’ tag requirement is 90% efficient. The selected candle sample dimuon invariant mass is shown in figure 4.15 overlaid with the one using the Monte Carlo truth.

Note that a comparison of the Monte Carlo truth boson E_T , the boson E_T using Monte Carlo truth E_T vectorial sum of the muons, the vectorial sum of the E_T of the two reconstructed muons in the Z -mass window, and the E_T^{miss} in the candle sample are all in agreement as expected, and in agreement with the E_T^{miss} of the $Z \rightarrow \nu\nu + \geq 2$ jet sample. Considering both the electron and muon decays of the Z boson, a statistically adequate (5% precision) ‘‘candle’’ sample that can be used to normalize the $Z \rightarrow \nu\nu + \geq 2$ jet predictions for $E_T^{miss} > 200 \text{ GeV}$ will be obtained with $\sim 1.5 \text{ fb}^{-1}$.

4.2.8 Analysis Results

The signal to background ratio is further enhanced in the final steps of the analysis (shown in table 13.5) by requiring the two leading jets E_T be above 180 and 110 GeV respectively. Furthermore the H_T in the event is required to be $H_T \equiv E_{T(2)} + E_{T(3)} + E_{T(4)} + E_T^{miss} > 500 \text{ GeV}$. The global signal efficiency for the analysis is 13% while the signal to background ratio is ~ 26 . The results are shown in table 13.6.




Figure 4.14: E_T^{miss} in $Z \rightarrow \mu\mu + \geq 2$ jets candle sample and normalized E_T^{miss} in $Z \rightarrow \nu\bar{\nu} + \geq 2$ jets sample.

Table 4.3: Selected SUSY and Standard Model background events for 1 fb^{-1}

Signal	$t\bar{t}$	single t	$Z(\rightarrow \nu\bar{\nu}) + \text{jets}$	$(W/Z, WW/ZZ/ZW) + \text{jets}$	QCD
6319	53.9	2.6	48	33	107

Due to the QCD Monte Carlo limited statistics to derive the QCD background component the analysis path is run without the topological QCD clean-up requirements and ILV requirements. The estimate is conservative and is based on factorizing the clean-up and efficiency and assuming them uncorrelated with the rest of the analysis requirements. **FIXME:** One sentence on parameterizing topological cuts efficiency as a function of MET – will reduce the QCD tails.

4.2.9 Systematic Uncertainties

4.2.9.1 E_T^{miss} shape systematic uncertainty due to tails in the jet resolution

A bootstrap-like study is performed to estimate the positive systematic uncertainty of the E_T^{miss} due to the non Gaussian tails in the jet resolution. The study uses the inclusive $t\bar{t}$ sample. The events are re-weighted according to a grading of the mismeasured jets, and on a jet-by-jet basis. The grading of a jet being considered mismeasured is derived from the jet resolution shape of jets in three p_T bins. Jets are considered mismeasured when they fall in the non-Gaussian tails of the jet resolution. The event weight is derived using each jet's weight and for three different scenarios that involve one, two or three jets being simultaneously mismeasured and positively contributing to the enhancement of the E_T^{miss} tail.

FIXME: explain the 15% studies as per IN-NOTE

The ratio of the E_T^{miss} distribution resulting from the one, two and three under-measured jets scenarios study over the nominal E_T^{miss} is shown in figure 4.16 and it shows graphically




Figure 4.15: Reconstructed and generator level Z di-muon invariant mass for $Z \rightarrow \mu\mu + \geq 2$ jets and $E_T^{miss} > 200$ GeV.

the positive systematic uncertainty band as a function of the E_T^{miss} due to jet tails in the resolution.

The positive systematic uncertainty due to one mismeasured jet in the high E_T^{miss} tails is estimated over the bins where in the nominal distribution we have enough statistics, namely between 180 and 240 GeV (statistical uncertainty $< 5\%$). The result is 8.5%. For the scenario with the two undermeasured jets, and assuming that 50% of the times the simultaneous under-measurement results in the overestimate of the E_T^{miss} the result is 6% and for the case of the three under-measured jets it is also 6%. We take the weighted average of these three scenarios, namely 7%, as an index of the positive systematic uncertainty due to the tails of the jet resolution in the tails of the E_T^{miss} above 180 GeV. The result in the method presented is bound to overestimate the increase in the tails, since by design positive interference of all under-measured jets in the event is considered (in reality there is some combinatorial compensation in the E_T^{miss} vector given the jet topology). It is noted that the ultimate measurement of the shape of the high E_T^{miss} tails and its systematic should be done using a candle physics process in the real data such as the Z -jets and the $t\bar{t}$ data sample.

4.2.9.2 Jet Energy Scale

To determine the effect of the JES uncertainty each jet four-vector is scaled with the uncertainty value α as follow:

$$\begin{aligned}
 p_{scaled}^{\mu,jet} &= (1 \pm \alpha) \cdot p_{meas}^{\mu,jet} \\
 &= (1 \pm \alpha) \cdot (p_x, p_y, p_z, E)
 \end{aligned}
 \tag{4.10}$$

The JES uncertainty for the high E_T jets that enter this analysis is taken to be about 7% for 1 fb^{-1} ; The resulting uncertainty in $t\bar{t}$ and QCD events is 22%.




Figure 4.16: Ratio of E_T^{miss} weighted distribution for one, two and three under-measured jets (described in the text) over the corresponding nominal E_T^{miss} distribution.

4.2.9.3 Luminosity uncertainty

Since the W/Z +jets background is taken to be normalized with real data, the estimate carries the luminosity uncertainty on it. Hence a $\pm 5\%$ uncertainty is taken on the background estimates due to the luminosity measurement.

4.2.9.4 ALPGEN- PYTHIA ILV

As discussed in section 4.2.6 a 5% positive systematic on the background estimate is taken due to the variation in efficiency of the ILV requirement between ALPGEN and PYTHIA.

4.2.9.5 Total background systematic

In summary for the major background components the uncertainties are as follows:

- $t\bar{t}$ uncertainties: 7% E_T^{miss} shape, 22% JES, 13% statistical
- $Z \rightarrow \nu\bar{\nu}$ +jets, W/Z +jets: 5% Luminosity (direct candle normalization to the data)
- QCD: E_T^{miss} 7% shape, 22% JES, 10% statistical

The number of background events per background component and their uncertainties are tabulated in Table 13.7.

4.2.10 Discussion


In conclusion, based on the Standard Model background estimates and their uncertainties, a 5σ observation of low mass SUSY at LM1 (gluino mass $600 \text{ GeV}/c^2$ is achievable with ~ 6

Table 4.4: Standard Model background components and uncertainties for 1 fb^{-1}

$t\bar{t}, \text{single}top$	$Z(\rightarrow \nu\bar{\nu}) + \text{jets}$	$(W/Z, WW/ZZ/ZW) + \text{jets}$	QCD
$56 \pm 11(\text{sys}) \pm 7.5(\text{stat})$	$48 \pm 3.5 (\text{all})$	$33 \pm 2.5 (\text{all})$	$107 \pm 25(\text{sys}) \pm 10(\text{stat})$

pb^{-1} in events with large missing energy plus multijets. It is found that with $\sim 1.5 \text{ fb}^{-1}$ the $W/Z+\text{jets}$ background including the invisible decays of the Z boson which constitutes a large irreducible background component can be reliably normalized using the $Z \rightarrow \mu\mu$ and $Z \rightarrow ee + \text{multijet}$ data candle. Furthermore the global raw E_T^{miss} measurement from the calorimeter towers can be calibrated for multijet topologies using the superb tracking and muons systems of CMS and the $Z \rightarrow \mu\mu + \text{multijet}$ candle data sample. This analysis demonstrates that the E_T^{miss} measurement from the calorimeter towers can be used as such at the startup of the experiment provided that adequate strategies are in place to discard spurious instrumental backgrounds. It is also found that an indirect lepton veto makes possible the $t\bar{t}$ and $W/Z+\text{jets}$ background rejection, without compromising the inclusive nature of the search. In anticipation of data, there is no accurate way of accurately predicting the contribution of the QCD background tails; Although the ME Monte predictions are to-date far more complete, the experiment has in place proper prescaled QCD triggers in order to estimate this background component using directly the data.

Finally the comparison of the signal, total background estimated and its components for the $E_T^{\text{miss}}, H_T, N_{jet}$ and $M_{eff} \equiv E_{T(1)} + E_{T(2)} + E_{T(3)} + E_{T(4)} + E_T^{\text{miss}}$ are shown in figures 4.18 and 4.20. It is to be underlined that the slopes of the tails of the missing energy, H_T , and M_{eff} distributions are very similar between the Standard Model background and the SUSY

Figure 4.17: E_T^{miss} LM1 signal and Standard Model background distributions.




Figure 4.18: H_T LM1 signal and Standard Model background distributions.




Figure 4.19: Jet Multiplicity LM1 signal and Standard Model background distributions.




Figure 4.20: M_{eff} LM1 signal and Standard Model background distributions.

Chapter 5

Physics Studies with tracks, B mesons, and taus



Benchmark Channels: Study of the decay $B_s \rightarrow J/\psi\phi$

5.1.1 Introduction

The decay $B_s \rightarrow J/\psi\phi \rightarrow K^+K^-\mu^+\mu^-$ is of particular interest, since it allows to study many properties of the B_s system, such as the differences between the widths and the masses of the two weak eigenstates, B_s^H and B_s^L . Contrary to the B^0 system, the difference between the widths $\Delta\Gamma_s$ of the two weak eigenstates is expected to be large, with a relative difference $\Delta\Gamma_s/\Gamma_s$ predicted to be in the order of 10% in the Standard Model. The first measurement from CDF ($\Delta\Gamma_s/\Gamma_s = (65_{-33}^{+25} \pm 1)\%$ [122]) and the new preliminary result from DØ ($\Delta\Gamma_s/\Gamma_s = (15 \pm 10_{-4}^{+3})\%$ [123]) have large discrepancies between the two measured values themselves and with the Standard Model prediction. It is only very recently that a first measurement of the mass difference, Δm_s , has been performed at CDF. Time-integrated measurements are not possible, as the time-integrated mixing probability χ saturates at a value of 0.5 for large mass differences, and in time-dependent measurements, the high mass difference generates very rapid oscillations. As in the B_s system the ratio $\Delta m_s/\Delta\Gamma_s$ depends on the ratio $|V_{cb}V_{cs}|/|V_{tb}V_{ts}|$, which is quite well known, and on QCD corrections, a measurement of $\Delta\Gamma_s$ would therefore yield an independent measurement of Δm_s . With the measurement already performed in the B^0 system, the ratio between the mixing parameters of the B^0 and B_s could provide a measurement of the ratio $|V_{ts}|/|V_{td}|$.

Furthermore, this decay provides one of the best ways to determine the height of the Unitarity Triangle, η in the Wolfenstein parameterization. At first order of the Wolfenstein parameterization, the CP -violating weak phase $\phi_{CKM} = [\arg(V_{cs}^*V_{cb}) - \arg(V_{ts}^*V_{tb})]$, measured in the rate asymmetry, cancels, and higher order terms have to be taken, yielding a weak phase $\phi_{CKM} = 2\lambda^2\eta$. The weak phase is therefore expected to be very small, of the order of 0.03. The measurement of a significantly larger phase would indicate contributions from non-Standard Model processes.

Because of the relative orbital angular momentum between the decay products, the $J/\psi\phi$ final state is an admixture of CP -even and CP -odd states, and the total rate asymmetry suffers from a partial cancellation. As the CP -even and CP -odd components have different angular dependences, an analysis of the angular correlation of the decay will permit to separate the two states, thereby permitting to access the different parameters.

With a total B production cross section at $\sqrt{s} = 14$ TeV expected to be as high as $500 \mu\text{b}$, a substantial number of fully reconstructed B_s candidates can be expected. Nevertheless, a high background has to be contended with. The main sources of backgrounds identified are those containing a J/ψ decaying to two muons susceptible to satisfy the Level 1 trigger requirements.

The decay $B_s \rightarrow J/\psi\phi$ is chosen as a benchmark channel since it is representative of exclusive B -physics studies. It allows to study the capability of CMS to identify, select and reconstruct a fully reconstructed decay of the B_s , which presents a significant challenge due to its relatively low momentum and high background. In addition, the measurement of the width difference $\Delta\Gamma_s$ on a sample of untagged $B_s \rightarrow J/\psi\phi \rightarrow \mu^+\mu^-K^+K^-$ candidates using a maximum likelihood fit of the time dependent angular distribution can be attempted.

An analysis on tagged candidates is not yet possible, as flavour tagging tools need to be implemented in the CMS reconstruction software.

5.1.2 Event generation

In addition to the signal itself, the main backgrounds identified have been simulated with low luminosity pile-up ($\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$). Kinematic requirements were applied in order to ensure that a significant fraction of the generated events would fulfill the Level-1 trigger requirements and that the final state particles are within the acceptance of the tracker ($|\eta| < 2.5$). The transverse momentum of the muons is thus required to be above $3 \text{ GeV}/c$ for muons in the barrel ($|\eta| < 1.2$) and $2 \text{ GeV}/c$ elsewhere. For the signal, the momenta of the kaons is required to be above $0.8 \text{ GeV}/c$.

For the samples composed of events with decays of B hadrons, $b\bar{b}$ pairs were generated with PYTHIA 6.215 with the MSEL=1 card in order to correctly reproduce the three different contributions to the total cross section (parton fusion, flavour excitation, and gluon splitting). The fragmentation of the b quark is performed by PYTHIA and the subsequent decay of the B hadron is performed using the SIMUB generator [124], a dedicated B physics event generator.

The decay $B_s \rightarrow J/\psi\phi$ has to be performed with SIMUB, since PYTHIA does not take into account the angular distributions of the final decay products. One of the b quarks in the event is forced to hadronize to a B_s^0 or \bar{B}_s^0 meson and to decay through the complete decay chain. With the kinematic requirements, using the world-average branching ratios for the decays of the B_s , J/ψ and ϕ mesons[45], the cross section is predicted to be $\sigma(B_s^0 \rightarrow J/\psi\phi \rightarrow \mu^+\mu^-K^+K^-) = 74 \pm 27 \text{ pb}$.

The most important background for the measurement is foreseen to be the inclusive decays of B hadrons to final states with a J/ψ resonance. These were simulated using PYTHIA, since no detailed simulation of angular distributions of the final decay products is needed. In order to increase the number of events similar to the signal events, a pair of oppositely charged particles with $p_T > 0.5 \text{ GeV}/c$ and $|\eta| < 2.5$ forming a fake ϕ candidate is required for in a region ($|\Delta\eta| < 1.5$, $|\Delta\varphi| < 1.5$) around the J/ψ direction and with an invariant mass within $30 \text{ MeV}/c^2$ of the world-average ϕ mass. In addition, this fake ϕ candidate is required to form a fake B_s^0 candidate with an invariant mass within $300 \text{ MeV}/c^2$ of the world-average B_s^0 mass. The cross section, including the kinematic requirements and branching-fractions, is estimated to be $\sigma(b \rightarrow J/\psi X) = 3.20 \pm 0.3 \text{ nb}$.

Furthermore, a sample of $B^0 \rightarrow J/\psi K^{*0} \rightarrow \mu^+\mu^-K^+\pi^-$ events were simulated, since this final state can be misidentified as a $B_s \rightarrow J/\psi\phi$ decay. In addition, this decay has a similar

Table 5.1: Values used for the mixing parameters, decay amplitudes, strong and weak phases in the simulation of the $B_s \rightarrow J/\psi \phi$ and $B^0 \rightarrow J/\psi K^{*0}$ Monte Carlo sample.

Parameter	$B_s \rightarrow J/\psi \phi$	$B^0 \rightarrow J/\psi K^{*0}$
$\tau = 1/\Gamma$	1.405×10^{-12} s	1.528×10^{-12} s
$\Delta\Gamma/\Gamma$	-0.2	0
ΔM	17.8 ps^{-1}	0.509 ps^{-1}
$ A_0(0) ^2/\Gamma$	0.570	0.570
$ A_{\parallel}(0) ^2/\Gamma$	0.217	0.217
$ A_{\perp}(0) ^2/\Gamma$	0.213	0.213
δ_1	π	π
δ_2	0	0
ϕ	-0.04	0

differential decay rate [125, 126] to the studied B_s decay. The B^0 decay is simulated with SIMUB, where one of the b quarks in the event is forced to hadronize to a B^0 or \bar{B}^0 meson, and to decay through the complete decay chain. With the kinematic requirements, and using the world-average branching ratios, the cross section is predicted to be $\sigma(B^0 \rightarrow J/\psi K^{*0} \rightarrow \mu^+ \mu^- K^+ \pi^-) = 366 \pm 22 \text{ pb}$.

The uncertainties quoted on the estimates above do not include the uncertainties on the total $b\bar{b}$ cross section at LHC energies, the b fragmentation functions, the transverse momentum distribution of b quarks, and the uncertainties introduced by using the model of $b \rightarrow J/\psi X$ decays in PYTHIA. However, since both the signal and background are proportional to the same $b\bar{b}$ cross section, the signal-to-background ratio is unaffected by the latter uncertainty. The parameters used in the simulation of the $B_s \rightarrow J/\psi \phi$ and $B^0 \rightarrow J/\psi K^{*0}$ decays are given in Table 5.1.

The direct production of J/ψ mesons is an important background at trigger level. Measurements at the Tevatron [127] have shown that predictions of the color-singlet model, which is presently the one implemented in the PYTHIA generator, underestimate the measurements by several orders of magnitude. Perturbative QCD is used in this model to generate $c\bar{c}$ pairs, which then hadronize to a charmonium state in a non-perturbative way.

The observed discrepancy has led to a different approach, where $c\bar{c}$ pairs are first formed taking into account all perturbative QCD diagrams, regardless of the final color state. The $c\bar{c}$ state is then transformed into a color-singlet by non-perturbative processes, such as the emission of a soft gluon.

A modified version [128] of PYTHIA 6.225, tuned on Tevatron data, has been used to simulate a sample of J/ψ decaying to two muons for background studies. The J/ψ production cross section is calculated to be $141 \mu\text{b}$. Taking the $J/\psi \rightarrow \mu^+ \mu^-$ branching ratio and the kinematic requirements into account, a cross section of $310 \pm 5 \text{ nb}$ is expected. Only the statistical uncertainty is quoted and used; the large uncertainties on total cross section for J/ψ production on p_T distribution are not included.

5.1.3 Trigger Selection

5.1.3.1 The Level-1 trigger

The B_s decay chain is selected at Level 1 by the di-muon trigger stream. At low luminosity it is foreseen [68] that it will be possible to use a symmetric threshold of $3 \text{ GeV}/c$ on the transverse momentum of the two muons, still keeping a low bandwidth occupancy of 0.9 kHz . Such a low p_T threshold ensures a very high selection efficiency on this channel, with a rate low-enough to allow the use of lower quality muon candidates in the endcap region, recovering full geometrical acceptance of the muon detector up to $|\eta| < 2.4$. For this decay, two of the identified muons are required to have opposite charge.

5.1.3.2 The High-Level trigger

In the HLT, the signal events will be identified by doing a full reconstruction of the B_s decay, imposing invariant mass and vertex constraints. Indeed, at this stage, tracks can be reconstructed in the tracker in restricted (η, ϕ) regions via a partial reconstruction algorithm, where only the first 5 hits are used [7, Section 6.5.3.2]. To define the tracking regions, the primary (interaction) vertex is first identified and reconstructed using only hits in the Pixel detector, with the ‘‘Divisive Method’’ described in reference [129]. Since the primary vertex of $b\bar{b}$ events involve low momentum tracks, the three vertex candidates with the highest sum of the p_T^2 of the tracks, which is the default selection criterion, have to be retained to have a good efficiency.

For the muons, the tracking regions is chosen around the direction of the muons identified at Level-1. Since no link to the muon detectors can be done at this stage, all track pairs of opposite charge for which the invariant mass is within $150 \text{ MeV}/c^2$ of the world-average J/ψ mass are retained. The resolution on the invariant mass of the J/ψ meson is found to be $51 \text{ MeV}/c^2$. In addition, the p_T of each muon is required to be above $2.5 \text{ GeV}/c$ in $|\eta| < 1.2$ or $2 \text{ GeV}/c$ in $|\eta| \geq 1.2$, and the p_T of the J/ψ candidate above $4 \text{ GeV}/c$. To remove the prompt J/ψ background, the two muon candidates are then fitted to a common decay vertex. The χ^2 of the fit is required to be below 10 and the significance of the transverse decay length is required to be above 3. Furthermore, the transverse momentum of the J/ψ candidate is required to be nearly parallel to its flight path in the transverse plane, since the J/ψ mesons produced in the decays of B_s mesons are collimated around the direction of the B_s meson by the relativistic boost. The cosine of the angle between the reconstructed momentum vector and the vector pointing from the production to the decay vertex is thus required to be less than 0.9.

To reconstruct the kaons, the tracking region is chosen around the direction of each J/ψ candidate. Assigning the kaon mass to the reconstructed tracks, all oppositely charged track pairs for which the invariant mass is within $20 \text{ MeV}/c^2$ of the world-average mass of the ϕ meson are retained, for a resolution on the invariant mass of the ϕ meson of $4.5 \text{ MeV}/c^2$. The p_T of each of the kaon tracks is required to be above $0.7 \text{ GeV}/c$, the p_T of the ϕ candidate above $1 \text{ GeV}/c$ and the p_T of the B_s candidate above $5 \text{ GeV}/c$. With the two muon candidates, the four-track invariant mass is required to be within $200 \text{ MeV}/c^2$ of the world-average mass of the B_s meson. The resolution on the invariant mass of the B_s meson is found to be $65 \text{ MeV}/c^2$. Here as well, a vertex fit of the four tracks is performed, imposing similar requirements as above.

The distribution of the invariant mass of the candidates after the HLT requirements is shown in Figure 5.1 (left). The efficiencies for the different criteria, which include the respective

Table 5.2: Trigger selection efficiencies for the signal and background (defined with respect to the number of generated events) after each requirement, and estimated HLT rate.

Requirement	Signal $B_s \rightarrow J/\psi \phi$	Background		
		Prompt J/ψ	$b \rightarrow J/\psi X$	$B^0 \rightarrow J/\psi K^{*0}$
Level-1	45.76(6)%	36.91(12)%	38.25(13)%	46.91(13)%
HLT - J/ψ selection	28.69(7)%	0.65(2)%	21.91(11)%	30.28(12)%
HLT - ϕ selection	20.50(6)%	0.0007(7)%	1.23(3)%	0.961(26)%
HLT rate (Hz)	0.03034(8)	0.002(2)	0.0792(18)	0.0077(2)

reconstruction efficiencies, are given in Table 5.2 for the signal and the different background samples, together with the estimated rate. The total rate for this selection is well below 1 Hz, a yield of approximately 456'000 signal events can be expected within 30 fb^{-1} of data.

5.1.4 Offline Selection and Reconstruction

The first step in the offline selection is similar to HLT selection, with the difference that the complete information from the detector is available. Candidates are reconstructed by combining two muons of opposite charge with two further tracks of opposite charge. As CMS does not possess a particle identification system suitable for this measurement, all measured tracks have to be considered as possible kaon candidates, which adds a substantial combinatorial background. At this stage, only loose requirements are applied, which are tightened after a kinematic fit.

First, all muons in the event are reconstructed using the global muon reconstruction algorithm [7, Section 9.1.3]. This algorithm is not fully efficient for low- p_T muons from J/ψ decays, being more suited to the reconstruction of high- p_T muons. Therefore, all tracks are reconstructed with the standard track reconstruction algorithm [7, Section 6.5]. Track-pairs of opposite charge for which the invariant mass is within $120 \text{ MeV}/c^2$ of the world-average J/ψ mass are retained as a J/ψ candidate. The p_T of each muon is required to be above $3 \text{ GeV}/c$ in $|\eta| < 1.2$ or $2 \text{ GeV}/c$ in $|\eta| \geq 1.2$, and the p_T of the J/ψ candidate above $4 \text{ GeV}/c$. The muon identification algorithm which uses information from the muon detector [7, Section 9.2.1.2], is applied to both tracks forming the J/ψ candidate. A J/ψ candidate is confirmed if both tracks share more than half of their hits in the silicon tracker with the muon tracks reconstructed by the global muon reconstructor, or if their compatibility score returned by the muon identification algorithm is greater than 0.1.

To reconstruct the ϕ meson, all tracks reconstructed with the standard track reconstruction are used. Requiring the p_T of each track to be above $0.8 \text{ GeV}/c$ and assigning a kaon mass to the thus reconstructed tracks, all oppositely charged track pairs for which the invariant mass is within $20 \text{ MeV}/c^2$ of the world-average mass of the ϕ meson are retained. The p_T of the ϕ candidate is required to be above $1 \text{ GeV}/c$, and the p_T of the B_s candidate above $5 \text{ GeV}/c$.

A kinematic fit [130] is then made, where the four tracks are constrained to come from a common vertex and the invariant mass of the two muons is constrained to be equal to the mass of the J/ψ . Since the natural width of the ϕ meson is of the same order as the resolution due to the reconstruction, no mass constraint is applied to the two kaon tracks. With this fit, a resolution on the invariant mass of the B_s meson of $14 \text{ MeV}/c^2$ is found. The confidence level of the fit is required to be greater than 1×10^{-3} (seven degrees of freedom). The invariant

Table 5.3: Offline selection efficiencies for the signal and background (defined with respect to the number of generated events) after each requirement.

Requirement	Signal	Background		
	$B_s \rightarrow J/\psi\phi$	$b \rightarrow J/\psi X$	Prompt J/ψ	$B^0 \rightarrow J/\psi K^{*0}$
HLT selection	20.50(6) %	1.23(3) %	0.0007(7) %	0.937(14) %
Reconstruction + Basic p_T req.	18.15(5) %	0.63(2) %	0.0007(7) %	0.675(12) %
Muon Identification	17.89(5) %	0.585(19) %	0.0007(7) %	0.636(11) %
Kinematic fit χ^2 req.	16.58(5) %	0.282(14) %	0.0007(7) %	0.503(10) %
Pointing constraint	16.48(5) %	0.258(13) %	–	0.497(10) %
ϕ mass req.	14.65(5) %	0.113(13) %	–	0.202(10) %

Table 5.4: Expected cross sections for the signal and background, after each requirement, with number of expected events.

	Signal	Background		
	$B_s \rightarrow J/\psi\phi$	Inclusive $b \rightarrow J/\psi X$	$B^0 \rightarrow J\psi K^*(892)^0$	Prompt J/ψ
$\sigma \times \text{Br}$	$2.87 \pm 1.07 \text{ nb}$	$682 \pm 64 \text{ nb}$	$20.4 \pm 1.7 \text{ nb}$	$141 \mu\text{b}$
Kin. preselection	$74 \pm 27 \text{ pb}$	$3.20 \pm 0.3 \text{ nb}$	$366 \pm 22 \text{ nb}$	$176 \pm 2 \text{ nb}$
Level-1	$34 \pm 12 \text{ pb}$	$1.22 \pm 0.11 \text{ nb}$	$172 \pm 10 \text{ nb}$	$65 \pm 1 \text{ nb}$
HLT	$15.2 \pm 5.5 \text{ pb}$	$39.4 \pm 3.8 \text{ nb}$	$3.52 \pm 0.21 \text{ nb}$	$12 \pm 12 \text{ nb}$
Offline	$10.9 \pm 4.0 \text{ pb}$	$36.2 \pm 0.54 \text{ nb}$	$0.74 \pm 0.06 \text{ nb}$	–
Events per 30 fb^{-1}	327'000	108'500	22'200	-

mass of the two kaons is required to be within $\pm 8 \text{ MeV}/c^2$ of the world-average mass of the ϕ meson. Finally, the cosine of the angle between the reconstructed momentum vector of the B_s candidate and the vector pointing from the production to the decay vertex is required to be less than 0.95. The distribution of the invariant mass of the candidates after all selection requirements is shown in Figure 5.1 (right).

The primary vertex is not used at this stage, since the efficiency of the standard primary vertex finder [7, Section 6.6.4], which uses all fully reconstructed tracks, is 92%, and drops to 83% if the vertex is required to be within $500 \mu\text{m}$ from the simulated vertex. In order to prevent this unnecessary loss of efficiency, no use is made of the primary vertex, and all quantities of interest are evaluated in the transverse plane.

With this selection, a yield of approximately 327 000 signal events can be expected within 30 fb^{-1} of data, with a background of 39 000 events. The efficiencies for the different criteria, which include the respective reconstruction efficiencies, are given in Table 5.3 for the signal and the different background samples, and the number of expected in Table 5.4. These do not include a requirement on the four-track invariant mass of the candidates, since the sidebands will be used later in the analysis. However, only a small fraction of these events are directly under the B_s peak, and even a simple cut will reduce the number of background events by a significant factor.




Figure 5.1: Four-track invariant mass distribution after the HLT (left) and offline (right) requirements. The right distribution includes only combinatorial background and the left distribution the expected inclusive $b \rightarrow J/\psi X$ and $B^0 \rightarrow J/\psi K^{*0}$ background.

5.5 The maximum likelihood analysis

The final state of the decay of a pseudo-scalar B meson into two vector mesons $B \rightarrow V_1 V_2$ is an admixture of CP -even and CP -odd states [131, 125, 126]. The CP -odd states correspond to transitions in which the relative orbital momentum L between the two vector mesons is 1 and the CP -even states to transitions in which L is either 0 or 2. The amplitude of the decay can be decomposed in three independent decay amplitudes which correspond to the linear polarization states of the two mesons. The first, A_0 , describes states in which the linear polarization vectors are longitudinal and is CP -even. The other two describe states in which the linear polarization vectors are transverse, either parallel (A_{\parallel} - CP -even) or perpendicular (A_{\perp} - CP -odd) to each other.

The differential decay rate can be written as:

$$\frac{d^4\Gamma(B_s(t))}{d\Theta dt} = f(\Theta, \alpha, t) = \sum_{i=1}^6 O_i(\alpha, t) \cdot g_i(\Theta), \quad (5.1)$$

where O_i are the kinematics-independent observables, g_i the angular distributions and Θ generically denote the angles which defines the kinematics. The time evolution of the different observables is given by bilinear combinations of the polarization amplitudes, $|A_0(t)|^2$, $|A_{\parallel}(t)|^2$, $|A_{\perp}(t)|^2$, $\Im(A_{\parallel}^*(t)A_{\perp}(t))$, $\Re(A_0^*(t)A_{\parallel}(t))$ and $\Im(A_0^*(t)A_{\perp}(t))$. These are functions of the widths of the two light and heavy eigenstates, Γ_L and Γ_H , weak phase ϕ_{CKM} , the magnitudes of the amplitudes at $t = 0$ ($A_0(0)$, $A_{\parallel}(0)$ and $A_{\perp}(0)$) which describe all hadronization effects, and, for a flavour-tagged sample, the mass difference $\Delta m_s = m_H - m_L$. Since the overall phase of the polarization states is not observable, two strong phases are defined as $\delta_1 \equiv \arg |A_{\parallel}^* A_{\perp}|$ and $\delta_2 \equiv \arg |A_0^* A_{\perp}|$. These are CP conserving, and are expected to be 0 (mod π) in the absence of final-state interaction. Assuming $SU(3)$ flavour-symmetry, the magnitudes and the two strong phases are equal for the decay $B_s \rightarrow J/\psi \phi$ and $B \rightarrow J/\psi K^*$ in unmixed samples. The measurement of these parameters is of interest to study and improve the phenomenological models used to calculate all hadronic effects.

In such decays, the kinematics is uniquely defined by a set of three angle. The transversity base is used in this analysis, in which the set of variables is $\Theta = (\cos \theta, \phi, \cos \varphi)$. In this base,

(θ, φ) are the polar and azimuthal angles of the momentum of the μ^+ in the J/ψ rest-frame. This coordinate system is defined such that the ϕ moves in the positive x direction and the z axis is perpendicular to the decay plane of $\phi \rightarrow K^+K^-$. The angle ψ is defined in the rest frame of the ϕ as the negative angle between the K^+ direction and the J/ψ direction.

In order to measure the values of the different parameters, an unbinned maximum likelihood fit is performed on the observed time evolution of the angular distribution. In the absence of background and without distortion, the p.d.f. describing the data would be the original differential decay rate $f(\Theta, \alpha, t)$ (Eq. 5.1). The distortion of this distribution by the detector acceptance, trigger efficiency and the different selection criteria is taken into account by an efficiency term $\epsilon(t, \Theta)$. In addition, a term describing the background has to be added.

The resolution of the proper decay length is taken into account by convolving the p.d.f. with a Gaussian resolution function where the width is a free parameter. Since the uncertainty of the measured angles are found to be small, these will not be taken into account in the fit. A contribution will be added to the systematic uncertainty to reflect this omission.

It is assumed that the efficiency can be factorized in two functions, the first modeling the effect of the decay length requirements and the second the distortion of the angular distribution,

$$\epsilon(t, \Theta) = \epsilon(t) \cdot \epsilon(\Theta). \quad (5.2)$$

The angular efficiency is described by an expansion of products of spherical harmonics ([132]):

$$\epsilon(\Theta) = \sum_{LRM} T_{LRM}^\epsilon \cdot \mathcal{Y}_{LRM}(\Theta), \quad (5.3)$$

$$\text{with } \mathcal{Y}_{LRM}(\Theta) = \sqrt{2\pi} \cdot Y_{LM}(\theta, \varphi) \cdot Y_{RM}(\psi, 0), \quad (5.4)$$

where \mathcal{Y}_{LRM} are orthonormal basis functions and Y_{LM} are spherical harmonic functions. In principle, L and R run from 0 to infinity and the sum over M from $-\min(L; R)$ to $+\min(L; R)$, but it has been found that the expansion can be limited to $L, R \leq 8$. These \mathcal{Y}_{LRM} functions describe the partial waves involved in a scalar \rightarrow vector decay [133]. The moments of the efficiency are determined from a Monte Carlo simulation with full detector simulation:

$$T_{LRM}^\epsilon = \int \epsilon(\Theta) \cdot \mathcal{Y}_{LRM}^*(\Theta) d\Theta \quad (5.5)$$

$$\approx \frac{1}{N_{gen}} \sum_{i=1}^{N_{obs}} \frac{1}{f(\Theta_i)} \mathcal{Y}_{LRM}^*(\Theta_i). \quad (5.6)$$

where $f(\Theta_i)$ is the expected time-integrated angular distribution (Eq. 5.1).

The time-dependent efficiency describes mainly the effects of the requirements on the proper decay length distribution. After the initial turn-on and a stable plateau, a deficit of events can be observed. Initial studies attribute this decrease in efficiency to the restrictions imposed on the seeds by the tracking regions in the HLT, which causes an additional track reconstruction inefficiency for displaced tracks such as those originating from B decays. The tolerance on the transverse and the longitudinal direction imposed on the tracking regions in the HLT results in an implicit cut on the impact parameters. Further studies are needed to find solutions to alleviate this inefficiency. Without corrections, the main effect of this inefficiency would be to lower the estimated lifetime of the longer-lived eigenstate B_s^H .

The different features in this distributions can not easily be described by a simple function. We have thus chosen two sigmoidal functions combined with a quadratic function

$$\epsilon(t) = \begin{cases} c \cdot \left(1 + \tanh\left(\frac{t-t_0}{\Delta t_1}\right)\right) & t < t_0 \\ (a \cdot t^2 + b \cdot t + c) \cdot \left(1 + \tanh\left(\frac{t-t_0}{\Delta t_2}\right)\right) & t > t_0 \end{cases} . \quad (5.7)$$

The parameters are found by fitting this function to the distribution obtained by the full Monte Carlo simulation.

The best way to gauge our ability to account for all effects and our capacity to correct them through this time-dependent efficiency curve is by comparing the proper time distributions foreseen by the simulation and observed in the data for the different B mesons. The first obvious choice is again the decay $B^0 \rightarrow J/\psi K^{*0}$, which is very similar to the studied B_s decay, and for which the lifetime has been measured with a high precision. Any discrepancy between the efficiency determined by Monte Carlo and the reality will be reflected in a mismeasurement of the B^0 lifetime. Further studies would be needed to determine the sensitivity of the efficiency on the lifetime of the selected B meson. It is dubious whether the number B_s events recovered in other trigger streams such as the dimuon, which has no decay length requirement, would be enough to estimate the time-dependent efficiency.

The background can be divided in two different types of distributions. The first type arises from misidentified $B_d \rightarrow J/\psi K^* \rightarrow \mu^+ \mu^- K \pi$ events, which has a similar differential decay rate [125, 126] to the decay of interest. The width difference of the two eigenstates of the B^0 are assumed to be negligible, and no CP violation is present since the final state is flavour specific. To describe this background in our dataset, it is not possible to use its time dependent angular distribution, which is in principal well known, since all variables are mismeasured because of the misidentification of the π . In addition, the distortion of the distribution due to the various requirements is much more severe than in the case of the B_s . Indeed, due to its lower mass, the momentum of the π in the laboratory frame is lower than that of the corresponding K when the π is emitted in the direction opposite to the momentum of the K^{*0} .

The same set of functions $\mathcal{Y}_{LRM}(\Theta)$ (Eq. 5.4) are used to model this background, with the moments computed in the following way:

$$T_{LRM}^b = \int b(\Theta) \cdot \mathcal{Y}_{LRM}^*(\Theta) d\Theta \quad (5.8)$$

$$\approx \frac{1}{N_b} \sum_{i=1}^{N_b} \mathcal{Y}_{LRM}^*(\Theta_i) , \quad (5.9)$$

Here as well, the expansion would be done up to $L, R \leq 8$. These would be obtained by a Monte Carlo simulation and could be cross-checked by a fully reconstructed sample of well-identified $B^0 \rightarrow J/\psi K^{*0}$ decays misreconstructed as B_s candidates.

The time dependence of this background is modeled as a single exponential decay, again with a time-dependent efficiency. The lifetime is left as a free parameter, τ_d , since the mismeasurement of the proper decay length precludes using the well-measured lifetime of the B^0 .

The other sources of background are assumed to have no angular dependence. The distribution of their proper decay time is modeled by two exponential decays, the first describing

the short-lined prompt background and the second misidentified long-lived heavy-flavour hadrons.

A better separation of the signal and background is obtained by using the events in a wider invariant mass region between 5.219 and 5.559 GeV/ c^2 , and including in the fit the distribution of the invariant mass of the candidates. The distribution of the B_s candidates is modeled by a Gaussian $G_s(m; M_{B_s}, \sigma_s)$, where M_{B_s} is the mass of the B_s meson and σ_s the variance due to the reconstruction. The distribution of the misidentified $B^0 \rightarrow J/\psi K^{*0}$ decays can reasonably well be modelled in the chosen region by a Gaussian $G_d(m; M_d, \sigma_d)$. Because of the misidentification of the pion, M_d will not correspond to the true mass of the B^0 meson, and will be left as a free parameter in the fit. The other sources of background are assumed to have a flat mass distribution and will be modeled by a linear function $L(m)$.

The total p.d.f. to be fit is thus given by

$$\begin{aligned} \mathcal{P} = & (1 - b_d - b_c) \cdot \epsilon(t, \Theta) \cdot f(\Theta, \alpha, t) \cdot G_s(m; M_{B_s}, \sigma_s) \\ & + b_d \cdot b_d(\Theta) \cdot \epsilon(t) \cdot \frac{1}{\tau_d} e^{-t/\tau_d} \cdot G_d(m; M_d, \sigma_d) \\ & + b_c \cdot \epsilon(t) \cdot \left(\frac{1}{\tau_{cl}} e^{-t/\tau_{cs}} + \frac{1}{\tau_{cl}} e^{-t/\tau_{cl}} \right) \cdot L(m), \end{aligned} \quad (5.10)$$

where b_d , respectively b_c , are the fraction of misidentified B^0 background, respectively combinatorial background in the sample. These parameters are left free in the fit. The resolution of the proper decay length is taken into account by convolving the p.d.f. with a Gaussian resolution function. The standard deviation of the Gaussian is taken as the uncertainty of each candidates proper decay length measurement multiplied by a scale factor, which is left free in the fit. Since the uncertainty of the measured angles are found to be small, these will not be taken into account in the fit. A contribution will be added to the systematic uncertainty to reflect this omission.

5.1.6 Result

Due to the high production cross sections of the identified background, only limited samples could be generated and analyzed, which does not permit to have a final dataset with the foreseen signal-to-background ratio. Indeed, the signal sample corresponds to an integrated luminosity of 6.8 fb $^{-1}$, while the inclusive background corresponds to an integrated luminosity of barely 48 pb $^{-1}$. The situation is somewhat better for the decay $B^0 \rightarrow J/\psi K^{*0}$ for which the sample corresponds to an integrated luminosity of 1.3 fb $^{-1}$.

We have thus proceeded in two steps. A fit was first performed on the complete set of selected and associated B_s candidates only, using the efficiency functions determined in the previous section. The relative width difference $\Delta\Gamma_s/\Gamma_s$ can be determined with an uncertainty of 0.016 (Table 5.5), but no sensitivity on the weak phase and the strong phases is obtained.

Then, we consider a sample corresponding to an integrated luminosity of 1.3 fb $^{-1}$, which allows us to have a realistic ratio of $B^0 \rightarrow J/\psi K^{*0}$ and signal events. With the low number of background events which remain after all selection requirements, an accurate model through the described p.d.f. is not possible. In addition, the low number of $B^0 \rightarrow J/\psi K^{*0}$ events does not permit an accurate estimate neither of its angular distribution nor of its time-dependent efficiency. As such, the background events are simply added to the dataset and their expected

Table 5.5: Results of the maximum likelihood fit for 73813 signal events.

Parameter	Input value	Result	Stat.error	Rel.error
$ A_0(0) ^2$	0.57	0.57398	0.00267	0.4%
$ A_{ }(0) ^2$	0.217	0.21808	0.00473	2.1%
$ A_{\perp}(0) ^2$	0.213	0.20794	0.00396	1.9%
$\bar{\Gamma}_s$	0.712 ps^{-1}	0.712358 ps^{-1}	$0.00350643 \text{ ps}^{-1}$	0.5%
$\Delta\Gamma_s$	0.142 ps^{-1}	0.134645 ps^{-1}	$0.0108247 \text{ ps}^{-1}$	8.0%
$\Delta\Gamma_s/\Gamma_s$	0.2	0.189013	0.0157993	8.4%
δ_1	π	2.94405	0.632682	
δ_2	0	-0.109493	0.639713	
ϕ_{CKM}	-0.04	-0.0297427	0.0758856	

distribution is not included in the p.d.f. used in the fit. The p.d.f. would thus simply describe the B_s distribution:

$$\mathcal{P} = \epsilon(t, \Theta) \cdot f(\Theta, \alpha, t)$$

With such a fit in which the invariant mass of the candidates is not taken into account, a requirement on the invariant mass of the candidates would obviously be made, choosing a window of $\pm 36 \text{ MeV}/c^2$ around the world-average B_s mass. This reduces the number of B^0 background events by a further 59%, while reducing the number of signal candidates by 2.9%. The results of the fit without background is given in table 5.6 and with background in table 5.7. With the lower number of B_s candidates, the statistical error of the measurement is, as expected, markedly worse. As can be seen, the influence of the background is very small, with only a slight degradation of the width difference. The distributions of the proper decay length of the selected events with the fit projection is shown in Figure 5.2.

Table 5.6: Results of the maximum likelihood fit for an integrated luminosity of 1.3 fb^{-1} (signal only).

Parameter	Input value	Result	Stat.error	Rel.error
$ A_0(0) ^2$	0.57	0.5859	0.0062	1.1%
$ A_{ }(0) ^2$	0.217	0.2141	0.0078	3.6%
$ A_{\perp}(0) ^2$	0.213	0.2002	0.0064	3.2%
$\bar{\Gamma}_s$	0.712 ps^{-1}	0.7018 ps^{-1}	0.0081 ps^{-1}	1.2%
$\Delta\Gamma_s$	0.142 ps^{-1}	0.1470 ps^{-1}	0.0256 ps^{-1}	17.4%
$\Delta\Gamma_s/\Gamma_s$	0.2	0.2095	0.0371	18.1%



5.1.7 Systematics and detector effects

The list of systematic uncertainties which were considered are summarized in two tables. The first, Table 5.8, summarizes the uncertainties which affect the HLT rate and the number of foreseen events after all selection requirements. The second, Table 5.9, summarizes the uncertainties which affect the measurement of the various parameters.

- Signal and background statistics:
Among the various uncertainties listed in Section 5.1.2, the largest single source of

Table 5.7: Results of the maximum likelihood fit for an integrated luminosity of 1.3 fb^{-1} (signal and background).

Parameter	Input value	Result	Stat.error	Rel.error
$ A_0(0) ^2$	0.57	0.5823	0.0061	1.1%
$ A_{ }(0) ^2$	0.217	0.2130	0.0077	3.6%
$ A_{\perp}(0) ^2$	0.213	0.2047	0.0065	3.2%
$\bar{\Gamma}_s$	0.712 ps^{-1}	0.7060 ps^{-1}	0.0080 ps^{-1}	1.1%
$\Delta\Gamma_s$	0.142 ps^{-1}	0.1437 ps^{-1}	0.0255 ps^{-1}	17.7%
$\Delta\Gamma_s/\Gamma_s$	0.2	0.2036	0.0374	18.4%




Figure 5.2: Distributions of the proper decay length of the selected signal and background events with fit projection.

uncertainty in the estimate of the number of events is obviously the poor knowledge of the $B_s \rightarrow J/\psi\phi$ branching ratio. The uncertainties quoted on the estimates above do not include the uncertainties on the total $b\bar{b}$ cross section at LHC energies, the $b \rightarrow B^0$ fragmentation functions, the transverse momentum distribution of b quarks. However, since both the signal and background are proportional to the same $b\bar{b}$ cross section, the signal-to-background ratio is unaffected by the latter uncertainty.

- Track reconstruction efficiency:
A 1% uncertainty per track on the track reconstruction efficiency is assumed for all tracks.
- Muon reconstruction:
The selection relies heavily on the correct identification of muons. A 1% uncertainty per track on the combined muon identification procedure is assumed.
- Tracker and muon detector misalignment:
The study has been conducted with a perfectly aligned detector. To gauge the sensitivity of the analysis with respect to the alignment it has been repeated on a

Table 5.8: List of systematic uncertainties with effect (in %) on the predictions of the rates.

Source	HLT Uncertainty	Offline uncertainty	Common uncertainty
Branching ratio B_s			36.4 %
Branching ratio B^0			6 %
Branching ratio $b \rightarrow J/\psi X$			9 %
Traching inefficiency	2 %	2%	
Muon reco.	-	1.4%	
Misalignment	17%	-	

detector with the short-term alignment scenario, which is supposed to be typical of the misalignment of the initial data taking period [78]. The effects of misalignment of the tracker on various aspects of track and vertex reconstruction have been extensively studied and reported in [134, 135]. The degradation affect both the selection, mostly through the requirement on the significance of the transverse decay length of the J/ψ in the HLT, and the analysis, through the degradation of the proper decay length. The resolution of the latter is degraded from $24 \mu\text{m}$ for a perfectly aligned detector to $32 \mu\text{m}$ with the short-term alignment. The HLT efficiency is degraded by some 17% with respect to a perfectly aligned detector.

- Background distributions:

To gauge the influence of the background on the fit, the variation observed between the fits performed on the reduced 1.3 fb^{-1} dataset with and without these events is added to the systematic uncertainty (“Bckg. distrib.” in the table).

Since a the signal-to-background ratio has a significant uncertainty, the fit on the reduced 1.3 fb^{-1} sample is repeated varying the number of B_s signal events to match the uncertainty in the signal-to-background ratio. For this estimate, a different uncertainty for the B_s branching fraction has been chosen, since it is believed that it will be measured again in the current run of the Tevatron. Two main uncertainties plague the measurement done at CDF in the Run I, the low number of observed B_s candidates and the uncertainty on the fragmentation. Based on recent publications, it is estimated that approximately 30 times more $B_s \rightarrow J/\psi \phi$ decays than in Run 1 should already be collected in the current dataset of 1 fb^{-1} . The uncertainty of the branching fraction is therefore reduce to 20%. For the other uncertainties, the numbers listed in Table 5.8 are used. The variation observed on the fit is listed under the heading “S/B ratio”. In a larger dataset, where the full p.d.f. (Eq. 5.11) is used, the influence of the uncertainty on the signal-to-background ratio should be much smaller, since the fraction of signal and background events in the dataset is a free parameter in the fit.

- Distortion of the proper-time distribution (“ $c\tau$ distortion”):

Other fits were then performed where the parameters of the time dependent efficiency function are varied by one standard deviation. The mean variation of the fitted parameters was added to the systematic uncertainty. As already mentioned, the decay $B^0 \rightarrow J/\psi K^{*0}$ can be used to compare the accuracy of this model by comparing the Monte Carlo prediction with the efficiency function observed in the data.

- Distortion of the angular distributions (“Ang. distortion”):

The expansion used to model the distortion of the angular distributions (Eq. 5.3)

Table 5.9: List of systematic uncertainties with effect on the measurements.

Source	$ A_0(0) ^2$	$ A_{ }(0) ^2$	$ A_{\perp}(0) ^2$	$\bar{\Gamma}_s$	$\Delta\Gamma_s/\Gamma_s$
Bckg. distrib.	0.0034	0.0011	0.0045	0.0043	0.0059
S/B ratio	0.0037	0.0001	0.0024	0.0025	0.0055
Resolution	-	-	-	0.00025	0.0040
Ang. distortion	0.0143	0.0061	0.0082	0.00083	0.0010
$c\tau$ distortion	0.0016	0.00073	0.0023	0.0221	0.0146
Alignment	0.00012	0.00042	0.00055	0.00040	0.0014
Total	0.0152	0.0063	0.0099	0.0227	0.0173

is limited to $L, R \leq 8$. When limiting the expansion to $L, R \leq 6$ or $L, R \leq 10$, the result of the fit shows negligible differences. In addition, to account for the possibility that the efficiencies do not factorize and that the angular efficiency is grossly misestimated, the fit was also repeated without the angular efficiency, hence without correction of the distortion. While this has little influence on the estimated lifetimes, a large variation is found for the amplitudes. This variation is used as systematic uncertainty.

- Resolution on the angular variables (“Resolution”):

In order to estimate the influence of the uncertainties of the angles on the fit, a fully controlled toy Monte Carlo, in which only the proper time and angles were generated according to the expected p.d.f. and smeared with Gaussian resolution functions. The default standard deviations are taken to be equal to those measured in the Monte Carlo with full detector simulation. The simulation was then repeated without smearing and with a harsher smearing, where the resolution is taken to be five times larger than in the default simulation. The value of parameters found in both cases were very close to the values found with the default smearing, and the observed variation is added to the systematic uncertainty.



5.1.8 Conclusion

The present section describes a study on the selection of the $B_s \rightarrow J/\psi\phi$ decay and the measurement of the width difference through an untagged analysis. An example of trigger is given which would be efficient for this decay and reject a large fraction of the background. It is based on the identification of J/ψ and B_s candidates with a displaced decay vertex. Nevertheless, this trigger precludes the selection of other decays of the B meson, and should certainly evolve as a true precursor to a B physics trigger. Indeed, the strategy proposed for the Level 2 would select inclusive $b \rightarrow J/\psi$ decays with a high efficiency and a good purity with respect to the prompt J/ψ background. Large uncertainties plague nevertheless our estimates of the rates, since large uncertainties remain on the production cross section of the b -quark and prompt J/ψ , on their momentum distribution and on the $b \rightarrow B_s$ fragmentation function.

A first measurement of one of the main parameters of the B_s system, the difference of the width of the weak eigenstates could be measured with a statistical uncertainty of 0.011 in a sample corresponding to an integrated luminosity of 10 fb^{-1} . A first measurement undertaken on approximately 1.3 fb^{-1} of data could already yield a measurement with an uncertainty of 20% (Table 5.10). A natural extension of this study should be a tagged analysis, for which flavour tagging algorithms need to be developed.



Table 5.10: Results of the maximum likelihood fit for an integrated luminosity of 1.3 fb^{-1} (signal and background).

Parameter	Input value	Result	Stat. error	Sys. error	Total error	Rel. error
$ A_0(0) ^2$	0.57	0.5823	0.0061	0.0152	0.0163	2.8%
$ A_{ }(0) ^2$	0.217	0.2130	0.0077	0.0063	0.0099	4.6%
$ A_{\perp}(0) ^2$	0.213	0.2047	0.0065	0.0099	0.0118	5.8%
$\bar{\Gamma}_s$	0.712 ps^{-1}	0.7060 ps^{-1}	0.0080 ps^{-1}	0.0227 ps^{-1}	0.0240 ps^{-1}	3.4%
$\Delta\Gamma_s$	0.142 ps^{-1}	0.1437 ps^{-1}	0.0255 ps^{-1}	0.0113 ps^{-1}	0.0279 ps^{-1}	19%
$\Delta\Gamma_s/\Gamma_s$	0.2	0.2036	0.0374	0.0173	0.0412	20%

5.2 Associated production of MSSM heavy neutral Higgs bosons $b\bar{b}H(A)$ with $H(A) \rightarrow \tau\tau$

5.2.1 Introduction

Observation of the heavy neutral scalar accompanied by b jets and decaying into two τ leptons would be an important justification of the MSSM Higgs sector. The associated Higgs boson production $gg \rightarrow b\bar{b}H(A)$ in the MSSM is dominated at large values of $\tan\beta$. The cross section of $gg \rightarrow b\bar{b}H(A)$, $H(A) \rightarrow \tau\tau$ channel is proportional to $\tan^2\beta_{\text{eff}}$ and will be used in a global fit together with other relevant measurements to determine the SUSY parameters simultaneously.

This channel is an excellent benchmark for the b and τ -tagging, jet and missing E_T reconstruction. The final state with two τ jets requires τ tagging both at Level-1 and High Level trigger. Along with reconstruction and tagging issues, a big number of various Standard Model backgrounds including QCD multi-jet production must be understood from the real data to be able to declare discovery.

5.2.2 Event generation

The signal events were generated by PYTHIA using processes 181 ($gg \rightarrow b\bar{b}H$) and 152 ($gg \rightarrow H$) for three values of Higgs boson mass: 200, 500 and 800 GeV/c^2 . The backgrounds considered were QCD multi-jet events (for $\tau\tau \rightarrow jj$ mode), $t\bar{t}$, $b\bar{b}$, Drell-Yan production of Z/γ^* , W +jet, Wt and $\tau\tau b\bar{b}$. All background processes except $\tau\tau b\bar{b}$ were generated with PYTHIA. The $\tau\tau b\bar{b}$ process was generated by CompHEP.

In order to reduce CPU time for full detector simulation and event reconstruction the loose preselections were applied for some of the backgrounds at the level of generation. The description of preselections for each final state can be found in the following chapters.

The cross sections for the associated Higgs boson production $gg \rightarrow b\bar{b}H(A)$ and branching ratio $H(A) \rightarrow \tau\tau$ were calculated using FeynHiggs2.3.2 [136, 137, 138] (the code can be obtained from <http://www.feynhiggs.de>) in the m_h^{max} scenario with $\mu=200 \text{ GeV}/c^2$ (see Section 11.3.1).

The uncertainty of the measured cross section of the $b(\bar{b})A$, $A \rightarrow \tau\tau$ process will include the uncertainty of the Monte Carlo generation. The verification of the Monte Carlo generation

for the Higgs boson production with the associated b jets will be done with the real data using $b\bar{b}Z$ ($Z \rightarrow \ell\ell$) events [139].

5.2.3 Level-1 and High Level trigger selections

The $\tau\tau \rightarrow jj$ final state is triggered by single or double Level-1 Tau trigger with thresholds 93 GeV for single and 66 GeV for double trigger. It is followed by the double τ jet tagging at High Level trigger. Currently there are two selection strategy at HLT under consideration [140]. In the first strategy the calorimeter isolation with the electromagnetic calorimeter is applied to the first τ jet in order to reduce the Level-1 output rate by factor of 3. The tracker isolation is then applied on both jets using the tracks reconstructed with the pixel detector only. The second strategy performs tracker isolation right after Level-1 trigger decision and uses full tracker with the regional track finding and restricted number of hits required to reconstruct track. The first strategy is exploited in this analysis.

The $\tau\tau \rightarrow \mu j$ final state uses single muon trigger at Level-1 with threshold 14 GeV. At the High Level the combined muon-plus- τ -jet trigger is used with the thresholds 15 GeV for muon and 40 GeV for τ -jet.

The $\tau\tau \rightarrow e j$ final state uses Level-1 single electron trigger with threshold 23 GeV together with the combined electron-plus- τ -jet trigger with thresholds 14 GeV for electron and 52 GeV for τ jet. At High Level again the single electron trigger with threshold 26 GeV and combined electron-plus- τ -jet triggers with thresholds 16 GeV for electron is used. No threshold is applied for τ jet candidate.

At High Level trigger for both the $\tau\tau \rightarrow \mu j$ and the $\tau\tau \rightarrow e j$ final states the ECAL and pixel track isolation is applied on τ -jet candidate similar to what is used in the double τ -jet trigger. For the lepton (e and μ) the same selections are used as for the single electron and muon High Level triggers. The lepton and τ jet are required to be originated from the same vertex found with the pixel detector. The tracks from this vertex only are used in the tracker isolation.

The search strategy for τ -jet candidates at High Level trigger for combined muon-plus- τ -jet and electron-plus- τ -jet triggers is the following. Two calorimeter jets are always reconstructed with the regional jet finder in the regions given by the two highest E_T Level-1 Tau jets. For muon-plus- τ -jet trigger the first (highest E_T) jet is taken as τ -jet candidate. For electron-plus- τ -jet trigger the requirement of non collinearity of the jet and the HLT electron candidate, $\Delta R(e - \text{jet}) > 0.3$, is checked for each jet, where $\Delta R(e - \text{jet})$ is the distance in η - φ space between the electron and the jet. The first not collinear jet is taken as the τ -jet candidate.

5.2.4 Off-line event selection strategy

The first step in off-line analysis is the τ -jet identification. The calorimeter jet is reconstructed in the η - φ region of the High Level trigger τ -jet candidate with the iterative cone algorithm using the cone size of 0.4. A number of requirements for τ -jet identification [140] is applied in addition to the tracker isolation which is more tight in off-line than at HLT and uses the tracks reconstructed with the full tracker. The additional τ -jet identification criteria include requirements to have one or three tracks in the signal cone and opposite charge of two τ jets for $\tau\tau \rightarrow jj$ mode or lepton and τ jet for $\tau\tau \rightarrow \ell j$ modes, the cut on the transverse impact parameter and p_T of the leading track in the signal cone. Finally the electron rejection

criteria was applied. The τ -jet tagging reduces the QCD multi-jet (including $b\bar{b}$) and the W +jet backgrounds.

The associated $b\bar{b}H(A)$ production dominates at high values of $\tan\beta$, thus it is natural to apply b -jet tagging which must suppress Drell-Yan $\tau\tau$ production and eliminate further the QCD multi-jet and the W +jet backgrounds. Since b jets in the signal are very soft in E_T and have flat distribution in pseudorapidity only single b tagging is applied. Furthermore, it is possible to veto events with an additional jets to reduce $t\bar{t}$ background. The τ jets found in the first step are not considered for b tagging. Non τ -jet candidates are reconstructed with the iterative cone algorithm using cone size of 0.5.

The energy of τ jet is corrected with the dedicated calibration obtained from the Monte-Carlo sample of single τ jets at low luminosity. The energy of other jets in the event is corrected applying Monte Carlo calibration evaluated from the QCD multi-jet events at low luminosity.

5.2.5 Method of the Higgs boson mass reconstruction

Despite the escaping neutrinos, the Higgs boson mass can be reconstructed in the $H \rightarrow \tau\tau$ channels from the visible τ momenta (leptons or τ jets) and the missing transverse energy (E_T^{miss}) with the collinearity approximation for the neutrinos from highly boosted τ 's. The mass resolution depends on the angle $\Delta\varphi$ between the visible τ momenta as $1/\sin(\Delta\varphi)$ and is sensitive to the E_T^{miss} measurement, both in magnitude and particularly in direction. The measurement of E_T^{miss} is affected by the non-linear calorimeter response. A method to improve the E_T^{miss} scale using the jet energy corrections was used [141, 142]. The correction of the missing E_T scale improves the reconstruction efficiency by reducing the number of events with negative reconstructed τ lepton and neutrino energies. In particular, for the case of the $\tau\tau \rightarrow jj$ final state the efficiency is improved by factor of $\simeq 1.6$. The $\tau\tau$ mass reconstruction method will be verified with the real data using $Z \rightarrow \tau\tau \rightarrow e(\mu) + \text{jet}$ and $Z \rightarrow \tau\tau \rightarrow e + \mu$ channels [139, 143].

5.2.6 $H \rightarrow \tau\tau \rightarrow$ two τ jet analysis

5.2.6.1 Event generation and preselections

The $t\bar{t}$, Drell-Yan production of Z/γ^* , W +jet and Wt backgrounds were generated with forced in PYTHIA $W \rightarrow \tau\nu$ and $Z/\gamma^* \rightarrow \tau\tau$ decays. The TAUOLA package was used for τ -lepton decays into all possible decay modes.

The Z/γ^* generation was splitted into three bins of generated di- τ -lepton mass $m_{\tau\tau}$: 80-130 GeV/c^2 , 130-300 GeV/c^2 and $>300 \text{GeV}/c^2$. The $\tau\tau b\bar{b}$ generation was divided into two bins of generated di- τ -lepton mass $m_{\tau\tau}$: 60-100 GeV/c^2 and $>100 \text{GeV}/c^2$. The $\tau\tau b\bar{b}$ CompHEP background was propagated to PYTHIA for the showering and hadronization and τ leptons decay into all possible modes.

The W +jet background was generated using PYTHIA processes 16 and 31 and with $\hat{p}_T > 65 \text{GeV}/c$. The QCD multi-jet background generation was done for four bins in \hat{p}_T : 50-80, 80-120, 120-170 and $> 170 \text{GeV}/c$.

The loose preselections at the level of generation were applied for all backgrounds (except $\tau\tau b\bar{b}$): the event was required to have at least two " τ -like" jets. The jets were reconstructed

5.2. Associated production of MSSM heavy neutral Higgs bosons $b\bar{b}H(A)$ with $H(A) \rightarrow \tau\tau$ 121

with PYTHIA PYCELL routine using cone size 0.5. The jet is selected as “ τ -like” if it has $E_T^{MC} > 50$ GeV, $|\eta^{MC}| < 2.4$ and transverse momentum of the leading stable charged particle in jet $p_T^{MC} > 30$ GeV/c. These cuts are smaller than ones applied at the trigger and off-line τ -jet selections. For Z/γ^* background no cut was applied on p_T^{MC} .

For the signal events the Higgs boson was forced to decay into two τ leptons and τ lepton was decayed hadronically using TAUOLA. No preselections were applied for the signal events.

5.2.6.2 Event selections

The calorimeter τ -jet candidates are reconstructed in the η - φ regions of the High Level trigger τ -jet candidates, thus no “volunteers” are searched for. It is due to the high ($\simeq 100\%$) purity of the HLT τ -jet candidates (fraction of true τ jets matched with τ -jet candidates)

The cut on the uncalibrated transverse jet energy for each of the two τ -jet candidates was required. It was $E_T > 50$ GeV for the $M_A = 200$ GeV/ c^2 . For higher Higgs boson masses the asymmetrical cuts were used: 100, 50 GeV for $M_A = 500$ GeV/ c^2 and 150, 50 GeV for $M_A = 800$ GeV/ c^2 . It allows more effective rejection of the QCD multi-jet background. The following τ -jet identification criteria were then used:

- tracker isolation with parameters: $R_m = 0.1$, $R_S = 0.04$, $R_i = 0.5$, $p_T^i = 1$ GeV/c.
- cut on the transverse momentum of the leading track $p_T > 35$ GeV/c
- number of tracks in the signal cone N_{tr}^S is one or three for $M_A = 200$ GeV/ c^2 . For higher Higgs boson masses the effective background rejection is only possible requiring only one track in the signal cone.

Finally, the two τ -jet candidates were required to have an opposite charge. The charge was calculated as the sum of charges of the tracks in the signal cone.

After identification of two τ jets the other jets in the event are considered. It was required to have only one additional jet with not calibrated energy $E_T^{raw} > 20$ GeV and $|\eta| < 2.4$ and it must be tagged as b jet. The b-jet identification was performed using the impact parameter tagging in 3D space [144]. The jet has to have at least three tracks with the impact parameter significance more than two. The purity of the b-tagged jet for the signal is very high ($> 95\%$).

The di- τ -jet mass reconstruction efficiency is determined by the requirements to have the positive reconstructed energy of both neutrinos, $E_T^{\nu_1, \nu_2} > 0$. In the missing E_T corrections the jets with raw energy $E_T^{raw} > 25$ were used.

5.2.6.3 Expected number of selected events

The section summarizes the event selections, the corresponding cross sections and expected number of events for the signal and the background processes after the selections. The efficiency of the every selection shown in the tables of this section was evaluated relative to the previous selection.

The signal.

The Table 5.11 summarizes the expectations for the signal of $M_A = 200, 500$ and 800 GeV/ c^2 . The signal cross sections and the branching ratios were obtained for the m_h^{max} scenario with $\mu = 200$ GeV (see Section 11.3.1).

Table 5.11: The summary table of the selections for the signal of $M_A = 200, 500$ and 800 GeV/c^2 .

	$m_A = 200 \text{ GeV}/c^2$ $\tan(\beta) = 20$	$m_A = 500 \text{ GeV}/c^2$ $\tan(\beta) = 30$	$m_A = 800 \text{ GeV}/c^2$ $\tan(\beta) = 40$
cross sections and branching ratios			
$\sigma(\text{gg} \rightarrow \text{bb}(A+H))$ (fb)	45795+44888	2741+2744	677+677
$\text{BR}(H/A \rightarrow \tau\tau)$	0.1	0.082	0.087
$\text{BR}(\tau \rightarrow \text{hadrons})^2$	0.65 \times 0.65		
$\sigma \times \text{BR}$ (fb)	3831	190	49.8
Experimental selection efficiencies			
Level 1 Trigger	0.506	0.854	0.896
HLT	0.289	0.319	0.314
Two off-line calo τ jets	0.997	0.999	0.999
cuts on E_T τ jets	0.430	0.755	0.780
Two off-line τ candidates	0.674	0.716	0.675
$p_T^{\text{ltr}} > 35 \text{ GeV}/c$	0.326	0.616	0.713
Tracker Isolation	0.859	0.950	0.954
N_{tracks} in signal cone	0.81	0.67	0.78
$Q_{\tau 1} \times Q_{\tau 2} = -1$	0.98	0.94	0.94
≥ 1 extra jet, $E_T^{\text{raw}} > 20 \text{ GeV}$, $ \eta < 2.4$	0.21	0.27	0.31
only 1 extra jet, $E_T^{\text{raw}} > 20 \text{ GeV}$, $ \eta < 2.4$	0.83	0.82	0.78
$M_{\tau\tau}$ reconstruction efficiency			
$E_{\tau 1, \tau 2} > 0$	0.93	0.93	0.92
$E_{\nu 1, \nu 2} > 0$	0.56	0.67	0.67
Total mass reco	0.52	0.62	0.62
b tagging of the extra jet	0.36	0.44	0.41
$M_{\tau\tau}$ mass window	150-300 GeV/c^2	400-700 GeV/c^2	600-1100 GeV/c^2
mass window efficiency	0.81	0.73	0.81
total efficiency	2.5×10^{-4}	2.4×10^{-3}	3.6×10^{-3}
σ after selections (fb)	0.96	0.46	0.19
N ev. at 60 fb^{-1}	58.0	27.0	11.0

The QCD multi-jet background

Despite of the huge amount of generated events (more than one million) and generation preselections, the statistics of the QCD multi-jet background events is not enough to ensure a large number of Monte Carlo events passing all the selections. In order to decrease the statistical uncertainties the factorization of the selections were applied. All selections were combined in three groups as shown in Table 5.12. The Group1 includes the Level 1 trigger and the calorimetric reconstruction of the τ jets (at HLT and offline). It includes also the cut on the transverse energy of the jets. After the event passed the Group1 selections the two other selection groups (Group2 and Group3) were applied independently. The Group2 is essentially the τ jet identification part of the analysis, i.e. the tracker isolation (at HLT and off-line), cut on the p_T of the leading track, selection on the number of tracks inside the signal cone. The Group3 describes the selections on the one extra jet in the event, the b tagging and the di- τ -jet mass reconstruction. The choice of the second and third selection groups was made minimizing the correlation among them. The further factorization was done for some selections inside the groups itself. Table 5.12 summarizes the selections and the QCD multi-jet background estimates for the signal of $M_A=200 \text{ GeV}/c^2$. The requirement to have the opposite charge τ -jet candidates ($Q_1 \times Q_2 = -1$) is not included in the Table 5.12. It reduces the QCD multi-jet background by another factor of two, leading to 104 events of the QCD multi-jet background expected with 60 fb^{-1} . With the selections applied to search for the signal of $M_A=500 \text{ GeV}/c^2$ and $M_A=800 \text{ GeV}/c^2$ the expected number of the QCD multi-jet background with 60 fb^{-1} is 25.0 and 4.0 correspondingly.

The irreducible background.

The irreducible background after all selections were applied is the small part of the total background dominated by the QCD multi-jet events. The Table 5.13 summarizes the expected number of events from the irreducible backgrounds with 60 fb^{-1} for the selections used to search for the signal of $M_A=200 \text{ GeV}/c^2$. In total, 6.0 events is expected. The efficiencies of some of the selections are also shown in the Table. With the selections applied to search for the signal of $M_A=500 \text{ GeV}/c^2$ and $M_A=800 \text{ GeV}/c^2$ the expected number of the irreducible background with 60 fb^{-1} is 4.0 and 1.0 correspondingly.

5.2.6.4 The detector effects, the experimental systematics and the evaluation of the background from the data.

The effect of the E_T^{miss} and the jet energy scale uncertainties.

The effect of the E_T^{miss} and the jet energy scale uncertainty on the Higgs boson mass reconstruction efficiency was estimated. The E_T^{miss} is reconstructed with the Type 1 corrections in the following form:

$$E_{T(x(y))}^{\text{miss}} = -(E_{T(x(y))}^{\text{raw}} + \sum_{\text{jets}} (E_{T(x(y))}^{\text{corr.jet}} - E_{T(x(y))}^{\text{rawjet}})) \quad (5.11)$$

where $E_{T(x(y))}^{\text{raw}}$ is the sum over the raw calorimeter tower energies from calorimeter towers and the jet sum in the equation is over jets with a reconstructed $E_T^{\text{raw}} > 25 \text{ GeV}$. The formula can be rewritten in the form:

$$E_{T(x(y))}^{\text{miss}} = -([E_{T(x(y))}^{\text{raw}} - \sum_{\text{jets}} E_{T(x(y))}^{\text{rawjet}}]_{\text{low } E_T} + [\sum_{\text{jets}} E_{T(x(y))}^{\text{corr.jet}}]_{\text{high } E_T}) \quad (5.12)$$

Table 5.12: The summary table of the selections for the QCD multi-jet background. The selections are factorized as explained in the text. The requirement to have the opposite charge τ -jet candidates ($Q_1 \times Q_2 = -1$) is not included.

	QCD di-jet background in bins of generated \hat{p}_T			
	>170 GeV/ c	120-170 GeV/ c	80-120 GeV/ c	50-80 GeV/ c
σ (fb)	1.33×10^8	5.03×10^8	2.94×10^9	2.08×10^{10}
$\varepsilon_{\text{kine pres.}}$	2.12×10^{-1}	4.19×10^{-2}	5.77×10^{-3}	2.44×10^{-4}
Group1 cuts: L1 trigger + L2 and offline calo reco + E_T cut				
Level 1 trigger	0.562	0.726	0.715	0.461
Two Level 2 calo jets with $\Delta R_{JJ} > 1.0$	0.927	0.959	0.982	0.987
Two off-line calo τ jets	0.975	0.975	0.982	0.994
cuts on E_T τ jets	0.753	0.804	0.774	0.343
$\varepsilon_{\text{Group1}}$	0.383	0.547	0.534	0.155
Group2 cuts: τ-jet identification at HLT and off-line				
HLT Calo+Pxl τ trigger	7.15×10^{-4}	1.81×10^{-3}	4.44×10^{-3}	1.12×10^{-2}
Two off-line τ candidates	0.86	0.84	0.825	0.84
$p_T^{\text{thr}} > 35$ GeV/ c	0.47	0.41	0.42	0.38
Tracker isolation	0.24	0.21	0.25	0.35
<i>Factorized inside group 2</i>				
1 or 3 prongs in 1 st τ jet	0.66	0.92	0.63	0.72
1 or 3 prongs in 2 nd τ jet	0.48	0.54	0.65	0.72
$\varepsilon_{\text{Group2}}/\varepsilon_{\text{Group1}}$	2.30×10^{-5}	6.33×10^{-5}	1.63×10^{-4}	6.54×10^{-4}
Group3 cuts: extra jet reco and b tagging plus $M_{\tau\tau}$ reco and mass window				
≥ 1 extra jet, $E_T^{\text{raw}} > 20$ GeV, $ \eta < 2.4$	0.463	0.235	0.127	0.090
only 1 extra jet, $E_T^{\text{raw}} > 20$ GeV, $ \eta < 2.4$	0.661	0.817	0.863	0.855
<i>Factorized inside group 3: $M_{\tau\tau}$ reco and b tagging</i>				
$E_{\tau_1, \tau_2} > 0$	0.921	0.898	0.882	0.834
$E_{\nu_1, \nu_2} > 0$	0.701	0.683	0.657	0.625
Total mass reco	0.646	0.613	0.579	0.522
b tagging of the extra jet	0.098	0.050	0.033	0.016
$M_{\tau\tau}$ window: 150-300 GeV/ c^2	0.142	0.295	0.433	0.430
$\varepsilon_{\text{Group3}}/\varepsilon_{\text{Group1}}$	2.77×10^{-3}	1.75×10^{-3}	9.15×10^{-4}	2.28×10^{-4}
$\varepsilon_{\text{Group1}} \times \varepsilon_{\text{Group2}} \times \varepsilon_{\text{Group3}}$	2.44×10^{-8}	6.07×10^{-8}	7.98×10^{-8}	2.84×10^{-8}
σ after selections (fb)	0.69	1.28	1.35	0.144
N_{ev} with 60 fb^{-1}	41.4	76.7	81.2	8.7

5.2. Associated production of MSSM heavy neutral Higgs bosons $b\bar{b}H(A)$ with $H(A) \rightarrow \tau\tau$ 125

Table 5.13: The number of expected events with 60 fb^{-1} and efficiencies of some of the selections for the reducible backgrounds.

process	$N_{\text{exp. at}} \text{ at } 60\text{fb}^{-1}$	$Q_{\tau 1} \times Q_{\tau 2} = -1$	only one extra jet	b tag. jet	$M_{\tau\tau}$ window
$t\bar{t}$	0.64	0.96	0.36	0.42	0.11
W+j	0.33	0.81	0.15	0.06	0.12
Wt	0.26	0.96	0.49	0.44	0.23
$Z/\gamma^* \rightarrow \tau\tau$ in bins of generated $m_{\tau\tau}$					
$130 < m_{\tau\tau} < 300 \text{ GeV}/c^2$	3.80	0.96	0.23	0.06	0.61
$m_{\tau\tau} > 300 \text{ GeV}/c^2$	0.18	0.95	0.27	0.05	0.04
$\tau\tau b\bar{b}, m_{\tau\tau} > 100 \text{ GeV}/c^2$	0.86	0.98	0.39	0.44	0.38

consisting from the low and the high E_T parts. For the low E_T part the scale uncertainty of 10% was applied, while for the high E_T part 3% uncertainty was used. The variation of the scale is applied independently for the two parts to obtain the maximal upper and low deviations from the case with no uncertainty. It was found that the missing E_T^{miss} scale uncertainty brings the largest contribution to the uncertainty of the Higgs boson mass reconstruction efficiency. In the worst case the uncertainty reaches 3 %. The mean fitted value of the $M_{\tau\tau}$ distribution for the signal of $M_A=500 \text{ GeV}/c^2$ is varied from $-10 \text{ GeV}/c^2$ to $+16 \text{ GeV}/c^2$ relative to the mean value evaluated without the scale uncertainty taken into account.

tracker misalignment.

The effect of the tracker misalignment on the rate of the fake τ jets from the QCD multi-jet background was studied for the first data taking scenario (Scenario 1) and the long term data taking scenario (Scenario 2). The tracker isolation efficiency and the efficiency of the track counting in the signal cone (one or three tracks requirement) was compared with the performance of the perfect tracker alignment (Scenario 0).

It was found that in the Scenario 2 the QCD multi-jet background can be increased by $\simeq 11\%$ due to the change of the tracker isolation efficiency. The efficiency of the requirement to have one track in the signal cone is increased by $\simeq 10\%$ in the Scenario 2 relative to the perfect alignment.

The measurement of the QCD multi-jet background from the data.

Figure 5.3 (left plot) shows the expected $M_{\tau\tau}$ distribution for the signal of $M_A=200 \text{ GeV}/c^2$ (thick solid histogram), the QCD multi-jet background (dashed histogram), the irreducible background (thick dashed-dotted histogram) and the sum of the signal and background (normal solid histogram) with 60 fb^{-1} in the m_h^{max} scenario with $\tan\beta=20$. The right plot of Figure 5.3 presents the case of $M_A=500 \text{ GeV}/c^2$, $\tan\beta=30$. The QCD multi-jet background is the biggest background in this analysis. The following way to evaluate this background from the data is proposed. The control sample must be used where all signal selections are applied except the mass window and the requirement to have an opposite charge of the two τ -jet candidates. It is proposed to select, instead, the sample with the same charge of the two τ -jet candidates (SS sample). The contamination of the signal events and irreducible background is negligible in the SS sample, thus giving possibility to predict from the data the QCD multi-jet background in a given mass window from the number of event and the




Figure 5.3: The expected $M_{\tau\tau}$ distributions for the signal of $M_A=200 \text{ GeV}/c^2$, $\tan\beta=20$ (left plot) and $M_A=500 \text{ GeV}/c^2$, $\tan\beta=30$ (right plot) and the background with 60 fb^{-1} . Thick solid histogram - signal in the m_h^{max} scenario; dashed histogram - the QCD multi-jet background; thick dashed-dotted histogram - the irreducible background; normal solid histogram - signal plus background.

measured shape of di- τ -jet mass in SS sample. The expected number of the QCD multi-jet SS events after all selections, but the mass window, used for the signal of $M_A=200 \text{ GeV}/c^2$ is 380 with 60 fb^{-1} . Neglecting the uncertainty of the measured shape of the di- τ -jet mass, it leads to 5% statistical uncertainty of the QCD multi-jet background estimates under the signal mass window. For the $M_A=500$ (800) GeV/c^2 selections about of 80 (28) the SS QCD multi-jet events is expected, thus giving $\simeq 10$ (20) % statistical uncertainty.

5.2.6.5 Discovery reach in M_A - $\tan\beta$ plane

The Table 5.14 shows the lowest value of the $\tan\beta$ for three considered in the analysis the Higgs boson masses, where the 5σ discovery is possible with 60 fb^{-1} . It is shown with and without QCD multi-jet background systematic uncertainty taken into account. The significance of the discovery is calculated with the S_{cP} method.

Table 5.14: The low limit of $\tan\beta$ where the 5σ discovery is possible with 60 fb^{-1} .

low $\tan\beta$ limit for 5σ discovery	Higgs boson mass		
	$m_A = 200 \text{ GeV}/c^2$	$m_A = 500 \text{ GeV}/c^2$	$m_A = 800 \text{ GeV}/c^2$
no systematics	20	32	46
with systematics	21	34	49

The extension of the discovery reach to the lower values of $\tan\beta$ would be possible with the lower threshold on the energy of the additional jet in the event, provided that the fake jets will be then suppressed with the jet-tracks matching criteria. Another improvement is expected from the increase of the Higgs boson mass reconstruction efficiency using the improved missing E_T measurement from the energy-flow like algorithms. Finally, the im-

Improved b-jet tagging performance is expected to extend the discovery reach to the lower $\tan\beta$ region.

5.2.7 $H \rightarrow \tau\tau \rightarrow \mu$ +jet analysis

5.2.7.1 Event generation and preselections

For the irreducible Drell Yan (DY) $\tau\tau$ background the $\tau_{1(2)} \rightarrow \mu\nu\nu$, $\tau_{2(1)} \rightarrow \text{hadrons} + \nu$ decays were forced in PYTHIA. The events containing b quarks were rejected to avoid the double counting with the $\tau\tau b\bar{b}$ background.

For the other background processes, $t\bar{t}$, Wt , W +jet and $b\bar{b}$ no specific decay mode was forced.

The DY $\tau\tau$ background was produced in two ranges of the $\tau\tau$ invariant mass: $40 < m_{\tau\tau} < 120 \text{ GeV}/c^2$ and $m_{\tau\tau} > 120 \text{ GeV}/c^2$. For $\tau\tau b\bar{b}$ the following mass bins were used: $60 < m_{\tau\tau} < 100 \text{ GeV}/c^2$ and $m_{\tau\tau} > 100 \text{ GeV}/c^2$. The W +jet background was generated with $\hat{p}_T > 20 \text{ GeV}/c^2$.

The SUSY background has been estimated using the events for the LM2 mSUGRA test point (see Section `refsusy:scope.testpts`) with the total NLO SUSY cross section of 9.4 pb. For this point $\tan(\beta)=35$, which makes the stau and tau production rate potentially dangerous. The number of events after all selection has been estimated to be less than one, therefore the SUSY background has been considered negligible, and was not studied in detail.

For the signal generation Higgs boson was forced to decay into τ pair. The τ leptons were decayed using TAUOLA and events with the $\tau_{1(2)} \rightarrow \mu\nu\nu$, $\tau_{2(1)} \rightarrow \text{hadrons} + \nu$ decays were selected.

The preselections at generation level were chosen in a way that selected events likely to pass the trigger selection. The requirements were: isolation of the muon defined as no charged particles with $p_T > 1 \text{ GeV}/c$ within cone of radius $r=0.2$ in the $\eta - \varphi$ space around the muon momentum direction. Isolation for the τ -like jet allowed for at most one charged particle with $p_T > 1 \text{ GeV}/c$ in the ring with the inner radius $R_{in}=0.1$, and the outer radius $R_{out}=0.4$ around the highest p_T charged particle in the jet. The leading track was required to have $p_T > 3 \text{ GeV}/c$. The $\tau\tau b\bar{b}$ events were generated without the preselection requirements.

Details on $b\bar{b}$ generation are explained in [145].

5.2.7.2 Event selection

The off-line τ -jet identification uses the parameters of the pixel HLT τ isolation, but with the full reconstructed tracks instead of the pixel tracks. Additionally one or three tracks in the signal cone are required. For the τ jet direction, the sum of the momenta of the signal tracks was used, improving the direction resolution. The leading τ -jet track is required to have $p_T > 10 \text{ GeV}/c$ in case of the one track in the signal cone, and $p_T > 20 \text{ GeV}/c$ for the three tracks, in order to suppress the $b\bar{b}$ and DY $\tau\tau$ backgrounds.

To select events with the associated $b\bar{b}H(A)$ production, the one b-tagged jet with calibrated $E_T > 20 \text{ GeV}/c$ was required. For the b tagging, the track counting method was used [144]: the jet is b tagged if it has at least two tracks with 2D transverse impact parameter significance ($S = d_{IP}/\sigma_d$) greater than two. The b tagging efficiency, including the jet finding, for the signal is 0.17 for $M_A = 200 \text{ GeV}/c^2$ and 0.27 for $M_A = 500 \text{ GeV}/c^2$. For the backgrounds

with the real b jet it is 0.67 for $t\bar{t}$ and 0.46 for Wt processes. For the backgrounds without the real b jet the mis-tag efficiency is 0.01 for the W +jet and 0.03 for the DY $\tau\tau$ processes. The b tagging purity for the signal and the $t\bar{t}$ background is 95% and it is 90% for the Wb and the $\tau\tau b\bar{b}$ processes.

Events containing W bosons decaying into $\mu + \nu_\mu$ are suppressed using the cut on the transverse mass of the muon and the missing transverse energy: $m_T = \sqrt{2 \cdot p_T^\mu \cdot \cancel{E}_T (1 - \cos(\vec{p}_T^\mu, \vec{\cancel{E}}_T))}$, where \cancel{E}_T is the missing transverse energy. The m_T has a Jacobian peak near the W mass. Rejecting events with $m_T > 60$ GeV largely reduces the $t\bar{t}$, Wt and W +jet backgrounds retaining good fraction of the signal events.

The additional selection against the $t\bar{t}$ background is the central jet veto. All events containing the additional jet (to the τ jet and the b -tagged jet) in the central region: $|\eta| < 2.5$ and with the calibrated $E_T > 20$ GeV were rejected.

The electrons from the W boson decays in the $t\bar{t}$ and Wt backgrounds can be misidentified as the τ jets. For the electron rejection a cut on the ratio of the τ -jet energy measured in the HCAL (E^{HCAL}) to the leading track momentum (p^{ltr}), $f = E^{\text{HCAL}}/p^{\text{ltr}}$, was used for the events with one track in the signal cone. The cut $f > 0.2$ retains 90% of the signal events, while rejects 95% of the events with the real electrons. The cut on the upper value of the ratio is efficient against the jets with the large fraction of the neutral hadrons. The requirement $f < 1.1$ rejects 50% of W +j and $b\bar{b}$ events and only 20% of signal events. Figure 5.4 shows the integrated distribution of the parameter f for the signal and the background events selected by High Level trigger. The labels on the right part of the figure are ordered by the decreasing of the rejection efficiency in the acceptance region of $0.2 < f < 1.1$, marked by the arrows.




Figure 5.4: The integrated distribution of the parameter $f = E^{\text{HCAL}}/p^{\text{ltr}}$. The acceptance region of $0.2 < f < 1.1$ is marked by the arrows (more explanations see in the text).

The Higgs boson mass reconstruction requires the rejection of the events with μ and τ jet back-to-back topology, therefore the cut $\cos(\Delta\varphi(\vec{p}_T, \vec{E}_T^{\text{jet}})) > -0.9962$ was applied. In addition, the upper cut on $\cos(\Delta\varphi(\vec{p}_T, \vec{E}_T^{\text{jet}})) < -0.5$ was used, retaining most of the signal events, while visibly reducing a fraction of the background events. Finally, the events with

5.2. Associated production of MSSM heavy neutral Higgs bosons $b\bar{b}H(A)$ with $H(A) \rightarrow \tau\tau$ 129

the negative reconstructed neutrino energy were rejected.

5.2.7.3 Expected number of selected events

The Table 5.15 presents the production cross sections in fb and the individual selection efficiencies for the signal of $M_A = 200$ and $500 \text{ GeV}/c^2$. The signal cross sections and the branching ratios were obtained for the m_h^{max} scenario with $\mu=200 \text{ GeV}$ (see Section 11.3.1). The Tables 5.16-5.18 summarize the cross sections and the individual selection efficiencies for the background processes. The total efficiency of all selections and the cross sections after all selections are also presented at the end of the tables. The events were counted

Table 5.15: The production cross sections, in fb, and the individual selection efficiencies for signal.



	$gg \rightarrow b\bar{b}(A + H), A, H \rightarrow \tau\tau$	
	$M_A=200 \text{ GeV}/c^2$ $\tan(\beta)=20$	$M_A=500 \text{ GeV}/c^2$ $\tan(\beta)=30$
$\sigma \times \text{BR} [\text{fb}]$	$9.12 \cdot 10^3$	$4.51 \cdot 10^2$
kine presel	$9.47 \cdot 10^{-2}$	$1.65 \cdot 10^{-1}$
Level-1	$8.99 \cdot 10^{-1}$	$9.09 \cdot 10^{-1}$
HLT	$4.17 \cdot 10^{-1}$	$4.99 \cdot 10^{-1}$
Offline τ -jet isolation	$9.54 \cdot 10^{-1}$	$9.60 \cdot 10^{-1}$
1 or 3 tk. in τ -jet signal cone	$9.12 \cdot 10^{-1}$	$9.19 \cdot 10^{-1}$
$p_T^{\text{tr}} > 10 \text{ GeV}/c$	$9.05 \cdot 10^{-1}$	$9.55 \cdot 10^{-1}$
$Q_\mu \cdot Q_{jet} = -1$	$9.61 \cdot 10^{-1}$	$9.60 \cdot 10^{-1}$
Single b tagging	$1.73 \cdot 10^{-1}$	$2.56 \cdot 10^{-1}$
No jet with $E_T > 20, \eta < 2.5$	$8.53 \cdot 10^{-1}$	$7.72 \cdot 10^{-1}$
$m_T(l, MET) < 60 \text{ GeV}$	$8.33 \cdot 10^{-1}$	$7.01 \cdot 10^{-1}$
$-0.996 < \cos(\Delta\varphi) < -0.5$	$8.05 \cdot 10^{-1}$	$7.51 \cdot 10^{-1}$
electron veto: $0.2 < f < 1.1$	$8.22 \cdot 10^{-1}$	$8.54 \cdot 10^{-1}$
$E_{\nu 1} > 0, E_{\nu 2} > 0$	$6.84 \cdot 10^{-1}$	$7.68 \cdot 10^{-1}$
Total efficiency:	$1.66 \cdot 10^{-3}$	$4.53 \cdot 10^{-3}$
σ after selections [fb]:	$1.52 \cdot 10^1$	2.05

in the $M_{\tau\tau}$ mass windows with the width taken to be $\pm\sigma$, where σ is given by the Gaussian fit of the signal $M_{\tau\tau}$ distributions. For the $M_A = 200 \text{ GeV}/c^2$ the sigma is $41 \text{ GeV}/c^2$, and it is 83 for the $m_A = 500 \text{ GeV}/c^2$. It is expected 146 signal and 127 background events for the $m_A = 200 \text{ GeV}/c^2, \tan(\beta) = 20$, and 21 signal with 61 background events for the $m_A = 500 \text{ GeV}/c^2, \tan(\beta) = 30$ with the integrated luminosity of 20 fb^{-1} . Figure 5.5 shows the expected $\tau\tau$ mass distribution for the total background and for the signal plus background for $M_A = 200 \text{ GeV}/c^2, \tan(\beta)=20$ and $M_A = 500 \text{ GeV}/c^2, \tan(\beta)=30$ with 20 fb^{-1} .

5.2.7.4 Background estimates and uncertainty

After all off-line selections the main background is represented by the $\tau\tau b\bar{b}$, DY $\tau\tau$ and the $t\bar{t}$ production processes. The contribution of the non Z/γ^* background, mainly the $t\bar{t}$ events,

Table 5.16: The production cross sections, in fb, and the individual selection efficiencies for irreducible background processes.

	$t\bar{t}$	$W + jet$	Wt	$b\bar{b}$
σ [fb]	$8.40 \cdot 10^5$	$4.15 \cdot 10^7$	$6.20 \cdot 10^4$	$2.29 \cdot 10^{10}$
kine presel	$9.01 \cdot 10^{-2}$	$1.44 \cdot 10^{-2}$	$6.58 \cdot 10^{-2}$	$7.56 \cdot 10^{-4}$
Level-1	$9.06 \cdot 10^{-1}$	$8.40 \cdot 10^{-1}$	$8.91 \cdot 10^{-1}$	$2.26 \cdot 10^{-2}$
HLT	$9.61 \cdot 10^{-2}$	$4.16 \cdot 10^{-2}$	$1.05 \cdot 10^{-1}$	$2.36 \cdot 10^{-4}$
Offline τ -jet isolation	$8.51 \cdot 10^{-1}$	$6.70 \cdot 10^{-1}$	$8.79 \cdot 10^{-1}$	$8.69 \cdot 10^{-1}$
1 or 3 tk. in τ -jet signal cone	$8.92 \cdot 10^{-1}$	$6.30 \cdot 10^{-1}$	$9.07 \cdot 10^{-1}$	$7.19 \cdot 10^{-1}$
$p_T^{\text{tr}} > 10 \text{ GeV}/c$	$9.42 \cdot 10^{-1}$	$8.58 \cdot 10^{-1}$	$9.37 \cdot 10^{-1}$	$7.17 \cdot 10^{-1}$
$Q_\mu \cdot Q_{jet} = -1$	$9.18 \cdot 10^{-1}$	$7.31 \cdot 10^{-1}$	$9.52 \cdot 10^{-1}$	$5.45 \cdot 10^{-1}$
Single b tagging	$6.73 \cdot 10^{-1}$	$1.09 \cdot 10^{-2}$	$4.56 \cdot 10^{-1}$	$9.42 \cdot 10^{-2}$
No jet with $E_T > 20, \eta < 2.5$	$3.43 \cdot 10^{-1}$	$8.17 \cdot 10^{-1}$	$8.60 \cdot 10^{-1}$	$4.30 \cdot 10^{-1}$
$m_T(l, MET) < 60 \text{ GeV}$	$3.53 \cdot 10^{-1}$	$3.76 \cdot 10^{-1}$	$3.62 \cdot 10^{-1}$	1.00
$-0.996 < \cos(\Delta\varphi) < -0.5$	$4.95 \cdot 10^{-1}$	$6.56 \cdot 10^{-1}$	$4.51 \cdot 10^{-1}$	$4.16 \cdot 10^{-1}$
electron veto: $0.2 < f < 1.1$	$1.65 \cdot 10^{-1}$	$4.76 \cdot 10^{-1}$	$1.27 \cdot 10^{-1}$	$2.98 \cdot 10^{-1}$
$E_{\nu 1} > 0, E_{\nu 2} > 0$	$4.08 \cdot 10^{-1}$	$2.00 \cdot 10^{-1}$	$4.15 \cdot 10^{-1}$	$3.60 \cdot 10^{-1}$
Total efficiency:	$1.54 \cdot 10^{-5}$	$3.31 \cdot 10^{-8}$	$1.66 \cdot 10^{-5}$	$7.86 \cdot 10^{-11}$
σ after selections [fb]:	$1.30 \cdot 10^1$	1.37	1.03	1.80

Table 5.17: The production cross sections, in fb, and the individual selection efficiencies for irreducible background processes.

	$Z/\gamma^* \rightarrow \tau\tau \rightarrow \mu + \text{jet}$	
	$40 < m_{\tau\tau} < 120 \text{ GeV}/c^2$	$m_{\tau\tau} > 120 \text{ GeV}/c^2$
$\sigma \times \text{BR}$ [fb]	$4.63 \cdot 10^5$	$4.88 \cdot 10^3$
kine presel	$6.56 \cdot 10^{-2}$	$2.14 \cdot 10^{-1}$
Level-1	$8.00 \cdot 10^{-1}$	$8.28 \cdot 10^{-1}$
HLT	$1.03 \cdot 10^{-1}$	$2.77 \cdot 10^{-1}$
Offline τ -jet isolation	$9.12 \cdot 10^{-1}$	$9.40 \cdot 10^{-1}$
1 or 3 tk. in τ -jet signal cone	$9.03 \cdot 10^{-1}$	$8.93 \cdot 10^{-1}$
$p_T^{\text{tr}} > 10 \text{ GeV}/c$	$8.12 \cdot 10^{-1}$	$9.00 \cdot 10^{-1}$
$Q_\mu \cdot Q_{jet} = -1$	$9.47 \cdot 10^{-1}$	$9.33 \cdot 10^{-1}$
Single b tagging	$2.68 \cdot 10^{-2}$	$2.51 \cdot 10^{-2}$
No jet with $E_T > 20, \eta < 2.5$	$7.77 \cdot 10^{-1}$	$6.98 \cdot 10^{-1}$
$m_T(l, MET) < 60 \text{ GeV}$	$9.41 \cdot 10^{-1}$	$7.74 \cdot 10^{-1}$
$-0.996 < \cos(\Delta\varphi) < -0.5$	$3.75 \cdot 10^{-1}$	$6.57 \cdot 10^{-1}$
electron veto: $0.2 < f < 1.1$	$6.46 \cdot 10^{-1}$	$7.29 \cdot 10^{-1}$
$E_{\nu 1} > 0, E_{\nu 2} > 0$	$6.45 \cdot 10^{-1}$	$6.46 \cdot 10^{-1}$
Total efficiency:	$1.31 \cdot 10^{-5}$	$1.75 \cdot 10^{-4}$
σ after selections [fb]:	6.08	$8.53 \cdot 10^{-1}$

5.2. Associated production of MSSM heavy neutral Higgs bosons $b\bar{b}H(A)$ with $H(A) \rightarrow \tau\tau$ 131

Table 5.18: The production cross sections, in fb, and the individual selection efficiencies for irreducible background processes.

	$bb(Z \rightarrow \tau\tau)$	
	$60 < m_{\tau\tau} < 100 \text{ GeV}/c^2$	$m_{\tau\tau} > 100 \text{ GeV}/c^2$
$\sigma \times \text{BR}$ [fb]	$2.61 \cdot 10^4$	$1.05 \cdot 10^3$
kine presel	1.00	1.00
Level-1	$1.41 \cdot 10^{-1}$	$1.64 \cdot 10^{-1}$
HLT	$4.10 \cdot 10^{-3}$	$1.21 \cdot 10^{-2}$
Offline τ -jet isolation	$9.05 \cdot 10^{-1}$	$9.34 \cdot 10^{-1}$
1 or 3 tk. in τ -jet signal cone	$9.12 \cdot 10^{-1}$	$9.17 \cdot 10^{-1}$
$p_T^{\text{tr}} > 10 \text{ GeV}/c$	$8.60 \cdot 10^{-1}$	$8.98 \cdot 10^{-1}$
$Q_\mu \cdot Q_{jet} = -1$	$9.41 \cdot 10^{-1}$	$9.48 \cdot 10^{-1}$
Single b tagging	$2.73 \cdot 10^{-1}$	$2.75 \cdot 10^{-1}$
No jet with $E_T > 20, \eta < 2.5$	$7.20 \cdot 10^{-1}$	$7.72 \cdot 10^{-1}$
$m_T(l, MET) < 60 \text{ GeV}$	$9.68 \cdot 10^{-1}$	$8.80 \cdot 10^{-1}$
$-0.996 < \cos(\Delta\varphi) < -0.5$	$4.23 \cdot 10^{-1}$	$5.84 \cdot 10^{-1}$
electron veto: $0.2 < f < 1.1$	$6.98 \cdot 10^{-1}$	$5.11 \cdot 10^{-1}$
$E_{\nu 1} > 0, E_{\nu 2} > 0$	$4.32 \cdot 10^{-1}$	$5.62 \cdot 10^{-1}$
Total efficiency:	$6.64 \cdot 10^{-5}$	$2.76 \cdot 10^{-4}$
σ after selections [fb]:	1.74	$2.89 \cdot 10^{-1}$




Figure 5.5: The reconstructed $\tau\tau$ mass distribution. The signal and the background contributions are shown with 20 fb^{-1} . The mass windows in which the events are counted for the significance calculations are shown.

can be estimated applying the inversion of the electron veto: $f < 0.1$ instead of $0.2 < f < 1.1$. All other cuts must be the same, including the $M_{\tau\tau}$ mass window. A relatively pure sample of the $t\bar{t}$ can be selected, since the requirement $f < 0.1$ rejects more than 95% of all processes except the $t\bar{t}$ and Wt as shown in Fig. 5.4. The number of the non Z/γ^* background events in the signal region can be then predicted using the ratio of the $t\bar{t}$ events in the signal region of $0.2 < f < 1.1$ and in the the region of $f < 1$. This ratio can be obtained from the Monte-Carlo (or from the real $t\bar{t}$ data). The systematic uncertainty on the number of the non Z/γ^* background

events predicted using this method has two contributions:

- the uncertainty of the HCAL energy scale, since the variable $f = E^{\text{HCAL}}/p^{\text{ltr}}$ includes the HCAL part of the τ -jet candidate energy measured by the calorimeter. It is taken as 3%.
- the uncertainty of the shape of the variable f . The shape is obtained from the $t\bar{t}$ events only, however the small fraction of the events from the other processes is presented in the "normalization" region of $f < 0.1$. It leads to the uncertainty of $\simeq 12\%$.

The contribution from the other systematic uncertainties, e.g. b tagging is expected to be small, due to the cancellation in the efficiency ratio. The total uncertainty on the number of the non Z/γ^* background events is thus 12.4 %.

The Z/γ^* background consists of the two parts: the $\tau\tau b\bar{b}$ process and the DY $\tau\tau$ process without the genuine b quarks in the event. The DY $\tau\tau$ background can be predicted using the DY $\ell\ell$ ($\ell = e, \mu$) cross section measured with the high precision at LHC and the selection efficiency obtained from the Monte-Carlo. The systematic uncertainty on the number of the DY $\tau\tau$ events has two main contributions due to:

- the jet scale uncertainty. The number of the events in the $M_{\tau\tau}$ signal window varies by $\pm 6\%$ for the jet scale variations of $\pm 3\%$ and the missing transverse energy scale variations of $\pm 5\%$.
- the b -misstaging uncertainty. The conservative estimate of 5% is taken.

The total uncertainty on the number of the DY $\tau\tau$ events with the jet misstaged as the b jet is therefore 8%.

For the $\tau\tau b\bar{b}$ background estimates the systematic uncertainty has the following main contributions:

- the uncertainty of the $\mu\mu b\bar{b}$ cross section measurement (without the luminosity uncertainty) is 14% [139].
- the jet scale uncertainty. It is assumed the same as for the DY $\tau\tau$ events.

The total uncertainty on the number of the $\tau\tau b\bar{b}$ events is 15%.

5.2.7.5 Discovery reach in $M_A - \tan(\beta)$ plane

The CMS discovery reach in the $M_A - \tan(\beta)$ plane with 30 fb^{-1} in the m_h^{max} scenario is shown in the Fig. 5.6. The 5σ discovery curves are shown without (lower curve) and with (upper curve) the uncertainty on the background taken into account.

5.2.8 $H \rightarrow \tau\tau \rightarrow e + \text{jet}$ analysis

5.2.8.1 Event generation

The signal process $gg \rightarrow bbH/A$, $H/A \rightarrow \tau\tau$, $\tau_1 \rightarrow e\nu_e\nu_\tau$, $\tau_2 \rightarrow \tau \text{ jet} + \nu_\tau$ leads to a final state of one isolated electron, an isolated τ jet and one or two detectable b jets. The background with genuine τ 's is due to two types of events, Z/γ^* events decaying into $\tau\tau$, and the $t\bar{t}$




Figure 5.6: The 5σ discovery region in the $M_A - \tan(\beta)$ plane with the 30 fb^{-1} of the integrated luminosity for the m_h^{max} MSSM scenario. The regions are shown without (lower curve) and with (upper curve) the uncertainty on the background taken into account.

events, where the $e + \tau$ jet final state can come from direct W decays to an electron and a τ or through $W \rightarrow \tau\nu_\tau \rightarrow e\nu_e\nu_\tau\nu_\tau$ decays:

- $Z/\gamma^* \rightarrow \tau\tau \rightarrow e + \tau \text{ jet} + X$
- $b\bar{b}Z/\gamma^*, Z/\gamma^* \rightarrow \tau\tau \rightarrow e + \tau \text{ jet} + X$
- $t\bar{t}$ with $W_1 \rightarrow \tau\nu_\tau$ ($\tau \rightarrow \text{jet}$), $W_2 \rightarrow e\nu_e$ or $W \rightarrow \tau\nu_\tau \rightarrow e\nu_e\nu_\tau\nu_\tau$.
- Wt , with $W_1 \rightarrow \tau\nu_\tau$ ($\tau \rightarrow \text{jet}$), $W_2 \rightarrow e\nu_e$ or $W \rightarrow \tau\nu_\tau \rightarrow e\nu_e\nu_\tau\nu_\tau$.

Background can arise also from the processes where a hadronic jet or an electron leads to a fake τ :

- $W + \text{jet}$, with $W \rightarrow e\nu_e$
- $Z/\gamma^* \rightarrow e^+e^-$
- $b\bar{b}Z/\gamma^*, Z/\gamma^* \rightarrow e^+e^-$
- $t\bar{t}$ with $W \rightarrow jj$ or $W \rightarrow e\nu_e$.

The QCD multi-jet production is a large potential background through hadronic jets faking both the electron and the τ jet.

For the inclusive Z/γ^* production the events containing b quarks in the final state were removed to avoid double counting with the $\tau\tau b\bar{b}$ background. The single top (Wt) events were generated with TopREX [35]. The τ decays in the signal were performed with the TAUOLA package [146].

5.2.8.2 Event selection

In the offline reconstruction an isolated electron from the decay of one of the τ 's is first searched for. On the average ~ 1.3 reconstructed electron candidates were found in the signal events. The reconstructed electrons were first required to be isolated in the tracker demanding that no track with $p_T > 1 \text{ GeV}/c$ was found in a cone of $\Delta R = 0.4$ around the electron

candidate direction. The further electron identification was performed following the algorithm of Ref. [147]. The largest contribution to the identification efficiency and purity was obtained from the ratio of hadronic cluster energy to the electromagnetic energy of the cluster ($E^{\text{hadronic}}/E^{\text{elm}} < 0.2$) and from the ratio of the supercluster energy to the track momentum ($E^{\text{super cluster}}/p^{\text{track}} > 0.8$). The identification efficiency, including the tracker isolation, was found to be 64.2%. A good purity of 97.5% was obtained for the selected electrons.




Figure 5.7: Distribution of transverse mass reconstructed from the electron and the missing transverse energy for the signal of $M_A = 200 \text{ GeV}/c^2$ and $\tan\beta = 20$ (filled histogram), for the $t\bar{t}$ (solid line) and for the $Z/\gamma^* \rightarrow e^+e^-$ (dashed line) background. Histogram normalization is arbitrary.




Figure 5.8: Reconstructed Higgs boson mass for $m_A = 200 \text{ GeV}/c^2$ and $\tan\beta = 20$.

The off-line τ -jet identification was applied to the jets with $E_T^{\text{jet}} > 40 \text{ GeV}$ reconstructed in the calorimeter with the cone of 0.4. The leading track with $p_T^{\text{tr}} > 10 \text{ GeV}/c$ was searched for in a cone of $R_m=0.1$ around the τ -jet direction. For an efficient isolation against the hadronic jets in the W +jet and QCD multi-jet backgrounds, a small signal cone, $R_S=0.04$, around the leading track was used. About 83% of the $\tau^\pm \rightarrow \text{hadron}^\pm + n\pi^0 + \nu_\tau$ decays were found to be reconstructed as one prong τ 's. Due to the small signal cone selected, about 50% of the $\tau^\pm \rightarrow 3 \text{ hadrons}^\pm + n\pi^0 + \nu_\tau$ decays were reconstructed as one or two prong τ jets. The cut $p_T^{\text{tr}} > 20 \text{ GeV}/c$ was found to be optimal for the suppression of the hadronic jets, in the presence of the QCD multi-jet background. The isolation was performed counting tracks with $p_T^i > 1 \text{ GeV}/c$ in the area between the signal cone and the isolation cone, which was taken to be then same as the jet reconstruction cone, $R_i = 0.4$. Following to the method described in [140], at least eight hits were required in the full silicon tracker and an upper bound of 0.3 mm on the transverse impact parameter was set on the leading track in order to suppress the background from the fake tracks.

The $Z/\gamma^* \rightarrow e^+e^-$ and $b\bar{b}Z/\gamma^*, Z/\gamma^* \rightarrow e^+e^-$ backgrounds contain an isolated genuine electron to pass the electron cuts and are not significantly suppressed with the τ selection

5.2. Associated production of MSSM heavy neutral Higgs bosons $b\bar{b}H(A)$ with $H(A) \rightarrow \tau\tau$ 135

cuts. These electronic τ -candidates were suppressed requiring a large energy deposition in the hadron calorimeter. A cut in the E_T of the most energetic HCAL cell in the τ jet, $E_{T(\text{max HCAL cell})} > 2 \text{ GeV}$, was found to suppresses the electrons with a factor of ~ 7 . A further reduction was obtained comparing the HCAL energy and the leading track momentum of the τ jet. The cut $E^{\text{HCAL}}/p^{\text{ltr}} > 0.35$, applied on the one-prong τ candidates only, was found to suppress further the electronic τ candidates by a factor of ~ 1.8 . The W+jet events show a tail at large values of $E^{\text{HCAL}}/p^{\text{ltr}}$ due to the neutral hadron component of the hadronic jets and was suppressed with the cut $E^{\text{HCAL}}/p^{\text{ltr}} < 1.5$.

Efficiencies of the τ -jet selections are shown in Tables 5.19, 5.20 and 5.21. The purity of $\sim 97\%$ is obtained for the signal events. A rejection factor of ~ 400 was obtained for the QCD multi-jet events generated with $50 < \hat{p}_T < 80 \text{ GeV}/c$ when the τ -jet selections described above were applied.

Finally, the charges of the electron and τ jet were required to be opposite. The charge of the τ jet was calculated as the sum of charges of the tracks in the signal cone.

The missing transverse energy, E_T^{miss} , measurement can be exploited to suppress the $t\bar{t}$ background with an upper bound on the transverse mass $m_T(e, E_T^{\text{miss}})$ reconstructed from the electron and the missing transverse energy. Figure 5.7 shows the $m_T(e, E_T^{\text{miss}})$ distribution for the signal events with $M_A = 200 \text{ GeV}/c^2$ and for the $t\bar{t}$ and $Z/\gamma^* \rightarrow e^+e^-$ backgrounds with the electron and τ -jet selections. An upper bound $m_T(e, E_T^{\text{miss}}) < 40 \text{ GeV}/c^2$ reduces the $t\bar{t}$ background with a factor of ~ 4 .

The events were further selected when at least one jet (in addition to the τ jet) with calibrated $E_T^{\text{jet}} > 20 \text{ GeV}$ and $|\eta| < 2.5$ was found and tagged as the b jet. A probabilistic secondary vertex algorithm with a discriminator cut from Ref. [148] was used for b tagging. The cut in the discriminator was set to 0.8, which suppresses efficiently the Z/γ^* , W+jet and the potential multi-jet background. The efficiency to tag at least one jet, including the jet finding efficiency, is between 13 and 19% for the signal, below 1% for the Z/γ^* backgrounds and 1.3% for the W+jet background. For the signal events the purity of the b-tagged jets is very high (99%).

The $t\bar{t}$ background, with a genuine electron, τ and b jets, can not be significantly suppressed with the cuts described above. This background, however, was suppressed applying the jet veto: the event must contain only the one b-tagged jet with calibrated $E_T^{\text{jet}} > 20 \text{ GeV}$ and $|\eta| < 2.5$. The fake jets, which generally do not contain tracks from the signal vertex, were suppressed with a cut in the fraction of the track p_T sum to the jet E_T , $\alpha = \Sigma p_T^{\text{track}}/E_T^{\text{jet}}$. The cut $\alpha > 0.1$ was found to improve the veto efficiency for the signal by about 10%. The jet veto efficiency is around 60% in the signal and $\sim 5\%$ for the $t\bar{t}$ background.

For the reconstruction of the di- τ -lepton mass the events with the back-to-back e- τ -jet configuration were removed applying the the upper cut on the angle between the τ jet and the electron in the transverse plane, $\Delta\varphi(e, \tau \text{ jet})$. The further cut on the reconstructed neutrino energies, $E_{\nu_1} > 0$ and $E_{\nu_2} > 0$, was applied leading to the reduction of $\simeq 40\%$ of the signal events. This requirement, however, rejects $\simeq 60\%$ of the $t\bar{t}$, tW and W+jet backgrounds. Figure 5.8 shows the reconstructed Higgs boson mass for the signal events with $M_A = 200 \text{ GeV}/c^2$. The Gaussian fit yields a mass resolution of 25%.

Table 5.19 shows the numbers of signal events with $M_A = 130\text{-}500 \text{ GeV}/c^2$ and $\tan\beta = 20$ for 30 fb^{-1} and the efficiency for all the event selection cuts described above. For $M_A = 130$

and $140 \text{ GeV}/c^2$, the mass of the lighter scalar Higgs boson h is only 4.4 and $11.2 \text{ GeV}/c^2$ smaller than m_A . With the mass resolution, which can be reached in the $H \rightarrow \tau\tau$ decay channels, the lighter scalar contributes to the signal and is added in the cross sections for $m_A = 130$ and $140 \text{ GeV}/c^2$. The contribution is 31 and 11% of the total production rate, respectively. Table 5.20 shows the number of events and efficiencies for the backgrounds originating from $Z/\gamma^* \rightarrow \tau\tau$ and $Z/\gamma^* \rightarrow e^+e^-$ decays in the inclusive and in the associated $b\bar{b}Z/\gamma^*$ production. The efficiency of removing the $b\bar{b}Z/\gamma^*$ component from the inclusive Z/γ^* samples is also shown. Table 5.21 shows the same for the backgrounds involving W 's from $t\bar{t}$, Wt and W +jet events. The cross section times branching fraction, trigger efficiency and the efficiency of the primary vertex reconstruction are also shown in the tables. The QCD multi-jet background after all selections was estimated to be 8.4 events for 30 fb^{-1} in the mass window around $M_A = 200 \text{ GeV}/c^2$, which is $\simeq 10\%$ of the all other backgrounds.

Figures 5.9 to 5.10 show the reconstructed Higgs boson mass distributions of the $H/A \rightarrow \tau^+\tau^- \rightarrow \text{electron} + \text{jet} + X$ signal and the total background for 30 fb^{-1} for $M_A = 200 \text{ GeV}/c^2$, $\tan\beta = 20$ and for $M_A = 300 \text{ GeV}/c^2$, $\tan\beta = 25$. The sum of the $Z/\gamma^* \rightarrow e^+e^-$ and $b\bar{b}Z/\gamma^* \rightarrow e^+e^-$ backgrounds is shown separately in the figures.

5.2.8.3 Systematic uncertainties for the background determination

The background uncertainty was evaluated using the cross-section uncertainties (measured or predicted from the theory) and the experimental uncertainties for the event selections.

The uncertainty of the event selection efficiency is related to the uncertainty of the electron and τ identification, the absolute calorimeter scale and the b -tagging efficiency. The systematic uncertainty due to the energy scale was estimated varying the jet energy and the E_T^{miss} values with the expected energy scale uncertainties yielding an average 5.1% uncertainty on the number of Z/γ^* events, 3.8% uncertainty on the number of $b\bar{b}Z/\gamma^*$ events, 7.3% uncertainty on the number of $t\bar{t}$ events, 11.3% uncertainty on the number of tW events and 11.8% uncertainty on the number of W +jet events passing the event selection cuts. The 5% uncer-

Table 5.19: The Higgs boson mass, production cross sections times branching fraction, efficiencies (%) for the selection cuts and numbers of events for 30 fb^{-1} for the signal with $\tan\beta = 20$.

m_A (GeV/c^2)	130	200	300	500
$\sigma \times \text{BR}$ (pb)	18.2	4.15	0.85	0.071
Level-1 and HLT	1.53 (8.4)	0.64 (15.4)	0.18 (21.6)	2.0×10^{-2} (28.7)
primary vertex	1.44 (94.1)	0.60 (94.2)	0.18 (97.2)	1.9×10^{-2} (93.6)
electron identification	1.11 (77.8)	0.48 (80.8)	0.14 (73.7)	1.4×10^{-2} (73.8)
one identified τ -jet	0.127 (11.4)	0.11 (23.4)	4.5×10^{-2} (32.9)	5.9×10^{-3} (41.7)
$Q^{\tau \text{ jet}} \times Q^e = -1$	0.127 (100.0)	0.11 (99.1)	4.5×10^{-2} (99.3)	5.8×10^{-3} (99.0)
$m_T < 40 \text{ GeV}/c^2$	9.9×10^{-2} (77.6)	3.8×10^{-2} (73.7)	3.1×10^{-2} (69.3)	3.9×10^{-3} (66.7)
≥ 1 jet, $E_T > 20 \text{ GeV}$	4.5×10^{-2} (45.9)	3.8×10^{-2} (46.6)	1.5×10^{-2} (48.6)	2.1×10^{-3} (53.5)
b tagging	1.3×10^{-2} (29.7)	1.2×10^{-2} (32.2)	5.0×10^{-3} (32.9)	7.6×10^{-4} (36.5)
jet veto	8.1×10^{-3} (60.2)	7.2×10^{-2} (62.5)	3.1×10^{-3} (63.2)	4.6×10^{-4} (61.0)
$\Delta\varphi(\tau_1, \tau_2) < 175^\circ$	7.6×10^{-3} (94.8)	6.8×10^{-3} (93.9)	2.7×10^{-3} (85.7)	3.4×10^{-4} (74.5)
$E_{\nu_1, \nu_2} > 0$	4.1×10^{-3} (54.1)	4.2×10^{-3} (61.7)	1.7×10^{-3} (64.3)	2.4×10^{-4} (70.6)
N_{ev} at 30 fb^{-1}	93.0	92.4	51.9	7.3

5.2. Associated production of MSSM heavy neutral Higgs bosons $b\bar{b}H(A)$ with $H(A) \rightarrow \tau\tau$ 137

Table 5.20: Background process, production cross sections times branching fraction, cross sections and efficiencies (%) for the selection cuts and number of events for 30 fb^{-1} .

	$Z/\gamma^* \rightarrow \tau\tau$	$b\bar{b}Z/\gamma^* \rightarrow \tau\tau$	$Z/\gamma^* \rightarrow e^+e^-$	$b\bar{b}Z/\gamma^*e^+e^-$
$\sigma \times \text{BR}$ (pb)	331.8	27.0	1890	26.3
preselection	173.5 (41.4)		811.2 (42.9)	
Level-1 and HLT	17.3 (10.0)	0.818 (3.1)	617.4 (76.1)	18.2 (67.2)
primary vertex	16.5 (95.4)	0.796 (97.3)	591.9 (95.9)	17.7 (97.3)
no b's in DY Z/γ^*	15.6 (94.6)		561.8 (94.9)	
electron identification	11.6 (74.4)	0.585 (80.2)	278.1 (50.1)	9.31 (52.6)
one identified τ -jet	0.13 (1.2)	1.0×10^{-2} (1.8)	3.40 (1.2)	9.0×10^{-2} (1.0)
$Q^{\tau \text{ jet}} \times Q^e = -1$	0.13 (96.3)	1.0×10^{-2} (100)	3.31 (97.4)	8.8×10^{-2} (97.8)
$m_T < 40 \text{ GeV}/c^2$	9.8×10^{-2} (76.3)	8.0×10^{-3} (80.0)	2.26 (68.3)	5.5×10^{-2} (62.5)
≥ 1 jet, $E_T > 20 \text{ GeV}$	4.0×10^{-2} (40.6)	5.6×10^{-3} (70.0)	0.85 (37.6)	3.0×10^{-2} (54.2)
b tagging	8.0×10^{-4} (2.0)	2.6×10^{-3} (46.4)	1.5×10^{-2} (1.8)	9.6×10^{-3} (32.2)
jet veto	5.2×10^{-4} (65.0)	1.5×10^{-3} (57.7)	6.0×10^{-3} (41.4)	5.9×10^{-3} (67.4)
$\Delta\varphi(\tau_1, \tau_2) < 175^\circ$	4.9×10^{-4} (94.2)	1.4×10^{-3} (90.7)	4.8×10^{-3} (80.0)	5.1×10^{-3} (85.7)
$E_{\nu_1, \nu_2} > 0$	2.0×10^{-4} (40.2)	7.6×10^{-4} (55.9)	1.7×10^{-4} (39.0)	1.9×10^{-3} (50.0)
N_{ev} at 30 fb^{-1}	5.9	22.8	51.3	57.9

tainty of the b tagging and mistagging efficiencies and the 2% uncertainty of the electron reconstruction and identification were used.

The uncertainty of the Z/γ^* cross section at LHC is the order of 1% [149]. For the $t\bar{t}$ background the theoretical NLO cross section uncertainty derives from the scale uncertainty, taken to be 5% according to Ref. [150], and the PDF uncertainty, $\sim 2.5\%$, yielding 5.6% for the total uncertainty. The same uncertainty is used for the cross sections of the Wt and W +jet processes. The uncertainty of the $b\bar{b}Z/\gamma^*$ cross section measurement is estimated to be 14.2% in [139]. With these estimates, the total systematic uncertainty, including the luminosity uncertainty of 3% [7], was found to be 8.1%, 15.9%, 11.1%, 14.0% and 14.5% for the Z/γ^* , $b\bar{b}Z/\gamma^*$, $t\bar{t}$, Wt and W +jet backgrounds, respectively.

5.2.8.4 Discovery reach in M_A - $\tan(\beta)$ plane.

Table 11.3 shows the number of signal plus background events and the number of background events for 30 fb^{-1} in the selected mass windows and the signal significance calculated according to Poisson statistics, with and without the background systematics taken into account. The mass windows were selected to optimize the significance. The prediction for the signal were obtained in the m_h^{max} scenario.

Figure 5.11 shows the 5σ discovery region in the M_A - $\tan\beta$ plane for 30 fb^{-1} for the m_h^{max} scenario. The lower (upper) curves was evaluated without (with) the effect of background systematics taken into account.

Table 5.21: Background process, production cross sections times branching fraction (pb), cross sections and efficiencies (%) for the selection cuts and number of events for 30 fb^{-1} . The QCD multi-jet events are generated with $50 < \hat{p}_T < 80 \text{ GeV}/c$.

	$t\bar{t}$	Wt	W+jet
$\sigma \times \text{BR}$ (pb)	840	6.16	673.2
preselection			315.0 (46.8)
Level-1 and HLT	94.4 (11.3)	2.00 (32.5)	145.6 (46.2)
primary vertex	93.9 (99.5)	1.97 (98.5)	143.9 (98.8)
electron identification	66.7 (71.0)	1.43 (72.6)	114.2 (79.4)
one id. τ -jet	0.66 (0.95)	4.10×10^{-2} (2.87)	0.57 (0.5)
$Q^{\tau \text{ jet}} \times Q^e = -1$	0.57 (89.8)	4.00×10^{-2} (97.6)	0.47 (82.7)
$m_T(e, E_T^{\text{miss}}) < 40 \text{ GeV}/c^2$	0.14 (24.3)	8.0×10^{-3} (20.0)	0.12 (25.2)
$\geq 1 \text{ jet}, E_T > 20 \text{ GeV}$	0.14 (98.6)	6.9×10^{-3} (86.3)	5.5×10^{-2} (46.2)
b tagging	9.4×10^{-2} (68.6)	4.1×10^{-3} (59.4)	1.6×10^{-3} (2.9)
jet veto	5.1×10^{-3} (5.4)	2.38×10^{-3} (58.1)	6.6×10^{-4} (41.9)
$\Delta\varphi(\tau_1, \tau_2) < 175^\circ$	4.9×10^{-3} (96.4)	2.33×10^{-3} (98.0)	5.6×10^{-4} (83.9)
$E_{\nu_1, \nu_2} > 0$	2.0×10^{-3} (40.9)	9.60×10^{-4} (41.2)	2.1×10^{-4} (38.5)
N_{ev} at 30 fb^{-1}	60.3	28.8	6.4

Table 5.22: Number of signal plus background events and the number of background events in the selected mass windows for 30 fb^{-1} and the signal significance without ($S_{\text{no syst.}}$) and with ($S_{\text{syst.}}$) the background systematics taken into account.

	$\Delta m_{\tau^+\tau^-}$	$N_S + N_B$	N_B	$S_{\text{no syst.}}$	$S_{\text{syst.}}$
$m_A = 130 \text{ GeV}/c^2, \tan\beta = 20$	120 - 200 GeV/c^2	176	83	8.9	6.4
$m_A = 140 \text{ GeV}/c^2, \tan\beta = 15$	130 - 220 GeV/c^2	136	76	9.1	6.7
$m_A = 200 \text{ GeV}/c^2, \tan\beta = 20$	140 - 280 GeV/c^2	175	83	8.8	6.3
$m_A = 300 \text{ GeV}/c^2, \tan\beta = 20$	240 - 480 GeV/c^2	78	39	5.4	4.3
$m_A = 500 \text{ GeV}/c^2, \tan\beta = 50$	360 - 780 GeV/c^2	57	22	6.2	5.3



Benchmark Channels: $t\bar{t}H, H \rightarrow b\bar{b}$

5.3.1 Introduction

The Higgs boson decay to $b\bar{b}$ is the dominant mode for the Higgs mass range up to $m_H \sim 135 \text{ GeV}/c^2$. Direct Higgs production is almost impossible to detect via this decay as a result of the combination of an overwhelming QCD cross section for $b\bar{b}$ production and the inability to reconstruct the Higgs mass very precisely. While the latter is still true in the case of Higgs production in association with a $t\bar{t}$ or $b\bar{b}$ pair, these channels hold promise because they entail substantially lower backgrounds. The separation of these events into 3 salient topologies follows as a result of the ways in which the two W bosons in the event decay. Thus, in addition to the four b jets, roughly 49% of these events also contain four hadronic jets (the all-hadron channel), while some 28% have two hadronic jets together with an isolated electron or muon and missing E_t (the semi-leptonic channel), with a further 5% of events containing two oppositely-charged leptons (either of which can be an electron or muon) and missing E_t (the di-lepton channel). The remaining 14% of events correspond to those cases where one or both of the W bosons decay to a tau lepton and neutrino and are not easily




Figure 5.9: Reconstructed Higgs boson mass for the signal of $M_A = 200 \text{ GeV}/c^2$ and $\tan\beta = 20$ and for the total background for an integrated luminosity of 30 fb^{-1} . The dashed line shows the sum of the $Z/\gamma^* \rightarrow e^+e^-$ and $b\bar{b}Z/\gamma^*e^+e^-$ backgrounds.




Figure 5.10: Reconstructed Higgs boson mass for the signal with $M_A = 300 \text{ GeV}/c^2$ and $\tan\beta = 25$ and for the total background for an integrated luminosity of 30 fb^{-1} . The dashed line shows the sum of the $Z/\gamma^* \rightarrow e^+e^-$ and $b\bar{b}Z/\gamma^*e^+e^-$ backgrounds.

distinguishable as such, as a result of the rich decay repertoire of the tau meson. In fact, these events do make a small contribution to the three other classes of events in the actual analyses. Additional hadronic jets can appear in these events and originate from initial and final state QCD radiation (IFSR).

A detailed description of the $t\bar{t}H$ analysis strategies and the results can be found in Reference [151]. All the results presented here are for an integrated luminosity of 60 fb^{-1} .

5.3.2 Event Generation and Simulation

As the identification of the signal relies upon the presence of top quark decay products, it comes as no surprise that the most significant backgrounds are those associated with $t\bar{t}$ events themselves. The main backgrounds are: $t\bar{t}jj$, $t\bar{t}b\bar{b}$ and $t\bar{t}Z$ with $Z \rightarrow b\bar{b}$.

These processes are studied in detail and are presented here. Secondary background sources include pure QCD multijet events in the case of the all-hadron channel, and W/Z plus jets or diboson plus jets events in the case of the semi-leptonic and di-lepton channels. With the exception of QCD multijets, the latter have substantially lower production cross-sections than $t\bar{t}$ events but very similar topologies. They are therefore not studied in detail.

Details about the primary Monte Carlo data samples used in this analysis are available in Reference [151]. The semi-leptonic and all-hadron $t\bar{t}H$ signal samples were generated using CompHEP (version 41.10) and PYTHIA (version 6.215), while the di-lepton samples used PYTHIA only. Though a leading order Monte Carlo, PYTHIA is known to do a very good job of reproducing IFSR as well as parton shower effects. This is adequate for the signal samples. For the $t\bar{t}$ plus jets backgrounds, greater care must be exercised. In particular,




Figure 5.11: The 5σ discovery region in the M_A - $\tan(\beta)$ plane with 30 fb^{-1} of the integrated luminosity for the m_h^{\max} scenario. The lower (upper) curves was evaluated without (with) the effect of background systematics taken into account.

PYTHIA alone cannot be expected to do a realistic job since the relevant processes are not leading order. On the other hand, there is not currently a full next-to-leading order (NLO) MC for $t\bar{t}$ plus jets production. As a result, higher order matrix elements are used including additional radiated partons in conjunction with the parton showering of PYTHIA to produce the appropriate event topologies.

ALPGEN and PYTHIA are used for the matrix elements and parton showering, respectively, for the $t\bar{t}$ plus n jets background samples. The matching of the two generators is done in ALPGEN as discussed in Ref. [152]. In particular, all of the matrix elements for $t\bar{t}$ plus n additional hard partons are included and properly combined at each order taking into account the interference between amplitudes.

QCD events were generated with PYTHIA (version 6.215) in the \hat{p}_t ranges from 120 to 170 GeV/c and greater than 170 GeV/c .

For the simulation of the interaction with the detector, the CMS tools, providing GEANT3 and GEANT4 based simulation of the CMS detector have been used.

The NLO signal cross-sections for different Higgs mass hypotheses are given in Table 5.23 together with the branching ratios for $H \rightarrow b\bar{b}$ [153].

The leading order CompHEP cross-sections for the different background processes together with the effective cross-sections after the application of the generator filters are listed in Table 5.24. The ALPGEN cross sections for the different jet multiplicity processes are listed in Table 5.25. A detailed comparison of ALPGEN versus CompHEP for the $t\bar{t}jj$ background is

Table 5.23: NLO signal cross-sections and $H \rightarrow b\bar{b}$ branching ratios for different Higgs mass hypotheses

m_H	115 GeV/ c^2	120 GeV/ c^2	130 GeV/ c^2
σ_{NLO} (pb)	0.747	0.664	0.532
$BR(H \rightarrow b\bar{b})$	0.731	0.677	0.525

Table 5.24: LO CompHEP cross-sections and effective cross-sections after the generator filters of the considered background processes.

	QCD $\hat{p}_t=120-170$ GeV/ c	QCD $\hat{p}_t > 170$ GeV/ c	$t\bar{t}b\bar{b}$	$t\bar{t}Z$
σ_{LO} (pb)	$3.82 \cdot 10^5$	$1.05 \cdot 10^5$	3.28	0.65
$\sigma_{LO} \times \epsilon$ (pb)	76.4	336.0	2.82	0.565

available in [151]. All the results that are presented here for the $t\bar{t}N_j$ backgrounds are based on the ALPGEN samples, where available.

5.3.3 L1 and High Level Trigger Selections

A dedicated $t\bar{t}H$ trigger was not available and therefore was not implemented in the analysis. As a result, it is assumed in what follows that the signal is recorded by the CMS Level 1 (L1) and High Level Triggers (HLT) as described in [68]. Wherever possible, the cleaner signature of at least one isolated lepton in the final state is exploited. The semi-leptonic channels thus use the single muon (stream #43) or single electron (stream #2) triggers.

A logical “OR” of the single muon, single electron and single tau streams is used for the di-lepton channel. The same trigger setups as for streams #43 and #2 were used, except that the p_T threshold was lowered to 15 GeV/ c to permit selection of 20 GeV/ c leptons later in the analysis. The tau trigger is the official stream (bit #91). Jet triggers are used to select all-hadron events. In particular, the single-jet, 3-jet and 4-jet triggers with low luminosity thresholds [68][154] are combined (stream #120 or #122 or #123).

Efficiencies for the various HLT and L1 triggers that were used are presented in Table 5.26. The efficiencies quoted are determined by counting the numbers of accepted events relative to the total numbers of events in each sample. In order to streamline the various studies that were performed, the analyses used different MC samples, produced with different final state constraints. Thus, efficiencies for single muon, single electron and fully hadronic final states were defined with respect to exclusive signal samples and inclusive background samples, as described in the preceding section. The di-lepton channel efficiency on the other hand, was defined with respect to samples containing at least one leptonic top decay for the signal and inclusive samples for the backgrounds.

Table 5.25: LO ALPGEN cross-sections for the different jet multiplicity samples.

	exclusive $t\bar{t}+1j$	exclusive $t\bar{t}+2j$	exclusive $t\bar{t}+3j$	inclusive $t\bar{t}+4j$
σ_{LO} (pb)	170	100	40	61

Table 5.26: Signal and background efficiencies of the Level 1 and High Level Triggers.

	Single μ	Single e	Single e OR μ OR τ	Jets
$H \rightarrow b\bar{b}$ (%) with $m_H = 120\text{GeV}/c^2$	63.5	52.4	76.7	24.93
$t\bar{t}b\bar{b}$ (%)	19.0	16.1	83.6	18.31
$t\bar{t}1j$ (%)	13.9	11.3	53.0	2.89
$t\bar{t}2j$ (%)	14.0	11.1	59.8	6.24
$t\bar{t}3j$ (%)	14.0	11.1	68.5	11.39
$t\bar{t}4j$ (%)	13.4	11.1	78.6	31.35
$t\bar{t}Z$ (%)	20.4	18.8	84.4	25.31
QCD 120-170 GeV/ c (%)	0.08	0.8	4.3	1.67
QCD >170 GeV/ c (%)	0.07	2.1	4.4	10.28



5.3.4 Reconstruction

5.3.4.1 Muon Reconstruction

The process of muon reconstruction begins in the Muon Chambers and is then extended to the tracking system, as described in Ref. [155]. For the studies presented here it is important to identify muons coming from W decays. To this end, additional selection criteria are applied to distinguish these muons, which will be referred to as *signal* muons, from the muons coming from other sources such as b decays. The latter will be referred to as *background* muons, even though they arise in signal events as well as background events. The desired discrimination between *signal* and *background* muons is achieved by constructing a discriminator that is based upon probability density functions (PDF) for the following observables associated with muon candidates:

- Transverse momentum, p_t
- Track isolation, $IsoTk$
- Calorimeter isolation, $IsoCalo$
- Significance of track impact parameter, $S_{ip} = d/\sigma_d$

The PDF's associated with these variables for *signal* and *background* muons are obtained by matching to generator-level muons.

The PDF's are combined into the following likelihood ratio:

$$L = \prod_i \frac{P_i^{sig}(x_i)}{P_i^{sig}(x_i) + P_i^{bkg}(x_i)} \quad (5.13)$$

where P_i^{sig} and P_i^{bkg} are the PDF's of an observable x_i for *signal* and *background* muons, respectively.




Figure 5.12: On the left: Performance of the muon likelihood discriminator for the semi-leptonic muon $t\bar{t}H$ channel. On the right: *Signal* versus *background* electron efficiencies for likelihood values ranging from 0.006 (the upper point) with a step size of 0.006, (i.e. approximately in the range $1.0 < -\text{Log}(L_e) < 2.0$).

The performance for *signal* and *background* muon discrimination are shown in Figure 5.12. For a *signal* muon efficiency of 90%, only 1% of background muons are selected. The PDF's are constructed using a sample of $t\bar{t}H$ events with $m_H = 120 \text{ GeV}/c^2$ in which one and only one of the W bosons decays to a muon and neutrino, while the other one decays hadronically.

If the likelihood selection is used after the HLT, a dramatic improvement in QCD ($\hat{p}_t > 170 \text{ GeV}/c$) rejection is possible with little or no loss in signal efficiency. For example, a small drop in signal efficiency from 60% to 60% reduces the QCD efficiency by more than a factor of 3 (i.e. from 0.07% to 0.02%).

5.3.4.2 Electron Reconstruction

A full description of the electron reconstruction in CMS can be found in Ref. [37]. Electrons coming from W boson decays are typically characterized by isolated high transverse energy clusters. These electrons are thus efficiently identified by means of an isolation requirement applied to the electron candidate with respect to other reconstructed tracks in the event.

In analogy to the muon reconstruction and equation 5.13, a likelihood method is used to identify the signal electrons, making use of the following observables:

- the p_t sum of tracks inside an *isolation cone* of radius $\Delta R = 0.3$ around the candidate electron direction
- the ΔR distance between the electron candidate and the closest track
- the transverse momentum of the electron candidate, p_t
- the ratio between the cluster energy and the track momentum, E/p
- the ratio between the hadronic and electromagnetic energies of the cluster, H/E

An appropriate choice of likelihood cut value has been studied by comparing *signal* versus *background* electron efficiencies as shown in Figure 5.12.

For a $-\text{Log}(L_e)$ cut value of 1.27, *signal* electrons are selected with an efficiency of 84% and *background* electrons with an efficiency of 1.5%. This value was chosen for the analyses described in subsequent sections.

Concerning the efficiency of the likelihood cut with respect to background rejection in $t\bar{t}jj$

events in which there were no isolated electrons coming from W decays, only 6% of these events were accepted for a likelihood cut of 1.27.

As in the case of the muon selection, the likelihood approach can be used to augment the HLT selection efficiency. Maintaining a roughly constant signal efficiency, the likelihood cut in combination with the HLT trigger yields an order of magnitude reduction in the QCD background selection efficiency.

5.3.4.3 Jet and Missing E_t Reconstruction

Jets are reconstructed using the iterative cone algorithm. A cone with $\Delta R = 0.5$ is used when at least one W boson decays into leptons, while a smaller cone size was found to be more suitable for the more dense jet environments associated with the all-hadron channel (see below).

A calorimetric-tower energy threshold of 0.8 GeV and a transverse-energy threshold of 0.5 GeV are used. Calorimeter towers that exceed 1 GeV are considered as jet seeds. For the leptonic channels, the jet energy is calibrated using MC calibrations [156] provided by the JetMET group for the corresponding set of reconstruction parameters.

The single lepton analyses, as described in more detail below, make use of an event likelihood to help select and properly reconstruct events and decay chains. This is facilitated, in part, by making use of the various invariant mass constraints associated with the top quark decays. The corresponding likelihoods thus rely upon the resolutions that are obtained for the invariant masses of the hadronically decaying W boson and the two top quarks. The “best-case” invariant mass distribution for the hadronically decaying top quark is reconstructed by matching to generator-level parton information and shown in Figure 5.13. The distributions for the leptonically decaying top quark and the hadronically decaying W boson (Ref. [151]) have similar shapes but different RMS (25.7 GeV/ c^2 and 15.7 GeV/ c^2 , respectively) since the longitudinal momentum of the leptonically decaying top quark has to be calculated from missing E_t . A reconstructed jet is considered as matched to the corresponding parton if their separation, ΔR_{j-p} , is less than 0.3.

The missing transverse energy of the event E_t^{miss} is computed as

$$E_t^{miss} = \sum_i E_t^{tower} - \left(\sum_j E_t^{RawJet} - \sum_k E_t^{CaliJet} \right) + \sum_m E_t^{Muon} \quad (5.14)$$

where the sum with index i runs over calorimeter towers, that with index j runs over raw jets, k runs over calibrated jets, and m runs over the reconstructed muons of the event. Equation 5.14 thus takes into account the corrections due to jet calibration and the contributions of muons that are not measured in the calorimeter.

The choice of the jet reconstruction algorithm is an important step in the event selection optimization for the all-hadron $t\bar{t}H$ channel, where at least 8 jets are expected in the final state. For this reason, an optimization is obtained by means of a simple “proto” analysis as described in Reference [151].

A dedicated $t\bar{t}H$ calibration [157] is applied to help recover the original transverse energy of the associated parton. Reconstructed jets with a b-tagging discriminator value higher than 0.4 are calibrated using a separate b-jet calibration procedure.

Figure 5.13 shows the significance with respect to the S/N ratio for a range of b-tag discriminator values for each of the several cone sizes indicated. Lower discriminator values yield

higher significance but only at the cost of low S/N while, on the contrary, higher discriminator values give lower significance but higher S/N . A good compromise is in the middle range of each of the curves where neither S/N nor significance are unreasonably low. With this in mind, the best choice for the jet cone is seen to be $\Delta R = 0.40$.




Figure 5.13: Left: Invariant mass of the hadronically decaying Top quark using jet-parton matching with $\Delta R_{j-p} < 0.3$. Right: Change in significance and S/N resulting from variations in the b-tagging discriminator for the various cone sizes indicated in the legend.

5.3.4.4 b-Tagging

The identification of jets from b-quarks is done with the *Combined Secondary Vertex* algorithm. This algorithm exploits secondary vertex and track properties to calculate a discriminator value which separates b-jets from non b-jets. A detailed description is published in Ref. [148] which also presents results of detailed studies of the performance of the b-tagging algorithm as applied to Monte Carlo $t\bar{t}$ and QCD samples.

In the $t\bar{t}H$ analyses, a fixed cut value for the b-tagging discriminator is applied, and four jets are required to pass this cut in the semi-leptonic and all-hadron channels, while only 3 jets are required to be tagged in the di-lepton analysis. The misidentification rate of charm and light flavour jets as a function of the b-tagging efficiency is shown in Fig. 5.14 for the $t\bar{t}H$ and the $t\bar{t}jj$ samples, respectively. It can be seen that the efficiencies are similar in these samples.

This fixed-cut b-tagging approach gives reasonable results, but is not necessarily optimal. Some potential improvements are possible such as the combination with a soft lepton tag or a discriminator cut which depends on p_t and η of the jets. Studies have shown that they have the potential to improve the results at the order of some percent. These improvements were not used in the current analyses.

5.3.5 Event Selection

In this section the event selection for the different channels under consideration is described. In order to be able to combine the results from all the $t\bar{t}H$ search channels, the different channels use mutually exclusive event samples. This is most easily facilitated by coordinating how high p_t electrons and muons from the W decays (previously referred to as *signal leptons*) are either selected or vetoed by the different analyses.

For the analyses reported here, the different data samples used were separated using selection and/or veto criteria based on the lepton likelihood value, as described in Ref. [151].




Figure 5.14: On the left: Non-b jet mistagging efficiency versus b-jet tagging efficiency for c-jets (triangles), and uds-jets (stars) for the $t\bar{t}H$ sample with $m_H = 120 \text{ GeV}/c^2$ and jets with a minimum transverse momentum of $20 \text{ GeV}/c$. For this plot the “physics definition” of the original jet flavour has been used. In this definition there are no original gluon jets in the $t\bar{t}H$ sample. On the right: The corresponding plot for the $t\bar{t}jj$ sample, where gluon jets are represented by crosses.

5.3.5.1 Semileptonic Channel: $t\bar{t}H \rightarrow b\bar{b}b\bar{b}q\bar{q}'\mu\nu_\mu$ and $b\bar{b}b\bar{b}q\bar{q}'e\nu_e$

The strategy for selecting $t\bar{t}H$ events with one isolated muon or electron in the final state can be summarized in the following three steps: preselection, choice of jet pairing and finally, selection. The preselection requires the HLT stream for a single muon or a single electron, one isolated lepton using the likelihood information as described in section 5.3.4.1 and 5.3.4.2, and 6 or 7 jets in the pseudorapidity region $|\eta| < 3.0$ with a calibrated transverse energy larger than 20 GeV . In order to recover some efficiency, jets with $10 \text{ GeV} < E_t < 20 \text{ GeV}$ are also accepted if they have at least two associated tracks pointing to the signal primary vertex¹ within a distance along the beam (z) axis of $(|z_{PV} - z_{track}| < 1 \text{ mm})$. The latter condition is required to reject low transverse energy fake jets, (i.e. jets that are not associated with any of the signature partons in the signal event). For the single electron channel, the misidentification of the jet with the isolated electron has been excluded by imposing a veto on the jet if the electron lies inside a jet cone radius of 0.1 .

At least 4 jets are required to be tagged as b-jets with a minimal discriminator value corresponding to a b-efficiency of about 70%.

To decrease the contamination from the di-lepton channel, a double muon, double electron and muon-electron veto is applied, in which events with the second lowest $-\text{Log}(L_\mu) < 1.4$ and events with $-\text{Log}(L_e) < 1.2$ are rejected from the analysis. In the case of the semileptonic electron channel the previous cuts are applied respectively to the first muon likelihood candidate and to the second electron likelihood candidate. The application of these vetoes results in a lowering of the signal efficiency by about 2%, while the total background rejection is increased by 13%.

In order to perform a complete reconstruction of the event, the longitudinal momentum of the neutrino has to be computed from four-momentum conservation for the W boson:

¹The signal interaction is generally the one which allows the event to be triggered.




Figure 5.15: $t\bar{t}H$ ($W \rightarrow qq', W \rightarrow \mu\nu$): Signal Significance (left) and Signal to Background ratio (right) as function of the cut on L_{bSele} .

$m_W^2 = (E^\mu + E^\nu)^2 - (\vec{p}^\mu + \vec{p}^\nu)^2$. This equation gives 2 real solutions for p_z^ν in 66% of the cases, while in the remaining 34%, the neutrino is assumed to be collinear with the lepton: $p_z^\nu = p_z^l$. This leads to a small degradation in the longitudinal momentum resolution, but the reconstruction efficiency of the leptonic W boson decay is increased to 100%.

In order to choose the jet combination that does the best job of reconstructing the two top quarks, a likelihood, L_{Event} , is defined using masses, b-tagging and kinematic information from the whole event:

$$L_{Event} = L_{Mass} \times L_{bTag} \times L_{Kine}. \quad (5.15)$$

The mass information considered in the likelihood L_{Mass} is the probability returned by the kinematic fit with invariant mass constraints (top quarks and hadronic W) that is described in Reference [158].

The b-tagging function L_{bTag} is defined as the product of the b-tag discriminators: $L_{bTag} = D_{TopHad} \times D_{TopLep} \times D_{H1} \times D_{H1} \times (1 - D_{W1}) \times (1 - D_{W2})$; where $TopHad$ and $TopLep$ are expected to be the two b jets from the hadronic and leptonic top, respectively, while H_1 and H_2 are expected to be the two b jets coming from Higgs and W_1 and W_2 are the two jets from the hadronically-decaying W boson.

The kinematic function takes into account the observation that the b-jets coming from top quarks tend to be slightly more energetic than b-jets coming from the Higgs boson (see [151] for a definition).

Among all possible combinations of jet-parton assignments, the one with the highest value of L_{Event} is chosen for use in the final reconstruction of the top quarks and the two remaining jets with highest b-tagging discriminator values are used to reconstruct the Higgs mass.

After the jet assignment is complete, additional criteria are applied to improve background rejection. In particular, a stronger b-tag requirement is applied on the event variable $L_{bSele} = D_{TopHad} \times D_{TopLep} \times D_{H1} \times D_{H2}$.

The signal significance as a function of the selection cut L_{bSele} is shown in Figure 5.15.

The distributions of reconstructed Higgs mass for the final selected events are shown in Figure 5.16 for signal only (left) and for the combination of the different backgrounds (right) for the muon channel only (similar results for the electron channel can be found in [151]). The fraction of signal events where the two b-jets are correctly assigned to the Higgs boson (i.e.




Figure 5.16: $t\bar{t}H$ ($W \rightarrow qq'$, $W \rightarrow \mu\nu$). Left: Invariant $b\bar{b}$ mass for signal only (combinatorial background is shaded grey). Right: The sum of the reconstructed $m_{b\bar{b}}$ spectra for backgrounds with a value of $L_{bSele} > 0.55$. The distributions are normalized to an integrated luminosity of 60fb^{-1} .

the pairing efficiency) is roughly 31% in the muon channel and about 29% for the electron channel.

5.3.5.2 Results

The Selection efficiencies with the corresponding numbers of expected events and signal significances are reported in Table 5.27 for the channels with a muon or an electron in the final state. The number of expected events is computed for an integrated luminosity of 60fb^{-1} in the Standard Model Higgs mass range from 115 to 130 GeV/c^2 .

5.3.5.3 Di-lepton Channel: $t\bar{t}H \rightarrow b\bar{b}b\bar{b}l'l\nu\nu$

Di-lepton $t\bar{t}H$ events are selected by requiring two reconstructed leptons (e, μ) accompanied by significant missing transverse energy and at least four but no more than seven jets, at least three of which have been b-tagged according to the *Combined Secondary Vertex* b-tagging algorithm.

Lepton identification is performed using the electron and muon likelihoods described in Section 5.3.4. In the semi-leptonic analyses, events with more than one identified lepton are vetoed, but in the di-lepton analysis those events are retained. The likelihood acceptance cuts used for leptons in the di-lepton channel are therefore chosen to be the same as the second-lepton veto cuts for both semi-leptonic channels. In this way, the sample of events for the di-lepton $t\bar{t}H$ analysis is by construction strictly complementary to those used in the semi-leptonic channels.

The details of the di-lepton $t\bar{t}H$ selection are summarized below:

- 2 oppositely-charged leptons (e, μ) passing identification criteria ($-\text{Log}(L_\mu) < 1.4$ for muons, $-\text{Log}(L_e) < 1.2$ for electrons)
- corrected $E_T^{\text{miss}} > 40$ GeV
- 4 to 7 jets with calibrated $E_T > 20$ GeV and $|\eta| < 2.5$
- ≥ 3 selected jets b-tagged with discriminator $D > 0.7$

Table 5.27: Selection efficiency for $L_{bSele} > 0.55$ (ϵ_{loose}) and for $L_{bSele} > 0.75$ (ϵ_{tight}), number of expected events and signal significance in 60 fb^{-1} for the muon and electron $t\bar{t}H$ channel. The numbers refer to the complete Higgs mass range.

muon channel						electron channel					
	Analyzed Ev.	ϵ_{loose} (%)	N_{loose}^{ev} 60fb^{-1}	ϵ_{tight} (%)	N_{tight}^{ev} 60fb^{-1}		Analyzed Ev.	ϵ_{loose} (%)	N_{loose}^{ev} 60fb^{-1}	ϵ_{tight}	N_{tight}^{ev} 60fb^{-1}
$t\bar{t}H$ (115)	27768	2.00 ± 0.08	96 ± 4	0.80 ± 0.05	38 ± 3	Total Background			1840		< 352
$t\bar{t}H$ (120)	41929	1.90 ± 0.07	75 ± 3	0.74 ± 0.04	29 ± 2	S/\sqrt{B} (115)	27692	1.39 ± 0.07	66 ± 3	0.52 ± 0.04	25 ± 2
$t\bar{t}H$ (130)	19466	2.23 ± 0.11	55 ± 3	0.84 ± 0.07	21 ± 2	S/B (115)	42228	1.42 ± 0.06	56 ± 2	0.53 ± 0.04	21 ± 1
$t\bar{t}b\bar{b}$	372737	0.247 ± 0.008	419 ± 14	0.0877 ± 0.0048	148 ± 8	S/\sqrt{B} (120)	19127	1.57 ± 0.09	39 ± 2	0.61 ± 0.06	15 ± 1
$t\bar{t}l_j$	393000	0.0051 ± 0.0011	520 ± 120	0.00076 ± 0.00044	78 ± 45	S/B (120)	372737	0.176 ± 0.007	297 ± 12	0.0641 ± 0.0041	109 ± 7
$t\bar{t}2j$	568999	0.0105 ± 0.0014	633 ± 82	0.00070 ± 0.00035	42 ± 21	S/\sqrt{B} (130)	393000	0.0038 ± 0.0010	390 ± 100	0.00025 ± 0.00025	26 ± 26
$t\bar{t}3j$	101000	0.0050 ± 0.0022	119 ± 53	0.	$< 27(68\%C.L.)$	S/B (130)	568999	0.0067 ± 0.0011	401 ± 65	0.00123 ± 0.00046	74 ± 28
$t\bar{t}4j$	86697	0.0035 ± 0.0020	126 ± 73	0.	$< 48(68\%C.L.)$	Total Background	101000	0.0040 ± 0.0020	95 ± 48	0.	$< 27(68\%C.L.)$
$Zt\bar{t}$	50000	0.068 ± 0.012	23 ± 4	0.026 ± 0.007	9 ± 2	S/\sqrt{B} (115)	86697	0.0023 ± 0.0016	84 ± 60	0.	$< 48(68\%C.L.)$
						S/B (115)	50000	0.064 ± 0.011	22 ± 4	0.022 ± 0.007	7 ± 2
						S/B (120)			1289		< 291
						S/\sqrt{B} (120)			1.8		1.5
						S/B (120)			5.1%		8.6%
						S/B (130)			1.6		1.2
									4.4%		7.2%
									1.1		0.9
									3.0%		5.2%

The above is termed the “loose” working point because there is evidence that it is possible to increase the purity (S/B) of the selection, by way of more stringent criteria:

- 4 to 6 jets with calibrated $E_T > 20 \text{ GeV}$ and $|\eta| < 2.5$
- ≥ 4 selected jets b-tagged with discriminator $D > 0.7$

The generated W^- was forced to decay leptonically (e, μ, τ), but the W^+ was allowed to decay freely. This “non-exclusive” dataset incurs a branching ratio of $1/3$, which has been factored into the selection efficiencies reported in Table 5.28. This choice allows us to obtain

a good estimate of the overlap of the contribution to the di-lepton sample arising from semi-leptonic top decays which are mis-reconstructed as di-lepton events; the same applies to tau decays which are mis-reconstructed as e, μ .

The background events have small efficiency to pass the selection criteria, so very large samples must be analyzed. To make these samples more manageable, a loose pre-selection requiring at least 3 b-tags with discriminator $D > 0.7$ is applied before analysis.

5.3.5.4 Results

The selection efficiencies for the two working points, with the corresponding number of expected events and the signal significance, are reported in Tables 5.28. The number of expected events is computed for an integrated luminosity of 60 fb^{-1} .

Table 5.28: Selection efficiency ϵ_{loose} (including branching fraction where applicable) and resulting number of expected events N_{loose}^{ev} in 60 fb^{-1} , for the di-lepton $t\bar{t}H$ channel. For a glimpse of possible improvements, the same for a tighter set of cuts is provided ($\epsilon_{tight}, N_{tight}^{ev}$). Also quoted are binomial errors arising from the finite sizes of processed datasets. The $t\bar{t}H$ datasets are labeled by the generated Higgs mass in GeV/c^2 (parentheses).

	#analyzed	$\epsilon_{loose}(\%)$	N_{loose}^{ev}	$\epsilon_{tight}(\%)$	N_{tight}^{ev}
$t\bar{t}H$ (115)	27900	0.511 ± 0.025	168 ± 8	0.0884 ± 0.01	29 ± 3
$t\bar{t}H$ (120)	26141	0.49 ± 0.025	132 ± 7	0.0701 ± 0.0094	19 ± 3
$t\bar{t}H$ (130)	25911	0.49 ± 0.025	82 ± 4	0.072 ± 0.0096	12 ± 2
$t\bar{t}b\bar{b}$	313894	0.637 ± 0.014	1080 ± 24	0.0939 ± 0.007	159 ± 12
$t\bar{t}1j$	280385	0.0125 ± 0.0021	1270 ± 220	0	< 42 (68% C.L.)
$t\bar{t}2j$	276917	0.0448 ± 0.004	2690 ± 240	0.00144 ± 0.00072	87 ± 43
$t\bar{t}3j$	90367	0.0553 ± 0.0078	1330 ± 190	0	< 31 (68% C.L.)
$t\bar{t}4j$	12281	0.0716 ± 0.0077	2620 ± 280	0.0025 ± 0.0014	92 ± 53
$t\bar{t}Z$	110156	0.304 ± 0.017	103 ± 6	0.0363 ± 0.0057	12 ± 2
all backgrounds			9090		< 422
S/\sqrt{B} (115)			1.8		1.4
S/B (115)			1.8 (%)		6.9 (%)
S/\sqrt{B} (120)			1.4		0.9
S/B (120)			1.5 (%)		4.5 (%)
S/\sqrt{B} (130)			0.9		0.6
S/B (130)			0.9 (%)		2.9 (%)

Since the event selection is quite simple for the di-lepton channel, it is possible to formulate simple equations predicting the selection efficiencies. This is detailed in Ref. [151], where some back-of-the-envelope calculations to estimate these efficiencies for both signal and backgrounds are presented including some of the backgrounds that were not taken into account in this analysis.

5.3.5.5 All-hadron Channel: $t\bar{t}H \rightarrow b\bar{b}b\bar{b}q'q''q'''$

A number of kinematic variables, together with the b-tagging discriminator, have been studied to optimize the signal selection with respect to background rejection. Moreover, in order to combine the results from the 4 different decay sub-channels, a veto on leptons has been

applied using the complementary cut developed within the semi and fully leptonic decays analyses: events are discarded if $-\text{Log}(L_\mu) < 1.4$ or $-\text{Log}(L_e) < 1.2$.

The final set of variables that are used in this analysis is the following:

- Jet Transverse Energy of the 8 most energetic jets in the tracker acceptance
- Combined b-Tag discriminator variable for each jet
- Centrality of the event defined as $\sum_{i=0}^8 E_T^i / E^i$
- Centrality of the Higgs defined similarly, with the sum restricted to the 2 jets paired to the Higgs

The jet-to-parton matching is performed using a χ^2 method as defined in [151].

Two working points have been chosen: the first uses loose cuts on the b-tagging discriminator to get higher statistical significance (but lower S/B), while the second uses a tighter cut on the b-tagging discriminator to obtain a higher S/B (but lower significance). For the first working point an event is selected if the following conditions are satisfied:

- $E_T^{7th} > 30$ GeV and $E_T^{8th} > 20$ GeV for the E_T ordered jets
- the χ^2 for each of the 2 W bosons and 2 t quarks are within 3 sigma of their expected values
- the 3 highest combined b-tagging discriminators for the 4 jets associated to the b -partons must satisfy $D_3 > 0.80$
- Higgs centrality higher than 0.55 and no cut on Event Centrality


For the tight working point, the b-tagging discriminator for the third highest jet is required to satisfy $D_3 > 0.85$ and the fourth one $D_4 > 0.70$, while the event and Higgs centrality are required to exceed 0.55 and 0.80, respectively.

All the applied cuts have been optimized to obtain the highest significance while keeping the S/B ratio as high as possible. All values chosen for $E_T^{7th}, E_T^{8th}, D_3, D_4$, Event and Higgs centrality have been varied simultaneously, thereby mapping out the complete set of combinations within the following limits:

- $20 \text{ GeV} < E_T^{8th} < 40 \text{ GeV}$
- $E_T^{8th} < E_T^{7th} < E_T^{8th} + 40 \text{ GeV}$
- $0.5 < D_3$ and $D_4 < 0.95$
- Event and Higgs Centrality in the range [0.50-0.95]

Variation of more than one cut has also been tested and the final implemented set of cut values is that for which significance and S/B are optimal.

5.3.5.6 Results

The number of analyzed events, selection efficiencies with the corresponding number of expected events and the signal significance are reported in Tables 5.29 for the all-hadron  channel. Both working points are considered.

5.3.6 Discussion of Systematic Uncertainties

Table 5.29: Analyzed events, selection efficiency, number of expected events and signal significance in 60 fb^{-1} for the all-hadron $t\bar{t}H$ channel for 2 different working points: ϵ_{loose} and ϵ_{tight} . The numbers refer to the full mass range.

	#analyzed	$\epsilon_{loose}(\%)$	$N_{loose}^{ev} 60\text{fb}^{-1}$	$\epsilon_{tight}(\%)$	$N_{tight}^{ev} 60\text{fb}^{-1}$
$t\bar{t}H$ (115)	49636	2.32 ± 0.07	347 ± 10	0.294 ± 0.015	44 ± 4
$t\bar{t}H$ (120)	163494	2.55 ± 0.04	314 ± 5	0.366 ± 0.024	45 ± 2
$t\bar{t}H$ (130)	43254	2.80 ± 0.08	214 ± 6	0.358 ± 0.029	27 ± 2
$t\bar{t}bb$	203135	0.702 ± 0.019	1190 ± 31	0.0645 ± 0.0056	109 ± 9
$t\bar{t}1j$	1031551	0.0084 ± 0.0009	860 ± 92	0.0005 ± 0.0002	49 ± 22
$t\bar{t}2j$	559111	0.0333 ± 0.0024	2000 ± 150	0.0009 ± 0.0004	54 ± 24
$t\bar{t}3j$	68015	0.079 ± 0.011	1910 ± 260	0.0015 ± 0.0015	35 ± 35
$t\bar{t}4j$	97334	0.182 ± 0.014	6660 ± 500	0.0021 ± 0.0015	75 ± 53
$Zt\bar{t}$	80226	0.358 ± 0.021	121 ± 7	0.0312 ± 0.0062	11 ± 2
qcd170	264310	0.0238 ± 0.0030	4810 ± 610	0.0004 ± 0.0004	76 ± 76
qcd120	55128	0.0018 ± 0.0018	83 ± 83	0 ± 0	$<95(68\%C.L.)$
Total Backgr.			17600		< 505
S/\sqrt{B} (115)			2.6		2.0
S/B (115)			2.0%		8.7 %
S/\sqrt{B} (120)			2.4		2.0
S/B (120)			1.8%		8.9 %
S/\sqrt{B} (130)			1.6		1.2
S/B (130)			1.2%		5.4 %

5.3.6.1 Estimation of “Standard” CMS Systematics

The uncertainties in various quantities, given the knowledge of the CMS experiment at the time of writing this note, are considered first. These differ from what they are expected to be after CMS has collected 60 fb^{-1} of data.

In keeping with other CMS analyses, the following “standard” sources of systematic error are considered:

- Jet energy scale (JES) (3% to 10% depending on p_t)
- Jet resolution (10%)
- b-jet and c-jet tagging efficiencies (4%)
- uds-jet tagging efficiencies (10%)
- Luminosity (3%)

It is assumed that the systematics listed above are uncorrelated. Each source is varied independently which produces a change in the selection efficiency $\Delta\epsilon$ and the corresponding change in expected event yields ΔN_X ($X = t\bar{t}H, t\bar{t}1j, \dots$) for the signal and background.

A very detailed breakdown of the various sources of systematic uncertainties and the methods of how they are computed for all the background and signal samples is available in Reference [151]. In Tables 5.30, the systematic uncertainties are propagated to the expected signal significance for “tight” and “loose” working points.


Table 5.30: Significance before and after taking into account the uncertainty dB in the total number of background events due to systematics.

muon	S/B	S/\sqrt{B}	$S/\sqrt{B + dB^2}$	di-lepton	S/B	S/\sqrt{B}	$S/\sqrt{B + dB^2}$
$L_{bS_{ele}} > 0.55 (\epsilon_{loose})$				4-7 jets, 3-4 b-tagged (ϵ_{loose})			
$t\bar{t}H$ (115)	0.052	2.2	0.20	$t\bar{t}H$ (115)	0.018	1.8	0.10
$t\bar{t}H$ (120)	0.041	1.8	0.15	$t\bar{t}H$ (120)	0.015	1.4	0.08
$t\bar{t}H$ (130)	0.030	1.3	0.11	$t\bar{t}H$ (130)	0.009	0.9	0.05
$L_{bS_{ele}} > 0.75 (\epsilon_{tight})$				4-6 jets, 4-6 b-tagged (ϵ_{tight})			
$t\bar{t}H$ (115)	0.108	2.0	0.44	$t\bar{t}H$ (115)	0.069	1.4	0.42
$t\bar{t}H$ (120)	0.082	1.6	0.34	$t\bar{t}H$ (120)	0.045	0.9	0.27
$t\bar{t}H$ (130)	0.060	1.1	0.24	$t\bar{t}H$ (130)	0.029	0.6	0.18
electron	S/B	S/\sqrt{B}	$S/\sqrt{B + dB^2}$	hadron	S/B	S/\sqrt{B}	$S/\sqrt{B + dB^2}$
$L_{bS_{ele}} > 0.55 (\epsilon_{loose})$				Working Point ϵ_{loose}			
$t\bar{t}H$ (115)	0.051	1.8	0.20	$t\bar{t}H$ (115)	0.020	2.6	0.07
$t\bar{t}H$ (120)	0.044	1.6	0.17	$t\bar{t}H$ (120)	0.018	2.4	0.07
$t\bar{t}H$ (130)	0.030	1.1	0.12	$t\bar{t}H$ (130)	0.012	1.6	0.05
$L_{bS_{ele}} > 0.75 (\epsilon_{tight})$				Working Point ϵ_{tight}			
$t\bar{t}H$ (115)	0.086	1.5	0.37	$t\bar{t}H$ (115)	0.087	2.0	0.22
$t\bar{t}H$ (120)	0.072	1.2	0.31	$t\bar{t}H$ (120)	0.089	2.0	0.22
$t\bar{t}H$ (130)	0.052	0.9	0.22	$t\bar{t}H$ (130)	0.054	1.2	0.13

5.3.6.2 Background Rates from Data

There are relatively large theoretical uncertainties in the cross-sections used to normalize the signal yields [153], and even larger theoretical uncertainties in those used for the $t\bar{t}$ +jets backgrounds [159]. These have not been included as part of the systematic errors considered above, because when the CMS experiment reaches maturity, estimating the $t\bar{t}$ +jets background directly from data ought to be possible. In this way, the uncertainty associated with Monte Carlo derived tagging rates are avoided entirely. For example, the number of mis-tagged $t\bar{t}$ +jets which can be factorized as follows:

$$N_{t\bar{t}jj}^{mistag} = N_{t\bar{t}jj}^{no-tag} \times Pr(uds \rightarrow b; E_T, \eta, \dots)$$

where $N_{t\bar{t}jj}^{no-tag}$ is a high purity (e.g. fully reconstructed with a mass window) top sample that has been obtained without requiring b-tagging and $Pr(uds \rightarrow b; E_T, \eta, \dots)$ is a parametrized “fake matrix” that is derived from some independent dataset (e.g dijet data) which yields the probability for a light quark jet to fake a secondary vertex. It may also be possible to derive this fake matrix from the top sample itself. If a high-purity (e.g. double-tagged and fully reconstructed) semi-leptonic top sample were selected, the jets belonging to the hadronic W would provide a source of both light quark and charm jets. From these data, a measurement of the  responding uds-tag and c-tag rates at the relevant energy could be directly obtained.

5.3.7 Combined Significance

Since the event samples for the channels studied in this note are strictly disjoint, the results can be combined by simply adding the individual signal yields (background yields) to obtain a summed $S(B)$.

For each of the considered systematics, the resultant error in background yields are added for all four channels, since they are by definition fully correlated. The summed errors are then added by quadratures to get a combined systematic uncertainty dB . One then calculates the significance, inclusive of systematic uncertainties in the background yield, according to the formula $S/\sqrt{B + dB^2}$.

It is of interest to see how much better the results have the potential to be at tighter working points for the various analyses. Since the systematic uncertainties are not well quantified at these “tight” working points, because of a lack in Monte Carlo Statistics, the same uncertainties as for the “loose” working points are used to reduce spurious statistical effects. This procedure can be justified by the observation that the impact of the b-tagging and uds-mistagging uncertainty is smaller at the “tight” working points and the JES uncertainty becomes dominant. Since the “tight” working points are defined by stronger b-tagging cuts, while keeping the E_T cuts constant, no major change in the relative systematic uncertainty is expected. A more detailed study of the systematic error at the “tight” working points for samples with enough Monte Carlo Statistics is available in Ref. [151].

It is difficult to predict at this time exactly what will be the level to which the backgrounds can be understood, because the tools required are not yet in existence and because this understanding requires real data. In view of this, it is interesting to consider how the combined significance of the measurements presented in this note would vary as a function of the fractional uncertainty in background cross-sections, i.e. as dB_{xsec}/B .




Figure 5.17: Expected range of combined significance (di-lepton + semi-leptonic + all-hadron, and includes the systematic uncertainties estimated in Section 5.3.6.1) versus an additional systematic uncertainty on the background cross-section as a fraction of total background. Left: Results for the “loose” working points. Right: Results for the “tight” working points.

The solid central line in Figure 5.17 shows how the combined significance $S/\sqrt{B + (dB_{sys} + dB_{xsec})^2}$ degrades as a function of dB_{xsec}/B . The signal and background yields for the tightest working points (N_3^{ev} in Table 5.27, N_{tight}^{ev} in Table 5.28, and N_2^{ev} in Table 5.29) are used in the right side of Figure 5.17, because these give the best results after inclusion of systematics.

Other than this “fundamental” cross-section uncertainty, there is also the “correctible” errors in the cross-sections used at the time of writing, which can be compensated for once data has been collected. The upper and lower dashed curves in Figure 5.17 show the maximum and minimum allowed excursions, should the signal and background cross-sections be off by 10% and 20% respectively. Thus the upper (lower) dashed line corresponds to the signal cross-section scaled up (down) by 10% while at the same time the background cross-section

is scaled down (up) by 20%.




Figure 5.18: Expected range of combined significance (di-lepton + semi-leptonic + all-hadron) versus the total systematic uncertainty in background as a fraction of total background. Left: Results for the “loose” working points. Right: Results for the “tight” working points.

It is also of interest to see how much better the analyses could do if the total systematic uncertainty can be reduced (i.e. the region left of zero in Figure 5.17). Hence, Figure 5.18 shows the full range of obtainable significances, with the dot marking the currently estimated value with no cross-section uncertainty ($dB = dB_{sys}$). The star corresponds to what one would obtain for 1% and 4% uncertainties on the $t\bar{t}N_j$ and $t\bar{t}b\bar{b}$ backgrounds, respectively, an arbitrarily chosen reference. It is interesting to note that it does not quite yield a substantial significance, even though background uncertainties of 1% and 4% for $t\bar{t}N_j$ and $t\bar{t}b\bar{b}$ are probably substantially better than what will be accessible in reality. This highlights the challenge that is faced in observing $t\bar{t}H$.

Chapter 6

Physics Studies with Heavy Ions



6.1 Benchmark Channel: $PbPb \rightarrow Q\bar{Q} + X \rightarrow \mu^+\mu^- + X$

The measurement of the charmonium ($J/\psi, \psi'$) and bottomonium ($\Upsilon, \Upsilon', \Upsilon''$) resonances in PbPb collisions at $\sqrt{s_{NN}} = 5.5$ TeV provides crucial information on the many-body dynamics of high-density QCD matter. First, the step-wise suppression of heavy quarkonia production is generally agreed to be one of the most direct probes of Quark-Gluon-Plasma formation. Lattice QCD calculations of the heavy-quark potential indicate that color screening dissolves the ground-state charmonium and bottomonium states, J/ψ and Υ , at $T_{\text{diss}} \approx 2 \cdot T_{\text{crit}}$ and $4 \cdot T_{\text{crit}}$, respectively. While the interest of charmonia production studies in heavy-ion collisions is well established from measurements done at the SPS and at RHIC, the clarification of some important remaining questions requires equivalent studies of the Υ family, only possible at the LHC energies. Second, the production of heavy-quarks proceeds mainly via gluon-gluon fusion processes and, as such, is sensitive to saturation of the gluon density at low- x in the nucleus (“Color Glass Condensate”). Measured departures from the expected “vacuum” (proton-proton) quarkonia cross-sections in PbPb collisions at LHC will thus provide valuable information not only on the thermodynamical state of the produced partonic medium, but also on the initial-state modifications of the nuclear parton (especially, gluon) distribution functions.

This first CMS heavy-ion physics analysis focuses on the measurement of the heavy-quarkonia cross-sections in PbPb collisions at $\sqrt{s_{NN}} = 5.5$ TeV, via their dimuon decay channel. The generation of realistic signals and backgrounds, the dimuon reconstruction algorithm and the trigger, acceptance and efficiency corrections are discussed. The obtained dimuon mass resolutions, the signal over background as well as the expected yields in one-month PbPb running are presented.

6.1.1 Simulation of Physics and Background Processes

The relatively low Υ production rates ($\sim 10^{-4}$ per PbPb event) and the large number of particles to track in heavy-ion collisions make it very expensive computationally to use a full nucleus-nucleus event generator (such as e.g. HIJING [160]) with detailed detector simulation and reconstruction to obtain a statistically significant sample of signal events. Instead, a combination of fast and slow simulations are used in this analysis. The input signal and backgrounds are obtained from realistic distributions: NLO pQCD for heavy-quark production processes, and HIJING for the soft background, constrained by extrapolations from lower energy heavy-ion data. A full detector and trigger simulation plus reconstruction are carried

out for a few 10^7 events with single and pair particles of the different types and the corresponding response functions (acceptances, resolutions, efficiencies, etc) are parametrized in a fast MC, used to obtain the final fully corrected yields. The response functions are cross-checked by comparing the final dimuon spectra obtained with the fast MC against 5×10^5 PbPb HIJING events fully simulated and reconstructed in the detector.

The quarkonium production cross sections in PbPb are obtained from NLO pp calculations at $\sqrt{s} = 5.5$ TeV made in the color evaporation model (CEM) [161], using MRST PDF modified with the EKS98 prescription for nuclear shadowing [162], with renormalization and factorization scales $\mu_R = \mu_F = m_Q$, and scaled by A^2 ($A = 208$ for Pb). The resulting (impact-parameter averaged) inclusive quarkonia production cross sections are: $B_{\mu\mu} \sigma_{Q\bar{Q}} = 49000, 900, 300, 80, 45 \mu\text{b}$ for $J/\psi, \psi', \Upsilon, \Upsilon',$ and Υ'' , respectively. The NLO double-differential $d^2\sigma/dp_T d\phi$ distributions of J/ψ and Υ are also used for the other states within each quarkonium family.

The two main sources of background in the dimuon invariant mass spectrum are:

1. Uncorrelated decays of **charged pions and kaons**, which represent about 90% of the produced charged particles. This source was simulated using input pion and kaon $d^2N/dp_T d\eta$ distributions from HIJING, absolutely normalized to give $dN_{ch}/d\eta|_{\eta=0} = 2500$ (*low*) and 5000 (*high*) multiplicities in central PbPb. Both cases are conservative (“pessimistic”) estimates, since extrapolations from RHIC data indicate that $dN_{ch}/d\eta|_{\eta=0} \approx 2000$ at the LHC.
2. The other source of background muons are **open heavy flavour** (D, B mesons) decaying a few mm away from the interaction vertex. The probability to produce at least one muon at the end of the decay chain of charm (bottom) quarks is $\sim 18\%$ (38%) according to PYTHIA 6.025. The double differential (p_T, η) cross-sections are obtained from pp NLO calculations (with CTEQ5M1 PDF, and $\mu_R = \mu_F = m_Q$), which give $\sigma_{c\bar{c}, b\bar{b}} = 7.5, 0.2 \text{ mb}$ [161], scaled by the nuclear overlap function, $\langle T_{PbPb}(b = 0 \text{ fm}) \rangle = 30.4 \text{ mb}^{-1}$, to obtain the expected yields in central PbPb collisions.

A fast MC simulation equivalent to $5 \cdot 10^7$ PbPb events has been carried out superimposing the decay dimuons from the five quarkonium resonances on top of the background from the combinatorial decays of π, K and open heavy flavour. Each muon track (with a given momentum, pseudorapidity, charge and origin) is weighted by a factor that takes into account the corresponding detector acceptance, as well as trigger and reconstruction efficiency for the two event multiplicities considered (see next section).

6.1.2 Reconstruction and Analysis

6.1.2.1 Dimuon Trigger and Acceptance

The response of the CMS detector to muons (as well as long-lived punchthrough pions and kaons reaching the muon chambers) is parametrized by 2-dimensional p, η acceptance and trigger tables. The particles are fully tracked in CMS using GEANT4 from the vertex to the chambers. Each track is accepted or rejected according to the Level-1,2 heavy-ion dimuon trigger criteria [7] and the corresponding efficiencies, $\epsilon_{trig}^{LV L1}(p, \eta)$ and $\epsilon_{trig}^{LV L2}(p, \eta)$,

are computed. Trigger efficiencies are of the order of $\sim 90\%$ for those μ reaching the muon chambers. The J/ψ and Υ acceptances are shown as a function of p_T in Fig. 6.1, for two η ranges: full detector and central barrel. Because of its relatively low mass, low energy J/ψ 's ($p_T \lesssim 4 \text{ GeV}/c$) cannot be detected since their decay muons don't have enough energy to traverse the calorimeters and they are absorbed due to ionization losses before reaching the muon chambers. For larger p_T values the J/ψ acceptance increases and flattens out at $\sim 15\%$ for $p_T \gtrsim 12 \text{ GeV}/c$. The Υ acceptance starts at $\sim 40\%$ at $p_T = 0 \text{ GeV}/c$ and remains constant at 15% (full CMS) or 5% (barrel) for $p_T > 4 \text{ GeV}/c$. The p_T -integrated acceptance is about $\frac{1}{3}$ for the J/ψ and 21% for the Υ as obtained from our input theoretical distribution.




Figure 6.1: J/ψ (top) and Υ (bottom) acceptances as a function of p_T , in the full detector (barrel and endcap, $|\eta| < 2.4$, full line) and in the barrel alone ($|\eta| < 0.8$, dashed line).

6.1.2.2 Dimuon reconstruction efficiency, purity and mass resolution

The dimuon reconstruction algorithm used in the heavy-ion analysis is a version of the regional track finder based on the muons seeded by the muon stations and on the knowledge of the primary vertex, as described in [163, 164]. It is adapted to deal with the high hit occupancy of the silicon tracker in PbPb collisions. It uses the muon tracks found in the innermost muon stations to identify hits in the outer CMS tracker layer that can form the starting points (seeds) for the matching muon candidate tracks. The propagation in the tracker is performed from the outer layer towards the primary vertex, using two-dimensional parametrizations in the transverse and longitudinal planes. The final fit of trajectories is performed with a Kalman-fitter. The efficiency of a given muon pair is: $\epsilon_{pair}(p, \eta) = \epsilon_{track_1} \times \epsilon_{track_2} \times \epsilon_{vertex}$. The dependence of the Υ reconstruction efficiency on the event multiplicity was obtained from a full GEANT simulation using Υ signal dimuons embedded in HIJING PbPb events. Fig. 6.2 shows the Υ efficiency and purity (where purity is defined as the ratio of true Υ reconstructed over all Υ reconstructed) as a function of charged-particle multiplicity. In the central barrel, the dimuon reconstruction efficiency is above $\sim 80\%$ for all multiplicities, whereas the purity decreases slightly with $dN_{ch}/d\eta$ but stays also above 80% even at multiplicities as high as $dN_{ch}/d\eta|_{\eta=0} = 6500$. If (at least) one of the muons is detected in the endcaps, the efficiency and purity drop due to stronger reconstruction cuts. Nonetheless, for the maximum $dN_{ch}/d\eta|_{\eta=0} \approx 2500$ multiplicities expected in central PbPb at LHC, the efficiency (purity) remains above 65% (90%) even including the endcaps.




Figure 6.2: Υ reconstruction efficiency (left) and purity (right) as a function of the PbPb charged particle rapidity density, $dN_{ch}/d\eta|_{\eta=0}$.

If we only consider muon pairs in the central barrel, $|\eta| < 0.8$, the dimuon mass resolution is $\sim 54 \text{ MeV}/c^2$ at the Υ mass, as obtained from a Gaussian fit of the reconstructed $\mu\mu$ m_{inv} distribution (using a detailed MC simulation but without background). In the full pseudorapidity range, the dimuon mass resolution amounts to $\sim 1\%$: $35 \text{ MeV}/c^2$ at the J/ψ mass, and $86 \text{ MeV}/c^2$ at the Υ mass. These dimuon mass resolutions (the best among the LHC experiments) allow for a clean separation of the different quarkonia states. These values are used to smear the dimuon mass distribution in the fast MC studies.

6.1.3 Results

About 5×10^7 PbPb collisions were simulated. Muons passing the acceptance tables are combined to form pairs and each pair is weighted according to the trigger and reconstruction efficiencies (dependent on the momentum, η , purity and event multiplicity). Their invariant mass is calculated and smeared as described in the previous section. The obtained dimuon mass distributions are then scaled to 0.5 nb^{-1} , corresponding to the PbPb luminosity integrated in one month with average luminosity $L = 0.4 \cdot 10^{27} \text{ cm}^{-2}\text{s}^{-1}$ and 50% machine operation efficiency. Fig. 6.3 shows the resulting opposite-sign mass distributions, for the *high* multiplicity case, $dN_{ch}/d\eta|_{\eta=0} = 5000$ and full acceptance ($\eta < 2.4$). The different quarkonia resonances appear on top of a continuum due to several combinatorial background sources, the main ones being identified in the upper plots (h , c and b stand for $\pi + K$, charm and bottom decay muons, respectively). Since the CMS trigger and acceptance conditions treat opposite-sign and like-sign muon pairs in the same way, the uncorrelated background can be subtracted using the like-sign pairs: $N^{\text{Sig}} = N^{+-} - 2\sqrt{N^{++} N^{--}}$, shown also in the bottom panels of Fig. 6.3.

Figure 6.4 shows the *signal* dimuon mass distributions, after background subtraction, for two different scenarios: $dN_{ch}/d\eta|_{\eta=0} = 5000$, $|\eta| < 2.4$ (“worst” case conditions); and $dN_{ch}/d\eta|_{\eta=0} = 2500$, $|\eta| < 0.8$ (“best” case). Except for the ψ' , all quarkonia states are clearly visible. The corresponding signal-to-background ratios and yields (counted within 1σ of the resonance peaks) are collected in the Table 6.1 for one month of PbPb running.

6.1.4 Conclusions

With its very broad muon acceptance and precise tracking, CMS will provide significant contributions to heavy ion physics at the LHC. Studies of quarkonium production in PbPb collisions at $\sqrt{s_{NN}} = 5.5 \text{ TeV}$, will provide crucial information on the thermodynamical state of QCD medium formed in these collisions, through the expected step-wise “melting” pat-




Figure 6.3: Dimuon mass distributions measured within $|\eta| < 2.4$ for PbPb events with $dN_{ch}/d\eta|_{\eta=0} = 5000$ in the J/ψ (left) and Υ (right) mass regions. The main contributions of the background are shown in the top panels (h, c, b stand for $\pi + K$, charm, bottom decay muons resp.), while the bottom panels also show the like-sign pairs (combinatorial background).




Figure 6.4: Signal dimuon mass distributions after background subtraction in the J/ψ (left) and Υ (right) mass regions expected after one month of PbPb running. Top panels for $dN_{ch}/d\eta|_{\eta=0} = 5000$ and $|\eta| < 2.4$ ("worst" case conditions); bottom panels for $dN_{ch}/d\eta|_{\eta=0} = 2500$ and $|\eta| < 0.8$ ("best" measurement conditions).

Table 6.1: Signal-to-background ratios and expected quarkonia yields in one month of PbPb running (0.5 nb^{-1} integrated luminosity) for two multiplicity scenarios and two η windows.

$dN_{ch}/d\eta _{\eta=0}, \Delta\eta$	S/B	N(J/ψ)	S/B	N(Υ)	N(Υ')	N(Υ'')
2500, $ \eta < 2.4$	1.2	180 000	0.12	25 000	7300	4400
2500, $ \eta < 0.8$	4.5	11 600	0.97	6400		
5000, $ \eta < 2.4$	0.6	140 000	0.07	20 000	5900	3500
5000, $ \eta < 0.8$	2.75	12 600	0.52	6000		

tern of the different $Q\bar{Q}$ states due to colour screening. These results will also be sensitive to modifications of the low- x nuclear parton distribution functions, as expected in case of gluon saturation.

CMS can reconstruct the charmonium and bottomonium resonances, via their dimuon decay channel, with high efficiencies ($\sim 80\%$), good purity ($\sim 90\%$) and a very good dimuon mass resolution ($54 \text{ MeV}/c^2$ at the Υ mass), when both muons are detected in the central barrel ($|\eta| < 0.8$), even in the case of exceptionally high multiplicities ($dN_{ch}/d\eta|_{\eta=0} \approx 5000$). When considering the full pseudorapidity region ($|\eta| < 2.4$), the mass resolution becomes $\sim 86 \text{ MeV}/c^2$ at the Υ , and $35 \text{ MeV}/c^2$ at the J/ψ , with $\sim 50\%$ dimuon reconstruction efficiencies. The Υ states can be measured all the way down to $p_T = 0 \text{ GeV}/c$ with acceptances as large as 40%, while the lower rest mass of the J/ψ state and the large amount of material in the calorimeters absorbs “low” energy decay muons and prevents from measuring J/ψ 's below $p_T \approx 4 \text{ GeV}/c$. At high p_T (above $\sim 12 \text{ GeV}/c$ for the J/ψ and $\sim 4 \text{ GeV}/c$ for the Υ) the dimuon acceptance flattens out at 15%.

The large aperture of the muon detectors and the precise tracking result in a very good separation between the $Q\bar{Q}$ states in the dimuon mass distributions, and in relatively high statistics and good signal to background ratios ($S/B \approx 1(5)$, $S/B \approx 0.1(1)$ for J/ψ and Υ resp. in the full (central) rapidity range). After one month of PbPb running (0.5 nb^{-1}) we should collect $\sim 180\,000$ J/ψ and $\sim 25\,000$ Υ dimuons, enough to compare central and peripheral PbPb collisions, and to carry out some differential studies (dN/dy , dN/dp_T) which will surely contribute significantly to clarify the physics mechanisms behind the production (and “destruction”) of quarkonia states in nucleus-nucleus collisions at the LHC.

Part II

CMS Physics Reach

Chapter 7

Physics of Strong Interactions



7.1 The structure of the proton



QCD and jet physics

7.2.1 Introduction

With the start-up of LHC, a new domain of energy will be explored and an extrapolation of our current knowledge in the form of the Standard Model may not be sufficient to describe the new measurements. Even in a first data-taking phase with a rather low luminosity, studies of jet physics in the framework of quantum chromodynamics (QCD) will allow to check our current theory against the new data.

Figure 7.1 presents the decomposition of the total jet cross section into the partonic processes for $p\bar{p}$ collisions at the Tevatron and pp collisions at the LHC in dependence of the scaling variable $x_T = 2p_T/\sqrt{s}$, and illustrates the differences in cross section contributions of the PDFs compared to measurements possible today. In Figure 7.2 the expected statistical uncertainties on differential cross sections for all rapidities are presented for a pilot run with 0.1 fb^{-1} and for a first physics run with 10 fb^{-1} . Trigger pre-scales are taken into account. The figure demonstrates that already in the pilot high statistics will be available up to 1.5 TeV of transverse jet energy.

On the one hand, the measured data have to be corrected for detector effects using fully simulated events. Also, an energy calibration has to be performed on the reconstructed jets which ideally is extracted from data as well, but can also be done employing Monte-Carlo methods. On the other hand, for the theory predictions, which are most precise with respect to the hard parton-parton scattering amplitudes, effects of soft physics modelled in the form of parton showers and hadronization models with subsequent decays have to be taken into account. Once this is done, parameters of the current theory can be cross-checked or improved in precision by comparing the measured hadronic final state with the corrected theoretical predictions.

7.2.2 Jet algorithms

In order to re-establish a link between the observed particles that appear as collimated streams of hadrons in the detector and the hard process, algorithms are defined to group particles that are supposed to come from the same hard parton into jets. The required ingredients of such




Figure 7.1: Decomposition of the total jet cross section into the partonic processes for $p\bar{p}$ collisions at the Tevatron (left) and pp collisions at the LHC (right). The fractional contributions shown versus the scaling variable $x_T = 2p_T/\sqrt{s}$.




Figure 7.2: Expected statistical uncertainties on differential cross sections for all rapidities; left: for a pilot run with 0.1 fb^{-1} , and right: for a physics run with 10 fb^{-1} . The central cross section values are taken from a leading-order calculation in dependence of the transverse momenta of the hard interaction.

a jet algorithm are a distance measure to define the separation between objects, a procedure how to decide when objects are to be combined and a recombination scheme explaining how to combine objects. In addition, it has to be specified how the list of input objects has been determined.

Two principal types of algorithms are in common use: Cone type algorithms [165] that traditionally have been employed in hadron-hadron collisions where objects are clustered together that are close in angle around a high-energetic seed, and clustering algorithms where iteratively objects are combined that have the smallest distance of all pairwise combinations possible. The latter have predominantly been used in e^+e^- and $e^\pm p$ collisions, first in the form of the Jade algorithm [166, 167] and nowadays as k_T algorithm [168].

Both algorithms applied in this study use an angular distance measure based on the azimuthal angle Φ and, instead of the pseudo-rapidity η , the true rapidity $y = 0.5 \ln((E + p_z)/(E - p_z))$ which has become an established standard in recent publications [169, 170]. The distance between two objects i and j hence reads

$$\Delta R_{ij} = \sqrt{(\Delta_{ij}\Phi)^2 + (\Delta_{ij}y)^2}. \quad (7.1)$$

In addition, the most frequently used recombination scheme, the E scheme, implying a simple four-momentum addition, is employed in both cases.

Two types of jet algorithms are used here. The main results have been achieved with the k_T algorithm defined below, some cross checks have been performed with the midpoint cone algorithm:

1. Iterative clustering-type: Inclusive k_T algorithm [171] with
 - Distances are evaluated according to the ΔR scheme, i.e. $d_{ij} = \min(p_{T,i}^2, p_{T,j}^2) \frac{\Delta R_{ij}^2}{D^2}$ with R_{ij} as in eq. 7.1
 - Jet resolution parameter $D = 1.0$
2. Cone-type: Midpoint cone algorithm [172, 173] with:
 - Cone radius $R = 0.7$, all objects within a cone have to fulfill $R_{ic} \leq R$ with c labelling the four-vector of the current cone.
 - Overlap threshold $f_{\text{merge}} = 0.50$, i.e. overlapping cone jets are merged when they share more than 50% of the energy in the less energetic cone
 - Search-cone radius fraction $f_{\text{search}} = 0.5$, i.e. the first step to find the stable cones (before any splitting/merging is done) is performed with a smaller radius of $f_{\text{search}} * R$

Concerning the k_T algorithm, a jet resolution parameter of $D = 1.0$ is, from a theoretical point of view, best comparable to a cone algorithm with $R = 0.7$. In order to reduce the sensitivity to the underlying event it is advantageous to reduce the jet resolution parameter D or the cone radius R , respectively.

Note that primarily due to the limited choice of available jet energy calibrations the definition of the midpoint algorithm above has been selected. It does not exactly correspond to the definition given in [172] but to a modified one [173] that is in use by the CDF collaboration [169]. There have been indications that this algorithm leads to an infrared sensitive behaviour [174], so it is recommended to use the original definition of the midpoint algorithm without extra search cone radius.

7.2.3 Trigger scheme, event selection and phase space

The level one (L1) and the high level triggers (HLT) required for this analysis are the single-jet triggers which are described in more detail in section E.4.3.2. QCD jet production has, by several orders of magnitude, the largest cross section, but in contrast to most other analyses QCD jet events are the signal here. Therefore the sole other selection requirement for this study demands all jets to have a transverse momentum larger than 50 GeV. The available phase space is then subdivided into 17 ranges in transverse momentum p_T and five ranges in rapidity y , where the focus is mostly on the central region up to 2.5 in rapidity.

7.2.4 Input data

The analyzed events were generated with PYTHIA [175] and subsequently subjected to the full GEANT-4 based CMS detector simulation and reconstruction programs. Following the analysis setup presented in the introduction 7.2.1, four classes of input objects to the jet algorithms have been considered: The initial partons of the hard interaction, partons after parton shower (partonic final state, PFS), all stable particles of the hadronic final state (HFS) other than muons or neutrinos and calorimeter towers. The calorimeter towers fulfilling the requirements $E > 0.8 \text{ GeV}$ and $E_T > 0.5 \text{ GeV}$ were subjected to the same jet algorithms as the generator particles. If necessary a matching of generator and calorimeter jets was performed by looking for the pairs closest to each other in distance $d = \sqrt{(\Delta\Phi)^2 + (\Delta\eta)^2}$.

7.2.5 Jet energy calibration

The jet energy calibration has been performed with a MC calibration method implying calibration factors that are applied on a jet by jet basis to the calorimeter jets depending on pseudo-rapidity η and transverse momentum p_T . The alternative data based technique of gamma-jet calibration where jet transverse energies are measured against recoiling high energetic photons could not yet be employed for this study.

7.2.6 NLO calculation

In order to compare to theoretical predictions of perturbative QCD, calculations of at least next-to-leading order (NLO) precision are required. Here, the program NLOJET++ [176] is employed for the NLO calculation. However, since precise computations in NLO are very time consuming, a more efficient set-up in the form of the fastNLO project [177] is used which allows the fast rederivation of the considered cross section for arbitrary input PDFs and α_S values. This is done by separating the PDF dependency from the hard matrix element calculation by interpolating the PDFs between fixed support points in fractional proton momentum x so that the PDF dependency can be evaluated a posteriori from one complete calculation.

Note that neither PYTHIA nor NLOJET++ contain electroweak corrections which may change high p_T cross sections from 1 TeV onwards by up to 30% [178]. Insofar this study is consistent, but before comparing to real data this has to be taken into account.

7.2.7 Experimental and theoretical uncertainties

From the experience at the Tevatron [169, 179, 180], it is known that the jet energy scale with an uncertainty of 3% represents by far the dominant source of uncertainty for high p_T jet cross sections. Similarly, PDF uncertainties lead to the dominant uncertainty of the jet cross sections from the theoretical side.

According to CMS studies the jet energy scale in this analysis has been varied by $\pm 3\%$ in order to estimate the impact on the cross section determination. Figure 7.3 presents on the left hand side the corresponding relative experimental uncertainty on the jet cross section for three regions in rapidity. Starting at about 15% at low p_T it rises up to about 50% at high p_T for central rapidity. In the two non-central rapidity regions the uncertainties are of

comparable size below about 1 TeV of transverse momentum, but get considerably larger for higher p_T . In general, a similar behaviour as expected from Tevatron results is observed.

By evaluating the cross section calculation for the error set of the CTEQ6M [29] PDFs the ensuing theoretical uncertainty as shown in figure 7.3 on the right hand side could be derived. It is of the same order of magnitude as the energy scale uncertainty and rises from about 5% for low transverse momenta with a minimum of 3% at ≈ 200 GeV up to +65% and -30% at highest transverse momenta for central rapidity.




Figure 7.3: Relative systematic uncertainties of the jet cross sections for the k_T algorithm versus p_T due to a change in energy scale of $\pm 3\%$ for three bins in rapidity (left). The error bars represent the statistical uncertainty. On the right hand side, the relative uncertainties due to an evaluation of the error sets of the CTEQ6M [29] PDFs are shown for the same regions in rapidity.

7.2.8 Summary and Outlook

The dominant experimental and theoretical uncertainties on the differential inclusive cross sections of jets with high transverse momentum ranging from 80 GeV up to 4000 GeV have been investigated. A variation of $\pm 3\%$ in the jet energy scale results in an uncertainty of the derived jet cross sections of 15% at low transverse momenta, increasing up to about 50% at the highest p_T for central rapidity. The theoretical uncertainty due to the parton density functions of the proton has been found to be of the same order of magnitude and rises from about 5% for low transverse momenta with a minimum of 3% at ≈ 200 GeV up to +65% and -30% at the highest transverse momenta. For higher rapidities both uncertainties are considerably larger. The results shown have been derived with the k_T jet algorithm, similar values were obtained with the midpoint cone algorithm.

For transverse momenta below about 500 GeV further sources of uncertainties may give significant contributions to the total uncertainty, e.g. corrections due to pile-up, the underlying event and multiple interactions or hadronization. Theoretical contributions due to scale variations are of the order of 5% (10% for transverse momenta larger than 3 GeV) for rapidities y below 1.5. Above a rapidity of 1.5 they might be larger especially at the edge of the phase

space. In addition, contributions due to α_S and electroweak corrections have to be included before comparing to real data.

In the future, it will be possible to run simultaneous fits of α_S and the parton density functions, especially the gluon density at high x , to the data. To be less sensitive to the jet energy scale other jet related quantities, e.g. jet rates, will be considered. By including other processes into the fit procedure, like W/Z production as a luminosity measure or Drell-Yan reactions to fix the low x gluon density, powerful combined PDF fits to the data of one experiment will become possible.



Underlying Event Studies

7.3.1 Definition of the physics process and status of the art

The “Underlying Event” (UE) in a hard scattering process is everything accompanying an event but the hard 2-to-2 scattering component of the collision. A CDF analysis [181, 182] showed that the density of particles in the UE of jet events is about a factor of two larger than the density of particles in a typical Minimum Bias (MB) collision. At the LHC the difference might be even larger.

Hard scattering collider events have a distinct topology and one can use the topological structure of the collision to define regions of the η - ϕ space that are sensitive to the UE components of the interaction. By comparing different processes such as high transverse momentum jets, “back-to-back” dijet production, or Drell-Yan, one can partially isolate the various components contributing to the UE.

Multiple parton interaction (MPI) models [183], extending the QCD perturbative picture to the soft regime, turn out to be particularly adequate to describe the physics of the UE. In the framework of these models one can regard the observed differences between the UE in a hard scattering process and a MB collision as the effect of the increased probability of partonic interactions for small impact parameter hadron-hadron collisions: one hard scattering implies a small impact parameter collision which makes it more likely that an additional parton-parton interaction will occur. Also, a hard scattering promotes initial and final state gluon radiation which inevitably contributes to the UE.

Examples of MPI models are implemented in the general purpose simulation programs PYTHIA [62], JIMMY [184], and SHERPA [185]. Other successful descriptions of UE and MB at hadron colliders are achieved by alternative approaches like PHOJET [186], which rely on both perturbative QCD and the Dual Parton Models (DPM). The purely phenomenological description available in HERWIG [187] provides a very useful reference of a model not implementing multiple interactions.

The QCD models considered in this study are different settings, called tunes, of relevant parameters in HERWIG and PYTHIA 6.2. One of the PYTHIA tunes is the ATLAS tune [188] and the other (PY Tunes DW) is a tune by R. Field which is similar to PYTHIA Tune A [189]. All these tunes use the CTEQ5L parton distribution functions. Details of the settings are given in reference [190].

Both Tune A and Tune DW fit the CDF Run 1 and Run 2 UE data [181, 182]. Tune DW also fits the CDF Run 1 Z -boson transverse momentum distribution [191]. Both Tune A and

Tune DW use the same multiple parton interaction energy dependence parameter $\text{PARP}(90) = 0.25$, while the ATLAS tune uses the default value of 0.16.

The analyses summarised in this section are described in detail in reference [190].

7.3.2 Underlying event observables discussed for charged jet events

Charged jets are constructed from the charged particles using a simple clustering algorithm and then the direction of the leading charged particle jet is used to isolate regions of η - ϕ space that are sensitive to the UE. As illustrated in Figure 7.4, the direction of the leading charged particle jet, chgjet1 , is used to define correlations in the azimuthal angle, $\Delta\phi$. The angle $\Delta\phi = \phi - \phi_{\text{chgjet1}}$ is the relative azimuthal angle between a charged particle and the direction of chgjet1 . The “transverse” region is almost perpendicular to the plane of the hard 2-to-2 scattering and is therefore very sensitive to the UE. We restrict ourselves to charged particles in the central region $|\eta| < 1$ and consider two p_T thresholds, the nominal CMS cut $p_T > 0.9 \text{ GeV}/c$ and a lower threshold with $p_T > 0.5 \text{ GeV}/c$.




Figure 7.4: Illustration of correlations in azimuthal angle ϕ relative to the direction of the leading charged particle jet ($R = 0.7$) in the event, chgjet1 . The angle $\Delta\phi = \phi - \phi_{\text{chgjet1}}$ is the relative azimuthal angle between charged particles and the direction of chgjet1 . The “transverse” region is defined by $60^\circ < |\Delta\phi| < 120^\circ$ and $|\eta| < 1$. We examine charged particles in the range $|\eta| < 1$ with $p_T > 0.5 \text{ GeV}/c$ or $p_T > 0.9 \text{ GeV}/c$.

Figure 7.5 shows the QCD Monte Carlo models predictions for the average density of charged particles, $dN_{\text{chg}}/d\eta d\phi$, and the average charged PT_{sum} density, $dPT_{\text{sum}}/d\eta d\phi$, respectively, in the “transverse” region for $|\eta| < 1$ with $p_T > 0.5 \text{ GeV}/c$ and $p_T > 0.9 \text{ GeV}/c$ versus the transverse momentum of the leading charged particle jet. The charged particle density is constructed by dividing the average number of charged particles per event by the area in η - ϕ space (in this case $4\pi/3$). The charged PT_{sum} density is the average *scalar* p_T sum of charged particles per event divided by the area in η - ϕ space.

Due to the multiple parton interactions the PYTHIA tunes rise rapidly and then reach an approximately flat “plateau” region. At very high $P_T(\text{chgjet1})$ they begin to rise again due to initial and final state radiation which increases as the Q^2 scale of the hard scattering increases. HERWIG has considerably fewer particles in the “transverse” region and predicts a steady rise resulting from initial and final state radiation. The ATLAS tune predicts a larger charged




Figure 7.5: QCD Monte Carlo models predictions for charged particle jet production at 14 TeV. *Left:* Average density of charged particles, $dN_{chg}/d\eta d\phi$, with $|\eta| < 1$ in the “transverse” region versus the transverse momentum of the leading charged particle jet for $p_T > 0.5$ GeV/c (A) and $p_T > 0.9$ GeV/c (B). *Right:* Average charged PT_{sum} density, $dPT_{sum}/d\eta d\phi$, with $|\eta| < 1$ in the “transverse” region versus the transverse momentum of the leading charged particle jet for $p_T > 0.5$ GeV/c (C) and $p_T > 0.9$ GeV/c (D). The QCD models are HERWIG and two PYTHIA 6.2 tunes described in the text.

particle density than PYTHIA Tune DW for $p_T > 0.5 \text{ GeV}/c$. However, the ATLAS tune and Tune DW have similar charged particle densities for $p_T > 0.9 \text{ GeV}/c$. This is because the ATLAS tune has a “softer” charged particle p_T distribution than Tune DW.

7.3.3 Feasibility studies

Here we concentrate on the UE measurement that will be performed in nominal CMS conditions at low luminosity [190]. All the studies presented in this section have been obtained applying the GEANT-4 based simulation and reconstruction chain of the CMS experiment.

Events corresponding to Drell-Yan di-muon pairs and leading QCD processes with superimposed low luminosity pile-up have been generated with PYTHIA 6.2 in different \hat{p}_T regions. The relevant PYTHIA 6.2 parameters adopted by CMS in simulation production are documented in [192]. The triggers used to collect Jet and Drell-Yan samples are described in reference [68].

Charged track reconstruction relies on the Combinatorial Track Finder described elsewhere [193]. The default algorithm allows to reconstruct tracks with p_T above $0.9 \text{ GeV}/c$. However, the same algorithm can be used in special conditions (with reduced thresholds for the seeds) achieving reasonable performances down to $0.5 \text{ GeV}/c$ [190]. For $|\eta| < 1$, a reconstruction efficiency better than 90% and a fake rate below 1% are quoted for charged tracks with p_T above $0.7 \text{ GeV}/c$.

7.3.3.1 The Underlying Event as Observed in Charged Jet Events

The track-based measurement for the scale of the leading interaction allows to keep an acceptable resolution for jet energies below 20 GeV , where the calorimetric measurement is dominated by large systematic uncertainties.

In principle MB could be studied from any data selection, getting rid of the leading pp interaction and performing the reconstruction of all the primary vertices from all the other piled-up pp interactions. However this methodology turns out to be challenging as the resolution on the position of the pp vertices degrades when lowering the total p_T of the associated charged tracks. In this study an MB trigger is defined requiring at least a calorimetric jet of $p_T > 20 \text{ GeV}/c$. In order to combine the measurements performed at different leading charged jet scales, on top of the MB trigger, two additional triggers based on the p_T of the leading high level trigger jet are adopted: $p_T > 60 \text{ GeV}/c$ and $p_T > 120 \text{ GeV}/c$, which will be referred to as JET60 and JET120. Jets are reconstructed with an iterative cone algorithm of radius 0.5 in the pseudorapidity-azimuth space.

Tracks arising from the piled-up interactions are suppressed requiring the extrapolated coordinate along the beam axis to be inside 1 mm with respect to the primary vertex associated to the leading charged jet. The selection of the pp interaction with the highest p_T charged jet tends to create a small bias on the MB sample, reducing the statistics available at very low $P_T(\text{chgjet1})$.

The definition of the main UE observables have been introduced in section 7.3.2. The density of charged particles, $dN_{\text{chg}}/d\eta d\phi$, and the charged PT_{sum} density, $dPT_{\text{sum}}/d\eta d\phi$, with $p_T > 0.9 \text{ GeV}/c$ and $|\eta| < 1$ in the “transverse” region are reported in Figure 7.6. Bins of $2 \text{ GeV}/c$ are used up to $P_T(\text{chgjet1}) = 20 \text{ GeV}/c$ and bins of $10 \text{ GeV}/c$ above.




Figure 7.6: Charged jet production at 14 TeV. Charged tracks with $|\eta| < 1$ in the “transverse” region. Density of charged particles, $dN_{chg}/d\eta d\phi$ (A) and PT_{sum} density, $dPT_{sum}/d\eta d\phi$ (B), with $p_T > 0.9 \text{ GeV}/c$ versus the transverse momentum of the leading charged particle jet. Ratio between density of charged particles with $p_T > 0.9 \text{ GeV}/c$ and $p_T > 0.5 \text{ GeV}/c$ (C) and ratio between PT_{sum} density with $p_T > 0.9 \text{ GeV}/c$ and $p_T > 0.5 \text{ GeV}/c$ (D) versus the transverse momentum of the leading charged particle jet. Data from different triggers collected for an arbitrary integrated luminosity are superimposed: (*squares*) = Minimum Bias; (*circles*) = JET60; (*triangles*) = JET120. The lines show the generator level distributions; the points with error bars correspond to the raw (uncorrected) reconstruction level distributions.

The shapes of uncorrected reconstruction level distributions basically agree with the corresponding generator level ones. The difference in absolute scale (about -20% for both $dN_{chg}/d\eta d\phi$ and $dPT_{sum}/d\eta d\phi$) turns out to be compatible with charged track inefficiencies and fake rates. Further details on these systematic effects, including the calibration and resolution of the leading charged jet have been studied in [190].

Figure 7.6 shows also the ratio between the observables for $p_T > 0.9 \text{ GeV}/c$ and $p_T > 0.5 \text{ GeV}/c$ in the “transverse” region. These ratios, which are sensitive to the differences between the models and/or to the choice of the tuning for a given model, are also nicely free from the systematic effects enumerated above, and basically do not need to be corrected when comparing to the corresponding generator level observables.

7.3.3.2 The Underlying Event as Observed in Drell-Yan Muon-Pair Production

Drell-Yan muon pair production provides an excellent way to study the UE. Here one studies the outgoing charged particles (excluding the $\mu^+\mu^-$ pair) as a function of the muon-pair invariant mass. After removing the muon-pair everything else is the UE. As for the charged jet production, we restrict ourselves to charged particles in the central region $|\eta| < 1$ and

consider the two p_T thresholds $p_T > 0.5 \text{ GeV}/c$ and $p_T > 0.9 \text{ GeV}/c$.

Single muon and muon-pair CMS triggers ensure very high efficiencies for the studied process. The relative mass shift and the corresponding resolution of the reconstructed muon-pair are studied in detail in reference [190]. Tracks arising from the piled-up interactions are suppressed requiring the extrapolated coordinate along the beam axis to be inside 1 mm with respect to the primary vertex associated to the leading muons.




Figure 7.7: Muon-pair production at 14 TeV with two isolated muons. Density of charged particles, $dN_{chg}/d\eta d\phi$ (left), PT_{sum} density, $dPT_{sum}/d\eta d\phi$ (right), with $p_T > 0.9 \text{ GeV}/c$ and $|\eta| < 1$ versus the muon-pair invariant mass. (full circles) correspond to the generator level distributions; (empty circles) correspond to the raw (uncorrected) reconstruction level distributions. Arbitrary luminosity.

In our study, we require “isolated muons”, not to have charged tracks with $p_T > 0.9 \text{ GeV}/c$ in a cone of radius $R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} = 0.3$ in the azimuth-pseudorapidity space centered along the direction of the muon. Selecting isolated muons turns out to be essential to reduce the QCD background to negligible levels for $p_T > 15 \text{ GeV}/c$, while keeping an efficiency of 76.9% for Drell-Yan muon-pairs in the same p_T region.

The charge particle density, $dN_{chg}/d\eta d\phi$, and the charged PT_{sum} density, $dPT_{sum}/d\eta d\phi$ with $p_T > 0.9 \text{ GeV}/c$ and $|\eta| < 1$ in muon-pair production with isolated muons versus the muon-pair invariant mass are shown in Figure 7.7. Correlations between isolation and UE activity have been studied in references [190, 56].

7.3.4 Conclusions

Predictions on the amount of activity in UE at the LHC based on extrapolations from the lower energy data differ greatly. In this study we have demonstrated the feasibility of reference UE measurements at CMS under nominal conditions, assessing our capability to distinguish between the predictions of different models. The UE is studied by examining charged particles in the “transverse” region in charged particle jet production and in the central region of Drell-Yan muon-pair production (after removing the muon-pair).

7.4 Physics of b-quarks and hadrons



1 Inclusive b-quark production

7.4.1.1 Introduction

At the LHC new opportunities to improve our understanding of the physics of b quarks will become available because of the high statistics data samples and the high center-of-mass energy. A study [194] has been performed to investigate methods in CMS of identifying b jets (b “tagging”) in an inclusive sample of events containing jets and at least one muon. Here we present the capability to measure the inclusive b quark production cross section as a function of the B -hadron transverse momentum and pseudorapidity. An important result of our study is an estimate for the B -hadron p_T range reachable at LHC.

Inclusive b -quark production has been studied at other proton and electron colliders. The observed shapes of distributions and correlations are reasonably well explained by perturbative QCD. However, the observed cross-sections at the Tevatron (Run I) are larger than QCD predictions [195, 196, 197, 198, 199, 200, 201, 202] which is confirmed by Run II data. Similar effects are observed in γp collisions at HERA [203, 204, 205, 206, 207, 208, 209] and in $\gamma\gamma$ interactions at LEP [210, 211].

The agreement between experiment and theory has improved due to more precise parton density functions and proper estimates of fragmentation effects [212, 213, 214, 215, 216, 217]. But the agreement is not complete and the improvement of the phenomenological description is required using also experimental input.

7.4.1.2 Analysis

This study of the CMS capability to measure the inclusive b production is based on full detector simulation. The generated events are passed through the GEANT4 simulation of CMS. Pile-up corresponding to low-luminosity LHC running conditions ($\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$) is also generated.

7.4.1.2.1 Event selection About 4 million signal and background events were processed, mainly with high transverse momentum of the partons ($p_T > 50 \text{ GeV}/c$). Samples of QCD jets were used. Jets in those samples cover the full geometrical acceptance in pseudorapidity of the tracking detector, $|\eta| < 2.4$. The measurement of the differential cross sections is studied for B -hadrons of $p_T > 50 \text{ GeV}/c$ and within the fiducial volume of $|\eta| < 2.4$. First, the events are required to pass the Level-1 (L1) trigger selection for the single muon trigger stream which accepts events with muons having $p_T > 14 \text{ GeV}/c$. The most energetic B -hadron inside the phase space defined above is selected. The trigger efficiency is flat as a function of the B -hadron pseudorapidity within the L1 trigger acceptance of $|\eta| < 2.1$. It increases with transverse momentum of the B -particle. The average L1 trigger efficiency corresponds to the expected value of the branching fractions for the semileptonic b quark and c quark decays, about 19% [45]. At level-1, the single muon trigger is used. At the High Level Trigger (HLT) we require the “muon + b -jet” trigger, fired by non-isolated muons with $p_T > 19 \text{ GeV}/c$ and by jets with $E_T > 50 \text{ GeV}/c$, $|\eta| < 2.4$ and compatible with b tagging.

The event selection requires a b -tagged jet in the fiducial volume to be present in the event. B tagging is based on inclusive secondary vertex reconstruction in jets [148]. The tagging algorithms combine several topological and kinematic secondary vertex related variables into a single tagging variable to discriminate between jets originating from b quarks and those from light quarks and gluons.

To measure differential cross sections for inclusive B -particle production as a function of its

transverse momentum p_T and pseudorapidity η , $d\sigma/dp_T$ and $d\sigma/d|\eta|$, we select as the reconstructed B -particle candidate the most energetic b tagged jet. Good correspondence between the generated B -particle and the reconstructed b -tagged jet is observed. The corresponding p_T and pseudorapidity relative resolutions are shown in Figure 7.8 for B -particles with $p_T > 170$ GeV/ c . The resolutions are 13% and 6% for p_T and pseudorapidity, respectively.




Figure 7.8: Relative resolution, $(\text{Reconstructed} - \text{True}) / \text{True}$, for p_T and pseudorapidity of b tagged jets in CMS.

The efficiency of the b tagging by secondary vertices in jets is shown in Figure 7.9 as function of the B -particle transverse momentum and pseudorapidity. The b tagging efficiency is defined with respect to events passing the L1 trigger and with a single muon of $p_T > 19$ GeV/ c selected. The efficiency decreases with increasing transverse momentum, while being rather flat as function of pseudorapidity. The slow degradation for larger transverse momenta is caused by the worsening of the tracking resolution with increasing p_T , an increased track multiplicity from fragmentation and more difficult pattern recognition in dense jets. The average b tagging efficiency is 65% in the barrel region, while the efficiency is about 10% less for the endcap region. The muon plus b -jet cross-channel trigger has a 4.3 Hz rate for the signal and a 6.1 Hz total event rate [194]. This trigger rate corresponds to a low-luminosity LHC run at $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$.

To measure the cross section one needs to know the number of selected events, the integrated luminosity, the event sample purity (signal fraction) and the signal efficiency. The signal fraction can be determined from the simulated prediction of the background contribution to the selected event sample. In order to rely less on the absolute prediction for the background one can extract the signal fraction using the prediction of the signal and background shapes for some sensitive variables. A fit to the data distribution using the simulated shapes for the signal and background is performed. To do so we apply a lepton tag by selecting inclusive muons.

7.4.1.2.2 Muon tag Muons are reconstructed in the muon chambers, matched to the inner tracker information and refitted using both subdetectors information. This provides the most precise muon track measurement. Each reconstructed muon is associated to the most energetic b tagged jet. The muon must be closer to this b tagged jet than to any other jet




Figure 7.9: The b tagging efficiency versus p_T and pseudorapidity of the generated B -particle.

in the event. Otherwise the event is discarded.

In most cases the tagged muon is inside the b jet. The average efficiency of associating the muon with the b tagged jet is 75 %.

7.4.1.2.3 Results We calculate the transverse momentum of the muon with respect to the b -jet axis which effectively discriminates between b events and background. The slopes of the p_T spectra are very different and this is exploited in the fit of the selected events to determine the fractions of the muon sources in the sample.

Figure 7.10 shows an example of the fit of the distribution of the muon p_T with respect to the closest jet, using the expected shapes for the muons from b events, charm events and light quark events. The normalisation of the three contributions are free parameters in the fit. The events in this plot are from a sample of QCD events generated with the PYTHIA “ p_T -hat” parameter in the range $230 < \hat{p}_T < 300 \text{ GeV}/c$. In the fit the shapes of the distributions were fixed using an independent QCD sample generated with $170 < \hat{p}_T < 230 \text{ GeV}/c$. The fit results as well as the Monte Carlo input are quoted in Table 7.1. The event fractions are well reproduced within statistical errors. In the actual experiment the shapes will be verified using data at different selection stages. Also the background shape will be derived from the data itself by applying an anti-tag selection (b -suppressed event sample).

Table 7.1: Results of the fit to the distribution of the transverse momentum of the muon with respect to the nearest b tagged jet. The number of beauty, charm and light quark events in the Monte Carlo input are compared to the fit result.

	MC input, $230 < \hat{p}_T < 300 \text{ GeV}/c$	Fit result
$N_{b\bar{b}}$	5250	5222 ± 501
$N_{c\bar{c}}$	2388	2050 ± 728
N_{uds}	1740	1778 ± 341

In Table 7.2 the b purity, $c\bar{c}$ and light quark event fractions for the different QCD samples




Figure 7.10: Fit of the muon p_T spectrum with respect to the closest b tagged jet. The sample of generated QCD events with “ p_T -hat” parameter in the range $230 < \hat{p}_T < 300 \text{ GeV}/c$ is tested. The contributions of tagged muons from b events (dashed curve), c events (dot-dashed curve) and light quark events (dotted curve) as defined by the fit are shown. The solid curve is the sum of the three contributions.

are shown. The b purity decreases from about 70 % down to 55 % from low p_T events to the high transverse momentum events. The expected number of $b\bar{b}$ events after event selection is quoted for 10 fb^{-1} of integrated luminosity. For the phase space of $p_T > 50 \text{ GeV}/c$ and $|\eta| < 2.4$ the event selection will allow for a b event statistics of about 16 million events. We conclude that for B -hadrons a p_T range up to $1.5 \text{ TeV}/c$ will be accessible with the CMS detector at the LHC.

Table 7.2: B purity and expected number of events after final event selection. The expected number of $b\bar{b}$ events is quoted for 10 fb^{-1} of integrated luminosity.

\hat{p}_T , GeV/c	$N_{\text{generated}}^{\text{QCD}}$	$b\bar{b}$ purity, %	$c\bar{c}$ fraction, %	uds fraction, %	$N_{\text{expected}}^{b\bar{b}}$
50 – 80	198993	66	32	2	1.4 M
80 – 120	294986	66	32	2	6.1 M
120 – 170	291982	72	26	2	5.1 M
170 – 230	355978	71	26	3	2.4 M
230 – 300	389978	73	24	3	0.9 M
300 – 380	283983	70	25	5	0.3 M
380 – 470	191989	68	27	5	88 k
470 – 600	190987	64	29	7	34 k
600 – 800	94996	60	31	9	10 k
800 – 1000	89999	60	30	10	2.0 k
1000 – 1400	89998	55	31	14	0.5 k

The background contribution from $t\bar{t}$ events has been estimated from a sample of one million simulated events including all decay modes. The total number of $t\bar{t}$ events passing the selection amounts to 104 thousand events for 10 fb^{-1} of integrated luminosity, corresponding on average to a less than 1 % background contribution. The $t\bar{t}$ background becomes more pronounced for the high p_T part of the inclusive B spectrum. In the region $p_T > 500 \text{ GeV}/c$

it amounts to 2.4 %.

The total event selection efficiency is about 5 %. By correcting for the semi-leptonic branching ratio of b quarks and c quarks it amounts to about 25 % on average. It turns out that the total efficiency is almost independent of transverse momentum and angle of the B -particle. Therefore the measurement of the differential cross section is less affected by systematic uncertainties due to bin-by-bin efficiency corrections.

7.4.1.2.4 Systematics Uncertainties Several potential sources for systematic uncertainties are considered and their impact on the observed cross section is detailed in Table 7.3. The largest uncertainty arises from the 3 % error on the jet energy scale [218] which leads to

Table 7.3: Sources of systematic uncertainty in % on the inclusive b production cross section measurement. The total systematic uncertainty is calculated by adding all contributions in quadrature.

Source	uncertainty, %
jet energy scale	12
event selection	6
B tagging	5
luminosity	5
trigger	3
muon Br	2.6
misalignment	2
muon efficiency	1
$t\bar{t}$ background	0.7
fragmentation	9
total	18

a cross section error of 12 % at $E_T > 50 \text{ GeV}/c$. Other important uncertainties arise from the event-selection procedure and the Monte Carlo modeling of the detector response, including the lepton identification and the detector resolution on the energy and angular variables which identify the fiducial volume. The effect of these systematic uncertainties is estimated by varying the corresponding cuts and repeating the fits for the newly selected event samples. It results in an uncertainty of 6 %. The expected b -tag systematics for 10 fb^{-1} integrated luminosity is 5 % [7]. The luminosity uncertainty is also 5 % [7].

The trigger efficiency will be determined from the data themselves. We estimate its uncertainty from Monte Carlo studies to be 3.0 %. The experimental uncertainties on the semi-leptonic branching ratio of b quarks [45] is also propagated to the measurement. The impact of the detector misalignment on the CMS b tagging performance has been investigated in [148]. The effect has been found to be small (2 %). The muon detection efficiency can be determined with better than 1 % precision [7]. The $t\bar{t}$ background subtraction uncertainty is conservatively taken as absolute value of the expected $t\bar{t}$ contribution to the considered phase space.

A large contribution is expected from the fragmentation modeling. We estimate the magnitude of the effect from the $D\bar{O}$ b -jet production measurement at Tevatron [202]. This uncertainty propagates to the cross section as a 9 % effect independent of jet E_T .

7.4.1.3 Conclusion

The event selection for inclusive b production measurement at CMS will allow to study b production mechanisms on an event sample of 16 million b events for 10 fb^{-1} of integrated luminosity. The b purity of the selected events varies as function of the transverse momentum in a range from 70 % to 55 %. Our estimate shows that with the CMS detector we can reach $1.5 \text{ TeV}/c$ as the highest measured transverse momentum of B hadrons.



2 Study of B_c hadrons

7.4.2.1 Introduction

The B_c meson is the ground state of the $\bar{b}c$ system, which is doubly heavy flavored. This unique character provides a window for studying heavy-quark dynamics that is very different from the one of quarkonium. The experimental study of B_c will help us to understand heavy quark dynamics and to test the spin symmetry derived in non-relativistic quantum chromodynamics (NRQCD) [219, 220, 221, 222, 223, 224, 225, 226, 227, 228]. B_c mesons have been observed at the Fermilab Tevatron collider by the CDF collaboration through the decay channel $B_c \rightarrow J/\Psi \ell \nu$ [229]. The mass and lifetime are measured to be [230] $M(B_c) = 6.40 \pm 0.39(\text{stat}) \pm 0.13(\text{sys}) \text{ GeV}/c^2$ and $\tau(B_c) = 0.46_{-0.16}^{+0.18} \pm 0.03(\text{sys}) \text{ ps}$, in agreement with the non-relativistic potential model [231, 232, 233] and other approaches [234, 235, 236].

Because of the higher colliding energy, the production cross section at the LHC is about a factor of 16 [223] larger than at the Tevatron. As also the LHC luminosity will be higher, CMS has the potential to collect much more B_c mesons than the Tevatron experiments do. We propose to study the B_c meson through $B_c \rightarrow J/\Psi \pi$, $J/\Psi \rightarrow \mu^+ \mu^-$. The goal is to measure the mass and lifetime, and to compare the results with theoretical predictions which do have large uncertainties at the moment. More details on the analysis can be found in reference [237].



2.2 Monte Carlo data samples




Figure 7.11: Comparison of p_T distributions of B_c mesons for the generator BCVEGPY, Gouz and PYTHIA.

A large amount of Monte Carlo data were produced to study the feasibility for CMS to measure the B_c mass and lifetime with the first fb^{-1} . There are two dedicated B_c generators,

one is called BCVEGPY, developed at ITP, Beijing, by Chang *et al.* [228, 223], and the other is developed at IHEP, Protvino, by Berezhnoy *et al.* [231, 232]. Both packages are based on perturbative QCD, and have been integrated into the SIMUB package [124]. PYTHIA [238] can also generate B_c events, but it takes much more CPU time than the dedicated ones. For comparison, the p_T distribution of B_c mesons, generated by PYTHIA, BCVEGPY and the Protvino package (named Gouz in the plot), are shown in Figure 7.11. One can see that the Protvino package produces higher p_T , while PYTHIA agrees with BCVEGPY. In order to save CPU time, BCVEGPY is used to generate B_c events. During generation, only events were retained which contain within $|\eta| < 2$ a B_c with $p_T > 10 \text{ GeV}/c$, together with a muon of $p_T > 4 \text{ GeV}/c$ within $|\eta| < 2.2$. After the kinematic cuts, the cross section multiplied by the branching ratio is 1.78 pb. 52,000 B_c events were produced, corresponding to 29.2 fb^{-1} of integrated luminosity.

Important background sources are J/Ψ mesons from decays of other B hadrons and prompt J/Ψ mesons. Because of their large cross sections also QCD jets, in particular $b\bar{b} \rightarrow \mu^+\mu^-X$, $c\bar{c} \rightarrow \mu^+\mu^-X$, as well as $W + \text{jets}$ and $Z + \text{jets}$ have to be considered.

B hadrons that decay into J/Ψ were generated with PYTHIA6.228 with kinematic cuts similar to B_c production, and prompt J/Ψ events were generated by PYTHIA6.324, where the color-octet contribution is included.

The full CMS detector simulation and reconstruction was applied to the generated samples. The fast simulation package FAMOS was also used to produce the B_c events, B hadrons, prompt J/Ψ and $c\bar{c} \rightarrow \mu^+\mu^-X$ (Table 7.4).

Table 7.4: The cross section multiplied by the branching ratio after kinematic cuts and the number of events produced for B hadrons and prompt J/Ψ and $c\bar{c} \rightarrow \mu^+\mu^-X$.

channel	$\sigma \cdot \text{Br. (pb)}$	Nevents
B^0	70.3	740,000
B^+	70.7	740,000
B_s	14.8	190,000
Λ_b	19.4	200,000
prompt J/Ψ	240.3	500,000
$c\bar{c} \rightarrow \mu^+\mu^-X$	1690	210,000

Samples corresponding to 10 fb^{-1} of B hadrons, 2 fb^{-1} of prompt J/Ψ and 0.12 fb^{-1} of $c\bar{c} \rightarrow \mu^+\mu^-X$ events were produced for the analysis. Additional background samples of about 950,000 QCD, 880,000 $W + \text{jets}$, 710,000 $Z + \text{jets}$ and 100,000 $b\bar{b} \rightarrow \mu^+\mu^-X$ events were used.

7.4.2.3 Selection

Signal events should have a b -jet, a c -jet and a B_c meson which decays into a J/Ψ and a pion, with the subsequent $J/\Psi \rightarrow \mu^+\mu^-$ decay. The selection starts from 2 muon tracks. The p_T of both muons should be larger than 4 GeV/c and the absolute value of η less than 2.2. The two muons should have different charge and share the same vertex. To form a J/Ψ candidate the invariant mass of the muons should be in a window between 3.0 and 3.2 GeV/c^2 . An additional track must be found at the same vertex of the J/Ψ which is inconsistent with a muon or an electron. The p_T of it should be larger than 2 GeV/c and the absolute value of η less than 2.4.

The decay length L_{xy} , the proper decay length L_{xy}^{PDL} and the error of the decay length σ_{xy} are calculated from the J/Ψ vertex and the primary vertex in the xy -plane. The resolution of the proper decay length is $25 \mu\text{m}$. It is found that the resolution is almost independent of the proper decay length. In order to suppress the prompt backgrounds, the second vertex has to be displaced from the primary one. We require $L_{xy}/\sigma_{xy} > 2.5$ and $L_{xy}^{PDL} > 60 \mu\text{m}$. In addition, the condition $\cos\theta_{sp} > 0.8$ is applied where θ_{sp} is the opening angle between the second vertex (pointing from the primary vertex) and the reconstructed B_c momentum. Finally, the reconstructed B_c candidate must be in a mass window between 6.25 and $6.55 \text{ GeV}/c^2$.

The number of B_c and background events for 1 fb^{-1} after the selection are listed in Table 7.5. The total number of background events was estimated to be 2.6 ± 0.4 , mainly from B hadron decays into J/Ψ . So far tagging of the b jet is not used in the analysis.

Table 7.5: Estimated number of signal and background events for 1 fb^{-1} .

B_c	B^+	B_s	B_0	prompt J/Ψ	Λ_b	$c\bar{c}$	$b\bar{b}$	QCD
120 ± 11	0.7 ± 0.2	0.1	0.9 ± 0.3	0.1	0.1	0.01	0.01	0.7 ± 0.1

Because of the high cross section the number of produced QCD Monte Carlo events is not sufficient to directly determine the QCD background which is therefore estimated in three steps [237]. At first the efficiency to select two muons is obtained directly from the QCD sample, then the efficiency to reconstruct two muons into a J/Ψ candidate is calculated from the $c\bar{c} \rightarrow \mu^+\mu^-X$ sample, and finally the efficiency for the J/Ψ candidate to fake a B_c meson is obtained from the prompt J/Ψ sample. The probability of a QCD event to pass the selection cuts is then approximated as the product of the above three efficiencies. In this way, the total number of QCD background for 1 fb^{-1} is estimated to be 0.7 events.

This study which is aimed at the first fb^{-1} collected with the CMS detector assumes that in this initial phase the di-muon trigger threshold can be set at values such that the applied cut of $p_T > 4 \text{ GeV}/c$ on both muons does not introduce a significant inefficiency at trigger level. In case the available trigger bandwidth will prohibit this, more sophisticated High Level Trigger algorithms like a J/Ψ mass window could be invoked to restore the trigger efficiency. A detailed study is underway.

7.4.2.4 Mass and lifetime fitting

A kinematic fit was applied to the selected events imposing a J/Ψ mass constraint and forcing the two muon tracks as well as the pion track to share the same vertex. After the kinematic fit the invariant mass of the J/Ψ – pion system is shown in Figure 7.12. A Gaussian fit provides a mean value of $6406 \text{ MeV}/c^2$, close to the input of $6400 \text{ MeV}/c^2$, and a mass resolution of $22 \text{ MeV}/c^2$. The number of signal events in the plot for 1 fb^{-1} is 120. Backgrounds from B hadrons and prompt J/Ψ are included in the plot, while other backgrounds are neglected here.

A binned likelihood fit was done on the proper decay length distribution of the selected B_c events with the likelihood defined as $L = \prod P(n_i, \mu_i)$. $P(n_i, \mu_i)$ denotes the Poisson distribution with n_i events observed and μ_i events predicted in the i -th bin:

$$\mu = N \cdot \epsilon(x) \cdot \exp(-x/c\tau) \otimes G(x, \sigma)$$




Figure 7.12: Left: The invariant mass of the J/Ψ and pion candidate for the selected B_c . Right: The B_c proper decay length distribution. Both plots correspond to 1 fb^{-1} .

Here x represents the proper decay length, N and $c\tau$ are the parameters to be fitted and $G(x, \sigma)$ is a Gaussian smearing function with σ fixed to $25 \mu\text{m}$ which is the resolution of the proper decay length. The efficiency $\epsilon(x)$ is obtained from the large B_c sample.

The result of the fit is $c\tau = 148.8 \pm 13.1 \mu\text{m}$ which is consistent with the used input value of $150 \mu\text{m}$. The distribution of the proper decay length together with the fit result is shown in Figure 7.12.

7.4.2.5 Systematic uncertainty

The influence of imperfect detector alignment which is of particular importance at the beginning of the CMS experiment on the track and vertex reconstruction has been studied in [93, 134]. It will affect the study of B_c in three ways: the momentum scale of muons and pions, the mass resolution and finally the vertex precision. Taking the scale uncertainty to be $\Delta(1/p_T) = 0.0005/\text{GeV}/c$, the resulting uncertainties on the B_c mass is $11 \text{ MeV}/c^2$ and $0.2 \mu\text{m}$ on $c\tau$.

The effect of the muon momentum resolution was estimated following [93] and muon p_T -values of 10, 100 and 1000 GeV/c were studied for different η . The Δp_T to be smeared for a muon track from B_c was extrapolated from its p_T and η according to [93]. The resulting B_c mass uncertainty is $10 \text{ MeV}/c^2$, and $0.8 \mu\text{m}$ on $c\tau$. The error from the vertex uncertainty was determined according to [134] causing an uncertainty on $c\tau$ of $2.4 \mu\text{m}$.

The uncertainty on the efficiency as function of the proper decay length originates from the limited Monte Carlo statistics. By subtracting \sqrt{N} events from the sample ($N = 3600$ events), new efficiencies were calculated and the fit was repeated. The observed difference of $0.1 \mu\text{m}$ on $c\tau$ is taken as systematic uncertainty.

The theoretical uncertainty was estimated from Figure 7.11 which shows the p_T distributions from different generator packages. The B_c events, generated by BCVEGPY, were reweighted to agree with the Gouz distribution and the analysis was repeated. The difference on $c\tau$ was

found to be $1.5 \mu\text{m}$ which is taken as the error from this source.

To check the sensitivity on the cuts, the muon and pion p_T cuts were changed by one standard deviation of their resolution, about 1.5% depending on η . Other cuts like on $\cos \theta_{sp}$ and on the proper decay length were changed by 10%. The resulting mass uncertainty is $0.1 \text{ MeV}/c^2$ and $0.2 \mu\text{m}$ on $c\tau$.

In total the systematic uncertainties on the mass and on $c\tau$ are estimated to be $14.9 \text{ MeV}/c^2$ and $3.0 \mu\text{m}$, respectively.

7.4.2.6 Conclusion

With the first fb^{-1} of data CMS is expected to measure the B_c mass with an uncertainty of $22.0(\text{stat.}) \pm 14.9(\text{syst.}) \text{ MeV}/c^2$ and $c\tau$ with $13.1(\text{stat.}) \pm 3.0(\text{syst.}) \mu\text{m}$, corresponding to a lifetime uncertainty of $0.044(\text{fit}) \pm 0.010(\text{syst.}) \text{ ps}$. About 120 $B_c^+ \rightarrow J/\Psi\pi^+$, with $J/\Psi \rightarrow \mu^+\mu^-$, events would be observed. At the moment, the theoretical calculation is at the leading order without the color-octet contribution. Therefore the uncertainties on the total cross section and the p_T distribution are large. In the real data analysis, J/Ψ + one track with $J/\Psi \rightarrow \mu^+\mu^-$ will be selected as a control sample, $B^+ \rightarrow J/\Psi K^+$ will be used to estimate the efficiency, and the side band of the J/Ψ peak will be used to estimate the background to B_c .



Diffraction and forward physics

7.5.1 Introduction

This section outlines the diffractive and forward physics that CMS can do – together with the TOTEM experiment. The CMS and TOTEM detectors involved are presented in Chapter 7 of Volume 1 of the CMS Physics TDR [7].

The combined phase space coverage of the two experiments makes it possible to study many physics subjects in diffractive interactions – from QCD and the investigation of the low- x structure of the proton to the production of SM and MSSM Higgs bosons. Diffractive events are characterised by the fact that the incoming proton(s) emerge from the interaction intact, or excited into a low mass state, with only a small energy loss. Diffractive processes with proton energy losses up to a few per cent are dominated by the exchange of an object with vacuum quantum numbers, the so called Pomeron, now understood in terms of partons from the proton. For larger energy losses, mesonic exchanges – Reggeons and pions – become important. The topology of diffractive events is characterised by a gap in the rapidity distribution of final-state hadrons due to the lack of colour of the exchanged object.

Events with a fast proton in the final state can also originate from the exchange of a photon. In particular, forward tagging one leading proton allows the selection of photon-proton events with known photon energy; likewise, tagging two leading protons gives access to photon-photon interactions of well known centre-of-mass energy.

Triggering of diffractive/forward events is discussed in [239]. More details on the work presented here can be found in [240].

7.5.2 The interest of diffractive interactions

There have been major advances in the field of diffractive interactions recently, largely driven by the study of diffraction at HERA and Fermilab. The essential results are discussed in [241] and can be summarised as follows:

- Many aspects of hard diffractive processes are well understood in QCD: the presence of a hard scale allows the use of perturbative techniques and thus to formulate the dynamics in terms of quarks and gluons.
- A key to this success are factorisation theorems in electron-proton scattering, which render part of the dynamics accessible to calculation in perturbation theory. The remaining non-perturbative quantities are the so-called diffractive parton distribution functions (dPDFs) and generalised (or “skewed”) parton distributions (GPDs). They can be extracted from measurements and contain specific information about small- x partons in the proton that can only be obtained in diffractive processes.

Diffractive parton densities are determined from inclusive diffractive processes and can be interpreted as conditional probabilities to find a parton in the proton when the final state of the process contains a fast proton of given four-momentum. Generalised parton distributions can be accessed in exclusive diffractive processes; they quantify correlations between parton momenta in the proton. Their t -dependence is sensitive to the distribution of partons in the transverse plane.

- To describe hard diffractive hadron-hadron collisions is more challenging since factorization is broken by rescattering between spectator partons. These soft reinteractions can produce additional final-state particles which fill the would-be rapidity gap. When such additional particles are produced, a very fast proton can no longer appear in the final state because of energy conservation. The effect is often quantified in terms of the so called “gap survival probability”. These rescattering effects are of interest in their own right because of their intimate relation with multiple scattering effects, which at LHC energies are expected to be crucial for understanding the structure of events in hard collisions.
- A fascinating link has emerged between diffraction and the physics of heavy-ion collisions through the concept of saturation, which offers a new window on QCD dynamics in the regime of high parton densities.
- Perhaps unexpectedly, the production of a SM or MSSM Higgs boson in diffractive pp collisions is drawing more and more attention as a clean channel to study the properties of a light Higgs boson or even to discover it. The central exclusive reaction, $pp \rightarrow pHp$, appears particularly promising.

7.5.3 A survey of the accessible diffractive/forward processes

The accessible physics is a function of the integrated luminosity. We assume standard LHC optics with $\beta^* = 0.5$ m unless stated otherwise. We recall that, in this case, the TOTEM Roman Pots (RP) at 220 m from the CMS interaction point have coverage for $0.02 < \xi < 0.2$, where ξ is the proton fractional momentum loss. Near-beam detectors at 420 m from the interaction point, currently also being considered [242], would cover $0.002 < \xi < 0.02$.

Low-luminosity ($\sim 10^{28} - 10^{30} \text{ cm}^{-2} \text{ s}^{-1}$) studies could profit from running with $\beta^* > 0.5$ m, where the ξ coverage of the 220 m RPs would be wider and the t resolution would improve because of the lower transverse momentum spread of the beam.

7.5.3.1 Inclusive single diffraction and double Pomeron exchange at low luminosity

At modest instantaneous luminosities, up to $10^{32} \text{ cm}^{-2} \text{ s}^{-1}$, inclusive single diffractive (SD) events, $pp \rightarrow pX$, as well as inclusive double-pomeron exchange (DPE) events, $pp \rightarrow pXp$, can be studied by requiring the presence of one or two rapidity gaps in the event. In the ξ range given above, the scattered proton can be detected and the kinematics of the events fully measured.

The inclusive SD and DPE cross sections, as well as their M_X dependence, even in the absence of a hard scale, are important quantities to measure at the LHC. Here M_X indicates the mass of the system X . These cross sections amount to approximately 15% and 1% of the total proton-proton cross section, respectively; their energy dependence is a fundamental parameter of (non-perturbative) QCD. In addition, since diffractive events constitute a major fraction of the pile-up events, their measurement is mandatory to be able to properly simulate and understand high-luminosity data, where, at instantaneous luminosities of $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$, approximately 35 pile-up events are superimposed, on average, to any event.

7.5.3.2 SD and DPE production of dijets, vector bosons and heavy quarks

The study of SD and DPE events in which the diffractively excited state includes high- E_T jets, heavy quarks or vector bosons opens up the possibility of accessing dPDFs and GPDs. The comparison of the DPE and SD rates for these processes may also give information on the hard diffractive factorisation breaking at LHC (see Sect. 7.5.2). A few examples of these processes are given here.

Production of dijets

The measurement of the reaction $pp \rightarrow pXjj$ (j indicates a jet) has been used for the first time by CDF to measure the diffractive structure function in antiproton-proton collisions [243]. A similar measurement is possible at LHC with wider kinematic coverage (CDF: $\xi > 0.035$) and larger minimum jet E_T . For $E_T > 45 \text{ GeV}$, of the order of 10^8 events per fb^{-1} can be expected.

Production of heavy quarks

Inclusive DPE production of $t\bar{t}$ pairs has been studied in the case in which the final state contains one muon and four jets (i.e. with one top quark decaying to b plus lepton and neutrino, and the other to three jets). The analysis required the detection of both final-state protons. The expected number of events is of order 1 – 100 for 10 fb^{-1} , depending on the theoretical model assumed.

SD and DPE production of B -mesons has also been looked at, with $B \rightarrow J/\Psi X$ and $J/\Psi \rightarrow \mu^+ \mu^-$. Here the number of expected events is much larger, of the order of a few events per 10 pb^{-1} in the DPE case and thousands in the SD case.

Inclusive DPE production of W bosons

Inclusive DPE production of W bosons, $pp \rightarrow pXWp$, is also sensitive to the dPDFs of the proton and is a relatively abundant process that can be studied at instantaneous luminosities where pile-up is small. In these conditions, the requirement that two final state protons be measured in the 220 m RPs suppresses both the QCD background and the inclusive W production. Several thousand events with $W \rightarrow e\nu$ or $W \rightarrow \mu\nu$ are expected, after cuts, for an integrated luminosity of 1 fb^{-1} . This process, in conjunction with SD production of

W bosons, can be used to study hard diffractive factorisation breaking using the LHC data alone, as mentioned above.

7.5.3.3 SM and MSSM central exclusive Higgs production

As the delivered luminosity reaches tens of fb^{-1} , the central exclusive production process (DPE) becomes a tool to search for new physics, delivering signal to background ratios of order 0.1 – 1 for Standard Model (SM) Higgs production [244] and more than an order of magnitude larger for certain supersymmetric (MSSM) scenarios.

By central exclusive, we refer to the process $pp \rightarrow p\phi p$, where there are large rapidity gaps between the outgoing protons and the decay products of ϕ . There are three primary reasons why this process is attractive. Firstly, if the outgoing protons remain intact and scatter through small angles, then, under some general assumptions, the central system ϕ is produced in the $J_Z = 0$, C and P even state. Secondly, the mass of the central system can be determined very accurately from a measurement of the transverse and longitudinal momentum components of the outgoing protons alone. This means an accurate determination of the mass irrespective of the decay mode of the centrally produced particle such as $\phi \rightarrow b\bar{b}$. Thirdly, the process delivers excellent signal to background ratios, due to the combination of the $J_Z=0$ selection rules, the mass resolution, and the simplicity of the event in the central detectors. An additional attractive property of central exclusive production is its sensitivity to CP violating effects in the couplings of the object ϕ to gluons.

The left panel of Fig. 7.13 shows the cross section times the branching ratio for central exclusive production of a Standard Model Higgs, with $H \rightarrow b\bar{b}$ and $H \rightarrow WW$, as a function of the Higgs mass for different theoretical approaches. The $b\bar{b}$ mode is particularly interesting for masses close to the current exclusion limit. The right panel of Fig. 7.13 shows the acceptance assuming various combinations of RPs at 220 m and near-beam detectors at 420 m. Both protons can be detected in the 220 m stations only for Higgs masses larger than $280 \text{ GeV}/c^2$; this reflects the ξ range for which the 220 m RPs have acceptance, $0.02 < \xi < 0.2$ (the mass of the centrally produced Higgs is related to the ξ via $M_H^2 = \xi_1 \xi_2 s$, with ξ_1, ξ_2 the fractional momentum losses of the two protons). However, asymmetric events with one proton at low ξ and another at large ξ can be detected by the combination of the 220 m and 420 m detectors ($0.002 < \xi < 0.02$).

Central exclusive production is generally an attractive way of searching for any new particles that couple strongly to glue. An example studied in [245] is the scenario in which the gluino is the lightest supersymmetric particle. In such models, there should exist a spectrum of gluino-gluino bound states which can be produced in the central exclusive channel. Likewise, central exclusive production of radions, the fields introduced in the Randall-Sundrum model of five-dimensional quantum gravity, has been studied [246].

$H \rightarrow b\bar{b}$

The analysis is based on the requirement of two back-to-back central b -tagged jets in addition to the detection of both final-state protons yielding a mass of the central system consistent with that calculated from the protons alone. The event yield is very low, about 2 – 4 events per 30 fb^{-1} after all cuts, depending on the model. The non-resonant continuum b -jet background is largely suppressed by the $J_Z = 0$ rule. The residual background, mostly due to dijet production ($gg \rightarrow \text{dijets}$) and diffractive $gg \rightarrow b\bar{b}$ production, is a function of the mass resolution, which is about 1.6% for the '420+420' combination and 5.6% for the '220+420'




Figure 7.13: Left: The cross section for the exclusive production of the Higgs boson as a function of the Higgs boson mass for $H \rightarrow b\bar{b}$ and $H \rightarrow WW$. The different curves were obtained with the generators Exhume1.3 [247], DPEMC2.4 [248] and EDDE1.2 [249]. Right: Acceptance for the 420 m detectors alone and for the combination of the 220 m and 420 m detectors as a function of the Higgs boson mass.

combination (for $M_H = 120 \text{ GeV}/c^2$). The number of expected background events is of order 10 for 30 fb^{-1} .

$H \rightarrow WW$

In this case, the suppression of the background does not rely primarily on the mass resolution of the RPs. There are three main categories of WW events. Events in which at least one of the W bosons decays to an electron or a muon are the simplest, and pass the L1 trigger thanks to the high- p_T final-state lepton. This holds also if one of the W bosons decays into a tau, which subsequently decays leptonically. The four-jet mode occurs approximately half of the time; here, however, the RP information is necessary already at the L1 level. The expected event yields range between 1 and 7 events for 30 fb^{-1} , depending on the mass. Irreducible backgrounds are small and controllable.

MSSM Higgs

Double proton tagging is especially beneficial in the MSSM case. The b -jet channel is very important in the 'intense coupling regime' of MSSM ($M_h \approx M_A \approx M_H \approx 100 \text{ GeV}/c^2$) [250]: couplings of the Higgs to gg , WW^* , ZZ^* are strongly suppressed, making the discovery challenging by conventional means. Rates for central exclusive production of the two scalar (0^+) MSSM Higgs bosons (h, H) are more than a factor 10 larger than for the SM Higgs. The enhancement for $H \rightarrow b\bar{b}$ is by orders of magnitude in the M_h -max scenario for $M_H \approx 180 - 250 \text{ GeV}/c^2$; likewise for $h \rightarrow b\bar{b}$ and $h \rightarrow \tau\tau$ for $M_h \approx 90 - 130 \text{ GeV}/c^2$ [251]. In the small α_{eff} scenario, $h \rightarrow b\bar{b}$ and $h \rightarrow \tau\tau$ can be heavily suppressed for large $\tan\beta$ and for $M_h \approx 120 \text{ GeV}/c^2$ [251], whereas $h \rightarrow WW$ may be enhanced by up to a factor 4 compared to the SM predictions. Also, the pseudo-scalar (0^-) Higgs boson (A) is practically not produced in the central exclusive channel, yielding a clean separation of the scalar and pseudo-scalar Higgs bosons, impossible in conventional channels. The good missing mass resolution al-

lows to resolve h , H and, if enough statistics is available, measure their widths. This makes central exclusive production a possible discovery channel. Central exclusive production is also interesting in the '3-way mixing' scenario of CP-violating MSSM [252]: here the 3 neutral Higgs bosons are nearly degenerate, mix strongly and have masses close to $120 \text{ GeV}/c^2$.

Central exclusive production, with its good mass resolution via the scattered protons, may allow disentangling the Higgs bosons by studying the production lineshape. Explicit CP-violation in the Higgs sector causes an asymmetry in the azimuthal distributions of tagged protons (via the interference of P-even and P-odd amplitudes) – a measurement unique at the LHC [250, 253].

7.5.3.4 High-energy photon interactions

A significant fraction of events at the LHC involves photon interactions at energies above the electroweak scale [254]. The protons radiating the photon often survive the collision intact and are scattered at angles comparable to the beam angular divergence. Detection of such events at the LHC will open up a new field of high-energy photon physics, which is briefly outlined below. By requiring the detection of one or two forward protons like in diffractive interactions, photon-photon and photon-proton interactions can be selected. The photon fluxes, and the effective luminosities of photon-photon and photon-proton collisions are well known [255, 256]. The average proton energy loss is larger and the proton scattering angle smaller in photon exchanges than for the diffractive case. This can be used to establish relative contributions of these two processes.

Two-photon exclusive production of W and Z boson pairs

The cross section for the production of W pairs via photon-photon interactions, $pp \rightarrow ppWW$, is slightly above 100 fb ; in almost half of these events both forward protons are produced within the acceptance of the TOTEM RPs. About $100 \text{ events per } 10 \text{ fb}^{-1}$ with leptonic W decays can be detected in CMS. This allows a precise study of the gauge couplings, in particular of the $\gamma\gamma WW$ coupling. The expected sensitivity to anomalous quartic gauge couplings (QGCs) will surpass the LEP and Tevatron limits by orders of magnitude. A deviation from the Standard Model predictions would also allow a clean detection of anomalous WW production as predicted e.g. by A. White's theory of the supercritical Pomeron [257]. Two-photon production of Z pairs, $pp \rightarrow ppZZ$, is not allowed at the SM tree level, but yields similar sensitivities to the anomalous QGCs in this channel.

Two-photon exclusive production of pairs of SUSY particles

The cross sections for production of pairs of charginos, sleptons and charged Higgs bosons via photon-photon fusion at the LHC decrease rapidly with the masses of these particles [258]. This limits the scope of SUSY searches to particle masses below $150 - 200 \text{ GeV}/c^2$. However, the very clean environment of this reaction makes it attractive compared to other production mechanisms; the final state typically consists of two opposite-sign leptons and of missing p_T . The main background is due to the exclusive production of W pairs discussed above.

Two-photon production of doubly charged Higgs bosons (appearing in GUTs) is strongly enhanced, and leads to exclusive final states with two pairs of same-sign leptons.

Two-photon lepton pair production

Exclusive production of lepton pairs – a purely QED process at low $|t|$ – may serve for calibra-

tion of the pp luminosity; it may also be used for calibration of the momentum measurement of the scattered proton. Thousands of exclusive muon pairs are expected to be reconstructed in CMS for an integrated luminosity of 1 fb^{-1} . The striking signature of extremely small muon acoplanarity angles of less than about 10 mrad may be exploited already at the trigger level.

Single W and single top photoproduction

The cross section for single W photoproduction, $pp \rightarrow pWjX$, reaches almost 100 pb. This process can be therefore studied already at low luminosity. It also provides a means to study rescattering effects [256]. At higher luminosities, studies of high mass Wj states will be possible; for Wj invariant masses above 1 TeV, tens of events are expected to be detected in CMS (and tagged by TOTEM) per 10 fb^{-1} . This will allow to search for, as an example, an anomalous triple gauge coupling γWW . This process is the main background in the search for anomalous photoproduction of single top.

Associated WH and top pair photoproduction

The associated photoproduction of a SM Higgs boson and a W boson has a cross section of about 20 fb for Higgs mass below $180 \text{ GeV}/c^2$. About 50% of the forward protons are tagged by TOTEM, and events with leptonic W decay can be triggered efficiently in CMS. The cross section for photoproduction of top pairs is slightly above 1 pb. Top pair production is the main background for WH production, and in the photoproduction case the signal-to-background ratio for photoproduction of WH pairs is superior to the one in inclusive production.

7.5.3.5 Drell-Yan

The study of forward production of low mass Drell-Yan lepton pairs at the LHC provides a unique opportunity to directly access low- x partons in the proton. In this process, the lepton pair originates from the annihilation of a quark-antiquark pair whose fractional momenta, x_1 and x_2 , are related to the dilepton mass, M , and rapidity, y , through

$$M^2 = sx_1x_2; \quad x_{1,2} = \frac{M}{\sqrt{s}} \exp^{\pm y}, \quad (7.2)$$

with $\sqrt{s} = 14 \text{ TeV}$, the centre-of-mass energy of the colliding protons. In order to access low x , a large imbalance in fractional momenta is required, boosting the lepton pair to large rapidities.

The CASTOR calorimeter will cover the pseudorapidity range $5.3 < \eta < 6.6$, corresponding to Bjorken- x values down to 10^{-7} . With CASTOR alone, it may be possible to obtain a crude estimate of the dilepton mass. With the additional information provided by the T2 tracker, one can enhance the signal to background ratio by requiring tracks in association to the electromagnetic energy deposits. As T2 will measure both the azimuthal and polar angles of the tracks, a much more accurate measurement of the opening angle (and therefore of the dilepton mass) and a two-dimensional study in M^2 and x will become possible.

7.5.3.6 Validation of cosmic-ray generators

The correct simulation of the interaction of primary cosmic rays in the PeV energy range with the atmosphere is a key tool in the study of cosmic rays. Unfortunately, the available

generators differ significantly in their predictions for the energy flow, multiplicity, hadronic energy fraction etc., in particular at high rapidities. These models can be tested at the LHC: a 100 PeV fixed-target collision in air corresponds to the centre-of-mass energy of a pp collision at the LHC. Several generators were used to simulate inelastic and diffractive collisions at CMS: QGSjet [259], SIBYLL [260], DPMJet [261], neXus [259]. There are significant differences in the predictions, notably in the region covered by CASTOR, T1 and T2. A measurement of these features with CASTOR, T1 and T2 may thus be used to validate/tune these generators.



Physics with heavy ions

7.6.1 High-density QCD: heavy-ion physics

Quantum Chromodynamics (QCD) is the only existing quantum field theory whose collective behaviour, phase diagram and phase transitions, is accessible to study in the laboratory. High-energy nucleus-nucleus collisions offer the only experimental means known so far to concentrate a significant amount of energy ($\mathcal{O}(10 \text{ TeV})$ at the LHC) in a “large” volume ($\mathcal{O}(100 \text{ fm}^3)$) at thermalization times of $\tau_0 \approx 1 \text{ fm}/c$, allowing the study the many-body dynamics of strongly interacting matter. The programme of high-energy heavy-ion physics addresses several key open questions of the strong interaction:

- **Deconfinement and chiral symmetry restoration:** Lattice QCD calculations predict a new form of matter at energy densities above $\varepsilon \approx 1 \text{ GeV}/\text{fm}^3$ consisting of an extended volume of deconfined and bare-mass quarks and gluons: the Quark Gluon Plasma (QGP) [262]. The scrutiny of this new state of matter (equation-of-state, order of the phase transition, ...) promises to shed light on fundamental questions such as the nature of confinement, the mechanism of mass generation (chiral symmetry breaking, structure of the QCD vacuum) and hadronization, that still evade a thorough theoretical description due to their highly non-perturbative nature.
- **Non-linear parton evolution at small- x :** At high energies, hadrons consist of a very dense system of gluons with small (Bjorken) parton fractional momenta $x = p_{\text{parton}}/p_{\text{hadron}}$. At low- x , the probability to emit an extra gluon is large $\sim \alpha_S \ln(1/x)$ and non-linear gluon-gluon fusion processes start to dominate the parton evolution in the hadronic wave functions. Whereas at values of $x \gtrsim 10^{-3}$, the parton evolution with Q^2 (or $\ln(1/x)$) is described by the usual DGLAP (or BFKL) equations, at lower values of x and around $Q_s^2 \sim 3 \text{ GeV}^2/c^2$, such a saturated configuration is theoretically described in terms of the “Colour Glass Condensate” (CGC) picture [263]. Since the nonlinear growth of the gluon density depends on the transverse size of the system, the effects of gluon saturation are expected to set in earlier (at higher x) for heavy nuclei than for free nucleons.

In addition, the study of heavy-ion collisions has interesting connections to other research areas such as:

- **Early Universe cosmology:** The quark-hadron phase transition took place some $10 \mu\text{s}$ after the Big-Bang and was the most important event taking place in the Universe between the electro-weak (or SUSY) transition ($\tau \sim 10^{-10} \text{ s}$) and Big Bang

nucleosynthesis (BBN, at $\tau \sim 200$ s). Depending on the order of the QCD phase transition, several cosmological implications such as the formation of strangelets and cold dark-matter (WIMP) clumps or baryon fluctuations leading to inhomogeneous nucleosynthesis, have been postulated [264].

- **High-energy cosmic-ray physics:** The energy and mass of cosmic particles with energies above 10^{14} eV can only be measured via the ground-based detection of “extended air showers” (EAS) generated in upper-atmosphere interactions of cosmic rays (protons and ions up to Fe) with air (N,O nuclei). The interpretation of the EAS (and the related astro-particle phenomena) relies heavily on the accurate modeling of hadronic multi-particle production in proton-nucleus (p+N, p+O) and nucleus-nucleus (He+N, N+N, Fe+N) collisions in the TeV range. Direct measurements at LHC are needed in order to calibrate and tune the EAS models and correctly extrapolate their predictions to the highest cosmic-ray energies measured ($\sim 10^{20}$ eV).
- **Gauge/String duality:** Theoretical calculations based on the AdS/CFT correspondence permit to obtain results in strongly coupled ($g^2 N_c \gg 1$) gauge theories (QCD-like: SUSY $\mathcal{N} = 4$ Yang-Mills) in terms of a dual (10-D) gravity theory. Recent applications of this formalism have allowed, for the first time, to compute finite temperature QCD transport coefficients (such as the ratio of the QGP viscosity over entropy density, η/s) experimentally accessible, from black hole thermodynamics calculations [265].

7.6.2 Hard probes of QCD matter at LHC

Nucleus-nucleus collisions at the LHC offer a unique opportunity for studying strongly interacting matter at values of energy and particle densities never reached before. The factor of 30 increase in energy between RHIC and the LHC ($\sqrt{s_{NN}} = 5.5$ TeV for PbPb) leads to copious production of hard QCD probes: high- p_T hadrons, jets, quarkonia, direct photons, etc., arising from parton-parton scatterings with large squared momentum transfer, Q^2 . Such perturbative processes take place at time scales $\tau \approx 1/p_T \lesssim 0.1$ fm/c, and involve primary partons with fractional momenta of order $x \sim 10^{-3}$ (10^{-5}) at central (forward) rapidities. The produced hard probes are, thus, sensitive to initial-state modifications of the low- x parton distribution functions, as well as to final-state effects while propagating through the bulk matter formed in the collision. Thanks to the excellent detection capabilities of CMS in the perturbative sector (unparalleled $\eta - \phi$ acceptance for tracking and calorimetry, high granularity and resolution, fast detector technologies as well as sophisticated triggering), the heavy-ion physics reach of CMS will be extremely competitive at LHC. Among the various perturbative probes accessible to measurement, we focus in this report on the quarkonia detection via the $\mu^+ \mu^-$ decay channel. Other experimental capabilities, both in the hard (notably jet reconstruction in the heavy-ion environment) and “soft” (multiplicities, elliptic flow ...) sectors will be discussed in detail in CMS Physics TDR Volume III.

7.6.3 Gluon saturation and QGP colour screening via Quarkonia

The production of heavy-quarks at LHC proceeds mainly via gluon-gluon fusion processes and, as such, is sensitive to nuclear modifications of the gluon density at low- x . At $\sqrt{s_{NN}} = 5.5$ TeV, the average fraction of the nucleon momentum carried by the interacting parton

producing a J/Ψ at mid (forward) rapidity is $\langle x \rangle \approx 3 \cdot 10^{-3}(10^{-5})$. Such a kinematical domain is well in the regime where gluon saturation effects and departures from linear Q^2 (DGLAP) and $\ln(1/x)$ (BFKL) evolutions should be observable. In addition, the final-state formation of $Q\bar{Q}$ bound states is expected to be suppressed in a deconfined medium due to color screening of the heavy-quark potential. Recent finite-temperature lattice QCD calculations exhibit a substantial reduction of the heavy-quark internal energy $U_{Q\bar{Q}}$, with increasing temperature. The low-lying charmonia resonances (J/Ψ and η_c mesons) have been found to dissolve slightly below $2 \cdot T_{crit} \approx 330$ MeV, whereas much higher dissociation temperatures, $T_{diss} \approx 4 \cdot T_{crit}$ reachable at LHC, are needed to dissociate the Υ . Although J/Ψ suppression has been indeed observed in central A+A collisions both at CERN-SPS and RHIC energies, competing mechanisms to colour deconfinement (hadronic co-movers interactions and charm quark recombination) have been proposed to explain the observed cross-sections. At variance with charmonia states, the study of the much heavier bottomonia spectroscopy accessible at LHC is free from the distorting hadronic and coalescence contributions, and is directly sensitive to the temperature conditions of the produced partonic medium.

CMS has focused on the quarkonia detection through their decays to muon pairs. The good muon momentum resolution translates in an Υ mass resolution of $\sigma = 56$ MeV/ c^2 (in the central barrel region $|\eta| < 0.8$), the best of all the LHC detectors. This good resolution provides a clean separation between the members of the Υ family with a consequent improvement in the signal to background ratio, even in central PbPb with particle multiplicities as large as $dN_{ch}/d\eta = 5000$. The expected signal/background ratios are $S/B \approx 1(5)$, $S/B \approx 0.1(1)$ for J/Ψ and Υ respectively in the full ($|\eta| < 0.8$) rapidity range. In the absence of initial- or final-state medium effects, production cross sections of $B_{\mu\mu}\sigma = 50$ mb and $300 \mu\text{b}$ respectively will be measured in minimum bias PbPb collisions. The expected reconstructed yields for both charmonium and bottomonium resonances after background subtraction, in one-month data taking (with 50% overall efficiency) and nominal PbPb luminosity (0.5 nb^{-1}), are $\mathcal{O}(1.5 \cdot 10^5)$, $\mathcal{O}(2 \cdot 10^4)$ respectively. These statistics will allow detailed quantitative studies of quarkonia production as a function of p_T , rapidity and/or centrality. Any departure from the expected “vacuum” cross-sections will provide valuable information on the initial-state modifications of the nuclear parton (especially, gluon) distribution functions, as well as on the thermodynamical state of the produced medium from the predicted “melting” pattern of different quarkonia states due to colour screening.

Chapter 8

Physics of Top Quarks



8.1 Selection of $t\bar{t}$ events and measurement of the cross sections



8.1.1 Introduction

The goal of top physics at the LHC is to characterize the properties of this heaviest fermion of the Standard Model by measuring observables in its production and decay exploiting all possible decay channels. Important examples are the production cross section and the mass and spin properties of the top quark.

Most of the top quarks at the LHC will be produced as $t\bar{t}$ pairs. The $t\bar{t}$ production cross section is estimated to be 830 pb [266] at NLO and the dominant production mechanisms are gluon-gluon fusion ($\approx 90\%$) and quark-antiquark annihilation ($\approx 10\%$). Within the Standard Model the top quark decays almost exclusively to a W boson and a b quark. The decays of the $t\bar{t}$ system are then classified according to the decays of the W^+W^- system as di-leptonic, semi-leptonic or fully hadronic. The W can decay into leptons, $e^-\bar{\nu}_e, \mu^-\bar{\nu}_\mu, \tau^-\bar{\nu}_\tau$, or into quarks, $u\bar{d}', c\bar{s}'$, where the charge conjugate is implicit. Neglecting QCD corrections, branching fractions of 9/81 (11.1%) for the di-leptonic, 36/81 (44.4%) for the semi-leptonic and 36/81 (44.4%) for the fully hadronic decay channel are obtained.

For our studies we use PYTHIA for the simulation of signal and background events. As it includes spin correlation in $t\bar{t}$ production also samples generated with TOPREX are used for signal events.



8.1.2 Di-leptonic channel

8.1.2.1 Event Selection for 1 fb^{-1}

The very clean signature of this channel combined with a high signal-to-background ratio makes it possible to select $t\bar{t}$ -events with simple kinematic cuts. The selection is therefore suitable for the the expected early performance of the CMS detector and will allow to establish the signal as well as to measure the top mass at an early stage of the experiment.

For an integrated luminosity of 1 fb^{-1} about 54000 signal events are expected according to the leading-order estimate of PYTHIA. The main backgrounds with a final state mimicking the signal are Z, WW, WZ and ZZ production accompanied by jets. Furthermore, events from

semi-leptonic and fully-hadronic top-quark pair production with misidentified leptons and leptons from b -quark jets eventually constitute the dominating background. Here, dilepton events with W bosons decaying into τ -leptons are considered signal events if the τ lepton decays leptonically. Details of the analysis can be found in Reference [267].

Events are required to pass the Level-1 and High Level Trigger, in particular the single and di-lepton subtriggers. In addition to trigger criteria, events must contain at least two jets and two oppositely charged leptons. Electrons are identified using an electron likelihood method combining various electromagnetic shower variables and track-to-supercluster-matching criteria. After this preselection about 15000 signal events are left in a 1 fb^{-1} data set with a signal over background ratio of $S/B = 1/10$. The most important background at this stage consists of $Z + \text{jets}$ production with an accepted cross section of about 120 pb and a similar final state.

Isolation criteria reduce the contribution from misidentified leptons and leptons from b -jets. For a lepton candidate no other track or calorimeter hits amounting to 10% or more of the lepton p_T are allowed in a cone of $\Delta R < 0.2$. Two charged leptons are then chosen with a discriminant based on the likelihood ratio in case of an electron, the energy deposited in a cone of $\Delta R = 0.2$ around the lepton axis and the p_T of the lepton.

Both b -jets are selected with a discriminator based on the jet p_T , the invariant mass of tracks inside the jet and the output of the combined b -tagging algorithm [148]. Using this scheme the correct jets and leptons of the signal are selected for more than 90% of the events, if they could be reconstructed. It has been shown in reference [148] that, during the *first data taking phases* of the LHC, the degradation in b -tagging performance is still acceptable. This implies the b -tagging results presented here remain essentially correct.




Figure 8.1: Left: Invariant mass of the two lepton candidates indicating the cut window to remove $Z + \text{jets}$ events. Right: Most likely top mass after selection for 1 fb^{-1} .


Figure 8.1 shows the invariant mass of the two lepton candidates. The Z mass peak of the invariant mass distribution of two same type leptons is used to remove the contamination due to $Z + \text{jets}$ events. As a further improvement a cut on the b -tag discriminator is applied to the two selected jets.

The non-dilepton $t\bar{t}$ events usually contain more jets with a p_T greater than $30 \text{ GeV}/c$ but do

not contain two high p_T leptons. The second lepton candidate is considerably softer than the corresponding lepton from the signal decay channel. So a cut on the lower transverse momentum lepton is imposed with $p_T > 20 \text{ GeV}/c$. The two neutrinos in the decay of the W bosons lead to significant missing transverse energy E_T^{miss} whereas the decay of Z bosons into electrons or muons does not generate E_T^{miss} . The cut $E_T^{\text{miss}} > 40 \text{ GeV}$ further improves the signal to background ratio. At this stage about 1800 signal events are left with a signal over background ratio of $S/B = 7.3/1$.

The kinematics of the $t\bar{t}$ dilepton events yield an underconstrained equation system due to the two undetected neutrinos in the final state. However if all other kinematic quantities have been measured it is possible to make a fit imposing m_W and assuming a top mass parameter in the range of 100 to 300 GeV/c^2 . A weight can then be assigned to the different solutions obtained [267]. Figure 8.1 shows the distribution of the most likely top mass for signal and background events in the range $100 \text{ GeV}/c^2 < m_t < 300 \text{ GeV}/c^2$.

The event topology of most of the background events passing the previous cuts does not satisfy the dilepton kinematical constraints. Therefore considering only candidates which give a mass estimate in the range of 100 to 300 GeV/c^2 further reduces the background and raises the signal over background ratio to about $S : B = 12 : 1$. The remaining background essentially contains only non-dilepton $t\bar{t}$ events. In a dataset equivalent to 1 fb^{-1} , 657 signal events are selected with an overall efficiency of 1.2%.

We conclude that a measurement of the $t\bar{t}$ cross section and the top mass (see Section 8.2.1) in the dileptonic channel will be possible already with a modest amount of luminosity [267]. 

8.1.2.2 Event Selection for Higher Luminosities

The trigger is based on the presence of one muon or electron which covers with high efficiency all the possible final states in this channel. The selection of events in this channel then requires after trigger selection the presence of just two oppositely charged leptons with $E_T > 20 \text{ GeV}$ within pseudorapidity ranges of ± 2.4 and ± 2.5 for muons and electrons respectively. Details are available in [267].

The reconstruction efficiency is good for both for muons and electrons. More than 97% of the generated muons are correctly reconstructed in the considered range, as well as 90% of the electrons, with p_T above 20 GeV/c [267]. An electron is considered isolated if the total uncorrected E_T of the jets within a cone $\Delta R \leq 0.3$, minus the lepton E_T , is less than 30% of the lepton E_T . In a similar way a muon is considered isolated, if the sum of the p_T of all the tracks present in a cone of $\Delta R \leq 0.3$ minus p_T of the muon is less than 2 GeV/c . Candidate events must have $E_T^{\text{miss}} > 40 \text{ GeV}$. The analysis requires at least two jets with uncorrected $E_T > 20 \text{ GeV}$ detected within $|\eta| < 2.5$, where a jet is defined as a fixed-cone cluster with a cone size of $R = 0.5$. Jets produced by electrons are discarded before applying the previous selection by removing those which have an electromagnetic supercluster within $\Delta R = 0.2$ with a ratio between the electromagnetic energy of that supercluster and the uncorrected jet energy above 0.75.

b -tagging techniques based on the explicit reconstruction of a secondary vertex in a jet [148] are used to further suppress backgrounds in which no jets from b -quarks are present. The dominant backgrounds to dilepton $t\bar{t}$ events are those which have real leptons, real E_T^{miss} and jets originating from initial or final state radiation, arising mainly from diboson (WW , WZ , and ZZ) + jets production, and also from top quark decays, either from the semileptonic

Table 8.1: Cumulative effect of the different selection criteria applied to the simulated $t\bar{t}$ dilepton sample (electrons and muons) and simulated backgrounds. The column denoted as τ corresponds to $t\bar{t}$ dilepton sample in which at least one W decays into a τ lepton. The numbers correspond to LO accepted cross sections in pb.

	Signal	τ	WW	WZ	ZZ	$Z + \text{jets}$	other $t\bar{t}$
Before selection	24.3	30.4	7.74	0.89	0.11	3912	438
L1 + HLT	19.4	15.1	4.4	0.37	0.07	657	92
2 jets $E_T > 20 \text{ GeV}$	11.5	9.8	0.6	0.012	0.006	23.9	73.1
$E_T^{\text{miss}} > 40 \text{ GeV}$	9.6	8.1	0.5	0.01	0.003	5.8	53.6
Two opp. charged leptons	3.2	0.42	0.04	0.001	0.001	1.17	0.12
b-tag of two highest E_T jets	1.12	0.15	0.002	$\sim 10^{-4}$	$\sim 10^{-5}$	< 0.01	0.05

channel or from tau decays producing leptons. This kind of backgrounds are expected to be determined using MC simulation. Instrumental backgrounds, are characterized in general by their large cross sections but not having real E_T^{miss} , among them are: $Z + \text{jets}$, Drell-Yan ($Z/\gamma^* \rightarrow \ell^+\ell^-$) production, “fake” leptons in $W \rightarrow \ell\nu + \text{jet}$ events where a jet is falsely reconstructed as a lepton candidate. In principle it is harder to estimate their contribution to the final sample using MC simulation.

After this selection an efficiency close to 5% is obtained, with a very high rejection of all the backgrounds considered at the level of $10^{-3} : 1$ or better, as shown in Table 8.1. A S/B value of 5.5 is obtained, the main background being the one arising from the dilepton channel itself in which at least one of the W decays into $\tau\nu_\tau$ and with a subsequent leptonic tau decay.

Different sources of systematic uncertainties have been identified that affect event selection and background determination and thus the cross section measurement. Detailed studies [267] of these sources have been done based mainly on the results of the studies performed in [7] and [192]. Among the most important experimental sources are uncertainties on the jet energy scale and the b-tag efficiency. The impact of theoretical and phenomenological uncertainties such as those on hadron fragmentation and PDF have been studied using samples generated with different PYTHIA parameters and simulated and reconstructed with the CMS fast simulation and reconstruction program. The uncertainty in the cross section coming from the luminosity estimation was taken as 3% as expected for 10 fb^{-1} integrated luminosity. As the non- $t\bar{t}$ background is small it does not contribute significantly to the uncertainty. The results are summarized in Table 8.2 and lead to an estimated total error on the $t\bar{t}$ cross section measured in the di-leptonic channel using electrons and muons of $\Delta\sigma_{t\bar{t}}/\sigma_{t\bar{t}} = 11\% (\text{syst}) \pm 0.9\% (\text{stat}) \pm 3\% (\text{luminosity})$.

8.1.2.3 Top decays to tau leptons

In this section studies performed to select events with τ leptons in the final state are presented. We consider here di-leptonic $t\bar{t}$ decays with one tau lepton decaying into hadrons in the final state $t\bar{t} \rightarrow b\bar{b}\tau\nu_\tau\ell\nu_\ell$, ($\ell = e, \mu$). The measurement of the ratio $BR(t\bar{t} \rightarrow \ell\tau + X)/BR(t\bar{t} \rightarrow \ell\ell + X)$ will allow to set new limits on the presence of non-standard physics in top decays. Furthermore, this channel is a source of background for Supersymmetry and Higgs searches, as well as for the other dileptonic top channels.

Tau candidates are selected and identified following the method of the MSSM Higgs and HLT analyses [268], adapting the different selection criteria to the momentum range in which tau candidates are expected to be produced in top decays [267]. The hadronic tau identification

Table 8.2: Uncertainties in the $t\bar{t}$ di-lepton cross section determination for 10 fb^{-1} .

Effect	$\Delta\sigma_{t\bar{t} \text{ dil } e/\mu}/\sigma_{t\bar{t} \text{ dil } e/\mu}$
Jet Energy Scale	3.6%
b-tag efficiency	3.8%
Lepton reconstruction	1.6%
E_T^{miss}	1.1%
ISR and FSR	2.5%
Pile-Up	3.6%
Underlying Event	4.1%
Heavy quark fragmentation	5.1%
PDF uncertainties	5.2%
Statistical uncertainty	0.9%
Integrated luminosity	3%

Table 8.3: Cumulative effect of the different selection criteria applied to the simulated $t\bar{t}$ sample. Numbers correspond to LO accepted cross sections.

Cut	Efficiency times cross sections (pb)			
	$t\bar{t}$ (signal)	$t\bar{t}$ (other dilepton)	$t\bar{t}$ (semileptonic)	$t\bar{t}$ (hadronic)
Before selection	15.62	38.94	218.88	218.88
Trigger	8.61	25.40	85.90	2.08
2 jets	6.97	18.90	80.08	2.04
≥ 1 Iso lepton	4.27	13.11	34.93	0.11
$E_T^{\text{miss}} \geq 40 \text{ GeV}$	3.58	10.89	26.41	0.05
1 lepton	3.48	6.73	25.24	0.04
τ cand. with opp. Q	0.75	0.20	0.75	0.001
b-tagging	0.29	0.07	0.30	0.0005

efficiency obtained in the di-lepton samples is about 30% using this method as can be seen in Figure 8.2.

Event selection proceeds in a similar way as in section 8.1.2.2 but only one isolated lepton (electron or muon) is allowed. One isolated tau candidate separated from the isolated lepton has to be present, and the isolated lepton and the tau candidate must have opposite charges. The effect of these selections are described in detail for the $t\bar{t}$ sample in Table 8.3. b -tag for the two accompanying jets is also required. An efficiency close to 2% is obtained, with a very high rejection of all the backgrounds considered. A S/B value close to 1 is obtained, the main background being the one arising from the $t\bar{t}$ semileptonic channel. The majority of the systematic uncertainties are described in section 8.1.2.2. There is another systematic uncertainty intrinsic to this analysis due to the τ reconstruction and identification. Based on preliminary studies, we assigned a 12% uncertainty to the τ reconstruction and identification. Statistical uncertainty in the cross section determination is about 1.3% for an integrated luminosity of 10 fb^{-1} . Then the relative uncertainty in the estimation of the cross section is given by $\Delta\sigma_{t\bar{t} \text{ dil } \tau, e\mu}/\sigma_{t\bar{t} \text{ dil } \tau, e\mu} = 16\% \text{ (syst)} \pm 1.3\% \text{ (stat)} \pm 3\% \text{ (luminosity)}$.



3 Semi-leptonic channel

The semi-leptonic $t\bar{t}$ decay has a final state topology of four hadronic jets of which two originate from a b -quark, an isolated lepton and missing transverse momentum. In this section




Figure 8.2: Reconstruction efficiency of tau candidates as a function of p_T and η . Errors are statistical only.

we consider the measurement of the cross section of the semi-leptonic $t\bar{t}$ production where the lepton is a muon [269].

Both the Level-1 and the High-Level Trigger selection criteria are applied on the simulated events, resulting in the efficiencies shown in Table 8.4. The single-muon trigger stream was used. The jets are reconstructed from the combined electromagnetic and hadronic calorimeter energy deposits and clustered with the Iterative Cone algorithm using an opening angle of $\Delta R = 0.5$. A transverse energy threshold of 0.5 GeV is applied on the input objects before clustering. Optimization of the parameter settings of the clustering algorithms are considered in [270]. Only the jets in the vicinity of the primary vertex are considered in the analyses, rejecting in general those jets with a small transverse momentum. The energy scale of the reconstructed jets is calibrated using the methods described in [271]. Among the list of muon candidates identified flavour, the muon originating directly from the W boson decay is selected following the procedure described in [272]. The transverse momentum components of the unobserved neutrino are estimated via the missing transverse momentum which balances the vectorial sum of the energy deposits in the calorimeter above the transverse energy threshold mentioned.

The event selection consists of a series of sequential cuts on kinematic or topological variables. The event is required to have at least four jets after applying the primary vertex constraint with a calibrated transverse energy, E_T , exceeding 30 GeV and within a pseudorapidity in the range of the tracker, $|\eta| < 2.4$. If more than four jets match this criterion, the four leading jets are selected as those with the highest E_T . Of these four jets, two have to be b -tagged according to the method applying a combined b -tag variable described in [269, 273, 274]. The selected lepton is required to be within the tracker acceptance and to have a transverse momentum larger than 20 GeV/ c .

After classifying two of the four reconstructed jets as b -quark and the other two as light quark jets, only two jet combinations remain to reconstruct the hadronically-decaying top. A kinematic fit [275] was applied on the reconstructed event for both jet combinations forcing the reconstructed W boson mass to its precisely known value. Before applying the kinematic

Table 8.4: Overview of the selection criteria applied. The expected S/B values take into account the respective Leading-Order cross-sections of the processes.

	Semi-lept. $t\bar{t}$	Other $t\bar{t}$	W+4j	Wbb+2j	Wbb+3j	S/B
Before selection	365k	1962k	82.5k	109.5k	22.5k	5.9
L1+HLT Trigger	62.2%	5.30%	24.1%	8.35%	8.29%	7.8
Four jets $E_T > 30$ GeV	25.4%	1.01%	4.1%	1.48%	3.37%	9.9
$p_T^{lepton} > 20$ GeV/ c	24.8%	0.97%	3.9%	1.41%	3.14%	10.3
b-tag criteria	6.5%	0.24%	0.064%	0.52%	0.79%	25.4
Kinematic fit	6.3%	0.23%	0.059%	0.48%	0.72%	26.7
Selected cross section (pb)	5.21	1.10	0.10	0.08	0.05	26.7
Scaled $\mathcal{L} = 1$ fb $^{-1}$	5211	1084	104	82	50	26.7

fit the energy scale of the light quark jets is corrected for an overall bias in the reconstructed W boson mass. Following the method described in [276] after the event selection mentioned above, an inclusive jet energy scale correction of -9.7% was obtained and applied to light quark jet candidates. The event is finally selected if the fit converged for at least one of the combinations.

The selection efficiency for the signal events is estimated to be $6.28 \pm 0.04\%$. The fraction of $t\bar{t}$ signal events in the selected sample of inclusive $t\bar{t}$ decays is estimated to be $82.8 \pm 0.2\%$. The signal-to-background ratio after the event selection is 26.7, where all $t\bar{t}$ decay channels are considered as signal. Hence the systematic effect of the background contribution is minor. It is shown in [269] that after the event selection topological observables will not help much in differentiating between signal and background. The cross section is therefore estimated from counting events. The statistical uncertainty on the estimated cross section is 1.2%, 0.6% and 0.4% for integrated luminosities of 1 fb $^{-1}$, 5 fb $^{-1}$ and 10 fb $^{-1}$, respectively.

Systematic effects are introduced only on the signal events, changing the efficiency of the event selection. Similar effects on the background samples should be a second order effect on the inferred cross section. For the theoretical or phenomenological uncertainties the prescription of [192] was used as described in [269]. The list of systematic uncertainties is shown in Table 8.5. The dominant systematic effects are b-tagging, and in the early stage the uncertainty on the integrated luminosity. For an extended discussion on the studied systematic effects we refer to [269]. As a consequence of the kinematic fit, the uncertainty on both the light- and heavy-quark jet energy scale results in a limited systematic uncertainty, of about 1.6%.

The total relative systematic uncertainty on the cross section is 10.5% which can be compared to a relative statistical uncertainty of 0.6% at 5 fb $^{-1}$. The total uncertainty of 10.5% scales with the integrated luminosity as shown in Figure 8.3. In this plot it is assumed that the uncertainty on the determination of the integrated luminosity scale as the inverse square root of the integrated luminosity. At an integrated luminosity of about 5 fb $^{-1}$ the total uncertainty is dominated by the uncertainty on the b-tagging performance. For the uncertainty on the b-tagging efficiency a conservative 5% is taken according to [274] although the Tevatron experience shows that a value of 2% can be reached [277, 278].

Table 8.5: Overview of the systematic uncertainties on the cross section.

	$\Delta\hat{\sigma}_{t\bar{t}(\mu)}/\hat{\sigma}_{t\bar{t}(\mu)}$		
	1 fb^{-1}	5 fb^{-1}	10 fb^{-1}
Simulation samples (ϵ_{sim})	0.6%		
Simulation samples (F_{sim})	0.2%		
Pile-Up (30% On-Off)	3.2%		
Underlying Event	0.8%		
Jet Energy Scale (light quarks) (2%)	1.6%		
Jet Energy Scale (heavy quarks) (2%)	1.6%		
Radiation (Λ_{QCD}, Q_0^2)	2.6%		
Fragmentation (Lund b, σ_q)	1.0%		
b-tagging (5%)	7.0%		
Parton Density Functions	3.4%		
Background level	0.9%		
Integrated luminosity	10%	5%	3%
Statistical Uncertainty	1.2%	0.6%	0.4%
Total Systematic Uncertainty	13.6%	10.5%	9.7%
Total Uncertainty	13.7%	10.5%	9.7%



4 Fully hadronic channel

The fully hadronic final state, characterized by a six-jets topology $t\bar{t} \rightarrow WWb\bar{b} \rightarrow qqqb\bar{b}$, has the largest branching fraction (46%), and kinematics that can be fully reconstructed. However, this channel is affected by a large background from QCD multi-jet production, which makes the isolation of the signal rather challenging, and internal jet-parton permutation uncertainties. Improvements in the signal-to-background ratio are possible by requiring the presence of b -quark jets and by selecting central and very high-energy kinematic configurations which are expected for jets arising from the decay of a massive object like the top quark. A specific multi-jet trigger which uses b -tagging information has been devised for this analysis and an optimized selection has been applied. The analysis is described in detail in [267].

The signal sample consists of 500000 inclusive $t\bar{t}$ events, from which a subsample of 230000 fully hadronic $t\bar{t}$ events is extracted. The background consists of 1.5 million multi-jet events (QCD) generated with $50 < p_T < 470 \text{ GeV}/c$, where the p_T symbol indicates the transverse momentum of the most energetic parton of the hard scattering before the final-state radiation processes.

8.1.4.1 Trigger preselection and event selection

The trigger preselection uses the inclusive jet trigger envisaged in [68] and a special inclusive b -jet trigger [279]. The inclusive b -jet trigger combines in the first stage the b -tagging requirement with an inclusive jet trigger which applies tuned E_T thresholds of 350 GeV for single jets, 150 GeV for 3-jet and 55 GeV for 4-jet topologies; then a b -tagging based on pixel and regional track and vertex reconstruction is performed on the two most energetic jets. The trigger requires either multiple jets in the event or a b -tagged jet among the two highest- E_T jets. After the trigger preselection the QCD rate is reduced to 23 Hz, the signal efficiency is 16.8% and the signal to background ratio, S/B , amounts to 1/300.

The selection is designed to optimise the statistical significance $S/\sqrt{S+B}$ for an integrated




Figure 8.3: Statistical and total uncertainty on the inferred cross section of the process $pp \rightarrow t\bar{t} \rightarrow bq\bar{q}b\mu\nu_\mu$ as a function of the integrated luminosity.

luminosity of $\mathcal{L} = 1 \text{ fb}^{-1}$. The first step of the selection requires a topology of $6 \leq N_{jet} \leq 8$. For a jet to be counted, the jet pseudorapidity must satisfy $|\eta| < 2.4$ and its transverse energy must be greater than 30 GeV. Event shape variables, potentially able to separate the signal from the background are then taken into account. The useful ones are centrality, aplanarity and non-leading jet total transverse energy obtained removing the two most energetic jets ($\sum_3 E_T$) of which distributions are shown in Figure 8.4. After the selection b -tagging is applied to the surviving samples of $t\bar{t}$ fully hadronic and QCD events. Selection criteria of at one b -jet and two b -jets are considered.




Figure 8.4: Distributions of centrality, aplanarity and $\sum_3 E_T$ for $t\bar{t}$ and QCD events (normalized to the same area).

Table 8.6 summarizes the selection applied in cascade. The signal-to-background ratio amounts to 1/17 and 1/9 for the 1 and 2 b -tag samples, respectively, and resulting in signal efficiencies of 3.8% and 2.7%.

The signal efficiency relative to the total inclusive $t\bar{t}$ sample, to be used in the calculation of the total $t\bar{t}$ production cross section, becomes 2.3% (1.6%), respectively for the 1 (2) b -

Table 8.6: $t\bar{t}$ fully hadronic and QCD effective cross sections, signal-to-background ratio, statistical significance for 1 fb^{-1} and signal efficiency at each step of the selection.

Selection	Requirement	$\sigma\epsilon$ [pb]	$\sigma\epsilon_{\text{QCD}}$ [pb]	S/B	$S/\sqrt{S+B}$	ϵ (%)
Before Selection (PYTHIA LO)		225	25M	$1/10^5$	0.04	100
Trigger	HLT multi-jet+b-jet	38	11600	1/300	11.1	16.8
Event	$6 \leq N_{jet} \leq 8$	35	7900	1/225	12.4	15.5
	$E_T \geq 30 \text{ GeV}$	15	930	1/60	15.4	6.6
	centrality ≥ 0.68	9.9	324	1/33	17.1	4.4
	aplanarity ≥ 0.024	9.0	251	1/28	17.7	4.0
	$\sum_3 E_T \geq 148 \text{ GeV}$	9.0	229	1/25	18.4	4.0
b-tagging	1 b-tag	8.6	148	1/17	21.7	3.8
	2 b-tag	6.0	54	1/9	24.1	2.7

tag requirement. The estimated statistical uncertainty on the cross section is reported in Table 8.7.

Table 8.7: Number of $t\bar{t}$ and QCD events, $t\bar{t}$ efficiency, absolute and relative statistical uncertainties expected on the cross section measurement for an integrated luminosity of 1 fb^{-1} .

Requirement	$\mathcal{L} = 1 \text{ fb}^{-1}$				
	$t\bar{t}$ events	QCD events	ϵ (%)	$(\Delta\sigma)_{stat}$ [pb]	$(\Delta\sigma/\sigma)_{stat}$ (%)
1 b-tag	11500	148000	2.3	17	3.5
2 b-tag	8000	54000	1.6	15	3.0

Sources of systematic uncertainty are studied as described in detail in [192] and [7]. From the experience of CDF and $D\bar{O}$ experiments at Tevatron [280], one of the dominating systematic uncertainties arises from jet energy scale. The systematic uncertainty related with the trigger selection is calculated considering contributions from b-tagging and jet energy scale. Table 8.8 summarizes the contributions to the total uncertainty on the cross section, which combined lead to a relative uncertainty of $\Delta\sigma/\sigma = 3\%(stat) + 20\%(syst) + 5\%(luminosity)$.

Table 8.8: Contributions to the systematic uncertainty on the $t\bar{t}$ cross section measurement in the fully hadronic channel (cut based approach).

	$\Delta\sigma/\sigma$ (%)
HLT	5.9
Pile Up	10.0
Underlying Event	4.1
Fragmentation	1.9
PDF	4.2
IS/FS Radiation	7.9
Jet Energy Scale	11.2
b-tagging	2.0
Background	5.0
Integrated Luminosity	5.0

8.1.4.2 Event selection based on neural net

A more refined selection is based on a neural net exploiting the same variables considered so far. Such approach is attempted in order to investigate the possibility of improving the S/B ratio and/or the efficiency. The previous selection which will be called “early” selection could represent a more conservative approach for the first LHC analyses.

The most effective neural network configuration studied is applied to the $t\bar{t}$ and QCD events satisfying the topology request of $6 \leq N_{jet} \leq 8$ (jet pseudorapidity $|\eta| < 2.4$) after a cut on jet transverse energy of $E_T > 25$ GeV and consists of 6 input nodes: E_T of the first and sixth jet with the jets ordered in increasing E_T , centrality, aplanarity, $\sum_3 E_T$ and sphericity. The performance of the neural net is shown in Figure 8.5 which compares the output distributions for signal and QCD background. The S/B ratio as a function of the $t\bar{t}$ efficiency is also shown. With respect to the “early” selection, the request for a neural net output ≥ 0.77 improves the S/B ratio from 1/25 to 1/10 with same efficiency of about 4%.




Figure 8.5: Left: distribution of the neural net output for $t\bar{t}$ and QCD. Right: signal-to-background ratio as function of the signal efficiency. For comparison the result of the “early” selection is also shown.

As done after the “early” selection, a b -tagging is applied to the surviving samples of $t\bar{t}$ fully hadronic and QCD events, and selection criteria of at least one b -jet and two b -jets are considered. Improved signal-to-background ratio, amounting to 1/7 (1/3) respectively for 1 (2) b -tag samples, can be achieved using the neural net keeping the same signal efficiencies of 3.8% (2.7%). This means an estimated relative statistical uncertainty on the cross section of 2.3% (2.0%), with the same expected number of $t\bar{t}$ events for an integrated luminosity of $\mathcal{L} = 1 \text{ fb}^{-1}$.

8.2 Measurement of the top quark mass

1 Di-leptonic events

The di-lepton channel has the advantageous feature of the cleanest signature and the highest achievable signal to background ratio in the selection. However, concerning the measurement of the top mass the presence of the two neutrinos prohibits the direct reconstruction of any of the top quarks. But the kinematic properties of the di-leptonic events reveal a large

sensitivity to the top mass which can be exploited in various ways. The method presented here is discussed in more detail in [267].

The six unmeasured kinematic quantities corresponding to the momentum components of the two neutrinos are reduced assuming momentum balance in the transverse plane, imposing the m_W constraint and the equality of both top quark masses. This way the event kinematics can be cast into a fourth order polynomial with the top mass as parameter. For each candidate event we step through top mass values in the range $100 \text{ GeV}/c^2 \leq m_t \leq 200 \text{ GeV}/c^2$ and weight the kinematic solutions, including their fourfold ambiguity, for the neutrino momenta with the spectrum expected in the Standard Model. For each event the most likely solution, i.e. the solution with the highest weight, is retained and their mass distribution is shown in Figure 8.1. This kinematic event reconstruction was already used to improve the signal to background ratio in the selection for 1 fb^{-1} (see Section 8.1.2.1). The figure shows a clear mass peak at the expected value for the fully simulated and reconstructed events. A Gaussian fit to the signal in a range corresponding to 0.4 of the maximum yields $m_t = 178.5 \pm 1.5 \text{ GeV}/c^2$ for an input top mass of $175 \text{ GeV}/c^2$. Hence the expected statistical accuracy on the top mass exploiting this method in the di-lepton channel is $1.5 \text{ GeV}/c^2$. The background is small and essentially flat and does not affect the mass determination significantly.

The main systematic effects on the mass estimation in the di-lepton channel using this kinematical reconstruction are due to the assumptions used to reduce the complexity of the kinematic equation system and the detector effects. The dominating systematic effect in the first category is the uncertainty on the initial and final state radiation, changing e.g. the amount of transverse momentum of the $t\bar{t}$ -system influencing the solvability of equation system and the neutrino solutions. This results in an uncertainty on the top mass of $\Delta m_t = 0.3 \text{ GeV}/c^2$ determined using the recipe in [192]. The zero width approximation for both the W bosons and the top quarks in the equation system gives rise to another shift of about $0.1 \text{ GeV}/c^2$.

The expected uncertainty of the jet energy scale after startup (using source calibration and test beam data) amounts to 15% independent on the jet p_T . For this measurement this shift amounts to $\Delta m_t = 4.2 \text{ GeV}/c^2$ in case of an integrated luminosity of up to 1 fb^{-1} . With better calibration ($\gamma + \text{jet}$) and W fit from the other $t\bar{t}$ channels) this error does reduce to $2.9 \text{ GeV}/c^2$ in a measurement with $1 - 10 \text{ fb}^{-1}$. **QUESTION:** What does W fit from the other $t\bar{t}$ channels mean? Further improvements in the knowledge of the jet energy scale (e.g. $Z + \text{jets}$ calibration) are expected reducing the uncertainty to about $1 \text{ GeV}/c^2$ after 10 fb^{-1} of integrated luminosity.

We conclude that employing the kinematic reconstruction method sketched here the di-lepton channel will allow already at an early stage of the experiment a meaningful measurement of the top mass. Assuming that the goal for a precise jet energy scale determination for b -quarks can be achieved the expected precision on the top mass in this channel with 10 fb^{-1} is $\Delta m_t = \pm 0.5 \text{ GeV}/c^2$ (*stat*) $\pm 1.1 \text{ GeV}/c^2$ (*sys*).



2 Semi-leptonic events

The semi-leptonic $t\bar{t}$ decay is traditionally called the *golden channel* for measuring the top-quark mass. A measurement based on advanced analysis tools is described in detail in [281]. The event reconstruction and initial event selection follows the one of Section 8.1.3. For the event to be selected, exactly two out of the four leading jets are b -tagged and the other

two need to be anti- b -tagged. The four leading jets should not overlap in order to reduce ambiguities in the jet energy scale calibration procedure. The efficiency of each sequential cut is shown in Table 8.9.

Table 8.9: Overview of the selection criteria applied after the lepton cut $p_T^{\text{lepton}} > 20 \text{ GeV}/c$ in Table 8.4.

	Signal	Other $t\bar{t}$	W+4j	Wbb+2j	Wbb+3j	S/B
Before selection	365k	1962k	82.5k	109.5k	22.5k	0.032
b-tag criteria	5.5%	0.21%	0.052%	0.47%	0.70%	3.73
No jet overlap	3.0%	0.11%	0.027%	0.25%	0.44%	3.87
P_{χ^2} -cut 20%	1.4%	0.039%	0.0097	0.061	0.07	5.3
P_{sign} -cut 80%	1.2%	0.025%	0.0085	0.052	0.05	6.8
P_{comb} -cut 50%	0.7%	0.013%	0.0036	0.013	0.	8.2
Scaled $\mathcal{L} = 1 \text{ fb}^{-1}$	588	64	6	2	0	8.2

The amount of events produced via a different $t\bar{t}$ decay channel in the selected event sample is reduced by a likelihood-ratio method combining three kinematic observables resulting in a variable L_{sign} which is transformed into a probability P_{sign} for the selected event to be a semi-leptonic muon $t\bar{t}$ event. An extra sequential cut is applied by requiring this probability P_{sign} to exceed 80%.

Among the four reconstructed jets, three have to be chosen to form the hadronic decaying top quark. The efficiency and purity of this selection was significantly enhanced by applying a second likelihood ratio method combining the information from several sensitive variables. The jet combination with the largest L_{comb} value is taken as the best pairing. The L_{comb} value is transformed into a probability P_{comb} for the chosen combination to be the correct one. The event probability P_{comb} is used in the event selection where events are selected if their value for P_{comb} exceeds 60%, increasing the purity of the selected jet pairings to 81.6% in the mass window of $25 \text{ GeV}/c^2$ around the expected m_t of about $175 \text{ GeV}/c^2$.

For each jet combination a kinematic fit was applied as described which imposes the W-boson mass for the hadronically-decaying W boson in the event [275]. Only jet combinations are considered with a probability of the kinematic fit calculated from its χ^2/ndf exceeding 20%. For some events none of the jet combinations fulfill this criterium, therefore reducing the total event selection efficiency. The fraction of fully hadronic $t\bar{t}$ events selected is negligible (less than 0.05 events expected at 1 fb^{-1}). From this we conclude that the also influence of QCD produced jet events is minor.

When estimating m_t from the selected event sample by a simple Gaussian fit in a range of $20 \text{ GeV}/c^2$ in both directions around the modal bin, a value of $176.5 \pm 0.65 \text{ GeV}/c^2$ is obtained before applying the kinematic fit and $172.2 \pm 0.48 \text{ GeV}/c^2$ after applying the kinematic fit, for an input value of $175 \text{ GeV}/c^2$. The errors reflect the statistical precision of the available Monte Carlo signal sample. The top quark mass after the kinematic fit is shown in Figure 8.6.

Rather than developing m_t estimators on samples of events, an event-by-event likelihood approach is used to estimate m_t from the fitted kinematics of the three jets of the hadronically decaying top quark. The uncertainty on m_t for each event is determined from the covariance matrices of the kinematic fit. This uncertainty can either be assumed Gaussian or the full m_t range can be explicitly scanned with the kinematic fit.




Figure 8.6: Left: Distribution of the mass of the hadronic decaying top quark for the selected events after applying the kinematic fit. Right: Estimated shift in $M_t^{FullIdeo}$ versus a relative shift α applied on the inclusive heavy quark jet energy scale.

To obtain information about the true value of M_t we convolute the reconstructed resolution function or ideogram with the theoretical expected probability density function $P(m_t|M_t)$ in the reconstruction space

$$\mathcal{L}_i(M_t) = \int P(\{\bar{p}_j\}|m_t) \cdot P(m_t|M_t) dm_t \quad (8.1)$$

where one integrates over the kinematic relevant range of m_t to obtain a likelihood function $\mathcal{L}_i(M_t)$ for each event i . Several contributions are added in the expected density $P(m_t|M_t)$: a Breit-Wigner shape for the correct jet combinations $S(m_t|M_t)$, a parametrized combinatorial background contribution $B_{comb}(m_t)$ and a parametrized background contribution $B_{proc}(m_t)$. This results in a function

$$P(m_t|M_t) = P_{sign} \cdot [P_{comb} \cdot S(m_t|M_t) + (1 - P_{comb}) \cdot B_{comb}(m_t)] + (1 - P_{sign}) \cdot B_{back}(m_t) \quad (8.2)$$

where each contribution is weighted according to the probabilities extracted from the observed event. After combining the likelihoods $\mathcal{L}_i(M_t)$ from all selected events, a maximum likelihood method is applied to obtain the best value for the estimator \hat{M}_t .

The linearity of the estimators have been checked and the slopes are found to be compatible with unity. The width of the pull distribution of the top quark mass estimators \hat{M}_t are found to be 0.82 for \hat{M}_t^{fit} (simple fit on reconstructed mass spectrum), 1.04 for $\hat{M}_t^{ParIdeo}$ (convolution with the parametrized ideogram) and 1.02 for $\hat{M}_t^{FullIdeo}$ (convolution with the full scanned ideogram). The resulting top quark mass for the estimator \hat{M}_t^{fit} applied on the simulated events samples with a generated top quark mass of $175 \text{ GeV}/c^2$ is $174.16 \pm 0.59 \text{ GeV}/c^2$, hence reflecting a bias of $-0.84 \text{ GeV}/c^2$. For the convolution method this is $170.65 \pm 0.54 \text{ GeV}/c^2$ and $172.42 \pm 0.31 \text{ GeV}/c^2$ for respectively the $\hat{M}_t^{ParIdeo}$ and the $\hat{M}_t^{FullIdeo}$ estimator. Figure 8.7 illustrates the results.

Several systematic effects introduce an uncertainty on the top quark mass estimator. They originate from our understanding of the detector performance, the robustness of the reconstructed objects, for example jets, and the general description of the proton collisions in the




Figure 8.7: Distribution of the mass of the hadronic decaying top quark before the kinematic fit used for the \hat{M}_t^{fit} estimator (left) and the combined $\Delta\chi^2(M_t)$ function over all events for both ideogram based estimators $\hat{M}_t^{\text{ParIdeo}}$ and $\hat{M}_t^{\text{FullIdeo}}$ (right).

simulation. A full description can be found in [281]. The estimation of the systematic uncertainties follows that of the cross section measurement in Section 8.1.3. We conservatively conclude that a total precision on the top quark mass of $1.9 \text{ GeV}/c^2$ can be reached with 10 fb^{-1} of data. The uncertainty is dominated by systematic effects like pile-up collisions and the knowledge of the jet energy scale of b -quark jets (see Figure 8.6).

After achieving a better understanding of the accelerator settings and the detector performance, however, the total uncertainty will decrease. Our understanding of the underlying event model will improve in the future significantly when new tuning data become available. The magnitude of pile-up collisions could be monitored to the level of 10%. To take into account the overlap between the pile-up and the jet energy scale uncertainty, the systematic shift due to a 10% variation in the pile-up collisions is divided by two. The uncertainty on the energy scale of b -quark jets can be extrapolated to about 1.5% after a better understanding of the detector performance and with the application of advanced tools like energy flow algorithms or selecting jets only in well understood regions in the detector. The measurement of the b -tag efficiency [274] is dominated by systematic uncertainties of radiation effects. The experience at the Tevatron collider [277, 278] illustrates that an uncertainty of 2% could be reached.

Table 8.10 summarizes and combines the extrapolated systematic uncertainties on each of the top quark mass estimators. The uncertainty on the inferred top quark mass of about $1.2 \text{ GeV}/c^2$ is dominated by the uncertainty on the energy scale of the b -quark jets. This relative uncertainty is taken to be 1.5% which defines a goal for the performance of jet calibration methods.



3 Fully hadronic events

The kinematic selection described in Section 8.1.4.1, including the demand for the two b -tags, forms the basis for a selection of fully hadronic $t\bar{t}$ events suitable for a kinematic top-mass

Table 8.10: Overview of all uncertainty components on the top quark mass estimators, extrapolated to a better understanding of both the proton collisions at the LHC and the detector performance.

	Standard Selection		
	Gaussian Fit Δm_t (GeV/c ²)	Gaussian Ideogram Δm_t (GeV/c ²)	Full Scan Ideogram Δm_t (GeV/c ²)
Pile-Up (5%)	0.32	0.23	0.21
Underlying Event	0.50	0.35	0.25
Jet Energy Scale (1.5%)	2.90	1.05	0.96
Radiation (Λ_{QCD}, Q_0^2)	0.80	0.27	0.22
Fragmentation (Lund b, σ_q)	0.40	0.40	0.30
b-tagging (2%)	0.80	0.20	0.18
Background	0.30	0.25	0.25
Parton Density Functions	0.12	0.10	0.08
Total Systematical uncertainty	3.21	1.27	1.13
Statistical Uncertainty (10 fb ⁻¹)	0.32	0.36	0.21
Total Uncertainty	3.23	1.32	1.15

reconstruction. An additional cut on the two leading jets, $100 \text{ GeV}/c < p_T < 300 \text{ GeV}/c$, is effective against background from mis-reconstructed events and combinatorial background.

The six partons in $pp \rightarrow t\bar{t} \rightarrow bW^+\bar{b}W^- \rightarrow bq_1\bar{q}'_1\bar{b}q_2\bar{q}'_2$ are matched to six reconstructed jets by picking the matching which minimises the sum of the angular separation between reconstructed jet and matched parton. Only jets satisfying our initial jet-definition, $p_T > 30 \text{ GeV}/c$ and $|\eta| < 2.4$, as employed in the selection, are taken into account in the matching process. Based on the amount of the angular separation three disjunctive classes of signal events are defined: good (36%), half-good (45%) and bad jet-parton-matching (19%). The first class being the events where all six partons are matched well by jets, the second class where only the three partons from one top are matched well by jets. The reason for the mismatch can be traced to parton-level properties, like high $|\eta|$ and low p_T , described in more detail in [267].

In order to perform the correct jet pairing, a likelihood variable is constructed from the following event observables: (a) average of the two W -boson masses, (b) difference of the two W -boson masses, (c) sum of the inter-jet angles of the W -boson candidates $\angle(q_1\bar{q}'_1) + \angle(q_2\bar{q}'_2)$, (d) difference of the two top-quark masses, (e) sum of the inter-jet angles of the top quark candidates $\angle(bq_1) + \angle(b\bar{q}'_1) + \angle(q_1\bar{q}'_1)\angle(\bar{b}q_2) + \angle(\bar{b}\bar{q}'_2) + \angle(q_2\bar{q}'_2)$, (f) angle between the direction of the two top-quark candidates. Their distributions are shown in [267]. Taking for each event the pairing with the highest likelihood value yields pairing efficiencies of 71% for the good and 64% for the half-good jet-parton-matching.

Only one top per event is chosen for the kinematic mass determination, the choice is once again based on a likelihood variable constructed from the following event observables: (a) p_T of the softest of the three jets of each top-quark candidate (b) mass of the W boson as reconstructed in top decay (c) sum of the inter-jet angles of jets from top decay, $\angle(b_iq_i) + \angle(b_i\bar{q}'_i) + \angle(q_i\bar{q}'_i)$. Taking the top with the larger likelihood value yields a 72% efficiency, far greater than the 50% efficiency of a random choice.

The differentiation of the selected signal events into the now six classes is summarised in

Table 8.11, where the six classes are being mapped onto two labels, indicating whether the events are considered signal- or background-like.

Table 8.11: Distribution of the different signal event classes after jet-pairing and top-choice in the $t\bar{t}$ fully hadronic channel. The label column indicates whether the class is considered signal- or background-like.

reconstruction	pairing	[pb]	top-choice	[pb]	label
good	correct	0.62 (35%)	always correct	0.62 (35%)	sig.
	wrong	0.26 (14%)	always wrong	0.26 (14%)	bkg.
half-good	correct	0.46 (25%)	correct	0.33 (18%)	sig.
			wrong	0.13 (7%)	bkg.
	wrong	0.26 (15%)	always wrong	0.26 (15%)	bkg.
bad	always wrong	0.20 (11%)	always correct	0.20 (11%)	bkg.

With all the pieces in place a kinematic reconstruction of the top quarks is straightforward and the resulting invariant mass distribution of the chosen top, with the paired non- b -jets scaled such that they yield the W -mass, is shown in Figure 8.8.




Figure 8.8: Invariant mass distribution of the reconstructed and rescaled, chosen top for both signal classes with a Gaussian fit to the peak.

As expected the signal-like events form a narrow peak, while the wrongly-reconstructed events have a far broader shape. Fitting a Gaussian to the peak of the invariant mass distributions with a fit range corresponding to 0.4 of the peak maximum, as shown in Figure 8.8 serves as a simple mass estimator. The extracted top-mass is $m_t = 175.0 \pm 0.6$ (stat.) ± 4.2 (syst.) GeV/c^2 for an input top-mass of $175 \text{ GeV}/c^2$ and an integrated luminosity of $\mathcal{L} = 1 \text{ fb}^{-1}$.

Already with this amount of data the statistical error becomes negligible compared to the systematic uncertainties which are summarised in Table 8.12. By far the biggest systematic uncertainty is the QCD background. The S/B in the displayed mass window of Figure 8.8 is about $2/3$, although not shown since the currently available number of simulated events does not allow a determination of the QCD background shape and of the uncertainty it in-

Table 8.12: Summary of the systematics for the top-mass determination with fully hadronic events.

	$\Delta m_t [\text{GeV}/c^2]$
Pile Up	0.4
Underlying Event	0.6
PDF	1.4
IS/FS Radiation	2.3
Fragmentation	0.9
Jet Energy Scale	2.3
b-Tagging	0.3
Background	2.0

roduces into the top-mass determination. Experience from CDF at the Tevatron [282, 283] indicates that this uncertainty can be understood at the $\sim 2 \text{ GeV}/c^2$ level, when using data for background estimation.



4 Top quark mass from J/ψ final states

8.2.4.1 Introduction

At the LHC the measurement of the top quark mass via direct reconstruction will soon be limited by systematic errors. It is expected that the most severe systematic contributions will be linked to the modelling of the hadronic environment and the knowledge of the jet energies. It would be particularly desirable, therefore, to consider methods for the extraction of m_t from the data which could reduce the contribution from these uncertainties considerably. An alternative method, which is making use of exclusive b decays in semileptonic top-pair events with the presence of a J/Ψ decaying into an electron or muon pair was proposed in [284, 285].

The top quark mass is determined by its correlation with the invariant mass of the reconstructed J/Ψ and the lepton from the W decay coming from the same top decay, $m_{J/\Psi\ell}$. The correlation is present because the reconstruction of the J/Ψ gives an accurate measurement of the b quark flight direction and its momentum thanks to the relatively high mass of the meson. Moreover, this measure is expected to have an excellent resolution because of the very clean experimental reconstruction of the lepton three-vectors. Details on the analysis presented here can be found in [286].

8.2.4.2 Event generation and selection

Signal events are generated using the TOPREX generator [35] and consist of $t\bar{t}$ events where the presence of at least one J/Ψ in the final state from the hadronization of b -quarks is required. No distinction is made about the origin of the J/Ψ ; therefore the same samples also contains combinatorial background where the J/Ψ is coming from a b quark produced together with a W boson not decaying leptonically. Five samples corresponding to five different top masses are generated with a statistics of 200K events each. The event hadronization and the description of the underlying event and the minimum bias is realized with PYTHIA 6.227 [30].

All the signal samples are passed through full detector simulation (ORCA) [15] with a simulation of the minimum bias corresponding to high luminosity data taking. Indeed, the statistics is expected to be so low that the use of high luminosity data must be considered. The same signal samples, and several millions more for studies on systematics, are passed through the fast simulation of the detector (FAMOS) [287]. The shape of the variables used in the selections are fully compatible in both scenarios.

The studied physics backgrounds are generated with the ALPGEN [152] generator and include $W + \text{jets}$, $Zb\bar{b} + \text{jets}$, $Wb\bar{b} + \text{jets}$. In these cases the samples are not biased by requiring an explicit J/Ψ in the final state, therefore the separation from the signal is studied on the basis of cuts not involving the search for a J/Ψ and the contribution of the resulting background is then rescaled taking into account the proper branching fractions. The selection, in terms of signal efficiency, is also cross-checked against $t\bar{t} + \text{jets}$ signal generated with ALPGEN, and is found to be consistent.

The main difficulty of the analysis comes from the extremely low branching ratio for a $t\bar{t}$ event to give a final state with a leptonic J/Ψ . This can be written as:

$$BR(t\bar{t} \rightarrow (Wb)(Wb) \rightarrow (Xb)(\ell\nu J/\Psi X)) = 2 \cdot BR(W \rightarrow \ell\nu) \cdot BR(b \rightarrow X) \rightarrow B^{\pm,0}, B_s, B_{\text{baryon}} \rightarrow J/\Psi X) \cdot BR(J/\Psi \rightarrow \ell\ell) \quad (8.3)$$

where charge conjugation is implicit, ℓ indicates either an electron or a muon, and having assumed a $BR(t \rightarrow Wb)$ of 1. Replacing the branching ratios with up-to-date numbers [45] one gets for the global branching ratio the value $5.5 \cdot 10^{-4}$ that, in terms of event yield and assuming a cross section for $pp \rightarrow t\bar{t}$ of 830 pb, makes approximately 4500 events per 10 fb^{-1} . This number does not include neither the trigger and selection efficiency, nor the efficiency for the correct pairing of the J/Ψ to the correct lepton from the W decay.

Events are triggered using the inclusive lepton trigger with thresholds described in [68]. The efficiency for triggering signal events is reported in Table 8.13 and is included in all numbers presented here. In events passing the trigger thresholds a J/Ψ is searched for by looking for same-flavour, opposite-sign leptons with invariant mass in the range $[2.8, 3.2] \text{ GeV}/c^2$ and forming an angle greater than 2 and lower than 35 degrees. No isolation requirements must be imposed on these leptons. The efficiency for reconstructing a J/Ψ at this stage is (0.386 ± 0.007) and (0.114 ± 0.004) for the muon and electron channels, respectively. It is limited by the low momenta of the leptons and because they are produced inside a jet, making the reconstruction more difficult, particularly for electrons.

If a J/Ψ is found in an event, the isolated lepton with the highest p_T and higher than 20 GeV/c is considered as the lepton candidate from the W decay. The isolation discriminant is defined as the sum of the energies in the electromagnetic and hadronic calorimeters in a cone of opening angle $\Delta R = 0.3$ around the lepton candidate. The selection requires that the isolation energy is less than 15 GeV for electrons and less than 20 GeV for muons.

We define as background all contributions from processes not resulting in the decay chain $t \rightarrow Wb \rightarrow \ell\nu J/\Psi X$. We call physics background the contribution from processes other than $t\bar{t}$ (semi)leptonic and as combinatorial background the irreducible part of $t\bar{t}$ (semi)leptonic where the J/Ψ is wrongly associated to the lepton not coming from the W in the same top decay. Any physics background needs to mimic a final state with the presence of a J/Ψ and an isolated and energetic lepton. The obvious candidates are bosons in association with jets. It is important to distinguish between b jets and light jets, which produce J/Ψ at very

different rates, suppressing the contribution of processes with light jets very much. To remove these contributions the total scalar sum of the transverse jet momenta is required to be greater than $100 \text{ GeV}/c$. This cut is not applied if two isolated leptons are found, in order to preserve di-leptonic $t\bar{t}$ events. If the flavour of the two leptons is the same, an explicit cut to remove the presence of leptonic Z is made, vetoing events where the invariant mass of the two leptons is between 85 and $97 \text{ GeV}/c^2$. To further reduce soft background the cut on the transverse momentum of the isolated lepton is brought to $40 \text{ GeV}/c$, making the analysis less sensitive also to systematic effects involving soft QCD. Table 8.13 presents, in terms of predicted cross sections, efficiencies and events yields per 10 fb^{-1} , the performance of the analysis.

8.2.4.3 Reconstruction of $m_{J/\Psi\ell}$ and statistical performance

In order to estimate the correct invariant mass J/Ψ -lepton it would be necessary to efficiently discriminate between right pairings, where both particles are coming from the decay of the same top, and from wrong pairings where, in $t\bar{t}$ events, they come from the two different top decays. In the present analysis, in order to increase the available statistics, we propose not to attempt any separation of the combinatorial but, instead, to use the full distribution containing both signal and background.

Figure 8.9 shows the three-lepton invariant mass in $t\bar{t}$ events at generator level without selection and at full reconstruction after the selection described in the previous section. The distribution of the components of signal and background from $t\bar{t}$ are shown, where the Monte Carlo truth is used to judge when the correct pairing is made. No equivalent distribution can be done for non- $t\bar{t}$ backgrounds since no J/Ψ is present in those samples. To take this into account the pure background shape is scaled up according to the extra contribution of non $t\bar{t}$ background (Table 8.13), in the hypothesis that the shape of the two samples are the

Table 8.13: Selection performance on signal and expected backgrounds. The first column indicates the channel and its final state, the second the predicted cross section, where the branching ratio for producing at least a J/Ψ into leptons from either a b jet or a light jet is accounted for, the third the trigger efficiency, the fourth the selection efficiency, the fifth the expected number of events in 10 fb^{-1} , the sixth the classification of the contribution as signal (S), physics background (B) or combinatorial background (C).

Channel	BR· σ (fb)	ϵ_{trig} (%)	ϵ_{sel} (%)	Events in 10 fb^{-1}	Class
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)\ell\nu - b\ell\nu$	107	93.9	15.7 ± 0.4	158	S+C
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)\ell\nu - b\tau\nu$	53	61.1	11.0 ± 0.8	36	S
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)\ell\nu - bq\bar{q}$	320	55.3	10.9 ± 0.3	193	S
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)\tau\nu - b\ell\nu$	53	61.1	10.6 ± 0.8	34	C
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)\tau\nu - b\tau\nu$	27	14.2	2.8 ± 1.2	1	B
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)\tau\nu - bq\bar{q}$	160	7.9	1.5 ± 0.5	2	B
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)q\bar{q} - b\ell\nu$	320	55.3	10.7 ± 0.3	190	C
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)q\bar{q} - b\tau\nu$	160	7.9	1.5 ± 0.5	2	B
$t\bar{t} \rightarrow (b \rightarrow J/\Psi)q\bar{q} - bq\bar{q}$	959	0.1	0.2 ± 0.5	0	B
$W + N \text{ jets}, N > 1 \rightarrow J/\Psi X$	394	55.3	2.1 ± 0.1	43	B
$Wb\bar{b} + \text{jets} \rightarrow J/\Psi X$	196	55.3	1.6 ± 0.1	16	B
$Zb\bar{b} + \text{jets} \rightarrow J/\Psi X$	23	93.9	9.4 ± 0.1	20	B
$b\bar{b} \rightarrow J/\Psi X$	$1.3 \cdot 10^9$	$< 2 \cdot 10^{-8}$	< 1	< 2.6	B




Figure 8.9: Three-lepton mass distribution for $m_t = 175 \text{ GeV}/c^2$ at generator level (left) and after detector simulation and reconstruction (right). In the pictures the components coming from correct and wrong lepton pairing - from both combinatorial and physics backgrounds - are shown.

same. Uncertainty in the background description will then be translated into a systematic contribution on the measurement.

The observable most sensitive to the top mass is the position of the maximum of the three-lepton mass distribution. It is determined via a fit of the full shape with a polynomial function of fourth degree. The range chosen for the fit is centered around the maximum and goes from 20 to 120 GeV/c^2 . The error on the maximum of the fitted polynomial is determined by propagating the errors on the fitted coefficients and taking into account their correlation. As a cross check, an alternative way of fitting the signal with a gaussian was tried. In this case the background is first subtracted on a bin-by-bin basis making use of an average background distribution determined by using all the simulated samples. The results obtained are comparable.

The fitted maxima are expected to be correlated to the input value of the top mass. This correlation is proven and fitted by a line (Figure 8.10). The two results at fast and full simulation are in impressive agreement. The correlation curves can be used to estimate the expected statistical error on the top mass as a function of the available amount of data. This is done by using the number of events expected according to Table 8.13, and the result is presented in Figure 8.10. From the figure it can be concluded that the measurement of the top quark mass with this analysis can become, on the statistical footing, competitive already with other analyses' total error after the first years of data taking. Moreover the measurement is expected to be dominated by systematic errors in the long range, as explained in the next section.

8.2.4.4 Systematic errors

The sources of systematic errors can be divided into two main categories: theoretical and experimental. The former include the description of the hard process and the modelling of




Figure 8.10: Left: correlation between the reconstructed three-lepton invariant mass and the input top mass at full simulation. Right: expected statistical error on the top quark mass as a function of the integrated luminosity.

radiation, fragmentation and the underlying event in the simulation, whereas the latter includes all experimental sources coming from an imperfect detector description. The sources analyzed in what follows are considered as uncorrelated and the corresponding resulting errors on the top mass are summed in quadrature to form the total systematic error. To evaluate the effect of various sources the guidelines described in [192] and in Appendix B are followed.

With the exception of the PDF description, for each of the other sources of theoretical uncertainty and for each change in the simulation parameters an independent signal generation with TOPREX and PYTHIA has been performed, with statistics of a few 100K events each, and fast simulated. The variations on the resulting top masses are considered as systematics: when the mass difference with respect to the reference sample is smaller than the associated statistical error, this is conservatively quoted as the systematic error.

For all the experimental sources, smearings and shifts on the observed objects (leptons and jets) are applied after reconstruction and before selection in a consistent way. The observed difference on the top mass is taken as an estimation of the associated systematic uncertainty.

Table 8.14 presents the systematic breakdown on the top mass. The systematics error is dominated by theoretical sources, which are the ones affected by the larger statistical uncertainties, quoted here as systematics.

Putting together the systematic and the statistical error one can conclude that, with maybe exception for the first year of data taking, this measurement will be dominated by systematics, in turn dominated by our poor understanding of the theoretical sources. A total error on the top mass below $2 \text{ GeV}/c^2$ can be in reach from the first 20 fb^{-1} already. The present result suggests an uncertainty of $1.5 \text{ GeV}/c^2$ with full statistics, but this number is fully dominated by the theory systematics. A precision much better than this is not out of reach

Table 8.14: Systematic error breakdown. For each source either the maximum variation from a reference sample or the resulting statistical error on the difference is quoted as a systematic error.

Source	$\delta m_t (\text{GeV}/c^2)$
Λ_{QCD}	0.31
Q^2	0.56
Scale definition	0.71
b-quark fragmentation	0.51
Light jet fragmentation	0.46
Minimum bias/Underlying event	0.64
Proton PDF	0.28
Total theoretical	1.37
Electron E scale	0.21
Muon p scale	0.38
Electron E resolution	0.19
Muon p resolution	0.12
Jet E scale	0.05
Jet E resolution	0.05
Background knowledge	0.21
Total experimental	0.54
Total systematic	1.47

since, by the time this measurement will be made, the analysis will be hopefully repeated at (N)NLO and our understanding of the dominating systematics, for instance the minimum bias and the underlying event, will be drastically improved. Moreover, the results presented reflect the current status of the simulation and reconstruction tools in CMS. More dedicated reconstruction techniques and more sophisticated analyses will considerably improve the statistical treatment of the information.

This analysis reduces to a minimum those systematics which are expected to dominate in more traditional estimations of the top mass, especially the ones from direct reconstruction. Examples are given by the jet energy scale and the knowledge of the b-tagging both below $0.05 \text{ GeV}/c^2$.



5 Summary of top mass determinations

Measuring the mass of the top quark in different channels allows for a combination of the individual results [288]. As the statistical component in the total uncertainty on m_t in each channel is negligible, the correlation between the systematic uncertainties must be determined. The dominant uncertainty arises from the knowledge of the energy scale of b -quark jets, a component which is assumed to be fully correlated between decay channels. This uncertainty can however be subdivided in several components: detector understanding, clustering algorithms, related to the modelling of b - and light-quark fragmentation and decay and, finally, the statistical precision of the data-based estimates of the b -jet energy scale differentiated versus the pseudo-rapidity and the transverse momentum of the observed jet.

The measurement from the J/Ψ final states is however limited by other, mainly theoretical, sources of systematic uncertainties. Therefore a reduction of the uncertainty on m_t is ex-

pected when combining the direct measurements with the measurement from the J/ψ final states. The knowledge of the top quark mass can be improved by developing alternative methods which do not rely on the b -jet energy scale [289, 290]. Accounting for these future improvements an uncertainty of $1 \text{ GeV}/c^2$ on the top quark mass is feasible. The combination can be performed by applying techniques described in [291, 292].



Spin Correlation in Top-Quark Pair Production

8.3.1 Introduction

Because of its large width of $1.4 \text{ GeV}/c^2$ the top quark decays before either hadronization, governed by the scale Λ_{QCD} , or depolarization, governed by the scale Λ_{QCD}^2/m_t , can take place. This unique feature is used to investigate the spin of the top quark; such investigation is not possible in the case of light quarks, where the spin information is diluted by hadronization. Moreover, the top quark spin-flip time is much larger than its lifetime and the probability of a spin flip due to emission of one or several gluons via chromomagnetic dipole-transition is very small.

The angular distribution of a daughter particle in top quark decays can be written as [293, 294, 295]

$$\frac{1}{\Gamma} \frac{d\Gamma}{d \cos \theta_i} = \frac{1}{2} (1 + \kappa_i \cos \theta_i), \quad (8.4)$$

where the decay angle θ_i is defined as the angle between the direction of motion of the daughter particle i and the chosen spin quantisation axis. As gluon fusion is the dominant production mechanism at the LHC there is no well defined spin axis in the initial state. This leads to a choice of the helicity basis along the the top quark momenta in the in the partonic center-of-mass frame. The spin-analyzer quality κ_i of the top quark daughter particle is defined as the degree to which the daughter particle is correlated with the top-quark spin. The analysis presented here is based on the semi-leptonic $t\bar{t}$ decay channel with electrons or muons, which is considered to be the signal. Alternatively, the di-leptonic $t\bar{t}$ decay channel can also be considered. The κ values for the daughter particles used in this analysis [296], lepton, b quark and the lower energy quark from W decay, are 1, -0.41 and 0.51 , respectively.

The spin correlation in the semi-leptonic $t\bar{t}$ decay channel can be measured in terms of a double differential lepton and quark angular distribution, which, neglecting higher order QCD corrections, is given by

$$\frac{1}{N} \frac{d^2 N}{d \cos \theta_l d \cos \theta_q} = \frac{1}{4} (1 - \mathcal{A} \kappa_l \kappa_q \cos \theta_l \cos \theta_q). \quad (8.5)$$

Here, using the helicity basis the lepton and quark angles θ_l and θ_q are obtained by measuring the angle between the decay particle momentum in its parent top quark rest frame and the parent top quark momentum in the $t\bar{t}$ quark pair rest frame. The correlation coefficient

$$\mathcal{A} = \frac{N_{||} - N_X}{N_{||} + N_X} = \frac{N(t_L \bar{t}_L + t_R \bar{t}_R) - N(t_L \bar{t}_R + t_R \bar{t}_L)}{N(t_L \bar{t}_L + t_R \bar{t}_R) + N(t_L \bar{t}_R + t_R \bar{t}_L)}, \quad (8.6)$$

where $N_{||}$ and N_X give the number of events with parallel and anti-parallel top quark spins, respectively. Two angle combinations are considered: θ_l versus θ_b and θ_l versus $\theta_{q(\text{lower energy})}$; in the following description these two combinations are denoted as $b - t l - t$ and $q - t l - t$.

8.3.2 Simulation of $t\bar{t}$ With Spin Correlation

A $t\bar{t}$ sample of $3.1 \cdot 10^6$ containing $9.1 \cdot 10^5$ semi-leptonic signal events was generated with PYTHIA [30] and reconstructed using ORCA. As PYTHIA does not include spin correlations the events are weighted according to Formula 8.5 with $\mathcal{A} = 0.32$ [35] and appropriate values of κ . Then, this data sample is subdivided into two subsamples: one is regarded as the “reference” subsample (1.61M events), used for determination of the selection efficiency and backgrounds. The other is regarded as the “analysis” subsample (1.50M events), used for the measurement of \mathcal{A} . This sample provides 436K “signal” events. The double differential angular distributions obtained from the “analysis” sample are presented in Figure 8.11.




Figure 8.11: Double differential angular distributions obtained from the “analysis” sample, see text.

The distributions in Figure 8.11 are fitted according to the formula (8.5). The results are $\mathcal{A}_{b-tl-t} = 0.321 \pm 0.011$ (stat.) and $\mathcal{A}_{q-tl-t} = 0.319 \pm 0.009$ (stat.) which are statistically compatible with the input value of $\mathcal{A} = 0.32$.

8.3.3 Online and Offline Event Selection

The Level 1 and High Level triggers select events with a single isolated electron or muon; the overall trigger efficiency is 55%.

The following requirements are applied in the offline selection: missing transverse energy $E_T^{\text{miss}} > 20 \text{ GeV}$; at least one isolated lepton with $|\eta| < 2.5$, electron with $p_T > 27 \text{ GeV}/c$ or muon with $p_T > 20 \text{ GeV}/c$; at least four jets with $p_T > 30 \text{ GeV}/c$ and $|\eta| < 2.5$. Jets are reconstructed with a cone algorithm with $\Delta R = 0.5$ and at least two jets must be b -tag jets with a discriminant value larger than zero. This selection results in an overall efficiency of 12%.

The reconstruction of two top quarks includes the following requirements: Two jets that are not b -tagged and have an invariant mass in the range $50 - 135 \text{ GeV}/c^2$, consistent with the W mass, are found. A b -tag jet which combined with the above reconstructed W gives an invariant mass in the range $130 - 250 \text{ GeV}/c^2$, consistent with the t mass. In addition to the top quark reconstructed above, another top quark is required based on the other b -tag jet plus

lepton and missing energy combination. The neutrino components are determined by fitting the missing energy components, constrained with W and t quark masses. The azimuthal angle between the two top quarks is required to be greater than 2 rad. This selection results in an overall efficiency of 5% (Table 8.15).

A measure of the selection quality can be obtained by comparing the generated and reconstructed momentum directions expressed in terms of the cosine of the angles defined above. Figure 8.12 presents the differences between the generated and reconstructed cosines of the $b-ll-t$ and $q-ll-t$ systems. Quantifying this selection quality Q as the ratio of the number of events in the four central bins to all bins, one obtains: $Q_{b-tl-t} = 52\%$ and $Q_{q-tl-t} = 45\%$.




Figure 8.12: Selection quality: Difference between the generated and reconstructed cosine of the analysis angles in the $b-ll-t$ and $q-ll-t$ systems

The signal-to-background ratio is 4.5. The main background, detailed in Table 8.15, is $t\bar{t}$ production with decays different from those treated as the signal. It amounts to 88% of the total background and is used to model the shape of the total background.

Table 8.15: The physics processes considered for signal and background. The number of selected events for the non- $t\bar{t}$ processes are scaled to the same $t\bar{t}$ sample luminosity.

Process	Simulated events	$\sigma(\text{pb})$	Efficiency	Selected events
$t\bar{t}$ (signal)	436K	246	$5.0 \cdot 10^{-2}$	21589
$t\bar{t}$ (background)	1.07M	584	$4.0 \cdot 10^{-3}$	4236
$WW + \text{jets}$	310K	188	$4.5 \cdot 10^{-5}$	15
$W + \text{jets}$ ($\hat{p}_T = 20 - 400 \text{ GeV}/c$)	2.06M	43K	$3.4 \cdot 10^{-6}$	260
Wbt semileptonic decay	328K	63.1	$1.3 \cdot 10^{-3}$	144

8.3.4 Estimation of Correlation Coefficient

In order to correct for the selection efficiency, an efficiency (6×6) matrix is determined by taking the ratio of the reconstructed double differential angular distribution to the generated one, using the “reference” sample. The final double differential angular distribution is

obtained by subtracting, bin-by-bin, the background obtained from the “reference” sample from the total sample of signal plus background obtained from the “analysis” sample. The resulting distributions are corrected for the selection efficiency, Figure 8.13, and fitted using formula 8.5.




Figure 8.13: Background-subtracted and efficiency-corrected double-differential distribution of the cosine of the analysis angles in the $b - ll - t$ and $q - ll - t$ systems.

The correlation coefficients obtained from the fit are:

$$\begin{aligned} \mathcal{A}_{b-tl-t} &= 0.375 \pm 0.100 \text{ (stat.)} , \\ \mathcal{A}_{q-tl-t} &= 0.346 \pm 0.079 \text{ (stat.)} . \end{aligned}$$

These results agree, within statistical uncertainties, with those obtained from the generated events of Figure 8.11.

in \mathcal{A}_{q-tl-t} -12%.

The following sources of systematic uncertainties have been evaluated. The choice of the Parton Density Function in modeling $t\bar{t}$ production affects the number of $t\bar{t}$ events produced via gluon fusion and that via quark-antiquark annihilation. The variation in \mathcal{A} , determined using TOPREX with different PDFs (CTEQ6M, MRST2003), is found to be 4%. The mass of the top quark has an impact on the result of the kinematic fit and the selection. The nominal $m_t = 175 \text{ GeV}/c^2$ is varied by $\pm 5 \text{ GeV}/c^2$ [45] using TOPREX. The variation in \mathcal{A} is found to be comparable with statistical uncertainties of the fit.

The uncertainty on the $t\bar{t}$ cross section affects the shape of the final angular distribution after background subtraction; varying $\sigma(t\bar{t})$ by 10% results in 1% variation in correlation coefficients. The Uncertainty due to b -tagging efficiency is evaluated by varying the b -identification discriminant cut. The variation in \mathcal{A}_{b-tl-t} is found to be of -20%, and in \mathcal{A}_{q-tl-t} is +6.5%/-8.3%. The jet energy scale uncertainty is evaluated by varying the jet P_T . The variation in \mathcal{A}_{b-tl-t} is found to be +14%, and in \mathcal{A}_{q-tl-t} -12%.

Uncertainties in the initial and final state radiation, quark fragmentation, underlying event

and pile up rate could result in an underestimation of the number of non- $t\bar{t}$ jets (not originating from top decays). This possible underestimation of jet multiplicity is estimated to be 8%. To estimate the corresponding uncertainty, 10% additional jets per event are generated while processing the data sample. These jets are simulated randomly according to the η and p_T distributions of non- $t\bar{t}$ jets, obtained from the $t\bar{t}$ Monte Carlo. The variations in \mathcal{A}_{b-tl-t} and \mathcal{A}_{q-tl-t} are found to be -6.3% and -5.3%, respectively.

Summing up the systematic uncertainties and using the statistical uncertainties estimated for 10 fb^{-1} of integrated luminosity, the results are:


$$\begin{aligned}\mathcal{A}_{b-tl-t} &= 0.375 \pm 0.027 (\text{stat.})_{-0.084}^{+0.055} (\text{syst.}), \\ \mathcal{A}_{q-tl-t} &= 0.346 \pm 0.021 (\text{stat.})_{-0.055}^{+0.026} (\text{syst.}).\end{aligned}$$

In summary, the correlation coefficient of top quark spins in $t\bar{t}$ production is measured with a total uncertainty (dominated by systematic uncertainties) of 28% for \mathcal{A}_{b-tl-t} and of 18% for \mathcal{A}_{q-tl-t} .

Single top quark production

8.4.1 Introduction

The single top production cross section at the LHC is known to the NLO level for the tree production mechanisms (see Fig. 8.14, which are classified by the virtuality of the W-boson involved as: t -channel ($q_W^2 < 0$), s -channel ($q_W^2 > 0$), and associated tW production ($q_W^2 = M_W^2$) [297, 298, 299]. In all cases, the most dangerous background comes from $t\bar{t}$ process. Other dangerous background is multi-jet QCD events, but such background is reduced substantially by considering only leptonic decays of the W^\pm -bosons from top-quark decays.

 results presented in this Section were done for $10/\text{fb}$ of integrated luminosity.




Figure 8.14: Feynman diagrams for the three channels of single top production.

8.4.1.1 Details on the signal and background simulation

Two generators, SingleTop [300] (based on the CompHEP package [34]) and TOPREX [35] were used to generate events for all three single top production processes. The background

processes, namely, $Wb\bar{b}$, $Wb\bar{b} + j$, and $W + 2j$ were generated with CompHEP, TOPREX, MadGraph [73], and Alpgen[152] programs as indicated in the Table 8.16. The hard process events containing all needed information were passed to PYTHIA 6.227 [30] for showering, hadronization and decays of unstable particles. The $t\bar{t}$ and $W + \text{jets}$ background events were generated with the same PYTHIA version. All simulations were done with $M_t = 175 \text{ GeV}/c^2$ and $M_b = 4.7 - 4.8 \text{ GeV}/c^2$, proper considerations of the spin correlations, and the finite W -boson and t -quark widths. The list of the signal and background process cross sections as well as generators used are given in the Table 8.16. Both the full simulation chain (OSCAR [14] and ORCA [15]) and a fast simulation (FAMOS [287]) were used.

Table 8.16: Cross section values (including branching ratio and kinematic cuts) and generators for the signal and background processes (here $\ell = e, \mu, \tau$)

Process	$\sigma \times \text{BR}$, pb	generator	Process	$\sigma \times \text{BR}$, pb	generator
t -ch. ($W \rightarrow \mu\nu$)	18 (NLO)	SingleTop	$Wb\bar{b}$ ($W \rightarrow \ell\nu$)	100 (LO)	TOPREX
t -ch. ($W \rightarrow \ell\nu$)	81.7 (NLO)	TOPREX	$Wb\bar{b} + \text{jets}$ ($W \rightarrow \mu$)	32.4 (LO)	MadGraph
s -ch. ($W \rightarrow \ell\nu$)	3.3 (NLO)	TOPREX	$W + 2j$ ($W \rightarrow \mu\nu$)	987 (LO)	CompHEP
tW ($2W \rightarrow \ell\nu$)	6.7 (NLO)	TOPREX	$W + 2j$ ($W \rightarrow \ell\nu$)	2500 (LO)	ALPGEN
tW ($1W \rightarrow \ell\nu$)	33.3 (NLO)	TOPREX	$Z/\gamma^* (\rightarrow \mu^+\mu^-)b\bar{b}$	116 (LO)	CompHEP
$t\bar{t}$ (inclusive)	833 (NLO)	PYTHIA			

8.4.1.2 Reconstruction algorithms and triggers

Muons are reconstructed by using the standard algorithm combining tracker and muon chamber information as described in [301]; tracker and calorimeter isolation cuts are applied as described in [302]. The electrons are reconstructed by the standard algorithm combining tracker and ECAL information, see [303]. The jets are reconstructed by the Iterative Cone algorithm with the cone size of 0.5, see [304]; for the calibration both the Monte Carlo (in the t -channel analysis) and the $\gamma + \text{jets}$ (in the tW - and s -channel) methods are used, see [305]. For b -tagging a probability algorithm based on the impact parameter of the tracks is used, as described in [306].

The **transverse missing energy** is reconstructed as follows:

$$E_T^{\text{miss}} = - \left(\sum \vec{P}_T^\mu + \sum \vec{E}_T^{\text{tower}} + \sum (\vec{E}_{T,\text{jet}}^{\text{calib}}) - \sum (\vec{E}_{T,\text{jet}}^{\text{raw}}) \right) \quad (8.7)$$

where E_T^{tower} is the sum of transverse energy of towers, $E_{T,\text{jet}}^{\text{calib}}$ ($E_{T,\text{jet}}^{\text{raw}}$) is the transverse energy of calibrated (uncalibrated) jets. For the final states with one isolated lepton the neutrino (E_T^{miss}) longitudinal component, $P_{z,\nu}$, is extracted from the quadratic equation:

$$M_W^2 = 2 \left(E_\mu \sqrt{P_{z,\nu}^2 + (E_T^{\text{miss}})^2} - \vec{P}_{T,\mu} \cdot \vec{E}_T^{\text{miss}} - P_{z,\mu} P_{z,\nu} \right) \quad (8.8)$$

This equation has two solutions:

$$P_{z,\nu}^{(1,2)} = \frac{AP_{z,\mu} \pm \sqrt{\Delta}}{P_{T,\mu}^2}, \quad \text{where } A = \frac{M_W^2}{2} + \vec{P}_{T,\mu} \cdot \vec{E}_T^{\text{miss}}, \quad \Delta = E_\mu^2 (A^2 - (E_T^{\text{miss}})^2 P_{T,\mu}^2) \quad (8.9)$$

Among the two solutions of Eq. (8.8) the minimal value of $|P_{z,\nu}|$ is used for W -boson momentum reconstruction.

About 30% of the events have negative Δ values due to the finite detector resolution and to the presence of extra missing energy. In this case for t -channel analysis the parameter M_W in Eq. (8.9) is increased until Δ becomes zero. Using this value of M_W , $P_{z,\nu}$ is calculated from Eq. (8.9). For the tW and s -channels analyses, only the real part of $P_{z,\nu}$ is used for further analysis.

The **transverse mass of the W -boson** is defined as

$$M_T^W = \sqrt{2(P_{T,\mu} E_T^{\text{miss}} - \vec{P}_{T,\mu} \cdot \vec{E}_T^{\text{miss}})}. \quad (8.10)$$

The **sum of the transverse momentum vectors** of all reconstructed objects

$$\vec{\Sigma}_T \equiv \vec{P}_{T,\ell} + E_T^{\text{miss}} + \sum \vec{E}_{T,jet}, \quad (8.11)$$

is found to be very effective for signal/background separation. Note, at the partonic level this variable equals zero for signal events.

The **“jet charge”** (Q_j) is defined as the sum of the charges of the tracks inside the jet cone, weighted over the projections of the track momenta along the jet axis.

The **lepton isolation** criterion used is to sum the p_T of all the tracks in a cone of $\Delta R < 0.2$ around the lepton track, and to reject the event if this sum is greater than 5% of the lepton p_T .

The present study is based on leptonic decay channels ($e\nu_e$ or $\mu\nu_\mu$) of the W -boson. The signal is triggered by the trigger on leptons. The HLT p_T thresholds from the CMS DAQ-TDR [68] are assumed: 19 GeV/ c (29 GeV/ c) for the single muon (electron); with $|\eta_\mu| \leq 2.1$ and $|\eta_e| \leq 2.4$.

8.4.1.3 The contribution from multi-jet backgrounds

A special treatment is required for QCD events with jets, due to the huge cross section. The currently available samples have very small statistics and typically no events remain after the application of pre-selection cuts. Therefore, in order to estimate the impact of the QCD-background the cuts are applied separately, assuming they are uncorrelated.

For t -channel study these cuts are: (a) one isolated muon ($p_T > 19 \text{ GeV}/c$); (b) $E_T^{\text{miss}} > 40 \text{ GeV}$ and only two jets; one B -jet and one light forward jet. It was found a satisfactory suppression of the multi-jet events as compared to other background process ($N_{\text{QCD}}/N_{\text{bckg}} = 6924/(8.9 \times 10^4) = 0.078$ (see [307]) and the QCD-background was not considered in the analysis of the t - and s -channel single top production.

More detailed investigation of this problem was done for tW -channel [308]. The selection cuts are arranged into cut groups whose efficiencies are estimated with the Monte Carlo samples. The product of efficiencies is an indicator of the total efficiency.

Three cut groups are used in the di-leptonic channel: lepton, E_T^{miss} , jet. The same procedure is applied on signal sample to find the ratio of total efficiency to the product of efficiencies. The ratio is used to correct the product of efficiencies found in multi-jet sample and the result is 5.6 events. Four cut groups are used in the semi-leptonic channel: jets, leptons, kinematics and finally signal region and b tagging. The b tagging requirement is taken out from jets group to have reasonable statistics for the efficiency measurement. By comparing

the product of efficiencies with total efficiency of applying cut groups in series, the cut groups are found to be anti-correlated which would result in an over-estimate of the yield. The result of 508 events is kept to be conservative [308].

8.4.1.4 Systematic uncertainties

The following sources of systematic uncertainty are common for all three channels: (i) the **theoretical errors** to the total rates of the signal is $\Delta_{\text{th}} \approx 4\%$. The uncertainties in the background events are assumed to be: 5% for $t\bar{t}$ [36], 17% for $Wb\bar{b}j$, 7% for $W + \text{jets}$, 5% for Wjj [309], 10% for tW and 5% for $Wb\bar{b}$. (ii) the **jet energy scale (JES) uncertainty**: using a calibration method based on $t\bar{t}$ events [310], the JES uncertainty after 10 fb^{-1} integrated luminosity is expected to be $\pm 5\%$ ($\pm 2.5\%$) for jets with $p_T \approx 20 \text{ GeV}/c$ ($p_T > 50 \text{ GeV}/c$). In the region between 20 and 50 GeV/c a linear dependence is assumed. (iii) **b-tagging identification uncertainty**: of $\pm 4\%$ on the overall selection efficiencies is expected on the b -tagging efficiencies [148]. (iv) the **luminosity uncertainty**, expected to be 5% [311].



8.4.2 Selection and cross section - t -channel

The final state in t -channel includes one isolated muon, missing energy (neutrino), one or two jets from b -quark (B_{jet}), and one “forward” hadronic jet. A specific feature of single top events is production of a light jet in the forward/backward direction (see Figs. 8.15) providing an additional possibility for background suppression. The additional b -quark is produced with small transverse momentum, making the reconstruction of the associated low- p_T jet and its b -tagging very difficult. Therefore, in t -channel analysis [307] it is required to have only two hadronic jets in the final state. In this case, the most important background contribution arises from $t\bar{t}$ production and from W^\pm -boson production in association with $\gamma\gamma$ quarks ($Wb\bar{b}+\text{jet}$) or light quark jets ($W + \text{jets}$).




Figure 8.15: The distributions of pseudorapidity (η) of the light jet (left), and of $|\vec{\Sigma}_T|$ (right).

8.4.2.1 Analysis of the fully simulated events

The selection requires the presence of only one isolated muon with $p_T > 19 \text{ GeV}/c$ and $|\eta_\mu| < 2.1$ (HLT selection). Then, it is required: (i) $E_T^{\text{miss}} > 40 \text{ GeV}$; and (ii) at least two

hadronic uncalibrated jets, with $p_T > 20 \text{ GeV}/c$. For further analysis the following additional requirements are: at least one of the selected jets should have the b -tag; the second (light) jet should be in the forward region; only two jets (calibrated) with $p_T^{\text{calib}} \geq 35 \text{ GeV}$ and no other hadronic jets with $p_T^{\text{calib}} \geq 35 \text{ GeV}/c$ (jet veto). The GARCON program [55] is used for the final optimizations of the cuts. The signal-over-background ratio times significance is chosen as an optimization criterion. Finally, the optimal cut values found are:

- muon: $p_T(\mu) > 19.0 \text{ GeV}/c$ and $|\eta(\mu)| < 2.1$ and $E_T^{\text{miss}} > 40.0 \text{ GeV}$;
- b-jet: $p_T > 35.0 \text{ GeV}/c$, $|\eta| < 2.5$ and Discriminator > 2.4 ;
- the light forward: $p_T > 40.0 \text{ GeV}/c$ and $|\eta| > 2.5$;
- $|\vec{\Sigma}_T|$ cut window: $(0.0, 43.5) \text{ GeV}$; $50 < M_T^W < 120 \text{ GeV}/c^2$
- the reconstructed top mass window: $110 \text{ GeV}/c^2 < M_{\text{rec}}(bW) < 210 \text{ GeV}/c^2$

Table 8.17: Number of events (t -channel) and cumulative efficiencies for each cut used in the analysis of t -channel single top production. The symbol “ $p_{TB} \times p_{Tj} \times E_T^{\text{miss}}$ ” means: $p_{TB} > 35 \text{ GeV}/c$, $p_{Tj} > 40 \text{ GeV}/c$, $|\eta_j| > 2.5$, $E_T^{\text{miss}} > 40 \text{ GeV}$.

	signal	$t\bar{t}$	$Wb\bar{b}j$	Wj	Wjj
N(events) at 10 fb^{-1}	1.8×10^5	8.33×10^6	3.24×10^5	9.7×10^7	9.9×10^5
isolated muon	0.73	0.14	0.52	0.16	0.81
$p_{TB} \times p_{Tj} \times E_T^{\text{miss}}$	0.036	6.4×10^{-3}	3.4×10^{-3}	9×10^{-6}	3×10^{-3}
veto on 3 rd jet	0.021	5.8×10^{-4}	1.6×10^{-3}	4×10^{-6}	1.1×10^{-3}
$0.0 < \Sigma_T < 43.5 \text{ GeV}$	0.018	4.1×10^{-4}	1.2×10^{-3}	4×10^{-6}	6.8×10^{-4}
$50 < M_T^{W*} < 120$	0.015	2.2×10^{-4}	9.6×10^{-4}	1×10^{-6}	5.4×10^{-4}
$110 < M_{\text{rec}}(bW)^* < 210$	0.013	1.4×10^{-4}	5.8×10^{-4}	0	4.1×10^{-4}
Number of events	2389	1188	195	0	402

* in GeV/c^2

The efficiencies of these cuts and the resulting number of events are given in the Table 8.17. The resulting signal-to-background ratio and the significance are: $N_S/N_B = 1.34$ and $S_{\text{stat}} = N_S/\sqrt{N_S + N_B} = 37.0$. The final distribution of the reconstructed top mass is shown in Fig. 8.16. The cuts provide a satisfactory background suppression.

Table 8.18: Number of selected events (t -channel) at 10 fb^{-1} with uncertainties due to different sources. ΔN_{syst} represents the theoretical, JES and b -tagging uncertainties. ΔN_{stat} is expected statistical uncertainty.

sample	selected	ΔN_{th}	JES	$\Delta N_{b\text{-tag}}$	ΔN_{syst}	ΔN_{stat}
t -channel	2389	96	71	96	153	49
$t\bar{t}$	1188	59	73	48	105	34
$Wb\bar{b}j$	195	33	6	8	35	14
Wjj	402	20	0	16	26	20

The systematic uncertainties (see Section 8.4.1.4) evaluated for 10 fb^{-1} are given in Table 8.18. In summary, the statistical error is 2.7%, the total systematic error excluding the 5% luminosity uncertainty is 8%, resulting in a total error of 10%.




Figure 8.16: The distribution on the reconstructed top mass, for signal only (left) and with background included (right).

8.4.3 Selection and cross section - tW -channel

The $pp \rightarrow tW$ process contains two W -bosons and a b -quark in the final state. In this study only leptonic decays of the W 's are considered. The nominal final states are $\ell^+ \ell^- E_T^{\text{miss}} b$ and $\ell^\pm E_T^{\text{miss}} b j j$ for the di-leptonic and semi-leptonic modes, respectively. The dominant background arises from $t\bar{t}$ production. Other backgrounds are t - and s -channel single top production, $Wb\bar{b}$, $W + \text{jets}$, $WW + \text{jets}$, and to a lesser extent QCD multi-jet background.

8.4.3.1 Jet quality requirements and extra jet reduction

The most significant difference between tW events and $t\bar{t}$ events is the number of jets in the final state. However, most of the time there are also additional jets due to the underlying event, pile-up or calorimeter noise. These "extra jets" were identified and excluded from the counting by consideration of five jet quality variables (see [308]). It was found that the most discriminating variables are E_T^{max} (the maximum tower E_T in a cone of 0.5) and N_{track} (the number of associated tracks). A Fisher discriminant [312] (F) is constructed from the jet quality variables to separate real jets from extra jets. Each jet is classified value F into one of three categories: good ($F < -0.5$), loose ($|F| < 0.5$) and bad ($F \geq 0.5$) jets. This method yields 84.3% efficiency on true jets and rejects 86.9% of extra jets. Only "good" jets and "loose" jets are used in preselection and event reconstruction. The jet multiplicity after the extra jet reduction in semi-leptonic channels reveals that the number of good jets peaks at the 2 and 3 jet bins for signal events, and at the 3 and 4 jet bins for $t\bar{t}$ backgrounds.

8.4.3.2 Event selection and reconstruction

The kinematic cuts used for this study are presented in Table 8.19 and Table 8.20. For the semi-leptonic channel, two non- b -like jets with $m_{jj} < 115 \text{ GeV}/c^2$ are used for reconstruction of the W -boson (that decays hadronically). In events with a 4th jet that survives jet veto cuts, it is required that the invariant mass of the 4th jet with any of the selected non- b -like jets must be outside a window of $M_W \pm 20 \text{ GeV}/c^2$. For the leptonic decays of the W -boson it is required that $M_T^W < 120 \text{ GeV}/c^2$.

Table 8.19: Kinematic cuts used in the di-leptonic channel. The final electron and muon should have the opposite charges.

Leptons	Jets
$ \eta(e) < 2.4, \eta(\mu) < 2.1$ $p_T(e, \mu) > 20 \text{ GeV}/c$ no other lepton with $p_T > 5 \text{ GeV}/c$	leading jet: $ \eta < 2.4, p_T > 60 \text{ GeV}/c, \text{disc} > 0$ at most one extra jet No other jets with $p_T > 20 \text{ GeV}/c$
Missing $E_T: E_T^{\text{miss}} > 20 \text{ GeV}$	

Table 8.20: Kinematic cuts used in the semi-leptonic channel. The presence of a good fourth jet would veto the whole event.

Leptons
$p_T(e) > 30 \text{ GeV}/c, p_T(\mu) > 20 \text{ GeV}/c, \eta(e) < 2.4, \eta(\mu) < 2.1$ no other lepton $p_T > 10 \text{ GeV}/c$
Jets (after removing all bad quality jets)
b -like jet: good quality, $\text{disc} > 2, \eta < 2.5, p_T > 35 \text{ GeV}/c$ non- b -like jet: good quality, $ \eta < 3.0, \text{disc} < 0$ if $ \eta < 2.5, p_T > 35 \text{ GeV}/c$ Jet counting: one b -like jet and 2 non- b -like jets Jet veto: no other “good” or “loose” jets with $p_T > 20 \text{ GeV}/c$ and $ \eta < 3$
Missing $E_T: E_T^{\text{miss}} > 40 \text{ GeV}$

To find the correct pairing of b -jet and reconstructed W -boson (coming from top decay) the following variables were used: the p_T of (b, W) systems; the separation of the b -jet with each of the W in (η, ϕ) space; the “charges” of jets (see Section 8.4.1.2) and W -bosons (see Ref. [308] for details). A Fisher discriminant based on these variables is used for discriminating leptonic top events from hadronic top events. A cut of 0.56 is optimal in separating these 2 types of events, and 72% of the events are correctly paired.

To further enhance the signal to background ratio the following “global” cuts are applied:

- p_T of the reconstructed tW system: $|\vec{\Sigma}(t + W)| < 60 \text{ GeV}/c$.
- Scalar sum of transverse energies $H_T: H_T < 850 \text{ GeV}$.
- Reconstructed top quark mass: $110 \text{ GeV}/c^2 < m(t) < 230 \text{ GeV}/c^2$.
- p_T of the reconstructed top quark: $20 \text{ GeV}/c < p_T(t) < 200 \text{ GeV}/c$.

8.4.3.3 Efficiencies and expected yields

The efficiencies estimated with Monte Carlo samples are converted to the effective cross sections by multiplying the production cross sections of each process. The effective cross sections, as well as the expected yields with 10 fb^{-1} of data for all signal and background samples, are shown in Table 8.21 and 8.22. The signal to background ratio is found to be 0.37 for di-leptonic channel and 0.18 for semi-leptonic channel.

Table 8.21: Summary of cross section times branching ratio times efficiencies at each stage of the analysis for the di-leptonic channel. All values are in picobarns. The last row is the expected number of events for 10 fb^{-1} . Multi-jet background has been estimated separately (see Section 8.4.1.3). When only a limit on the number of events is stated, this is due to MC statistics.

	tW dil.	$t\bar{t}$ dil.	$t\bar{t}$ oth.	WW dil.	WW oth.	t ch. lept.
Production	6.667	92.222	737.778	11.111	88.889	81.667
HLT	4.865	74.090	346.151	7.674	27.259	41.409
2ℓ	1.944	25.150	21.012	2.574	0.226	2.309
Lepton p_T	0.675	7.919	0.703	0.543	0.012	0.098
≤ 1 extra jet	0.459	6.574	0.664	0.416	0.010	0.067
Jet p_T, η	0.307	5.234	0.556	0.339	0.004	0.033
≥ 1 b -jet	0.184	3.864	0.379	0.017	0.000	0.018
$E_T^{\text{miss}} > 20$	0.170	3.640	0.349	0.017	0.000	0.016
≤ 2 jet	0.150	2.734	0.221	0.015	0.000	0.012
Final select.	0.057	0.145	0.000	0.006	0.000	0.000
Expected events	567	1450	≤ 55	61	≤ 10	≤ 20

Table 8.22: Summary of cross section times branching ratio times efficiencies at each stage of the analysis for the semi-leptonic channel. All values are in picobarns. The last row is the expected number of events for 10 fb^{-1} .

	tW	$t\bar{t}$	t ch.	s ch.	Wbb	W2j	W3j	W4j	Multi-jet
Total cross section	60	833	245	10	300	7500	2166	522	9.73×10^9
HLT	18.9	263.9	39.5	1.52	34.0	1006	300	73	1.86×10^5
Presel. & isolation	9.05	179.4	12.0	0.54	2.15	52	35	12	1325
jet & lepton p_T , jet veto	1.28	18.5	1.31	0.046	0.061	0.60	4.9	1.0	4.23
b -tagging	0.669	6.13	0.476	0.013	0.016	0.10	0.99	0.26	0.85
kinematic cuts	0.223	0.999	0.047	0.002	0.003	0.017	0.101	0.008	0.105
Signal box cuts	0.170	0.771	0.035	0.001	0.001	0.013	0.054	0.008	0.051
Events in 10 fb^{-1}	1699	7709	351	14	10	130	539	80	508

8.4.3.4 The ratio method

The *ratio method* is developed to reduce systematic uncertainties related to the dominant $t\bar{t}$ background. We define a $t\bar{t}$ -rich control region and use ratio of efficiencies to estimate the yield of $t\bar{t}$ in the signal region. The kinematics of tW and $t\bar{t}$ are similar so tW is present in the control region, therefore the ratio of efficiencies for tW is also used. The signal and background yield is determined by the following equation:

$$S = \frac{R_{t\bar{t}}(N_s - N_s^o) - (N_c - N_c^o)}{R_{t\bar{t}} - R_{tW}}, \quad (8.12)$$

$$B = \frac{(N_c - N_c^o) - R_{tW}(N_s - N_s^o)}{R_{t\bar{t}} - R_{tW} + N_s^o}. \quad (8.13)$$

Here R_x is the ratio of efficiencies $R_x = \epsilon_x(\text{control region})/\epsilon_x(\text{signal region})$ for $x = t\bar{t}, tW$; N_s (N_c) is total number of events in the signal (control) region; N_s^o (N_c^o) is the estimated number of non- $t\bar{t}$ background events in the signal (control) region. With S measured with 2 regions and the ratio method, the cross section can be found by $S/\epsilon\mathcal{L}$.

For the ratio method to work it is important to find a control region with similar kinematics except with one more jet. It is expected that systematic uncertainties from PDF, JES and b tagging cancel to a large extend, while the luminosity uncertainty drops out for the $t\bar{t}$ background. The lepton selection and jet quality requirements in the control region is identical to the signal region. The differences are outlined below.

Di-leptonic. A second jet is required with $p_T = 20 - 80$ GeV, $|\eta| < 2.4$ and b -tagged ($\text{disc} > 0$). No other jets with $p_T > 20$ GeV are allowed. The background region is found to be filled by 97.9% di-leptonic $t\bar{t}$, 0.4% other $t\bar{t}$ decays, 1.6% di-leptonic tW , and 0.1% for leptonic t channel single top while WW +jets yield is negligible.

Semi-leptonic. It requires 2 jets with $p_T > 30$, 2 more jets with $p_T > 20$, and no bad jets with $p_T > 20$. It is required that one of the 2 high- p_T jets is b -tagged ($\text{disc} > 2$), and that both low- p_T jets be not tagged ($\text{disc} < 0$). The $b - W$ pairing is done in the same way, with a 72% correct pairing. It is found that the $t\bar{t}$ purity in the control region is 93.9%. The non- $t\bar{t}$ events are mainly composed of W +jets (2.8%), tW (2.0%) and t -channel single top (1.2%). The ratio of efficiencies are found to be $R_{tW} = 0.319$ and $R_{t\bar{t}} = 3.31$.

8.4.3.5 Systematic Uncertainties

- **Theoretical uncertainties** The $t\bar{t}$ cross section does not show up in the ratio method. The effect is 0.8% for t -channel single top and 3.1% for W +jets. It is found to be negligible for other background.

- **Pileup amount** A difference of 30% between normal pileup and no pileup is used as an estimate of the systematic uncertainty, as was done in [192] for the di-leptonic $t\bar{t}$ studies.

- ◊ *Dileptonic mode* The analysis is found to be rather sensitive to the pileup, as the relative shift of the “measured” cross section is +20.4% for no pileup, and -16.2% for double pileup, while is the difference between the check sample and the reference sample 4.6% (which has purely statistical origin). The value of 6.1% is used as the systematic uncertainty.

- ◊ *Semi-leptonic mode* The extracted cross section varies by +35% for no pileup and -63% for double pile-up so a systematic uncertainty of 10.3% is obtained. The results for both channels are shown in table 8.23.

The results from the ratio method were used in the significance calculation. In addition, the uncertainty on the background expectation, evaluated for di-leptonic ($\Delta_B/B = \pm 9.6\%$) and semi-leptonic ($\Delta_B/B = +3.6\% / -4.4\%$), was taken into account. The resulting significance is 4.2 for the di-leptonic channel and 5.1 for the semi-leptonic channel. Combining the two channels gives a total significance of 6.4.



8.4.4 Selection and cross section - s -channel

The present analysis of the s -channel single top production is based on leptonic channels, i.e. the top is identified and reconstructed by its semileptonic decays into $\ell\nu b$ final states, with $\ell = e, \mu$. For this study, a fast simulation of the CMS detector with FAMOS was used, see [307, 308] for details.

Table 8.23: Summary of uncertainties of cross section measurement.

Source	Uncertainty	$\Delta\sigma/\sigma$ (di-lept.)	$\Delta\sigma/\sigma$ (semi-lept.)
Statistical uncertainty	—	8.8%	7.5%
Integrated luminosity	5%	5.4%	7.8%
$t\bar{t}$ cross-section	9%	<i>negligible</i>	<i>negligible</i>
t -channel cross-section	5%	<i>negligible</i>	0.8%
W+jets cross-section	10%	<i>not applicable</i>	3.1%
WW+jets cross-section	10%	1%	<i>not applicable</i>
Jet energy scale	5%-2.5%	19.7 %	9.4%
b tagging efficiency	4% - 5%	8.7 %	3.6%
PDF	1σ	+4%/-6.0%	1.6%
Pileup	30%	6.1 %	10.3%
MC statistics	—	9.9%	15.2%
Total uncertainty		$\pm 23.9\%$ (syst.) $\pm 9.9\%$ (MC)	$\pm 16.8\%$ (syst.) $\pm 15.2\%$ (MC)

The signal events are triggered by the single lepton triggers. Since this production mode suffers from low statistics, one could envisage the introduction of a combined trigger $e \times jet$, with threshold 19 GeV for the electron (in order to make the electronic sample more coherent with the muonic sample) and 45 GeV for the jet. This value has been chosen to be the same as the threshold for the τ -jet in the already existing $e \times \tau - jet$ trigger.

8.4.4.1 Preselection

The preselection criteria are as follows:

- The event has to fire at least one of the previously described triggers (including the proposed $e \times j$).
- The event must contain one isolated lepton (μ or e) with $p_T \geq 19$ GeV/ c and $|\eta| \leq 2.1$ (≤ 2.4) for muons (electrons) and no other lepton above 10 GeV/ c .
- Exactly two uncalibrated jets must have $p_T \geq 30$ GeV/ c and $|\eta| \leq 2.5$ and no other jet has to be present with $p_T \geq 20$ GeV/ c .
- Both jets should have a positive b-tagging discriminator value.
- The event should have $E_T^{\text{miss}} > 30$ GeV.
- The transverse mass of the W -boson M_T^W should be less than 100 GeV/ c^2 .

Details on the effect of the preselection cuts are given in Table 8.24. Note, that as in Section 8.4.2, the multi-jet QCD contribution is neglected.

8.4.4.2 Genetic Algorithm analysis

The following observables have been chosen in order to further discriminate between signal and background after preselection: (i) the jet b -tagging discriminants; (ii) the calibrated jet transverse momenta; (iii) the mass of the reconstructed top; (iv) $|\Sigma(t, \bar{b})|$; (v) the scalar sum of the transverse momenta of all the reconstructed objects. The reconstructed top quark is formed by the reconstructed W and one of the two b -jets, chosen according to the value of

Table 8.24: Efficiencies of the preselection cuts, with respect to the initial number of events. For all process (except of $t\bar{t}$) the final W decays into charged lepton ($\ell = e, \mu, \tau$) and neutrino. “HLT” includes the $1\mu, 1e$ and $e \times j$ triggers. N_{ev} is the number of events surviving these cuts (the uncertainties are only those due to the limited Monte Carlo statistics).

Cut	s -ch.	t -ch.	$t\bar{t}$	$Wb\bar{b}$	Wt ($1 W \rightarrow l\nu$)
“HLT”	$37.5 \pm 0.2\%$	$42.5 \pm 0.1\%$	$30.1 \pm 0.1\%$	$29.4 \pm 0.1\%$	$46.5 \pm 0.1\%$
Isolation	$33.7 \pm 0.2\%$	$39.0 \pm 0.1\%$	$21.7 \pm 0.1\%$	$28.2 \pm 0.1\%$	$42.3 \pm 0.1\%$
E_T^{miss} cut	$27.3 \pm 0.2\%$	$31.9 \pm 0.1\%$	$17.4 \pm 0.1\%$	$22.6 \pm 0.1\%$	$34.4 \pm 0.1\%$
M_T^W cut	$23.2 \pm 0.2\%$	$26.3 \pm 0.1\%$	$13.6 \pm 0.1\%$	$18.4 \pm 0.1\%$	$29.2 \pm 0.1\%$
$N_j \geq 2j$	$11.9 \pm 0.1\%$	$11.5 \pm 0.1\%$	$11.9 \pm 0.1\%$	$0.88 \pm 0.03\%$	$18.5 \pm 0.1\%$
$N_j = 2j$	$8.9 \pm 0.1\%$	$8.2 \pm 0.1\%$	$1.84 \pm 0.04\%$	$0.76 \pm 0.03\%$	$7.09 \pm 0.05\%$
b -tag	$3.07 \pm 0.07\%$	$0.72 \pm 0.02\%$	$0.28 \pm 0.02\%$	$0.14 \pm 0.01\%$	$0.34 \pm 0.01\%$
N_{ev}	1010 ± 10	5880 ± 70	23300 ± 200	1400 ± 35	1150 ± 40

the “jet charge” (Q_j , see Section 8.4.1.2). Since in top decays the W and the original b quark have opposite sign of the charge, the jet with Q_j “most opposite” to the W is used for top reconstruction, leading to a probability of 67% to identify the correct pairing.

The cuts on these variables are optimized by means of the GARCON program [55]. The surviving events after these cuts are shown in cascade in Table 8.25. With this selection, after an integrated luminosity of 10 fb^{-1} one gets: $N_S/N_B \approx 0.13$.

Table 8.25: Final cuts and their efficiencies, with respect to the preselected samples, for the signal and the main backgrounds. For s - and t -channel and $Wb\bar{b}$ samples the final W -boson decays into lepton (e, μ, τ) and neutrino. $t\bar{t}$ samples includes all W -boson decay modes.

Cut	s -channel	t -channel	$t\bar{t}$	$Wb\bar{b}$
$b\text{-tag}(j_1) > 0.4, b\text{-tag}(j_2) > 0.1$	85%	75%	78%	85%
$p_T(j_1) > 50 \text{ GeV}/c, p_T(j_2) > 50 \text{ GeV}/c$	68%	53%	70%	37%
$120 < M(l\nu b) < 220 \text{ GeV}/c^2$	52%	34%	46%	26%
$25 < p_T(l\nu b) < 160 \text{ GeV}/c$	48%	32%	43%	26%
$\Sigma_T < 20 \text{ GeV}/c$	35%	15%	10.6%	12.5%
$H_T < 340 \text{ GeV}/c$	27%	10.7%	5.4%	11.1%
number of surviving events	273 ± 4	630 ± 14	1260 ± 60	155 ± 12

8.4.4.3 Systematic uncertainties

In addition to systematics described in Section 8.4.1.4 the following sources of systematic uncertainty are considered:

- **Top mass.** The variation of m_t within $\pm 2 \text{ GeV}/c^2$ around top mass $m_t = 175 \text{ GeV}/c^2$ leads to the relative systematic error on the selection efficiency $\sigma_{\text{syst}}^{m_t} = 0.5\%$ for the s -channel single top.
- **Parton Distribution Functions.** To extract the dependence on the PDF uncertainty, two different PDF sets were used: CTEQ61 and CTEQ6M [29]. The result is $\sigma_{\text{syst}}^{\text{PDF}} = 0.7\%$.
- **Initial/Final State Radiation modeling.** The model parameters were varied in the ranges $\Lambda_{\text{QCD}} = 0.25 \pm 0.1 \text{ GeV}$ and Q_{max}^2 from 0.25 to $4 \hat{s}$ (see [192]). The extreme values of the efficiencies are taken as systematic error: $\sigma_{\text{syst}}^{\text{rad}} = 0.5\%$.

Table 8.26: Number of selected events after 10 fb^{-1} and systematic uncertainties.

sample	selected	$\Delta\sigma$	JES	b-tag	M_{top}	PDF	ISR/FSR
<i>S</i> : <i>s</i> -channel	273	—	± 3	± 11	± 1.5	± 2	± 1.5
<i>B</i> : <i>t</i> -channel	630	± 25	± 8	± 25	—	—	—
<i>B</i> : $t\bar{t}$	1260	± 63	± 75	± 50	—	—	—
<i>B</i> : $Wb\bar{b}$	155	± 8	± 7	± 6	—	—	—

8.4.4.4 Background normalization

The $t\bar{t}$ events in Table 8.26 are, in 41% of the cases, $t\bar{t} \rightarrow l^+ \nu b l^- \bar{\nu} \bar{b}$ events with a lepton missed, and in the remain cases $t\bar{t} \rightarrow l^+ \nu b q \bar{q}' \bar{b}$ events with two jets missed ($t\bar{t} \rightarrow q \bar{q}' b q \bar{q}' \bar{b}$ events give a negligible contribution). These two categories of events are very differently affected by the Jet Energy Scale variation. In general, any variation going in the direction of more jets gives a better rejection of the $t\bar{t} \rightarrow l^+ \nu b q \bar{q}' \bar{b}$ component with respect to the signal, while the $t\bar{t} \rightarrow l^+ \nu b l^- \bar{\nu} \bar{b}$ events, having two quarks, are affected almost in the same way as the signal.

- $t\bar{t} \rightarrow \ell^\pm + X$ enriched control sample

In this case the difference with respect to Sec. 8.4.4.1 is the request of three jets instead of two and only the muon channel is used. The selection efficiency for $t\bar{t} \rightarrow \ell^\pm$ events is found to be 1.08%. The ratio R_{c1} between the efficiencies in the main sample and in this control sample is $R_{c1} = 0.0149$, whose variations under JES and b-tagging efficiency systematic shifts are $\Delta R_{c1} = \pm 0.0015(JES) \pm 0.0003(b - tag)$.

- $t\bar{t} \rightarrow \ell^+ \ell^- + X$ enriched control sample

This sample is obtained by the same selection as in Sec. 8.4.4.1, but two leptons with different flavours with the opposite sign are required. The selection efficiency for $t\bar{t} \rightarrow 2l$ events is found to be 0.822%. The ratio R_{c2} between the efficiencies in the main sample and in this control sample is $R_{c2} = 0.0681$, whose variations under JES and b-tagging efficiency systematic shifts are $\Delta R_{c2} = \pm 0.0010(JES) \pm 0.0004(b - tag)$.

8.4.4.5 Results

The number of the selected signal (N_S) and background (N_B) events and their estimated uncertainties are listed in Table 8.26. The cross section is extracted as

$$\sigma = \frac{N_{tot} - b^0 - R_{c1}(N_{c1} - b_{c1}^0) - R_{c2}(N_{c2} - b_{c2}^0)}{\epsilon L}, \quad (8.14)$$

where b^0 is the sum of the non-top backgrounds in the main sample, N_{c1} and N_{c2} are the total events selected in the two control regions, and b_{c1}^0 and b_{c2}^0 are their contamination by non-top backgrounds, single top and other $t\bar{t}$ decays. The statistical error is evaluated to be 18%. The total systematic uncertainty is 31%, where the largest contribution arises from the effect of the JES uncertainty, on the $t\bar{t}$ di-lepton background. The use of “Energy Flow” techniques, including the charged tracks information, is expected to significantly reduce this uncertainty. The total error, including also the 5% luminosity uncertainty, is 36%.



8.4.5 Conclusion

Selection strategies have been proposed for all the three single top production modes, and their effectiveness is shown, taking into account the expected statistics after 10 fb^{-1} . All

analyses will be systematics dominated. For the s -channel and tW -associated cases, control samples have been proposed in order to constrain the dominant $t\bar{t}$ background.

The resulting signal-to-background ratio and the significance for the t -channel are: $N_S/N_B = 1.34$ and $S_{stat} = N_S/\sqrt{N_S + N_B} = 37.0$, with a statistical error of 2.7%, and a systematic error excluding the 5% luminosity uncertainty of 8%, resulting in a total error of 10%.

For tW -channel we expect to reach the significance of 4.2 (5.1) for the di-lepton (semi-leptonic) channel, increasing to 6.4 after combining the two channels. The total uncertainty is $\pm 23.9\%$ (syst.) $\pm 9.9\%$ (MC) for di-lepton and $\pm 16.8\%$ (syst.) $\pm 15.2\%$ (MC) for semi-leptonic channels.

The total systematic uncertainty for the s -channel is 31%. The total error, including also the 5% luminosity uncertainty, is 36%.



Search for flavor changing neutral currents in top decays

8.5.1 Introduction

The study of Flavour Changing Neutral Current (FCNC) interactions plays an important role in testing the Standard Model (SM) and probing new physics beyond it. The top quark is regarded to be more sensitive to new physics than other fermions, due to its mass close to the electroweak scale. Owing to the GIM mechanism of the SM, top quark FCNC interactions are absent at tree level and extremely small at loop level.

In recent years a lot of work has been done to explore the top quark FCNC couplings. On the theoretical side, various FCNC top quark decays and top-charm associated production at high energy colliders were extensively studied in the SM [313, 314, 315, 316], the Minimal Supersymmetric Standard Model (MSSM) [317, 318, 319, 320, 321, 322, 323, 324] and other new physics models [325, 313, 326, 327, 328, 329, 330, 331, 332, 333, 334]. In models beyond the SM the top quark FCNC branching fractions may be significantly enhanced. Thus searching for top quark FCNC is a potentially powerful probe of new physics. The CDF and DØ collaborations have reported interesting bounds on the FCNC top quark decays [335, 336, 337]. The SM expectations for such top quark FCNC processes are far below the detectable level but the MSSM can enhance them by several orders of magnitude to make them potentially accessible at future collider experiments [338, 339, 340]. The theoretical branching ratios and the experimental limits are summarized in Table 8.27.

Details of this analysis can be found in [341].

Table 8.27: Theoretical branching ratios of FCNC top quark decays in various models and experimental limits

Decay	SM	two-Higgs	SUSY with \tilde{R}	Exotic Quarks	Exper. Limits(95% CL)
$t \rightarrow gq$	5×10^{-11}	$\sim 10^{-5}$	$\sim 10^{-3}$	$\sim 5 \times 10^{-4}$	< 0.29 (CDF+TH)
$t \rightarrow \gamma q$	5×10^{-13}	$\sim 10^{-7}$	$\sim 10^{-5}$	$\sim 10^{-5}$	< 0.0059 (HERA)
$t \rightarrow Zq$	$\sim 10^{-13}$	$\sim 10^{-6}$	$\sim 10^{-4}$	$\sim 10^{-2}$	< 0.14 (LEP-2)

8.5.2 Signal and background generation

Both the $t \rightarrow \gamma q$ and the $t \rightarrow Z^0 q$ decay channels are investigated, while the channel $t \rightarrow gq$ is not studied because of its very high QCD background. The $t\bar{t}$ signal is generated with TOPREX [35], while PYTHIA [175] is used for modelling of quark and gluon hadronization. The $t\bar{t}$ pair is generated through gluon-gluon and quark-antiquark annihilation, with subsequent SM decay for one top ($t \rightarrow Wb$) and FCNC decay of the other. Only leptonic decay channels of Z, W bosons are studied, where the lepton could be either e or μ . Hadronic decays as well as decays to tau leptons are not considered because of the large QCD background. On generator level both top quarks are produced on-shell, with a mass of $m_t = 175 \text{ GeV}/c^2$, including the effects of spin-state correlations to final decay products ($\gamma q, Z^0 q, Wb$). Both ISR and FSR are simulated with CTEQ5L PDFs. The generated events are passed through the full detector simulation and digitization, taking into account low luminosity pile-up.

Several SM processes contributing as background are studied: $t\bar{t}$ production, single top quark production (t-channel), $ZW + \text{jets}$, $WW + \text{jets}$, $ZZ + \text{jets}$, $W + \text{jets}$, $Z + \text{jets}$, Zbb and QCD multi-jets in various p_T bins.

8.5.3 Selections

The $t \rightarrow \gamma q$ channel is well identified by a high-energy isolated photon accompanying the FCNC top decay. One b-tagged jet and a light jet can be also used to distinguish from the $t\bar{t}$ standard decays. The considered background sources include multi-hadronic QCD processes having an outgoing parton with $50 < \hat{p}_T < 1000 \text{ GeV}/c$, $WW/Z+\text{jets}$, $W+\text{jets}$, $t\bar{t}$ inclusive and EW single top in the t-channel.

For the FCNC $t \rightarrow \gamma q$ channel our main selection cuts are: (a) ‘single electron or single muon’ trigger criteria at L1 and HLT levels; (b) one isolated e^\pm (with $p_T > 30 \text{ GeV}/c$) or μ^\pm (with $p_T > 20 \text{ GeV}/c$), and missing energy $E_T^{\text{miss}} > 25 \text{ GeV}$ that should form a transverse invariant mass $M_T(W) < 120 \text{ GeV}/c^2$; (c) only one jet compatible with b-jet with calibrated $p_T > 40 \text{ GeV}/c$ that in combination with the W candidate should give an invariant mass in the range $[110-220] \text{ GeV}/c^2$; (d) one single isolated photon with $p_T > 50 \text{ GeV}/c$ and one light-jet (not compatible with b-jet) with calibrated $p_T > 50 \text{ GeV}/c$, reconstructing as the FCNC top decay with invariant mass in the range $[150-200] \text{ GeV}/c^2$; (e) the two reconstructed top quarks recoil against each other ($\cos \phi(t\bar{t}) < -0.95$).

The SM backgrounds $t\bar{t}$ and EW single top (t-channel) are the only backgrounds giving some contribution. The total efficiency for the signal selection is calculated to be $\epsilon = 0.021 \pm 0.002$, while 54 ± 7 background events surviving at a luminosity of 10 fb^{-1} . The errors are statistical only. Adopting a factorization method, QCD background is proven to be not dangerous for the analysis. A set of independent cuts (hard jets, isolated hard lepton, isolated hard photon, b-tagging) is applied to both QCD and $t\bar{t}$ background, and efficiencies for single cuts are factorized together. b-tagging efficiency and mistagging are respectively 30% and 0.5%. Number of surviving QCD events for this preselection is found to be 43 and efficiency of last cuts on $t\bar{t}$ amounts to 2.5%. As a consequence, one can naively infer that multi-hadronic events surviving the whole selection will not exceed $42 \times 2.5\% \simeq 1$ under the most conservative assumptions.

For the FCNC $t \rightarrow Z^0 q$ channel our main selection cuts are: (a) ‘double electron or double muon’ trigger criteria at L1 and HLT levels; (b) two isolated e^\pm (each with $p_T > 20 \text{ GeV}/c$) or

μ^\pm (each with $p_T > 10 \text{ GeV}/c$), having an invariant mass $\pm 10 \text{ GeV}/c^2$ around the nominal Z^0 mass; (c) third lepton (e with $p_T > 20 \text{ GeV}/c$ or μ with $p_T > 15 \text{ GeV}/c$), which, in combination with the missing transverse energy ($E_T^{\text{miss}} > 20 \text{ GeV}$) should have a transverse mass less than $120 \text{ GeV}/c^2$; (d) only one jet compatible with b jet with calibrated $p_T > 40 \text{ GeV}/c$; (e) invariant mass of candidate W and b jet in the range $[110-220] \text{ GeV}/c^2$; (f) one light-jet (not compatible with b jet) with calibrated $p_T > 30 \text{ GeV}/c$, that in combination with the Z^0 should reconstruct a FCNC top with a mass in the range $[150-200] \text{ GeV}/c^2$, that recoils toward the SM one ($\cos\phi(t\bar{t}) < 0$).

The SM background $t\bar{t} \rightarrow (\nu lb)(\nu lb)$ is the only background that gives some contribution. The total efficiency for the signal selection is $\epsilon = 0.041 \pm 0.002$, with 1 ± 1 background events surviving the selection at a luminosity of 10 fb^{-1} . The errors are statistical only.

8.5.4 Sensitivity estimation

For the FCNC sensitivity estimation, it is assumed that new physics is observed when the signal significance is 5 at least. When dealing with a small number of background (B) events with respect to signal ones (S), an appropriate definition of significance is [40]:

$$S_{12} = 2 \left(\sqrt{B + S} - \sqrt{B} \right) \quad (8.15)$$

S_{12} evaluates the probability (in number of sigmas) that a background with expected value B fluctuates above observed number of events $S + B$ with Poisson statistics. Equation 8.15 can be used to evaluate the minimum number of signal events S claiming the FCNC discovery for a given level of significance. On the other hand, the number of signal events for the $t \rightarrow Zq$ and $t \rightarrow \gamma q$ can be expressed respectively as:

$$\begin{aligned} S(t \rightarrow Zq) &= 2 \times BR(t \rightarrow Zq) \times Br(W \rightarrow l\nu) \times Br(Z \rightarrow ll) \times \sigma(t\bar{t}) \times L \times \epsilon(t \rightarrow Zq) \\ S(t \rightarrow \gamma q) &= 2 \times BR(t \rightarrow \gamma q) \times Br(W \rightarrow l\nu) \times \sigma(t\bar{t}) \times L \times \epsilon(t \rightarrow \gamma q) \end{aligned} \quad (8.16)$$

where $L = 10 \text{ fb}^{-1}$, $\sigma(t\bar{t}) = 833 \text{ pb}$, $BR(W \rightarrow l\nu) = 0.2136$, $BR(Z \rightarrow ll) = 0.0673$ ($l = e, \mu$), and ϵ is selection efficiency for the signal. From this formula, the smallest observable $BR(t \rightarrow Zq)$ and $BR(t \rightarrow \gamma q)$ can be calculated for any significance level S_{12} .

8.5.4.1 Sensitivity estimation without systematics uncertainties

In the hypothesis of no systematics effects, the sensitivity can be estimated from the previous equations using various quantities given above. In Fig. 8.17 the sensitivity to FCNC is represented as a function of desired significance level: interpolating for $S_{12} = 5$, we obtain $BR(t \rightarrow Zq) = (11.4 \pm 2.6) \times 10^{-4}$ and $BR(t \rightarrow \gamma q) = (5.7 \pm 0.5) \times 10^{-4}$.

The statistical error on S is obtained from a simple propagation of the statistical error on B . In the branching ratio S and ϵ factorize and are uncorrelated, thus the error on BR is given adding quadratically the statistical errors on S and ϵ .

8.5.4.2 Effects of systematic errors

For all experimental sources of systematic errors, shifts and rescaling on the observed objects (leptons, jets and photons) are applied after the reconstruction in a consistent way and




Figure 8.17: Branching Ratios of FCNC signal versus signal significance, for qZ (right) and $q\gamma$ (left) channels, with $L=10 \text{ fb}^{-1}$. The estimated upper limits for the branching ratios for $S_{12} = 5$ are $BR(t \rightarrow Zq) = (11.4 \pm 2.6) \times 10^{-4}$ and $BR(t \rightarrow \gamma q) = (5.7 \pm 0.5) \times 10^{-4}$. Systematics uncertainties are not considered here.

their impact on the selection efficiency (ϵ) and the surviving number of background events (B) evaluated. Fluctuations are considered as the maximum shift from the central value without the systematic bias. Systematic error sources considered include: (a) lepton energy scale uncertainty, accounted for with relative increase/decrease of the reconstructed photon and electron 4-momenta by ± 0.005 ; (b) jet energy scale uncertainty, expected to lie in the range from $\pm 5\%$ at $p_T = 20 \text{ GeV}/c$ to $\pm 2.5\%$ at $p_T > 50 \text{ GeV}/c$, and totally correlated to missing energy uncertainty (assumed to be $\pm 5\%$) [310]; (c) b-tagging uncertainty (4% after 10 fb^{-1} integrated luminosity [273]); (d) uncertainty in anti-tagging b-jet instead of non- b ones.

From the point of view of hadronization the FCNC signal differs from the dominant $t\bar{t}$ background just by the flavour of the quark originating from top. We assume that different hadronization processes in light and b-jet do not induce a significant change in the signal and $t\bar{t}$ topology. As a consequence, selection efficiency will be barely affected by theoretical systematics.

Beside these effects, the machine luminosity has an uncertainty that, in the low luminosity phase, is expected to be 5% [342].

Additional fluctuations of ΔB on the background due to systematic errors have to be considered, to re-evaluate the number of signal for observability and thus the branching ratio. These fluctuations can be implemented with a convolution of the Poissonian with the background distribution, that is commonly considered Gaussian if only detector effects are at work. Fluctuations related to theoretical uncertainties in the background are assumed to follow a flat distribution. For a quick estimation of these values, a tool available for the CMS collaboration is adopedt [79], that for given S , B , ΔB uses the proper distributions to produce the value for S_{c12} . Quite large effects are induced by tagging inefficiency, which is of particular relevance for this analysis since only one b-tagged is allowed, and by constraints on the three leptons in qZ , that are mandatory to disentangle the $t\bar{t}$ contribution.

All the considered sources systematic uncertainties are assumed to be uncorrelated, thus their effects are added quadratically. With inclusion of systematics, the results are $BR(t \rightarrow Zq) = (11.4 \pm 2.6^{+8.0}_{-0.5}) \times 10^{-4}$ and $BR(t \rightarrow \gamma q) = (5.7 \pm 0.5 \pm 1.3) \times 10^{-4}$. We conclude that this analysis with an integrated luminosity of 10 fb^{-1} could set upper limits on FCNCs in top-decay branching ratios of $BR(t \rightarrow Zq) = 22.0 \times 10^{-4}$ and $BR(t \rightarrow \gamma q) = 7.5 \times 10^{-4}$.

Table 8.28: The shifts induced by systematic effects on upper limits of Branching Ratio.

	$t \rightarrow Zq$	$t \rightarrow \gamma q$
stat	11.4	5.7
JES	$^{+0.4}_{-0.1}$	+1.0
Mistag	0.2	+0.6
Antitag	+0.5	+0.6
LES	$^{+7.9}_{0}$	+0.2
Theory	< 0.1	+0.2
Luminosity	< 0.1	0.1
TOTAL	$^{+8.0}_{-0.1}$	+1.3

Under the assumption that the selection efficiency is unaffected by moderate instantaneous luminosity increases (i.e., pile-up) the decrease in the upper limit on the branching fraction with increasing luminosity can be evaluated in a straightforward way. An improvement in the branching ratio limits by a factor of 2 is expected for a luminosity increase by a factor of 5.

Chapter 9

Electroweak Physics



9.1 Production of W and Z bosons



9.1.1 Introduction

The reactions $pp \rightarrow W + X$ and $pp \rightarrow Z + X$ with subsequent leptonic decays of the massive electroweak vector bosons, $W \rightarrow \ell\nu$ and $Z \rightarrow \ell^+\ell^-$, have a large cross section and are theoretically well understood. Cross sections above 10 nb (1 nb) are expected at the LHC for the $W \rightarrow \ell\nu$ ($Z \rightarrow \ell^+\ell^-$) channel in the fiducial region of the CMS detector. Hence these reactions are useful for many purposes, including a precise luminosity monitor, a high-statistics detector calibration tool and to demonstrate the performance of the CMS experiment. These reactions will be among the first to be measured at the LHC.

Here we discuss prospects for precise measurements of the reactions $pp \rightarrow Z + X$ and $pp \rightarrow W + X$ at the LHC using the decays of the gauge bosons into electrons and muons. Studies have been performed based on Monte Carlo samples generated with PYTHIA including realistic detector simulation and addressing the most relevant systematic effects. The potentially most dangerous background in these analyses consists of QCD events with leptons from hadron decays or tracks misidentified as leptons. However, these lepton candidates are associated to jets and can be largely suppressed using isolation algorithms.

Robust criteria are developed which allow for a low-background event selection which is rather insensitive to detector inhomogeneities. This robust selection is considered as especially useful for the CMS startup phase. The results show that a determination of the W and Z rates with an experimental precision on the percent level is feasible already in the early phase of the experiment.

9.1.2 W/Z into Electrons

The process $pp \rightarrow ZX$ and $pp \rightarrow WX$ with subsequent decay of Z and W into electrons is studied using the full CMS detector simulation and analysis scheme. The aim is to define some baseline selection which is suppressing background to a very small level and detector inhomogeneities can be controlled. This selection can thus be considered as especially useful for the CMS startup phase. Details can be found in [343].

Electron (positron) candidates are selected with the following criteria [303]:

- The minimal E_T of the electromagnetic cluster has to be larger than 20 GeV with $|\eta_{\text{cluster}}| < 1.4$ for barrel electron candidates and $1.6 < |\eta_{\text{cluster}}| < 2.4$ for endcap

electron candidates.

- The cluster should be consistent with the shower shape expected for electromagnetic showers. The spread of the electromagnetic shower along the η direction is rather insensitive to bremsstrahlung, thus allowing a good separation of signal and background shower shapes. Therefore it is required that the spread of the electromagnetic shower in η with respect to η of the supercluster, $\sigma_{\eta\eta}$, is smaller than 0.01.
- The energy deposit in the associated hadron calorimeter cluster should be very small. For this selection the ratio $E_{\text{Had}}/E_{\text{EM}}$ has to be smaller than 0.05.
- In order to be identified as an electron, a reconstructed track has to be matched with the cluster such that $\Delta R < 0.15$ (where $\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2}$). Furthermore, it is required that the ratio of the cluster energy and the track momentum, E/P , is larger than 0.9 and that $|1/E - 1/P| < 0.02$.
- Finally, it is required that the electron candidate is isolated. The transverse momentum sum of all other tracks found within a cone radius ΔR of 0.35 divided by the electron candidate transverse supercluster energy has to be smaller than 0.2. Only tracks with a transverse momentum above 1.5 GeV/ c and with at least four hits in the central tracker which are close to the interaction vertex are considered.

9.1.2.1 $pp \rightarrow Z \rightarrow eeX$ Selection

We analyze events where one e^+e^- pair consistent with the Z mass is found (if more than two electrons pass the selection criteria, only those two with the highest transverse momenta are considered). The generated and reconstructed mass distribution are shown in Figure 9.1 left. For now, the “electron” clusters are not corrected for bremsstrahlung within the tracker and the reconstructed Z peak is found to be about 1 GeV lower than the generated one.

Using this selection, the rapidity distribution of the accepted Z events is shown in Figure 9.1 right. In addition, the rapidity distribution of the potentially accepted Z bosons, separated for the three cases where both decay electrons are within the acceptance of the barrel calorimeter (BB) $|\eta_{\text{BB}}| < 1.4$, both within the endcaps (EE) $1.6 < |\eta_{\text{EE}}| < 2.4$ or one within the barrel and the other one in the endcaps (EB) are also shown. In the case that both generated electrons are in the barrel, a Z detection efficiency of about 60% is reached.

Here the electron efficiency is defined by the ratio of reconstructed electrons from accepted Z events to the number of electrons from generated Z events, where the generated electrons fulfilled the condition $|\eta_{\text{gen}}^e| < 1.4$. Figure 9.2 left shows the efficiency distribution for all supermodules folded such that the local ϕ angle for all odd supermodules goes from 0-20 degrees and for all even supermodules from 20-40 degrees.

The efficiency drop of about 10% between the supermodules is clearly visible with the available sample of Z events corresponding to roughly 0.2 fb^{-1} . Similar inefficiencies were found in the η direction at supermodule boundaries. From the analysis of the reconstruction efficiency as function of the phi angle, we get an efficiency of $27.1\% \pm 0.4\%$ (if the inter-supermodule regions are excluded) while the average over the whole phi range is $26.5 \pm 0.4\%$.

The average Z efficiency, when both electrons are generated and reconstructed in the barrel calorimeter, is found to be $57.3 \pm 0.2\%$ (where the uncertainties are from the finite number of Monte Carlo events). Half the efficiency loss is caused by the shower-shape requirement, and




Figure 9.1: Left: Reconstructed and generated Z mass distribution with all cuts. Right: Generated rapidity distribution for all Z candidates and for those where both electrons were generated within the geometrical acceptance of the electromagnetic calorimeter. For comparison, the rapidity distribution of the finally accepted Z events is already shown here.

another quarter by the energy-momentum matching requirement. If events, where at least one electron is reconstructed within the gaps, are removed, the average efficiency is found to be $58.4 \pm 0.2\%$. Assuming that the produced electrons must be homogeneous in ϕ and that the effects from geometrical gaps can be monitored with some reasonable statistics, it should be straightforward to correct for the detector gaps. Already with the available statistics used for this study, the corrections for the efficiency loss in the gaps can certainly be determined with a relative accuracy smaller than about 25%. This number is estimated from comparing the minimal efficiency in the gap and the efficiency in the non-gap regions.

We conclude that already with a few 100 000 reconstructed Z events, collected at the early stage of the experiment, an efficiency determination with a systematic accuracy of better than 1-2% should be possible. Obviously, with the much larger statistics of a few million Z events, these uncertainties can be further reduced. Once data from the CMS detector becomes available, these cuts can be applied on one electron and varied on the other electron to compare the selection efficiency in data and Monte Carlo simulation. This can be used to further improve the detector simulation and to better access systematic uncertainties.

9.1.2.2 $pp \rightarrow W \rightarrow e\nu X$ Selection

In order to pass the $W \rightarrow e\nu$ selection, events must have exactly one electron candidate in the barrel fulfilling the requirements described above, and missing transverse energy associated with the neutrino: a cut on the transverse mass of the $e\nu$ system is applied. The transverse mass m_T is defined as follows:

$$m_T = \sqrt{2p_T^{(e)} p_T^{(\nu)} (1 - \cos \Delta\phi)} \quad (9.1)$$

where $p_T^{(e,\nu)}$ is the (reconstructed) transverse momentum of the electron and the neutrino respectively and $\Delta\phi$ is the azimuthal angle between the electron and the neutrino.




Figure 9.2: Left: The electron reconstruction efficiency in $Z \rightarrow e^+e^-$ events as a function of ϕ , all even and odd numbered supermodules are folded such that the odd (even) numbered supermodules always cover local ϕ angles from 0 to 20 degrees and from 20 to 40 degrees respectively. The dotted line corresponds to the average efficiency $57.3 \pm 0.2\%$ over the whole ϕ range and the solid lines correspond to the average efficiency $58.4 \pm 0.2\%$ with the gap regions excluded. Right: Generated (solid line) and reconstructed (dashed line) transverse W mass. The W transverse mass is reconstructed from the electron four-momentum and the missing transverse energy. In this plot, only events with no reconstructed jet above 20 GeV transverse energy are included.

The missing transverse energy can be determined in several ways, for example:

1. From the vector sum of all clusters in the calorimeter
2. From the vector sum of hard objects only

In the electromagnetic calorimeter, the electron transverse energy can be measured accurately. However, the reconstructed transverse missing energy shows a significant bias.

Suspecting that low energy objects (randomly distributed across the detector) are responsible for this bias, we follow the second approach: We select reconstructed jets with a transverse energy above 20 GeV and absolute pseudorapidity less than 2.4 and reconstruct the missing transverse energy only from these jets and the electron. Here we use uncalibrated jets, i.e. whenever we refer to the jet energy we mean raw jet energy.

To study this possibility in more detail, we split our sample into events without jets (as defined in the previous paragraph) and events with one or more jets. Note that in the case of zero accepted jets, only the electron is used to calculate the neutrino transverse energy which is then very close to the electron transverse energy (pointing into opposite directions in ϕ). The transverse mass is equal to twice the electron transverse energy in this case.

No systematic bias is found with this method and the mean value is close to zero. We thus use this method to reconstruct the neutrino transverse energy. The reconstructed W transverse

mass is shown in Figure 9.2 right. For the purpose of this analysis and the counting of resonant W events, we require the transverse mass to lie in the interval 60 to 100 GeV/ c^2 .

We consider two sources of systematic uncertainties here: The uncertainty due to inhomogeneities in the detector geometry and the uncertainty related to the jet veto. We expect that the uncertainty from the reconstruction efficiency as function of the electron azimuthal angle for the efficiency correction will be similar as for the Z selection.

To address the effect of the scale uncertainty of the absolute calibration on the jet definition, we investigated the changes in the selection efficiency when moving the threshold transverse energy for the jet definition. It follows that for a cut on the transverse jet energy at 20 GeV, the efficiency slope is roughly 0.1 % (absolute) per GeV, corresponding to a relative uncertainty of about 0.25% per GeV.

Assuming a jet energy scale uncertainty of 15% at the LHC startup we obtain an efficiency uncertainty of 0.75% relative. For 5% uncertainty in the jet energy scale (expected after the final detector calibration), this value reduces to 0.25%.

The efficiency change due to the jet veto can also be estimated directly from $Z \rightarrow e^+e^-$ events (applying a jet veto to these events). In the future, this can be done directly from the data recorded with the CMS detector. Thus with the expected large data samples of $Z \rightarrow e^+e^-$, remaining differences between data and Monte Carlo can be studied and corrected with very small uncertainties.

9.1.3 W/Z into Muons

Simple sets of cuts can be used in CMS to select large statistics samples of $Z \rightarrow \mu\mu$ and $W \rightarrow \mu\nu$ events with high purity. They are described in detail in Reference [344] and summarised here.

The $Z \rightarrow \mu\mu$ selection criteria have been chosen to minimize uncertainties from the muon chamber response and from the matching between the inner tracker and the muon spectrometer. The basic idea is to accept events in which one of the muons is reconstructed as an isolated track in the central tracker detector, even if no associated track in the muon spectrometer is present. This results in a more uniform efficiency as a function of the pseudo-rapidity, as observed in Figure 9.3 left. From the kinematics point of view only muons with $p_T > 20$ GeV and pseudorapidity $|\eta_\mu| < 2.0$ are considered in the present analysis. A dimuon mass window of $\pm 3\Gamma_Z = 7.5$ GeV around the reconstructed Z mass is used. Figure 9.3 right shows the efficiency of the HLT criteria on the selected sample as a function of the muon pseudo-rapidity. One can clearly observe two regions with smaller efficiency, around $|\eta| \approx 0.25$ and $|\eta| \approx 0.8$, where transitions between two muon wheels take place. The efficiency is dominated by the dimuon component, which represents a unique tool to study the performance of the single-muon subtrigger, which is of relevance for other selections, like $W \rightarrow \mu\nu$.

Even if the rate of $W \rightarrow \mu\nu$ events is expected to be larger than the $Z \rightarrow \mu\mu$ rate by an order of magnitude, the experimental context is more demanding due to a lower trigger efficiency, only moderate transverse missing energy in the event, the absence of a precise mass constraint and a full dependence on tracker and muon spectrometer behaviours. This will lead to larger experimental uncertainties, which can be studied with the $Z \rightarrow \mu\mu$ data samples. The selection of $W \rightarrow \mu\nu$ events uses the same η cut but a higher p_T threshold, 25 GeV, due to the higher threshold for the single-muon trigger. Figure 9.4 shows the transverse in-




Figure 9.3: Left: Muon efficiency as a function of pseudo-rapidity in the selected $Z \rightarrow \mu\mu$ sample. Two cases are considered: a selection using only muons reconstructed in the muon chambers (dashed histogram) and the selection described in the text (solid histogram), which also accepts isolated tracks in the inner tracker. For this test, no HLT trigger criteria have been applied. Right: HLT efficiency on the selected $Z \rightarrow \mu\mu$ sample as a function of the pseudorapidity of one of the muons. All but the HLT trigger criteria have been applied. The fitted line corresponds to the average HLT efficiency in the region $|\eta_\mu| < 2$. The regions at $|\eta| \approx 0.25$ and $|\eta| \approx 0.8$, with a slightly lower trigger efficiency, are visible. The fraction of events triggered by dimuon and single-muon triggers are also shown.

variant mass distribution of the muon- E_T^{miss} system in $W \rightarrow \mu\nu$ events, compared to QCD expectations.

Systematic uncertainties in the determination of $Z \rightarrow \mu\mu$ and $W \rightarrow \mu\nu$ acceptances are summarized in Tables 9.1 and 9.2. The various sources of uncertainties are discussed in detail in Reference [344]. Most of them are evaluated for a CMS detector calibrated with 1 fb^{-1} . The experimental components are well under control in the case of the $Z \rightarrow \mu\mu$ selection, with the limited knowledge on the track efficiency as the dominant source. In the $W \rightarrow \mu\nu$ case, many of them contribute at a similar level, with E_T^{miss} providing the largest uncertainty. Concerning theoretical sources, the boson p_T uncertainties are the dominant contribution. They are estimated from a comparison between LO and NLO CMS simulations using MC@NLO as event generator [345], as shown in Figure 9.5.

The results of the study can be summarized in terms of cross section measurement accuracies, for 1 fb^{-1} of integrated luminosity, as follows: $\Delta\sigma/\sigma(pp \rightarrow Z + X \rightarrow \mu\mu + X) = 0.13 \text{ (stat.)} \pm 2.3 \text{ (syst.)} \pm 10 \text{ (lumi) \%}$ and $\Delta\sigma/\sigma(pp \rightarrow W + X \rightarrow \mu\nu + X) = 0.04 \text{ (stat.)} \pm 3.3 \text{ (syst.)} \pm 10 \text{ (lumi)\%}$, where luminosity represents the dominant uncertainty which will eventually decrease to 5% with more integrated luminosity. QCD backgrounds seem to be under control, even if final checks with data will be necessary to determine the level of background with more precision.

Therefore, rates within the fiducial volume of the detector can be determined with high accuracy, even for the first stages of the LHC ($\approx 2.3\%$ for $Z \rightarrow \mu\mu$ and $\approx 3.3\%$ for $W \rightarrow \mu\nu$). These uncertainties will be significantly reduced with the use of the next generation of NLO Monte Carlos and final detector calibrations, and allow these reactions to be used to determine the luminosity.




Figure 9.4: Transverse invariant mass reconstructed in $W \rightarrow \mu\nu$ events. In order to observe the shape of the QCD background with more statistics, the HLT muon isolation criteria have not been applied to obtain the plot. The position of the lower cut ($M_{\mu\mu} > 40 \text{ GeV}/c^2$) is indicated with an arrow.

Table 9.1: Relative systematic uncertainties on the acceptance for the $Z \rightarrow \mu\mu$ sample.

Source	Uncertainty (%)
Tracker efficiency	1
Magnetic field knowledge	0.03
Tracker alignment	0.14
Trigger efficiency	0.2
Jet energy scale uncertainties	0.35
Pile-up effects	0.30
Underlying event	0.21
Total exp.	1.1
PDF choice (CTEQ61 sets)	0.7
ISR treatment	0.18
p_T effects (LO to NLO)	1.83
Total PDF/ISR/NLO	2.0
Total	2.3

Table 9.2: Relative systematic uncertainties on the acceptance for the $W \rightarrow \mu\nu$ sample.

Source	Uncertainty (%)
Tracker efficiency	0.5
Muon efficiency	1
Magnetic field knowledge	0.05
Tracker alignment	0.84
Trigger efficiency	1.0
Transverse missing energy	1.33
Pile-up effects	0.32
Underlying event	0.24
Total exp.	2.2
PDF choice (CTEQ61 sets)	0.9
ISR treatment	0.24
p_T effects (LO to NLO)	2.29
Total PDF/ISR/NLO	2.5
Total	3.3




Figure 9.5: Left: Comparison between LO and NLO predictions for the muon p_T distribution in $Z \rightarrow \mu\mu$ selected events. Both histograms have been normalized to the total number of events generated in the fiducial volume: $|\eta_\mu| < 2.5$, $p_{T\mu}^{max} > 20 \text{ GeV}/c$, $p_{T\mu}^{min} > 10 \text{ GeV}/c$ and $M_Z - 6\Gamma_Z < M_{\mu\mu} < M_Z + 6\Gamma_Z$ Right: Comparison between LO and NLO predictions for the muon p_T distribution in $W \rightarrow \mu\nu$ selected events. Both histograms have been normalized to the total number of events generated in the fiducial volume: $|\eta_\mu| < 2.5$.



Parton distribution functions and parton luminosities

The production of inclusive W and Z events is theoretically well understood and the couplings to quarks and leptons have been measured with accuracies of 1% or better. Thus, it follows from the previous sections that a precise counting of $W \rightarrow e\nu, \mu\nu$ and $Z \rightarrow ee, \mu\mu$ events is equivalent to a precise measurement of the quantity

$$\int_{q,\bar{q} \text{ partons}} dx_1 dx_2 \sigma_{q\bar{q} \rightarrow W,Z} \times L_{pp} \times PDF(x_1, x_2, Q^2), \quad (9.2)$$

where L_{pp} is the LHC integrated luminosity, $\sigma_{q\bar{q} \rightarrow W,Z}$ is the cross section for inclusive W or Z production at the partonic level and $PDF(x_1, x_2, Q^2)$ denotes the probability to produce quarks and antiquarks with proton fractions x_1 and x_2 at a scale Q^2 . The prospect studies of Reference [344], summarized in Table 9.3, show that uncertainties on the parton distribution functions (PDF) have a relatively small influence on the experimental acceptance for the rates, but a large effect on the global rate expectations.

Table 9.3: Estimated uncertainties in the rate and in the acceptance for the $pp \rightarrow Z + X \rightarrow \mu\mu + X$ and $pp \rightarrow W + X \rightarrow \mu\nu + X$ processes. The global rate is referred to the fiducial volumes used in Reference [344], which include a pseudorapidity cut of $|\eta_\mu| < 2.5$.

	$Z \rightarrow \mu\mu$	$W \rightarrow \mu\nu$
Global rate uncertainty (%)	+5.8 -7.9	+5.6 -7.4
Acceptance uncertainty (%)	+0.4 -0.7	+0.6 -0.9

On one hand we conclude from Table 9.3 that a comparison between theory and experiment with a 6 – 7% accuracy is possible. This comparison provides a measurement of the integrated luminosity L_{pp} with a similar level of precision. On the other hand, the small theoretical uncertainties on the experimentally measured rate (from the acceptance uncertainty) allow to significant improvements in many cross section rate measurements, like $\sigma(pp \rightarrow ZZ + X)/\sigma(pp \rightarrow Z + X)$, in which PDF and luminosity uncertainties cancel. Current studies within theoretical and experimental communities [346] aim to a further reduction of uncertainties associated to PDFs. Finally, PDF validity tests and further reductions in the acceptance uncertainty (below the percent level) will require dedicated studies of the lepton rapidity distributions observed in data, like those suggested in Reference [347].



Muon pairs from the Drell-Yan process

9.2.1 Introduction

In the Standard Model, the production of lepton pairs in hadron-hadron collisions, the Drell-Yan (DY) process [348], is described by s -channel exchange of photons or Z bosons. The parton cross section in the lepton-pair center-of-mass system has the form:

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4s} [A_0(1 + \cos^2 \theta) + A_1 \cos \theta] \quad (9.3)$$

where $\sigma = \frac{4\pi\alpha^2}{3s} A_0$ and $A_{\text{FB}} = \frac{3}{8} \frac{A_1}{A_0}$ are the total cross section and the forward-backward asymmetry, and θ is angle of lepton in the dilepton rest frame with respect to the quark direction. The terms A_0 and A_1 are fully determined by the electroweak couplings of the initial- and final-state fermions. At the Z peak the Z exchange is dominating and the interference term is vanishing. At higher energies both photon and Z exchange contribute and the large value of the forward-backward asymmetry is due to the interference between the neutral currents. Fermion-pair production above the Z pole is a rich search field for new phenomena at present and future high energy colliders. The differential cross section is sensitive to manifestation of new physics from a multi-TeV scale by adding new amplitudes or through their interference with the neutral currents of the SM. At hadron colliders the parton cross sections are folded with the parton density functions (PDF): $pp \rightarrow l_1 l_2$

$$\frac{d^2\sigma}{dM_{ll}dy} [pp \rightarrow l_1 l_2 + X] \approx \sum_{ij} (f_{i/p}(x_1) f_{j/p}(x_2) + (i \leftrightarrow j)) \hat{\sigma}, \quad (9.4)$$

where $\hat{\sigma}$ is the cross section for the partonic subprocess $ij \rightarrow l_1 l_2$, $M_{ll} = \sqrt{\tau s} = \sqrt{\hat{s}}$ the mass of the lepton-pair system, y the rapidity of the lepton pair, $x_1 = \sqrt{\tau} e^y$ and $x_2 = \sqrt{\tau} e^{-y}$ the parton momentum fractions, and $f_{i/p(\bar{p})}(x_i)$ the probability to find a parton i with momentum fraction x_i in the proton.

Table 9.4: x_1 and x_2 for different masses and rapidities.

y	0	2	4	0	2	4	0	2	4
	M = 91.2 GeV/ c^2			M = 200 GeV/ c^2			M = 1000 GeV/ c^2		
x_1	0.0065	0.0481	0.3557	0.0143	0.1056	0.7800	0.0714	0.5278	-
x_2	0.0065	0.0009	0.0001	0.0143	0.0019	0.0003	0.0714	0.0097	-

The total cross section and the forward-backward asymmetry are function of observables which are well measured experimentally for final states containing e^+e^- or $\mu^+\mu^-$: the invariant mass and the rapidity of the final-state lepton pair. This allows to reconstruct the center-of-mass energy of the initial partons, even if their flavors are unknown. For a ($x_1 \geq x_2$) pair of partons we have 4 combinations of up - or $down$ -type quarks initiating the interaction: $u\bar{u}, \bar{u}u, d\bar{d}, \bar{d}d$. In pp collisions the antiquarks come always from the sea and the quarks can have valence or sea origin. The x -range probed depends on the mass and rapidity of the lepton pair as shown in Table 9.4.

The results presented here extend the studies for the LHC SM workshop (see [149] and references therein), using more data and the CMS full detector simulation and reconstruction. More details can be found in [349].

9.2.2 Cross section measurements

Simulation of Drell-Yan events in proton-proton collisions at 14 TeV centre-of-mass energy is performed with PYTHIA 6.217 using the CTEQ5L parton distribution functions. The possible contributions from higher-order terms in the dimuon production cross section are taken into account by using a K factor of 1.3 PHOZPRMS [350]. Eleven samples of 10 000 events each with different cut-off values on the dimuon invariant mass are generated: $M_{inv} \geq 0.2, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5$ TeV/ c^2 . Only events with at least two muons in the pseudorapidity range $|\eta| \leq 2.5$, with transverse momentum $p_T \geq 7$ GeV are preselected. No cuts on isolation

of muons are made at the preselection stage. The total efficiency for dimuon preselection, ε , is about 87 % for a mass of 1 TeV/ c^2 and 96% for a mass of 5 TeV/ c^2 . To simulate the detector geometry, materials and particle propagation inside the detector, the GEANT 4-based simulation of the CMS detector is used.

The trigger simulation is based on the on-line reconstruction algorithms. Events are selected by the single- and double-muon triggers. This means that at least one muon candidate is within pseudorapidity region $|\eta| \leq 2.1$. The total efficiency of triggering including reconstruction and trigger selection efficiency is 98 % at 1 TeV. There is significant decrease in trigger efficiency after applying calorimeter isolation cuts (down by 15 %). The tracker isolation practically does not affect the trigger efficiency. Thus the additional cuts on calorimeter and tracker isolation of muon tracks are not applied in this analysis.

The off-line muon reconstruction algorithm is applied only to events which have passed trigger selection. At the off-line level two muons inside the CMS acceptance $|\eta| \leq 2.4$ are required. The overall efficiency of the full reconstruction procedure taking into account trigger and off-line reconstruction inefficiency is between 97% and 93% for a mass range of 0.2 to 5 TeV/ c^2 , as shown in Figure 9.6 left. In the case of an ideal detector the mass resolution smearing for fully-reconstructed events is between 1.8% and 6% for the same mass range, Figure 9.6 right. The effect of misalignment on the mass resolution varies from 1.1% up to 2.3% (1.3%) for the *First Data (Long Term)* scenarios at the Z and from 5% up to 25% (6%) for V/c^2 .




Figure 9.6: Left: dimuon reconstruction efficiency, and right: invariant mass resolution; both as function of the invariant mass cut.

The cross sections of Drell-Yan production for the simulated CMS runs are shown in Table 9.5. The non-reducible backgrounds considered are vector boson pair production ZZ , WZ , WW , $t\bar{t}$ production etc. The simulation and preselection of background events is done with the same cuts as for the signal above. In the SM the expected leading-order cross section of these events is negligible in comparison with the Drell-Yan one, see Table 9.5.

The $\tau\tau$ background (from τ decaying to μ and neutrinos) is 0.8 % at the Z pole and 0.7 % for masses above 1 TeV/ c^2 . The background from Drell-Yan production of $q\bar{q}$ pairs (mostly semileptonic b or c decays) is 0.3 % at the Z pole without applying any isolation cuts and below 0.1 % for masses above 1 TeV/ c^2 . The other background sources are negligible. If the need arises they can be further suppressed by acoplanarity and isolation cuts in the tracker.

The main experimental systematic effects in the cross section measurement arise from the

Table 9.5: Leading-order cross sections of Drell-Yan, preselected Drell-Yan, di-bosons (ZZ , ZW , WW) and $t\bar{t}$ events in fb. The CTEQ5L parton distributions are used.

$M_{\mu^+\mu^-}$, TeV/ c^2	≥ 1.0	≥ 1.5	≥ 2.0	≥ 2.5	≥ 3.0	≥ 4.0
Drell-Yan	6.61	1.04	$2.39 \cdot 10^{-1}$	$6.53 \cdot 10^{-2}$	$1.97 \cdot 10^{-2}$	$2.09 \cdot 10^{-3}$
Pre-sel. D-Y	5.77	$9.53 \cdot 10^{-1}$	$2.24 \cdot 10^{-1}$	$6.14 \cdot 10^{-2}$	$1.87 \cdot 10^{-2}$	$2.00 \cdot 10^{-3}$
Di-bosons	$2.59 \cdot 10^{-4}$	$1.51 \cdot 10^{-4}$	$5.6 \cdot 10^{-5}$	$2.26 \cdot 10^{-5}$	$9.06 \cdot 10^{-6}$	$1.66 \cdot 10^{-6}$
$t\bar{t}$	$2.88 \cdot 10^{-4}$	$2.58 \cdot 10^{-4}$	$1.55 \cdot 10^{-4}$	$7.02 \cdot 10^{-5}$	$2.93 \cdot 10^{-5}$	$3.65 \cdot 10^{-6}$

total muon inefficiency and momentum resolution. The latter is very important at high mass as smearing from lower masses from the steeply falling Drell-Yan spectrum can contaminate the high mass measurements, especially if the tails of the momentum resolution are not under control. The main sources of systematic uncertainties on the momentum resolution come from the alignment of the muon chambers and the central tracker, both at start-up and high luminosity.

Table 9.6: Relative errors of the Drell-Yan muon pairs cross section measurements in the fiducial volume.

$M_{\mu^+\mu^-}$, TeV/ c^2	Detector smearing	Statistical 1 fb $^{-1}$	Statistical 10 fb $^{-1}$	Statistical 100 fb $^{-1}$	Theor. Syst.
≥ 0.2	$8 \cdot 10^{-4}$	0.025	0.008	0.0026	0.058
≥ 0.5	0.0014	0.11	0.035	0.011	0.037
≥ 1.0	0.0049	0.37	0.11	0.037	0.063
≥ 2.0	0.017		0.56	0.18	0.097
≥ 3.0	0.029			0.64	0.134

The statistical errors for 1, 10 and 100 fb $^{-1}$ runs, the systematic uncertainty due to smearing in the detector and from theory side are given in Table 9.6. The modification of the measured cross section due to uncertainty of the mass resolution does not exceed 2.9% which is reached. for a mass of 3 TeV/ c^2 , see Table 9.6. This has been estimated by applying an additional smearing to the dimuon mass (see [349, 93]. The misalignment does not affect the efficiency of dimuon reconstruction for any masses [93]. Taking into account the trigger efficiency changes from 98.5 % to 97% for masses from 0.2 to 5 TeV/ c^2 , very conservatively we may assign half of this change with mass i.e. 0.75 % as a systematic uncertainty.

An important ingredient in the cross section measurement is the precise determination of the luminosity. A promising possibility is to go directly to the parton luminosity [347] by using the W^\pm (Z) production of single (pair) leptons. New estimates show that in this way the systematic error on $\sigma_{DY}^{high Q^2}$ relative to σ_Z can be reduced to $\approx 5 - 12\%$ [351].

On the theory side we consider several sources of systematic uncertainties. Higher order QCD corrections are often taken into account with K -factor of 1.3 as calculated with the program PHOZPRMS [350]. It is expected that the total value of additional NNLO contributions does not exceed 8 % .

A full-scale analysis of experimental data (comparison data with theory, taking into account acceptance corrections for precise measurement of σ and A_{FB} at large centre-of-mass energies \hat{s}) requires good knowledge of the different types of genuine electroweak (EW) radiative corrections to the DY process: vertex, propagator, EW boxes. A complete one-loop parton cross section calculation has been included in [149] and confirmed in [352]. The EW corrections change the cross section by 10-20%. The calculation [99] of the weak radiative corrections to the Drell-Yan processes due to additional heavy bosons contributions shows that these corrections are about 2.9% to 9.7% for mass region between 0.2 TeV/ c^2 and 5 TeV/ c^2 .

The phenomenological origin of PDF gives one additional systematic error. First of all, estimates of cross section obtained by using different sets of structure functions do not give exactly the same values. The results vary within $\pm 7\%$ for $M_{ll} \geq 1$ TeV. The internal PDF uncertainties are estimated using the LHAPDF library [88, 353]. The PDF-dependence of the acceptance efficiency is estimated by using the PDF sets CTEQ5L, CTEQ6L and MRST2001E. The changes in the acceptance efficiency are up to 0.5%. The ambiguity in the acceptance efficiency due to internal PDF uncertainties is larger, but less than 1.4% for any mass region.




Figure 9.7: Size of the EW corrections and the cross section uncertainties from PDFs, hard process scale and detector understanding as a function of the dimuon invariant mass cut.

The summary of the estimated systematic uncertainties as function of the di-lepton mass is given in Figure 9.7. The CMS experiment has excellent potential to measure the cross section for dimuon pairs up to the highest masses that will be accessible at the LHC, and to test the Standard Model up to very high momentum transfers in a new and unexplored energy range. Current uncertainties from theory are larger than the experimental uncertainties. The statistical errors will dominate for invariant masses larger than 2 TeV/ c^2 even for 100 fb^{-1} .

9.2.3 Prospects on the measurement of the forward-backward asymmetry

To measure the forward-backward asymmetry we need the original quark and anti-quark directions of the initiating partons, but these are not known in the case of pp experiments, where the initial state is symmetric. In Ref. [89, 106] it is shown that it is possible to approximate the quark direction with the boost direction of the dimuon system with respect to the

beam axis. This is due to the fact that the valence quarks have on average larger momentum than the sea antiquarks, and therefore the dimuon boost direction approximates the quark direction. The most unambiguous tagging occurs for large dimuon rapidity.

The approximation of the original quark direction for pp collisions leads to a flattening out of the original asymmetry (≈ 0.61 for Drell-Yan events) by a factor of almost 2. However, using multi-dimensional fits [105] or reweighting techniques depending on the mistag and acceptance which are under development, we can measure the original asymmetry.

The accuracy of asymmetry measurements depends on:

- statistical uncertainty which grows with rising mass cut value, as the number of events for integrated luminosity of e.g. $\int \mathcal{L} dt = 100 \text{ fb}^{-1}$ decreases with mass
- systematic uncertainty from the variation of the mistag probabilities for various PDF sets, typically below 10 %.

We expect the systematic uncertainty to dominate the statistical one for integrated luminosity of $\int \mathcal{L} dt = 100 \text{ fb}^{-1}$ and dimuon masses around $500 \text{ GeV}/c^2$, while the statistical one to be more important for dimuon mass cuts above $1000 \text{ GeV}/c^2$.



Determination of the W mass

9.3.1 Introduction

The precise measurement of the mass of the W boson constitutes an important consistency check of the Standard Model and, together with the top quark mass, is sensitive to supersymmetric corrections. Such a precision measurement of the W mass at the LHC becomes feasible because a huge sample of data available at the LHC will guarantee a nearly negligible statistical uncertainty and a good control of the systematic effects. Extrapolating from traditional approaches based on the reconstruction of the transverse mass $m_T = \sqrt{2p_T^l p_T^\nu (1 - \cos(\theta_{l\nu}))}$ in leptonic W decays, the most relevant contributions to the systematic uncertainties come from the lepton energy or momentum scale, the lepton energy or momentum resolution, the modelling of the system recoiling against the W boson, the parton distribution functions, the W intrinsic width, from radiative decays and from backgrounds. To accomplish a competitive measurement of the W boson mass, new strategies must be considered [354]. The most promising one consists in predicting the distribution of experimental observables sensitive to the W mass, such as the transverse momentum of the charged lepton (p_T^l) and the transverse mass of the boson from the corresponding distribution measured in Z boson decays into two charged leptons. The concept of transverse mass measurement can be applied to Z boson events by regarding one of the reconstructed leptons as missing energy. The theoretical description of both decays is very similar and the resulting distributions in transverse mass are comparable for a wide range in kinematics.

The advantage of this approach, conceptually discussed in [355], is that most of the experimental and theoretical uncertainties, being common between W and Z , cancel in the comparison, leading to a global reduction of the systematic uncertainty. The drawback is a larger statistical uncertainty due to the smaller production rate of Z bosons decaying to charged leptons. Yet a statistical precision of order $10 \text{ MeV}/c^2$ and $30 \text{ MeV}/c^2$ for an integrated luminosity of 10 fb^{-1} and 1 fb^{-1} respectively is anticipated. In order not to be limited by

statistics, the analyses are performed using large data samples produced with the fast simulation of the CMS experiment [287]. Smaller samples of fully simulated events are used for cross checks.

Two different ways to relate Z to W boson events are considered. One is based on the comparison of the same experimental observables in W - and Z -events scaled to the boson masses. The sensitivity of this method, which can take advantage of the precision calculation of the theoretical ratio of the W and Z boson differential production cross-sections, is fully addressed in the analysis of transverse energy distribution of the electrons from $W \rightarrow e\nu$ decays. An alternative approach considered in the analysis of $W \rightarrow \mu\nu$ events consists of predicting W boson distributions from Z -events by means of kinematic transformations of measured Z events, parameterized as a function of the boson masses and widths. This more phenomenological approach is exploited in the analysis of the transverse mass distributions, and relies less on the theoretical prediction of the boson p_T .

9.3.2 Event selections

In order to obtain a clean signal of $W \rightarrow l\nu$ decays, events that passed the High Level Trigger (HLT) for single leptons are required to satisfy the following selection cuts: one isolated muon with $p_T > 25$ GeV/ c within the pseudo-rapidity region $|\eta| < 2.3$ or one isolated electron with $p_T > 25$ GeV/ c and within $|\eta| < 2.4$; missing transverse energy $E_T^{\text{miss}} > 25$ GeV; no jets in the event with $p_{T,jet} > 30$ GeV/ c ; the transverse momentum of the system recoiling against W has to be lower than 20 GeV/ c , measured from the lepton p_T and the missing transverse energy.

The difference in minimum p_T of the charged lepton is determined by the single lepton trigger threshold. The last two selection cuts are intended to select W bosons produced with a small transverse momentum. The selection efficiency is about 15% for the electron channel and 25% for the muon channel, with a background at the percent level, dominated by leptonic Z decays with one lepton outside the acceptance, as shown in Figure 9.8.

Z events used to predict the W distribution are also selected from the sample of events passing the HLT for single leptons. Z candidates contain a pair of identified charged leptons consistent with the Z mass hypothesis [354]. One of the two leptons, randomly chosen, is removed from the event to mimic a W decay. The same selections discussed above are then applied, with the cut values on the lepton quantities (minimum lepton p_T and event missing transverse energy) scaled by the ratio M_Z/M_W . This choice is intended to minimize kinematic and acceptance differences in Z and W events and thus the theoretical uncertainties implied by the above mentioned approaches.

9.3.3 $W \rightarrow e\nu$

The analysis strategy is based on the prediction of the experimental distribution of the electron transverse energy in W events scaled to the boson mass from the corresponding distribution measured for Z bosons decaying into e^+e^- pairs, along with the theoretical ratio between the W and Z cross-sections, calculated at a fixed perturbative order. Ideally, the differential cross section for the W boson can be predicted from the one measured for Z boson by scaling the lepton transverse momenta with the boson masses, $p_T^{\text{lept},Z} = M_Z/M_W p_T^{\text{lept},W}$,




Figure 9.8: W events and main backgrounds for 1 fb^{-1} . Left: Electron scaled transverse energy distribution in $W \rightarrow e\nu$ decays and the backgrounds from $Z \rightarrow e^+e^-$, from $Z \rightarrow \tau^+\tau^-$ and from $W \rightarrow \tau\nu$ for 1 fb^{-1} . Right: Transverse mass distribution in the muon channel with the fractions of $Z^0/\gamma^* \rightarrow \mu^+\mu^-$ (red/grey) $W \rightarrow \tau\nu$ (blue/dark), and $W \rightarrow \mu\nu$ (yellow/light) events.

as:

$$\left. \frac{d\sigma^W}{dp_T^{\text{lept},W}} \right|_{\text{pred}} = \frac{M_Z}{M_W} R(X) \left. \frac{d\sigma^Z}{dp_T^{\text{lept},Z}} \left(p_T^{\text{lept},Z} = \frac{M_Z}{M_W} p_T^{\text{lept},W} \right) \right|_{\text{meas}}, \quad (9.5)$$

where $R(X) = \frac{d\sigma^W}{dX^W} / \frac{d\sigma^Z}{dX^Z}$ is the ratio, deduced from theoretical calculations, between the differential cross sections in terms of the scaled variable $X^V = \frac{p_T^{\text{lept},V}}{M_V}$, with $V=W,Z$. The parameter M_W can be extracted by fitting this prediction to the distribution for W events observed in the experiment. In practice, additional corrections to $R(X)$ are needed to account for the acceptance to Z and W events and for the experimental resolution. This calls for a detailed understanding of the detector response by means of Monte Carlo simulations compared to control samples. Clearly, the definition of $R(X)$ is the most critical aspect and must include both detector effects and theoretical predictions.

The results for 1 fb^{-1} of integrated luminosity using the technique just described are shown in Figure 9.9. The statistical precision of the method is determined from the resulting χ^2 distribution. The evaluation of the systematic uncertainties affecting the measurement of the W mass is performed by determining the distortions implied by the different systematic effects mentioned above. The effects of instrumental origin have been studied by fixing $R(X)$ to the theoretical prediction exactly describing the samples of generated events (i.e. an exact knowledge of the theory is assumed) and by introducing distortions and biases in the detector response. The resulting shift in M_W is assumed as the systematic uncertainty associated to the effect. The detector response to electrons, the largest source of systematic uncertainty of instrumental origin with this method, can be determined with the required precision from $Z \rightarrow ee$ events.

The prediction of the lepton transverse spectrum is plagued by large radiative QCD corrections. Yet, in the method adopted, large cancellations occur and $R(X)$ can be reliably pre-




Figure 9.9: Comparison of the scaled electron E_T spectra for Z (dots) and W boson (line) events (left) and χ^2 dependence on M_W (right) for 1 fb^{-1} of integrated luminosity.

dicted. The uncertainty related to the missing orders in the perturbative expansion can be quantified by the dependence of the available NLO prediction on the choice of the renormalisation and factorisation scales. A conservative figure of $30 \text{ MeV}/c^2$ for the mass uncertainty is deduced. This will become the dominant error at 10 fb^{-1} . Yet the reduction of this error by extending the calculation one order higher in α_S is technically feasible [355].

9.3.4 $W \rightarrow \mu\nu$

As a complementary method, the transverse mass distribution of W events in the muon channel is modelled from $Z \rightarrow \mu^+ \mu^-$ events by a kinematic transformation. In the rest frame of the Z boson, the lepton momenta are scaled such that their invariant mass distribution represents that of the W boson [354]. After removing one randomly chosen muon to mimic a neutrino, the whole system is boosted back into the detector frame, thus obtaining a template for the expected distribution of W events, which depends on the W and Z boson masses and widths as parameters. By iterating the procedure for different W boson masses, the best agreement with the observed transverse mass distribution in W events is determined using a χ^2 criterion. In practice, weighting factors take into account unavoidable differences between the W and Z samples, such as the acceptance for the second lepton, photon radiation, and differences in η and p_T of W and Z bosons. Thus perfect agreement of the distributions at the nominal W mass and for the simulated detector is ensured, while systematic effects are studied by introducing distortions of experimental or theoretical origin. The resulting shifts in the extracted W mass are taken as the related systematic uncertainties.

The dominant systematic error arises from scale and resolution uncertainties in the missing energy determined from the calorimeters. These can be controlled by using the Z sample, where the boson p_T can be measured from the two charged leptons, as is shown in Figure 9.10. The observed differences of 2% on the scale and 5% on the resolution are taken as the systematic uncertainties.




Figure 9.10: Left: x -component of the calibrated missing transverse energy in the calorimeters using the reconstructed muon p_T in Z events, as a function of the transverse W boson momentum at generator level. The slope of a fitted straight line is 0.98. Right: Difference between the reconstructed missing energy in the calorimeters and the measured muon p_T in Z events (red/grey line) or the W boson p_T at generator level (black dashed line). The RMS of the distribution is 8.15 GeV for Z events and 8.65 GeV for W events.

9.3.5 Expected precision and systematic uncertainties

The expected size of various detector effects for the early detector operation, after the analysis of an initial integrated luminosity of 1 fb^{-1} , and for a better detector understanding expected after employing an integrated luminosity of 10 fb^{-1} , is shown in Table 9.7 for the scaled p_T -lepton method applied to the electron channel, and for the muon channel using the transformation method.

The measurements of the W mass by means of $W \rightarrow e\nu$ and $W \rightarrow \mu\nu$ decays are largely independent. Common experimental uncertainties arise from the systematics involving the missing transverse energy in the calorimeters.

Based on the estimated systematic errors, it is clear that the scaled p^T -lepton method suffers less from experimental systematic errors than the transformation method. If systematic uncertainties arising from the theoretical prediction of the transverse momenta of the Z and W bosons can be brought to a level of $\approx 10 \text{ MeV}/c^2$, the scaled p^T -lepton method is clearly the first choice. Using the scaled p_T -lepton method in the muon channel leads to a better statistical precision of $30 \text{ MeV}/c^2$ for 1 fb^{-1} due to the higher acceptance for muons compared to electrons. The total instrumental uncertainty of the p_T -lepton method applied to the muon channel is estimated from the findings in the electron channel and amounts to about $25 \text{ MeV}/c^2$ for the initial measurement with an integrated luminosity of 1 fb^{-1} . Uncertainties due to the recoil modelling are fully correlated with the electron channel. The component of the experimental error in common with the electrons amounts to about $20 \text{ MeV}/c^2$. Clearly, all theoretical uncertainties are of similar size and also correlated between the electron and muon channels.

The transformation method has the advantage of providing templates for observables in W events from measured observables in Z events. In particular, the measurement of the

Table 9.7: Expected systematic uncertainties on M_W for the scaled E^T -lepton method with electrons (upper part) and for the Z transformation method applied to the muon channel (lower part). The first column lists the systematic effect considered, the second and third columns show the assumed detector uncertainty for an initial integrated luminosity of 1 fb^{-1} and the resulting uncertainty on M_W . The last two columns show the extrapolation to an integrated luminosity of 10 fb^{-1} , when the detector understanding is assumed to have significantly improved.

Source of uncertainty	uncertainty with 1 fb^{-1}	ΔM_W [MeV/ c^2]	uncertainty with 10 fb^{-1}	ΔM_W [MeV/ c^2]
scaled lepton- p_T method applied to $W \rightarrow e\nu$				
statistics		40		15
background	10%	10	2%	2
electron energy scale	0.25%	10	0.05%	2
scale linearity	0.00006/ GeV	30	<0.00002/ GeV	<10
energy resolution	8%	5	3%	2
MET scale	2%	15	<1.5%	<10
MET resolution	5%	9	<2.5%	< 5
recoil system	2%	15	<1.5%	<10
total instrumental		40		<20
PDF uncertainties		20		<10
Γ_W		15		<15
p_T^W		30		30 (or NNLO)
transformation method applied to $W \rightarrow \mu\nu$				
statistics		40		15
background	10%	4	2%	negligible
momentum scale	0.1%	14	<0.1%	<10
$1/p^T$ resolution	10%	30	<3%	<10
acceptance definition	η -resol.	19	< σ_η	<10
calorimeter E_T^{miss} , scale	2%	38	$\leq 1\%$	<20
calorimeter E_T^{miss} , resolution	5%	30	<3%	<18
detector alignment		12	–	negligible
total instrumental		64		<30
PDF uncertainties		≈ 20		<10
Γ_W		10		< 10

transverse momentum of Z bosons and the cross checks on the modelling of the missing energy are of vital importance to quantify systematic uncertainties.

The combination of the electron and muon channels brings the statistical uncertainty to a final precision of better than $10 \text{ MeV}/c^2$ for an integrated luminosity of 10 fb^{-1} , and a systematic uncertainty of instrumental origin below $20 \text{ MeV}/c^2$ should be within reach.

Multi-boson production

FIXME: particle names are different here from rest of SM section Z^0 instead of Z . Needs anyway global PTDR fix.

9.4.1 Introduction

The study of multiple gauge-boson production at the TeV scale constitutes a unique opportunity to test the Standard Model of Electroweak interactions at the highest possible ener-

gies. The production of $W^\pm Z^0$ and $W^\pm \gamma$ events at the LHC probes the triple gauge-boson couplings and therefore the non-Abelian gauge symmetry of the Standard Model. On the other hand, no neutral gauge-boson couplings exist in the Standard Model, thus anomalies in $Z^0 Z^0$ and $Z^0 \gamma$ production, hinting at large s -channel contributions, could be the first indirect manifestation of New Physics. In the following, the selections of $W^\pm Z^0$ and $Z^0 Z^0$ events are described, their signal-over-background ratio discussed and the outlook for an early measurement of multiple gauge-boson production is assessed. Further details are given in Reference [356].

The multi-lepton final states of multiple gauge-boson production are an important background in the search for New Physics, in particular Supersymmetry. A sound understanding of their production process is therefore needed in the first phase of LHC data-taking before any discovery can be claimed. In particular, $Z^0 Z^0$ production is an irreducible background to the most-coveted discovery at the LHC: the Standard Model Higgs boson. Its early measurement is therefore important.

The cross sections for multiple gauge-boson production at the LHC are of about 50 pb for the $W^\pm Z^0$ channel and 20 pb for the $Z^0 Z^0$ channel [149]. These large cross sections and the clean signature of fully-leptonic final states make $W^\pm Z^0$ and $Z^0 Z^0$ production observable in the early LHC data. Final states where the gauge bosons decay into electrons and muons are considered: $e^\pm e^+ e^-$, $\mu^\pm e^+ e^-$, $e^\pm \mu^+ \mu^-$ and $\mu^\pm \mu^+ \mu^-$ for $W^\pm Z^0$ production and $e^+ e^- e^+ e^-$ for the $Z^0 Z^0$ channel. The competing background processes are the Standard Model production of gauge bosons and top quarks, which also yield leptonic final states.

9.4.2 Signal definition and modeling

Both the $W^\pm Z^0$ and $Z^0 Z^0$ analyses focus on on-shell gauge bosons. On-shell production of the $W^\pm Z^0$ final state proceeds mainly through the s -channel, involving a WWZ triple gauge-boson coupling. Additional contributions from the $W^\pm \gamma^*$ final state through a $WW\gamma$ coupling are effectively suppressed by constraining the mass of the observed lepton pair to be compatible with a Z^0 boson. The PYTHIA Monte Carlo generator [30] is used to model $W^\pm Z^0$ production and subsequent decay into fully-leptonic final states. Gauge-boson decays into tau leptons are also included. These tau leptons are left free to decay into either leptons or hadrons.

Four-electron final-states can originate from $Z^0 Z^0$ production as well as via either $Z^0 \gamma^*$ or $\gamma^* \gamma^*$ production. The requirement of on-shell boson is enforced by considering only electron-positron pairs with a mass between 70 and 110 GeV/ c^2 . The PYTHIA Monte Carlo is used to generate events of this process, with the additional requirement that the electrons have a rapidity $|\eta| < 2.7$ and a transverse momentum $p_T > 5$ GeV/ c . Of all generated events, 72% are classified as $Z^0 Z^0$ signal while 26% are ascribed to the $Z^0 \gamma^*$ process and 2% to the $\gamma^* \gamma^*$ process.

Taking into account the branching fraction into leptons, \mathcal{B} , and the kinematic requirements, ϵ_{KIN} , the relevant NLO cross sections using the MCFM [48] Monte Carlo are:

$$\begin{aligned} \sigma_{NLO} \times \mathcal{B} \times \epsilon_{KIN}(pp \rightarrow W^+ Z^0 \rightarrow \ell^+ \ell^+ \ell^-) &= 1034 \text{ fb} \\ \sigma_{NLO} \times \mathcal{B} \times \epsilon_{KIN}(pp \rightarrow W^- Z^0 \rightarrow \ell^- \ell^+ \ell^-) &= 630 \text{ fb} \\ \sigma_{NLO} \times \mathcal{B} \times \epsilon_{KIN}(pp \rightarrow Z^0 Z^0 \rightarrow e^+ e^- e^+ e^-) &= 18.7 \text{ fb} \end{aligned}$$

The NLO corrections correspond to k -factors of 1.9 and 1.4 for $W^\pm Z^0$ and $Z^0 Z^0$ production,

respectively. The NNLO box-diagram contribution to $Z^0 Z^0$ production is not taken into account.

Three-lepton final-states from $W^\pm Z^0$ and $Z^0 Z^0$ production are collected with high efficiency by the L1 and HLT electron and muon triggers. The L1 and HLT efficiencies for events retained by the selections discussed below is 100% [68].

9.4.3 Background processes

The background to the selection of $W^\pm Z^0$ and $Z^0 Z^0$ events comprises other processes with multiple leptons in the final states, some of which might be due to fake signals. The most copious sources of multiple leptons at the LHC are $t\bar{t}$ and $Z^0 b\bar{b}$ production. The cross section of these processes is large: 830 pb and 1492 pb, respectively, as calculated with MCFM at NLO. These processes may have two leptons in the final states from leptonic decays of the W bosons arising from $t \rightarrow Wb$ decays or of the Z^0 boson, respectively. The other leptons can be produced in the direct or cascade decays of the b quarks. The $Z^0 b\bar{b}$ process is modeled with the COMPHEP Monte Carlo generator [47, 34] and the $t\bar{t}$ process with the TOPREX Monte Carlo program [35]. In addition, the special case in which four electrons are produced in $t\bar{t}$ events is considered in detail and modeled with PYTHIA. Contributions from Wt and $Zc\bar{c}$ to the selected samples are negligible.

Events from $Z^0 Z^0$ production also constitute a background to the $W^\pm Z^0$ selection. Events from the $Z^0 \gamma^*$ and $\gamma^* \gamma^*$ processes are a background for both the $W^\pm Z^0$ and $Z^0 Z^0$ analyses.

9.4.4 $W^\pm Z^0$ selection

Events with three charged leptons, either electrons or muons, with $p_T > 10 \text{ GeV}/c$ and $|\eta| < 2.5$, are considered by the $W^\pm Z^0$ selection. All possible Z^0 -boson candidates from same-flavours opposite-charge lepton pairs are formed. Events are retained if the mass of the Z^0 candidate is within $20 \text{ GeV}/c^2$ of the Z^0 -boson mass, m_Z . These criteria effectively suppress Z^0 decays into tau leptons. The background from $Z^0 Z^0$ final states is reduced by rejecting events with a second Z^0 candidate with a mass within $40 \text{ GeV}/c^2$ of m_Z . The remaining lepton is associated to the W^\pm -boson decay; its transverse momentum must be larger than 20 GeV . This criterion results in lower efficiencies for the W^\pm boson decays in tau leptons. The highest- p_T lepton associated to the Z^0 boson must satisfy $p_T > 15 \text{ GeV}/c$. If the event contains more than three leptons, the lepton with highest p_T is chosen as originating from the W^\pm . The signal efficiency after these cuts is 9.2% while the $t\bar{t}$, $e^+e^-b\bar{b}$ and $\mu^+\mu^-b\bar{b}$ efficiencies are 0.7%, 0.4% and 0.6%, respectively.

Leptons from the decay of b quarks in the background processes are produced in a higher-multiplicity environment and isolation criteria suppress the background contamination. Electrons associated to the W^\pm boson must have no other charged track with $p_T > 2 \text{ GeV}/c$ within a $\Delta R = 0.3$ cone around their direction. All muon candidates must have an energy measured in the calorimeters within a $\Delta R = 0.3$ cone around their direction smaller than 5 GeV and the sum of the p_T of tracks within a $\Delta R = 0.25$ cone smaller than $2 \text{ GeV}/c$. The significance of the lepton impact parameter in the plane transverse to the beam, S_{IP} , discriminates against leptons from heavy-quark decays. This variable is defined as the ratio between the measured impact parameter and its uncertainty and is required to satisfy $S_{IP} < 3$. The signal efficiency after these cuts is 7.3% while the $t\bar{t}$, $e^+e^-b\bar{b}$ and $\mu^+\mu^-b\bar{b}$ efficiencies are 0.07%, 0.008% and 0.03%, respectively.

The $t\bar{t}$ and $Z^0 b\bar{b}$ final states are associated with one or more hard jets and their contribution is reduced by removing events containing at least a jet with $E_T > 25$ GeV. Only jets outside cones of $\Delta R = 0.3$ around the three leptons are considered. The reconstructed mass of the Z^0 boson is required to be within $10 \text{ GeV}/c^2$ of m_Z , leading to the total efficiencies presented in Table 9.8.

9.4.5 $Z^0 Z^0$ selection

The $Z^0 Z^0$ selection is based on events with four electrons, identified from superclusters in the electromagnetic calorimeter matched with a charged track. The transverse momenta of the electron candidates, ordered from the largest to the smallest, have to be above $30 \text{ GeV}/c$, $20 \text{ GeV}/c$, $15 \text{ GeV}/c$ and $10 \text{ GeV}/c$, respectively. This cut suppresses the contribution from the $Z^0 \gamma^*$ and $\gamma^* \gamma^*$ final states and reduces by 30% and 60% the $t\bar{t}$ and $Z^0 b\bar{b}$ backgrounds, respectively. Leptons from b quarks decays in the $t\bar{t}$ and $Z^0 b\bar{b}$ background processes are produced in association with hadrons. Their contribution is reduced by requiring the electrons to be isolated: the ratio between the energy deposited in the hadronic and the electromagnetic calorimeters must be below 8%; no more than two other charged track with $p_T > 2 \text{ GeV}/c$ must be within a $\Delta R = 0.3$ cone around the electron; $\sum_i (p_{T,i} - E_{T,i})/E_T < 0.34$, where E_T is the transverse energy of the electron candidate and the sum runs on all tracks with $p_T > 2 \text{ GeV}/c$ within a $\Delta R = 0.3$ cone around the electron.

Electron-positron pairs are combined to form Z^0 candidates. Pairs with reconstructed masses between 50 and $120 \text{ GeV}/c^2$ are retained. Of the two possible $Z^0 Z^0$ pairings, the one where the Z^0 candidate masses are closest to m_Z is chosen. This pairing is correct for almost all events with two on-shell Z^0 bosons. For 2.5% of the events, more than four electrons are present and only the $Z^0 Z^0$ pairing which contains the highest- p_T electron is retained. Table 9.9 presents the signal and background selection efficiencies.

Table 9.8: Yield of the $W^\pm Z^0$ selection for an integrated luminosity of 1 fb^{-1} . Signal efficiencies include gauge-boson decays into tau leptons.

	$e^\pm e^+ e^-$	$\mu^\pm e^+ e^-$	$e^\pm \mu^+ \mu^-$	$\mu^\pm \mu^+ \mu^-$	Total	Efficiency
$W^\pm Z^0 \rightarrow \ell^\pm \ell^+ \ell^-$	14.8	26.9	28.1	27.0	96.8	6.1%
$Z^0 Z^0$	0.63	1.54	1.50	1.51	5.18	4.7%
$t\bar{t}$	0.93	1.55	–	0.31	2.79	0.02%
$\mu^+ \mu^- b\bar{b}$	–	–	6.54	4.9	11.4	0.005%
$e^+ e^- b\bar{b}$	1.21	1.82	–	–	3.03	0.005%

Table 9.9: Yield of the $Z^0 Z^0$ selection for integrated luminosities of 1 fb^{-1} and 10 fb^{-1} . The last row indicates the signal significance, which include systematic effects.

	Efficiency	$N_{\text{events}}/1 \text{ fb}^{-1}$	$N_{\text{events}}/10 \text{ fb}^{-1}$
$Z^0 Z^0$	38%	7.1	71.1
$Z^0 \gamma^*$	4.5%	0.16	1.60
$Z^0 b\bar{b}$	0.07%	0.08	0.84
$t\bar{t}$	0.06%	0.12	1.22
S_L		4.8	13.1

9.4.6 Systematic uncertainties

For the first 1 fb^{-1} of integrated luminosity, the total systematic uncertainties on the $W^\pm Z^0$ and $Z^0 Z^0$ cross section measurements are 17.4% and 12.9%, respectively. These figures include a 10% uncertainties on the determination of the integrated luminosity.

The most important sources of systematic uncertainties are lepton identification and isolation, and background subtraction. A 2% uncertainty on the efficiency of each lepton propagates to an uncertainty on the cross section between 2.6% and 7.8%, according to the channel. Background subtraction dominates the $W^\pm Z^0$ systematics with an uncertainty of 12%, while it accounts for a 1.3% uncertainty in the $Z^0 Z^0$ channel. An additional uncertainty of 5% on the jet energy scale affects the $W^\pm Z^0$ channel, while an uncertainty of 1% on the trigger efficiency affects both channels.

The significance of the observation of the $W^\pm Z^0$ and $Z^0 Z^0$ signals in the first 1 fb^{-1} is not sensitive to the luminosity uncertainty. It is affected by all other sources of systematic uncertainty listed above, with a total effect of 14.8% and 14.2% on the two channels, respectively. These uncertainties include additional PDF and QCD uncertainties in the Monte Carlo modeling, contributing 3.7% and 6.4% for the $W^\pm Z^0$ and $Z^0 Z^0$ selections, respectively.

9.4.7 Results

Figure 9.11 left presents the mass distribution of the Z^0 candidates in the $W^\pm Z^0$ channel for an integrated luminosity of 1 fb^{-1} before the last requirement of a $\pm 10 \text{ GeV}/c^2$ window is applied. A large signal-over-background ratio is observed, as shown in Table 9.8.

Figure 9.11 right shows the mass distribution of the Z^0 candidates, two entries per event, selected by the $Z^0 Z^0$ selection for an integrated luminosity of 10 fb^{-1} . Table 9.9 lists the selection yield for 1 fb^{-1} and 10 fb^{-1} . The selection results into an almost background-free signal sample, which will constitute a valuable input to assess the background in the search for the Higgs boson.

Both the $W^\pm Z^0$ and $Z^0 Z^0$ final states can be selected with high purity. A significance of 12.8 and 4.8, respectively, is expected in the first 1 fb^{-1} of integrated luminosity, including systematic uncertainties. The $W^\pm Z^0$ channel can be observed with a significance of 5, including systematic effects, in an integrated luminosity of 150 pb^{-1} .

This study of multiple gauge-boson production and couplings at the LHC will be extended to include the $W^\pm \gamma$ and $Z^0 \gamma$ channels, as well as the other flavours of $Z^0 Z^0$ fully-leptonic decays.

In conclusion, the large signal-over-background ratios achieved by the $W^\pm Z^0$ and $Z^0 Z^0$ selections suggest that early observation of these channels will take place at the LHC start up. In addition, precise investigations of triple gauge-boson couplings will be possible with the first 10 fb^{-1} of LHC data.




Figure 9.11: Left: Distribution of the mass of the Z^0 candidates for events retained by the $W^\pm Z^0$ selection, for an integrated luminosity of 1 fb^{-1} . Right: Distribution of the mass of the Z^0 candidates, two entries per event, retained by the $Z^0 Z^0$ selection, for an integrated luminosity of 10 fb^{-1} .

Chapter 10

Standard Model Higgs Bosons



10.1 Introduction

The Higgs mechanism is a cornerstone of the Standard Model (SM) and its supersymmetric extensions. The introduction of the fundamental Higgs field [357, 358, 359, 360] renders the standard electroweak theory weakly interacting up to high energy scales without violating the unitarity bounds of scattering amplitudes [361, 362, 363, 364]. Due to spontaneous symmetry breaking in the Higgs sector the electroweak gauge bosons W, Z as well as the fermions acquire masses through the interaction with the Higgs fields. Since the gauge symmetry, though hidden, is still preserved, the theory of electroweak interactions is renormalizable [365, 366, 367, 368, 369]. In the Standard Model one weak isospin Higgs doublet is introduced and leads to the existence of one elementary Higgs particle after electroweak symmetry breaking. The Higgs couplings to the electroweak gauge bosons and all fermions grow with their masses. The only unknown parameter of the Higgs boson itself is the value of its mass M_H . Once this is known, all production and decay properties of the SM Higgs boson will be fixed [8, 370, 371]. The search for the Higgs boson will be a crucial endeavour at present and future experiments for establishing the standard formulation of the electroweak theory.

Although the Higgs mass cannot be predicted in the Standard Model, there are several constraints deduced from consistency conditions on the model [372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382]. Upper bounds can be derived from the requirement that the Standard Model can be extended up to a scale Λ , before perturbation theory breaks down and new non-perturbative phenomena dominate the predictions of the theory. If the SM is required to be weakly interacting up to the scale of grand unified theories (GUTs), which is of $\mathcal{O}(10^{16} \text{ GeV})$, the Higgs mass has to be less than $\sim 190 \text{ GeV}$. For a minimal cut-off $\Lambda \sim 1 \text{ TeV}$ a universal upper bound of $\sim 700 \text{ GeV}$ can be obtained from renormalization group analyses [372, 373, 374, 375, 376, 377, 378, 379] and lattice simulations of the SM Higgs sector [380, 381, 382]. This issue can be rephrased by stating that the Higgs sector has to be trivial, if the cut-off is extended to arbitrary magnitudes. Triviality means the absence of Higgs self-interactions.

If the top quark mass is large, the Higgs self-coupling can become negative and the Higgs potential deeply negative, thus rendering the SM vacuum unstable. The negative contribution of the top quark, however, can be compensated by a positive contribution due to the Higgs self-interaction, which is proportional to the Higgs mass. For a given top mass $m_t = 175 \text{ GeV}$ a lower bound of $\sim 60 \text{ GeV}$ can be obtained for the Higgs mass, if the SM remains weakly interacting up to scales $\Lambda \sim 1 \text{ TeV}$. For $\Lambda \sim M_{GUT}$ this lower bound is enhanced

to $M_H \gtrsim 130$ GeV. However, the assumption that the vacuum is metastable, with a lifetime larger than the age of the Universe, decreases these lower bounds significantly for $\Lambda \sim 1$ TeV, but only slightly for $\Lambda \sim M_{GUT}$ [379].

The direct search in the LEP2 experiments via the process $e^+e^- \rightarrow ZH$ yields a lower bound of 114.4 GeV on the Higgs mass [54]. After LEP2 the search for the SM Higgs particle is continued at the Tevatron for Higgs masses up to ~ 130 GeV [383] and the LHC for Higgs masses up to the theoretical upper limit [384, 385].

The Higgs decay modes can be divided into two different mass ranges. For $M_H \lesssim 135$ GeV the Higgs boson mainly decays into $b\bar{b}$ and $\tau^+\tau^-$ pairs with branching ratios of about 85% and 8% respectively (see Fig. 10.1, right plot). The decay modes into $c\bar{c}$ and gluon pairs, with the latter mediated by top and bottom quark loops, accumulate a branching ratio of up to about 10%, but do not play a relevant role at the LHC. The QCD corrections to the Higgs decays into quarks are known up to three-loop order [386, 387, 388, 389, 390, 391, 392] and the electroweak corrections up to NLO [393, 394, 395, 396]. The latter are also valid for leptonic decay modes. One of the most important Higgs decays in this mass range at the LHC is the decay into photon pairs, which is mediated by W , top and bottom quark loops. It reaches a branching fraction of up to 2×10^{-3} . The NLO QCD [397, 398, 399, 400, 401, 402, 403] and electroweak [404, 405, 406] corrections are known. They are small in the Higgs mass range relevant for the LHC. For Higgs masses above 135 GeV the main decay modes are those into WW and ZZ pairs, where one of the vector bosons is off-shell below the corresponding kinematical threshold. These decay modes dominate over the decay into $t\bar{t}$ pairs, the branching ratio of which does not exceed $\sim 20\%$ as can be inferred from Fig. 10.1 (right plot). The electroweak corrections to the WW, ZZ decays are of moderate size [393, 407, 408, 394]. The total decay width of the Higgs boson, shown in Fig. 10.1 (left plot), does not exceed about 1 GeV below the WW threshold. For very large Higgs masses the total decay width grows up to the order of the Higgs mass itself so that the interpretation of the Higgs boson as a resonance becomes questionable. This Higgs mass range coincides with the upper bound of the Higgs mass from triviality.

The dominant Higgs production mechanism at the LHC will be the gluon-fusion process [409]

$$pp \rightarrow gg \rightarrow H,$$

which provides the largest production cross section for the whole Higgs mass range of interest. This process is mediated by top and bottom quark loops (see Fig. 10.2a). Due to the large size of the top Yukawa couplings and the gluon densities gluon fusion comprises the dominant Higgs boson production mechanism for the whole Higgs mass range.

The QCD corrections to the top and bottom quark loops are known since a long time including the full Higgs and quark mass dependences [410, 411, 412]. They increase the total cross section by 50 – 100%. The limit of very heavy top quarks provides an approximation within $\sim 10\%$ for all Higgs masses [8, 370, 371, 410, 411, 412, 413]. In this limit the NLO QCD corrections have been calculated before [410, 411, 412, 414, 415, 416, 417] and recently the NNLO QCD corrections [418, 419, 420, 421] with the latter increasing the total cross section further by $\sim 20\%$. A full massive NNLO calculation is not available so far, so that the NNLO results can only be trusted for small and intermediate Higgs masses. The approximate NNLO results have been improved by a soft-gluon resummation at the next-to-next-to-leading log (NNLL) level, which yields another increase of the total cross section by $\sim 10\%$ [422]. Electroweak corrections have been computed, too, and turn out to be small




Figure 10.1: Left plot: total decay width (in GeV) of the SM Higgs boson as a function of its mass. Right plot: Branching ratios of the dominant decay modes of the SM Higgs particle. relevant higher-order corrections are taken into account




Figure 10.2: Typical diagrams for all relevant Higgs boson production mechanisms at leading order: (a) gluon fusion, (b) vector boson fusion, (c) Higgs-strahlung, (d) Higgs bremsstrahlung off top quarks.

[423, 424, 425, 426, 404]. The theoretical uncertainties of the total cross section can be estimated as $\sim 20\%$ at NNLO due to the residual scale dependence, the uncertainties of the parton densities and due to neglected quark mass effects.

At LO the Higgs boson does not acquire any transverse momentum in the gluon fusion process, so that Higgs bosons with non-vanishing transverse momentum can only be produced in the gluon fusion process, if an additional gluon is radiated. This contribution is part of the real NLO corrections to the total gluon fusion cross section. The LO p_T distribution of the Higgs boson is known including the full quark mass dependence [427, 428]. The NLO cor-

rections, however, are only known in the heavy quark limit, so that they can only be trusted for small and moderate Higgs masses and p_T [429, 430, 431, 432, 433]. In this limit a NLL soft gluon resummation has been performed [434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444], which has recently been extended to the NNLL level [445, 446, 447, 448, 449] thus yielding a reliable description of the small p_T range. It should be noted that these results are only reliable, if the top quark loops provide the dominant contribution and p_T is not too large. In the regions where the NLO and resummed results are valid the theoretical uncertainties have been reduced to $\mathcal{O}(20\%)$.

For large Higgs masses the W and Z boson-fusion processes [450, 451, 452] (see Fig. 10.2b)

$$pp \rightarrow qq \rightarrow qq + WW/ZZ \rightarrow qqH$$

become competitive. These processes are relevant in the intermediate Higgs mass range, too, since the additional forward jets offer the opportunity to reduce the background processes significantly. Since at NLO there is no color exchange between the two quark lines, the NLO QCD corrections can be derived from the NLO corrections to deep inelastic lepton-nucleon scattering. They turn out to be $\mathcal{O}(10\%)$ for the total cross section [453, 8, 370, 371]. Quite recently the NLO corrections to the differential cross sections have been computed, too, resulting in modifications of the relevant distributions by up to $\sim 30\%$ [454]. The residual uncertainties are of $\mathcal{O}(5\%)$.

In the intermediate mass range $M_H \lesssim 2M_Z$ Higgs-strahlung off W, Z gauge bosons [455, 456] (see Fig. 10.2c)

$$pp \rightarrow q\bar{q} \rightarrow Z^*/W^* \rightarrow H + Z/W$$

provides alternative signatures for the Higgs boson search. Since only the initial state quarks are strongly interacting at LO, the NLO QCD corrections can be inferred from the Drell-Yan process. They increase the total cross section by $\mathcal{O}(30\%)$ [457, 8, 370, 371]. Recently this calculation has been extended up to NNLO [458]. The NNLO corrections are small. Moreover, the full electroweak corrections have been obtained in Ref. [459] resulting in a decrease of the total cross sections by 5 – 10%. The total theoretical uncertainty is of $\mathcal{O}(5\%)$.

Higgs radiation off top quarks (see Fig. 10.2d)

$$pp \rightarrow q\bar{q}/gg \rightarrow Ht\bar{t}$$

plays a significant role for smaller Higgs masses below ~ 150 GeV. The LO cross section has been computed a long time ago [460, 461, 462, 463, 464]. During the last years the full NLO QCD corrections have been calculated resulting in a moderate increase of the total cross section by $\sim 20\%$ at the LHC [465, 153, 466]. These results confirm former estimates based on an effective Higgs approximation [467]. The effects on the relevant parts of final state particle distribution shapes are of moderate size, i.e. $\mathcal{O}(10\%)$, too, so that former experimental analyses are not expected to alter much due to these results. All SM Higgs production cross sections including NLO QCD corrections are shown in Fig. 10.3.

2 Higgs boson channels

10.2.1 Inclusive Higgs boson production with $H \rightarrow ZZ^{(*)} \rightarrow e^+e^-\mu^+\mu^-$




Figure 10.3: Higgs production cross sections at the LHC for the various production mechanisms as a function of the Higgs mass. The full QCD-corrected results for the gluon fusion $gg \rightarrow H$, vector-boson fusion $qq \rightarrow VVqq \rightarrow Hqq$, vector-boson bremsstrahlung $q\bar{q} \rightarrow V^* \rightarrow HV$ and associated production $gg, q\bar{q} \rightarrow Ht\bar{t}$ are shown.

10.2.1.1 Introduction

The $H \rightarrow ZZ^{(*)} \rightarrow 4\ell$ channel has a very clean signature with relatively small backgrounds and is therefore an important discovery channel for the Higgs boson for a large range of masses. This channel is also important for the measurement of the mass and width of the Higgs boson.

10.2.1.2 Event Generation

All Monte Carlo event samples used in the analysis were generated using the PYTHIA [62] event generator, except for the $Zb\bar{b}$ ($e^+e^-b\bar{b}$ and $\mu^+\mu^-b\bar{b}$) background samples which were generated with CompHEP [47].

Higgs-boson production was simulated through leading order gluon-gluon scattering and vector-boson fusion. Monte Carlo samples were produced for 18 values of the Higgs boson mass m_H ranging from 115 GeV/c^2 to 200 GeV/c^2 in 10 GeV/c^2 steps, and from 200 GeV/c^2 to 600 GeV/c^2 in 50 GeV/c^2 steps.

Three background processes which yield the same signature of two electrons and two muons in the final state, with significant cross-section times branching ratio, are considered:

1. $q\bar{q}/gg \rightarrow t\bar{t} \rightarrow W^+W^-b\bar{b} \rightarrow e^+e^-\mu^+\mu^-X$
2. $q\bar{q}/gg \rightarrow Zb\bar{b} \rightarrow e^+e^-\mu^+\mu^-X$
3. $q\bar{q} \rightarrow (Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*) \rightarrow e^+e^-\mu^+\mu^-$

For the $t\bar{t}$ and $Zb\bar{b}$ backgrounds, no restrictions are applied on b decays prior to the preselection. Only events with $|\eta_b| < 2.5$ were generated for the $Zb\bar{b}$ background. For the $Zb\bar{b}$ and $(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$ backgrounds, m_{γ^*} is required to be greater 5 GeV/c^2 .

For the $(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$ background, only the t -channel production through $q\bar{q}$ fusion is simulated. In order to account for contributions from all NLO diagrams and from the NNLO gluon fusion ($gg \rightarrow (Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$), all events are re-weighted at analysis level with an $m_{4\ell}$ dependent K-factor, calculated [468][469] using MCFM.

The potential background contribution from $Zc\bar{c} \rightarrow e^+e^-\mu^+\mu^-X$ was also investigated using fully simulated events and was shown to be negligible.

For all Monte Carlo samples, a preselection is applied at generator level with the following requirements:

1. Final state contains $e^+e^-\mu^+\mu^-$.
2. $p_T(e) > 5 \text{ GeV}/c$ and $|\eta(e)| < 2.5$ for both electrons
3. $p_T(\mu) > 3 \text{ GeV}/c$ and $|\eta(\mu)| < 2.4$ for both muons

The cross-section times branching ratio and the cross-section times branching ratio times preselection efficiency, are shown for the signal as a function of m_H in Figure 12.4. The NLO cross-section and the cross-section times branching ratio times preselection efficiency shown for each background process in Table 10.1.




Figure 10.4: Cross-section times branching ratio, and cross-section times branching ratio times preselection efficiency for $H \rightarrow ZZ^{(*)} \rightarrow 2e2\mu$.

Table 10.1: NLO cross-section and the cross-section times branching ratio times preselection efficiency for the three background process.

Process	σ_{NLO} (pb)	$\sigma_{\text{NLO}} \times \text{BR} \times \epsilon$ (fb)
$t\bar{t} \rightarrow W^+W^-b\bar{b} \rightarrow e^+e^-\mu^+\mu^-X$	840	744
$e^+e^-b\bar{b} \rightarrow e^+e^-\mu^+\mu^-X$	276	262
$\mu^+\mu^-b\bar{b} \rightarrow e^+e^-\mu^+\mu^-X$	279	128
$(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*) \rightarrow e^+e^-\mu^+\mu^-$	28.9	37.0

10.2.1.3 Online Selection

Events selected by the di-muon or the di-electron triggers are considered. This choice follows from the presence of an on-shell Z-boson in most events. The additional use of single-electron and single-muon triggers does not increase the significance of the results.

The efficiencies of the Level-1 and High Level Triggers are shown for the signal as a function of m_H in Figure 10.5. The corresponding trigger efficiencies for background processes are given in Table 10.2.




Figure 10.5: Efficiency of the Level-1 and High Level Triggers for the Higgs signal. Monte Carlo Statistical uncertainties are shown.

Table 10.2: Efficiency of the Level-1 and High Level Triggers for each of the three background processes. Monte Carlo Statistical uncertainties are shown.

	$t\bar{t}$	$Zb\bar{b}$	$(Z^{(*)}/\gamma^{*})(Z^{(*)}/\gamma^{*})$
Level-1 Trigger efficiency (%)	95.1 ± 0.1	92.3 ± 0.1	97.9 ± 0.2
HLT efficiency (%)	39.9 ± 0.1	65.8 ± 0.1	89.6 ± 0.4

10.2.1.4 Offline Event Selection

Offline reconstruction of electrons and muons is performed using standard algorithms. It is required that four leptons of type $e^+e^-\mu^+\mu^-$ are reconstructed.

The two largest backgrounds after the HLT, $t\bar{t}$ and $Zb\bar{b}$, are reducible, since unlike the Higgs signal, two of the leptons will be associated with b-jets and will therefore be displaced relative to the primary vertex and will not be isolated. These two considerations can be used to powerfully cut against these processes, whereas the $(Z^{(*)}/\gamma^{*})(Z^{(*)}/\gamma^{*})$ background is irreducible by such means. Kinematic cuts are then applied, which further reduce all three backgrounds.

Vertex and Impact Parameter

Three criteria are applied:

1. The transverse distance of the $\mu^+\mu^-$ vertex from the beam line is required to be less than 0.011 cm.
2. The three-dimensional distance between the $\mu^+\mu^-$ vertex and the e^+e^- vertex is required to be less than 0.06 cm.
3. The transverse impact parameter significance of all leptons required to be less than 7.

For events passing this selection, the primary vertex is reconstructed by performing a fit to the tracks of the four reconstructed leptons. The lepton tracks are then refitted using the reconstructed vertex position as an additional point, in order to obtain a more accurate measurement of the momentum at the primary vertex.

Isolation

A cut is applied on the sum of the p_T of reconstructed tracks with $p_T > 0.9 \text{ GeV}/c$ and at least five hits, which satisfy the following conditions:

1. The track lies within the region defined by the sum of cones of size $\Delta R = 0.25$ around each of the four leptons and lies outside veto cones of size $\Delta R = 0.015$ around each lepton.
2. The track is consistent with originating from the reconstructed primary vertex to within $|\Delta z| < 0.2 \text{ cm}$, where Δz is the difference between the z position of the point of closest approach of the track to the reconstructed vertex, and the z position of the reconstructed vertex.

Kinematic Cuts

The following kinematic cuts are applied:

1. Lower thresholds on the transverse momenta of each of the four reconstructed leptons.
2. Upper and lower thresholds on the invariant masses of the reconstructed e^+e^- and $\mu^+\mu^-$ pairs.
3. Upper and lower thresholds on the invariant mass of the four reconstructed leptons.

These kinematic thresholds, together with the threshold on Σp_T for tracker isolation are optimized simultaneously using MINUIT, such that the log-likelihood ratio:

$$S_L = \sqrt{2 \ln Q}, \quad \text{where} \quad Q = \left(1 + \frac{N_S}{N_B}\right)^{N_S+N_B} e^{-N_S} \quad (10.1)$$

is maximized. The optimization is performed separately for each Higgs mass.

10.2.1.5 Results

Tables 10.3 and 10.4 show the production cross-section, cross-section times branching ratio, cross-section times branching ratio times preselection efficiency and the cross-section times branching ratio times efficiency after each stage of the online and offline event selection, for Higgs masses of $140 \text{ GeV}/c^2$ and $200 \text{ GeV}/c^2$, respectively. Values are shown for signal and for each of the three background processes. For all values of m_H , the background after all selections is strongly dominated by $(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$. For low m_H $t\bar{t}$ and $Zb\bar{b}$ each contribute around 10-15% to the total residual background, whereas for $m_H > 200 \text{ GeV}/c^2$, $(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$ constitutes more than 99%.

Figure 10.6 shows the invariant mass of the four reconstructed leptons before and after the application of the offline selection, for signal events for $m_H = 130 \text{ GeV}/c^2$ (left) and $m_H = 200 \text{ GeV}/c^2$ (right), and for the three background processes.

Figure 10.7 shows the final cross-section times branching ratio times efficiency for selected events, for signal and background, as a function of the Higgs boson mass. The number of expected events passing all selections for 10 fb^{-1} of integrated luminosity is shown in Table 10.5 for several values of the Higgs boson mass.

Table 10.3: Production cross-section (NLO), cross-section times branching ratio, cross-section times branching ratio times preselection efficiency and cross-section times branching ratio times efficiency after each stage of the online and offline event selection, for $m_H=140 \text{ GeV}/c^2$, for signal and backgrounds. All values in fb, except for expected number of events. Uncertainties are statistical only.

	Signal	$t\bar{t}$	$Zb\bar{b}$	$(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$
Production cross-section (NLO)	33.6×10^3	840×10^3	555×10^3	28.9×10^3
$\sigma \times \text{BR}(4 \text{ lepton final state})$	11.6	-	-	367.5
Preselection: $\sigma \times \text{BR} \times \epsilon$	3.29 ± 0.04	743 ± 2	390 ± 1	37.0 ± 0.4
Level-1 trigger	3.24 ± 0.04	707 ± 2	360 ± 1	36.3 ± 0.4
High Level trigger	2.91 ± 0.03	282 ± 1	237 ± 1	32.5 ± 0.4
$e^+e^-\mu^+\mu^-$ reconstructed	2.23 ± 0.03	130 ± 1	141 ± 1	24.1 ± 0.3
Vertex and impact parameter cuts	2.01 ± 0.03	18.9 ± 0.3	18.4 ± 0.2	21.5 ± 0.3
Isolation cuts	1.83 ± 0.03	1.34 ± 0.07	5.8 ± 0.1	20.0 ± 0.3
Lepton p_T cuts	1.61 ± 0.03	0.40 ± 0.04	0.56 ± 0.03	17.6 ± 0.3
Z mass window cuts	1.35 ± 0.02	0.20 ± 0.03	0.23 ± 0.02	13.8 ± 0.3
Higgs mass window cuts	1.17 ± 0.02	0.02 ± 0.01	0.025 ± 0.007	0.15 ± 0.03
Expected events for $\int \mathcal{L} = 10 \text{ fb}^{-1}$	11.7 ± 0.2	0.2 ± 0.1	0.25 ± 0.07	1.5 ± 0.3

Table 10.4: Production cross-section (NLO), cross-section times branching ratio, cross-section times branching ratio times preselection efficiency and cross-section times branching ratio times efficiency after each stage of the online and offline event selection, for $m_H=200 \text{ GeV}/c^2$, for signal and backgrounds. All values in fb, except for expected number of events. Uncertainties are statistical only.

	Signal	$t\bar{t}$	$Zb\bar{b}$	$(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$
Production cross-section (NLO)	17.9×10^3	840×10^3	555×10^3	28.9×10^3
$\sigma \times \text{BR}(4 \text{ lepton final state})$	23.8	-	-	367.5
Preselection: $\sigma \times \text{BR} \times \epsilon$	7.39 ± 0.09	743 ± 2	390 ± 1	37.0 ± 0.4
Level-1 trigger	7.36 ± 0.09	707 ± 2	360 ± 1	36.3 ± 0.4
High Level trigger	6.82 ± 0.08	282 ± 1	237 ± 1	32.5 ± 0.4
$e^+e^-\mu^+\mu^-$ reconstructed	5.51 ± 0.07	130 ± 1	141 ± 1	24.1 ± 0.3
Vertex and impact parameter cuts	5.03 ± 0.07	18.9 ± 0.3	18.4 ± 0.2	21.5 ± 0.3
Isolation cuts	4.92 ± 0.07	5.1 ± 0.1	12.3 ± 0.2	21.3 ± 0.3
Lepton p_T cuts	4.78 ± 0.07	1.93 ± 0.09	1.78 ± 0.06	18.7 ± 0.3
Z mass window cuts	4.45 ± 0.07	0.15 ± 0.03	0.12 ± 0.02	14.4 ± 0.3
Higgs mass window cuts	3.64 ± 0.06	0.006 ± 0.005	0.006 ± 0.003	1.61 ± 0.09
Expected events for $\int \mathcal{L} = 10 \text{ fb}^{-1}$	36.4 ± 0.6	0.06 ± 0.05	0.06 ± 0.03	16.1 ± 0.9

Table 10.5: Expected number of events from signal and background processes after all selections for an integrated luminosity of 10 fb^{-1}

$m_H \text{ (GeV}/c^2)$	120	130	140	150	160	170	180	200	250	300	400	500
N signal for 10 fb^{-1}	1.9	4.6	11.7	14.1	7.8	3.8	8.7	36.4	29.1	19.4	18.0	9.6
N back for 10 fb^{-1}	1.5	0.6	2.0	2.1	2.0	2.9	4.0	16.2	13.6	4.1	3.7	2.6




Figure 10.6: Invariant mass of the four reconstructed leptons before (top) and after (bottom) the application of the offline selection, for signal events for $m_H = 130 \text{ GeV}/c^2$ (left) and $m_H = 200 \text{ GeV}/c^2$ (right), and for the three background processes.




Figure 10.7: Cross-section times branching ratio times efficiency after all selections

Significance

Figure 10.8 shows the S_{cP} significance after all selection cuts for integrated luminosities of 10 fb^{-1} and 30 fb^{-1} , with and without the systematic uncertainty on the background estimation taken into account. The background systematic uncertainty will be discussed in section 10.2.1.6. Figure 10.9 shows the integrated luminosity required to obtain a significance of 5σ using the $H \rightarrow ZZ^{(*)} \rightarrow 2e2\mu$ channel alone, with and without the background systematic uncertainty. It can be seen that a significance of 5σ can be achieved with less than 30 fb^{-1} of integrated luminosity for a Higgs boson with mass in the range $130 \leq m_H \leq 500 \text{ GeV}/c^2$, excluding a gap of about $15 \text{ GeV}/c^2$ close to $m_H = 170 \text{ GeV}/c^2$ for which close to 100 fb^{-1} is required. If the Higgs boson mass lies in the range $190 \leq m_H \leq 400 \text{ GeV}/c^2$, 5σ significance can be attained with less than 8 fb^{-1} of integrated luminosity.




Figure 10.8: ScP significance after all selection cuts for integrated luminosities of 10 fb^{-1} and 30 fb^{-1} , with and without the systematic uncertainty on the background estimation taken into account.




Figure 10.9: Integrated luminosity required to obtain a significance of 5σ using the $H \rightarrow ZZ^{(*)} \rightarrow 2e2\mu$ channel, with and without the systematic uncertainty on the background estimation taken into account.

10.2.1.6 Evaluation of Background from Data

The background normalization can be estimated from data by using the sidebands in the reconstructed four-lepton invariant mass distribution. Figure 10.10 shows the number of expected events from the signal and background Monte Carlo simulations for an integrated luminosity corresponding to a discovery significance of 5σ , for Higgs boson masses of 130 and 200 GeV/c^2 : 28.2 and 6.4fb^{-1} , respectively. Figure 10.10 also shows the results of a simulated experiment with these luminosities.




Figure 10.10: Number of expected events for signal and background for an integrated luminosity corresponding to a discovery significance of 5σ , for Higgs boson masses of 130 and 200 GeV/c^2 . The results of a simulated experiment are also shown to illustrate the statistical power of the analysis and the determination of the background normalization from data.

The number of background events measured from the data within the signal region, N_{Data}^{IN} , is calculated as:

$$N_{Data}^{IN} = \alpha_{MC} N_{Data}^{OUT}, \quad \text{where} \quad \alpha_{MC} = \frac{N_{MC}^{IN}}{N_{MC}^{OUT}} \quad (10.2)$$

N_{Data}^{OUT} is the number observed events lying outside the signal region and α_{MC} is the ratio of the number of background events inside the signal region (N_{MC}^{IN}) to outside the signal region (N_{MC}^{OUT}), as determined from the background Monte Carlos.

The uncertainty on the number of background events in the signal region measured using this method is given by:

$$\Delta B = \Delta B_{Stat} \oplus \Delta B_{Theory}, \quad \text{where} \quad \Delta B_{Stat} = \alpha \sqrt{N_{Data}^{OUT}}$$

ΔB_{Stat} provides the dominant contribution to the uncertainty. Taking N_{Data}^{OUT} as the expected number of events outside the signal region for an integrated luminosity corresponding to 5σ significance, the value ΔB_{Stat} varies between 2% and 13% for $m_H < 200 \text{ GeV}/c^2$ and increases to around 30% for high m_H where the statistics in the sidebands are low.

ΔB_{Theory} is the theoretical uncertainty on the shape of the $m_{4\ell}$ distribution for the $(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$ background. The value is taken from [468], which takes into account PDF and QCD scale uncertainties in the $(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$ production cross-section at NLO, and varies between 0.5 and 4.5% for the range Higgs boson masses considered.

10.2.1.7 Measurement of the properties of the Higgs Boson

The $H \rightarrow ZZ^{(*)} \rightarrow 4\ell$ channel can be used to evaluate the mass, width and production cross-section of the Higgs boson.

Mass Measurement

The statistical uncertainty on the Higgs boson mass measurement is given by $\Delta_{stat} = \sigma_{Gauss} / \sqrt{N_S}$, where σ_{Gauss} is the measured Gaussian width of the four lepton invariant mass peak from the signal Monte Carlo and N_S is the expected number of signal events passing all selections. The value, as a fraction of the true mass, is shown in Table 10.6, for an integrated luminosity and 30 fb^{-1} , as a function of m_H .

Width Measurement

Figure 10.11 shows the measured width of the Higgs boson mass peak, obtained from the Gaussian fit, as a function of m_H . The true width from theory Γ_H is also shown. The measured width is a convolution of the natural width and the experimental resolution. It can be seen that for m_H less than around $200 \text{ GeV}/c^2$, the measured width is completely dominated by the experimental resolution. The statistical uncertainty on the width measurement is given by $\Delta_{stat} = \sigma_{Gauss} / \sqrt{2N_S}$, where σ_{Gauss} is the measured Gaussian width of the peak and N_S is the expected number of signal events passing all selections. The value, as a fraction of the true width, is shown in Table 10.6, for an integrated luminosity and 30 fb^{-1} , as a function of m_H . The direct measurement the Higgs boson width is possible with $\Delta_{stat} < 30\%$ for $m_H \geq 200 \text{ GeV}/c^2$.




Figure 10.11: Measured width of the Higgs boson mass peak, obtained from a Gaussian fit to the peak, as a function of the true Higgs mass. The true width from theory is also shown.

Production Cross-Section Measurement

The Higgs boson production cross-section can be determined from the number of observed events N_{obs} after all selections, given the efficiency ϵ of the event selection and the integrated luminosity \mathcal{L} :

$$\sigma = \frac{N_{obs}}{\mathcal{L}\epsilon}$$

The total uncertainty on the cross-section measurement is given by:

$$\Delta\sigma^2 = \Delta_{stat}^2 + \Delta_{syst}^2 + \Delta\mathcal{L}^2 + \Delta B^2$$

where Δ_{stat} , Δ_{syst} , $\Delta\mathcal{L}$ and ΔB are the statistical uncertainty, the systematic uncertainty from the event selection, the uncertainty on the luminosity measurement and the background systematic uncertainty, respectively.

The statistical uncertainty Δ_{stat} is shown in Table 10.6 for an integrated luminosity 30 fb^{-1} , as a function of m_H .

The total systematic uncertainty arising from the offline reconstruction and event selection can be summarized as:

$$\Delta_{syst}^2 = 2\Delta\epsilon_e^2 + 2\Delta\epsilon_\mu^2 + \Delta\epsilon_{iso}^2$$

where $\Delta\epsilon_e$ is the uncertainty in the reconstruction efficiency for electrons, estimated to be around 1% per electron [470], $\Delta\epsilon_\mu$ is the uncertainty in the muon reconstruction efficiency, which has been shown to be measurable to be better than 1% per muon [468], and $\Delta\epsilon_{iso}$ is the uncertainty in the efficiency of the isolation cut, estimated in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ analysis [468] to be around 2% per event. This gives a total uncertainty $\Delta_{syst} = 3\%$.

The uncertainty on the measurement of the LHC luminosity $\Delta\mathcal{L}$ is expected to be around 3%. The background uncertainty ΔB is discussed in Section 10.2.1.6.

$m_H \text{ (GeV}/c^2)$	115	120	130	140	150	160	170	180	190
$\Delta_{Stat}(m_H)(\%)$	0.722	0.512	0.335	0.206	0.193	0.256	0.388	0.27	0.134
$\Delta_{Stat}(\Gamma_H)(\%)$	-	-	-	-	-	-	-	54.8	17.6
$\Delta_{Stat}(\sigma_H)(\%)$	75	55.6	28.6	18.2	16.5	23.1	39.2	23.7	11.5
$m_H \text{ (GeV}/c^2)$	200	250	300	350	400	450	500	550	600
$\Delta_{Stat}(m_H)(\%)$	0.145	0.207	0.328	0.408	0.588	0.896	1.25	1.62	2.43
$\Delta_{Stat}(\Gamma_H)(\%)$	14.4	7.38	8.2	5.43	5.8	5.91	6.52	6.61	8.36
$\Delta_{Stat}(\sigma_H)(\%)$	11.5	13	14.4	13.8	14.9	18	21.2	25.9	32.3

Figure 10.6: Statistical uncertainty on the measurement of the mass, width and production cross-section of the Higgs boson.

10.2.2 Inclusive Higgs boson production with $H \rightarrow WW^{(*)} \rightarrow 2\ell 2\nu$

The Higgs decay into two Ws and subsequently into two leptons ($H \rightarrow WW \rightarrow \ell\nu\ell\nu$) is the discovery channel for Higgs masses between $2m_W$ and $2m_Z$ [471]. In this mass range, the Higgs to WW branching ratio is close to one, leading to high statistics. The signature of this decay is characterized by two leptons and missing energy. However, since no narrow mass peak can be reconstructed, a good background control together with a high signal to background ratio is needed. The most important backgrounds, which give similar signature as the signal (i.e. two leptons and missing energy), are the continuum WW production and the $t\bar{t}$ production. To reduce these backgrounds, one has to require a small opening angle between the leptons in the transverse plane and apply a jet veto.

10.2.2.1 Signal and background generation

The signal samples were generated using PYTHIA. The two major Higgs production modes for the mass range studied, gluon and vector boson fusion were generated. The $p_t(H)$ spectrum predicted by PYTHIA was reweighted to the MC@NLO prediction, defining p_t dependent k-factors, as proposed in [472].

For the backgrounds, continuum vector boson production (WW, ZZ, WZ) was generated using PYTHIA. The p_t (WW) spectrum was reweighted using the same technique than for the signal. A NLO cross section of respectively 16 pb, 50 pb and 114 pb was taken for ZZ, WZ and WW. WW production via gluon box diagram, ggWW, was generated using a parton Monte Carlo provided by N. Kauer and linked to PYTHIA for the parton shower [63]. Top production ($t\bar{t}$ and tWb) was generated using TopReX. NLO cross sections of respectively 840 pb and 33.4 pb were used for $t\bar{t}$ and tWb [473].

10.2.2.2 Signal reconstruction

The signal signature is characterized by two leptons in the final state with opposite charge, missing energy and no jet. The leptons, either electrons or muons, are required to have $p_t > 20$ GeV and $|\eta| < 2$.

Muons candidates are asked to be isolated: The energy left in the calorimeters around the muon candidate within a $\Delta R = 0.3$ cone must be smaller than 5 GeV and the sum of the p_t of the tracks within a $\Delta R = 0.25$ cone around the muon candidate must be smaller than 2 GeV.

Electrons candidates are reconstructed combining tracks and ECAL clusters. They must fulfill in addition the following identification requirements:

- The electron must deposit few energy in the HCAL: $E_{\text{hcal}}/E_{\text{ecal}} < 0.05$
- The electron track and cluster must be precisely matched:
 - in direction: $|\eta_{\text{track}} - \eta_{\text{SC corr}}| < 0.005$ and $|\phi_{\text{track prop}} - \phi_{\text{SC}}| < 0.02^*$
 - in magnitude: $E/p > 0.8$ and $|1/E - 1/p| < 0.02$

The electron candidate must be also isolated by requiring, $\sum_{\text{tracks}} p_t(\text{track})/E_t(\text{SC}) < 0.05$, where the sum runs on all the tracks which have:

- $\Delta R_{\text{SC-track}} < 0.2$ (at vertex)
- $p_t^{\text{track}} > 0.9$ GeV
- $|z_{\text{track}} - z_{\text{electron}}| < 0.2$ cm

Finally a cut on the impact parameter significance in the transverse plane is applied in order to reduce the $b\bar{b}$ background. Each lepton is required to have $\sigma_{\text{IP}} < 3$ where σ_{IP} is the impact parameter significance. The two leptons are also required to come from the same vertex by asking $|z_{\text{lep1}} - z_{\text{lep2}}| < 0.2$ cm.

With this lepton selection, the contribution of reducible backgrounds like $W+\text{jet}$ where one jet is misidentified as a lepton or $b\bar{b}$ is expected to be less than 5 fb after all cuts applied.

Missing energy is reconstructed by summing the raw energy of all ECAL and HCAL towers, and correcting for muons. Since a jet veto is applied in the signal selection, further correction on the missing energy did not bring a significant improvement.

Jets are reconstructed using a Cone algorithm of size $\Delta R = 0.5$ and requiring its component calorimeter towers to have $E_T^{\text{tow}} > 0.5$ GeV and $E^{\text{tow}} > 0.8$ GeV. Since jets are reconstructed to be vetoed, no energy calibration was applied. For the events studied, $E_T(\text{jet}) \approx (1.5 - 2) \cdot E_T(\text{raw})$. To veto electrons and Bremsstrahlung photons, the jets are also required to be away from the leptons ($\Delta R_{\text{jet-lepton}} > 0.5$).

For jets with a raw energy between 15 and 20 GeV an additional cut on their track content

*Where $\phi_{\text{track prop}}$ is the track angle propagated in the magnetic field up to the ECAL cluster position

was applied in order to reduce the contamination from fake jets coming from the underlying event. For this, the so-called alpha parameter is defined, as the ratio of the sum of p_t of all tracks inside the jet over the transverse jet energy in the calorimeter. For a perfect detector, the alpha parameter of a jet would be around 0.66, as in mean two third of a jet are charged particles. This ratio is smeared and reduced by the detector energy resolution and the fact that particles need a minimal energy in order to be detected. In a fake jet, the alpha parameter looks different, as underlying events contain a lot of low p_t particles, which depose energy in the ECAL but are not seen in the tracker, leading to an alpha parameter around zero.

Alpha is determined using only tracks that are 'inside' the jet, i.e. with $\Delta R_{\text{track-jet}} < 0.5$ and coming from the event vertex † , fulfilling $|z_{\text{trk}} - z_{\text{vtx}}| < 0.4$ cm. Finally, these tracks should have more than 5 hits and $p_t > 2$ GeV. Alpha is then defined as $\text{alpha} = \frac{\sum p_t(\text{tracks})}{E_T(\text{jet})}$. If its raw energy lies between 15 and 20 GeV a jet is then required to have $\text{alpha} > 0.2$ to be kept.

10.2.2.3 Event selection and results

Events are first required to pass globally the L1 trigger and at least one of the following HLT triggers: single electron, double electron, single muon or double muon trigger.

Figure 10.12 shows the L1 trigger efficiency (blue dashed curve) and the combined L1+HLT trigger efficiencies (red dotted curve) as a function of the Higgs mass. Note that the W are allowed to decay into τ , reducing the trigger efficiency since no specific τ trigger is used. To estimate the numbers of 'useful events' rejected by this trigger requirement it is interesting to look at the trigger efficiency on events having exactly two leptons which fulfill the lepton selection cuts defined before. This is shown by the solid black curve on Figure 10.12. In this case, the trigger efficiency is higher than 95% on the full mass range and is around 100% for $\mu\mu$ final state, whereas for ee final state it is around 96%.




Figure 10.12: Trigger efficiencies (L1+HLT) as a function of the Higgs mass on all events (dotted line), on events where the W decays in electrons and muons (dashed line) and on events with exactly two leptons passing the lepton selection cuts (solid line).

Then each event has to contain exactly two opposite charge leptons with $p_t > 20$ GeV and

† The event vertex is defined as the mean z position of the two leptons.

Table 10.7: The expected number of events for the signal for three different Higgs masses and the different backgrounds given in fb. The first column shows the number of expected events after HLT requirement, the second one after having found two opposite charge leptons and the last one the number of events after all selection cuts are applied.

Reaction $pp \rightarrow X$	$\sigma_{\text{NLO}} \times \text{BR}$	L1+HLT	2 leptons	All cuts
$\ell = e, \mu, \tau$	pb	Expected event rate in fb		
$H \rightarrow WW \rightarrow \ell\ell, m_H = 160 \text{ GeV}$	2.34	1353 (58%)	359 (27%)	42 (12%)
$H \rightarrow WW \rightarrow \ell\ell, m_H = 165 \text{ GeV}$	2.36	1390 (59%)	393 (28%)	46 (12%)
$H \rightarrow WW \rightarrow \ell\ell, m_H = 170 \text{ GeV}$	2.26	1350 (60%)	376 (28%)	33 (8.8%)
$qq \rightarrow WW \rightarrow \ell\ell$	11.7	6040 (52%)	1400 (23%)	12 (0.9%)
$gg \rightarrow WW \rightarrow \ell\ell$	0.48	286 (60%)	73 (26%)	3.7 (5.1%)
$tt \rightarrow WWbb \rightarrow \ell\ell$	86.2	57400 (67%)	15700 (27%)	9.8 (0.06%)
$tWb \rightarrow WWb(b) \rightarrow \ell\ell$	3.4	2320 (68%)	676 (29%)	1.4 (0.2%)
$ZW \rightarrow \ell\ell\ell$	1.6	1062 (66%)	247 (23%)	0.50 (0.2%)
$ZZ \rightarrow \ell\ell, \nu\nu$	1.5	485 (32%)	163 (34%)	0.35 (0.2%)
Sum backgrounds	105	67600 (64%)	18300 (27%)	28 (0.2%)

$|\eta| < 2$ passing the cuts described before. The following kinematic selection was applied on the two lepton candidates:

- $E_t^{\text{miss}} > 50 \text{ GeV}$
- $\phi_{\ell\ell} < 45^\circ$ (angle between the leptons in the transverse plane)
- $12 \text{ GeV} < m_{\ell\ell} < 40 \text{ GeV}$ (the invariant mass of the two leptons)
- No jet with $E_t^{\text{raw}} > 15 \text{ GeV}$ and $|\eta| < 2.5$
- $30 \text{ GeV} < p_t^{\ell \text{max}} < 55 \text{ GeV}$ (lepton with the maximal p_t)
- $p_t^{\ell \text{min}} > 25 \text{ GeV}$ (lepton with the minimal p_t)

These cuts were optimized for a Higgs mass of 165 GeV. The expected number of events for the signal for three different Higgs masses and the different backgrounds in fb are given in Table 10.7. The first column shows the signal times branching ratio for the different processes, the second one shows the number of events passing the trigger requirement, the third one the number of events with two opposite charge leptons passing the lepton selection cuts and the last one the number of events after all selection cuts are applied. Figure 10.13, left shows the $\phi_{\ell\ell}$ distribution for the signal plotted on the top of the sum of all background when all selection cuts are applied except the one on $\phi_{\ell\ell}$.

10.2.2.4 Background normalization and systematics

The following procedure for background normalization is proposed.

- *Top background normalization:* Two procedures are proposed. A first possibility is to define a sample with the same lepton and missing energy cuts than the signal but requiring two b-tagged jets with $E_t > 20 \text{ GeV}$. A second possibility is to apply the same kinematic cuts on the leptons and require two additional jets with respectively $E_T^{\text{raw}} > 50 \text{ GeV}$ and $E_T^{\text{raw}} > 30 \text{ GeV}$. In this case, only $e\mu$ final states are considered in order to avoid a contamination from Drell-Yan. Both methods are expected to give an error of about 16% on $t\bar{t}$ estimate for a luminosity of 5 fb^{-1} .




Figure 10.13: The angle between the leptons in the transverse plane for the signal and the different background and a luminosity of 10fb^{-1} , (Left) For the signal cuts taking out the one on ϕ_{ll} . (Right) For the WW background normalization region where all signal cuts are applied except the one on the lepton invariant mass, which was set to $m_{ll} > 60\text{ GeV}$ and only electron-muon final states are kept.

- *WW background normalization:* A normalization region can be defined for WW by keeping the same cuts than the signal but requiring $\phi_{ll} < 140$ and $m_{ll} > 60\text{ GeV}$. Moreover only opposite flavor leptons are considered in order to reduce the Drell-Yan and WZ contribution. A systematic error of about 17% is expected with a luminosity of 5 fb^{-1} , dominated by statistical uncertainty. Figure 10.13 right shows the ϕ_{ll} distribution for the different process in this normalization region.
- *WZ background normalization:* WZ can be normalized by keeping the same signal cut and requiring an additional lepton in the final state. The cuts on ϕ_{ll} and m_{ll} are removed. An accuracy of about 20% is expected on this background with 5 fb^{-1} .
- *ggWW and tWb normalization:* The contribution of these backgrounds will be estimated using Monte Carlo prediction, since they represent only a small fraction of signal events. The error on ggWW is about 30% whereas the one on tWb is about 22%, both largely dominated by theoretical errors.

Taking into account the sum of the different backgrounds, an overall error of 10% is found on the total background. These results are calculated for a luminosity of 5 fb^{-1} . For luminosities of 1, 2 and 10 fb^{-1} , the total systematic errors scale to 19%, 16% and 11% respectively. Table 10.8 shows the signal to background ratio for the different Higgs masses together with the luminosity needed for a 5σ discovery, with and without the inclusion of background uncertainties. For Higgs masses of 120-140 GeV and 190-200 GeV, the background errors are too high to get a significant signal.

Figure 10.14 shows the signal to background ratio (left) and the luminosity needed for a 5σ discovery (right) as a function of the Higgs mass. A signal of more than 5σ significance could be already observed with a luminosity of 7 fb^{-1} for a Higgs mass between 150 and 180 GeV. For a Higgs mass of 165 GeV the luminosity needed for a 5σ discovery is expected to be less than 1 fb^{-1} .

Table 10.8: The signal to background ratio for the different Higgs masses together with the luminosity needed for a 5σ discovery, with and without the inclusion of background uncertainties. Also the statistical errors due to the restricted MOnte Carlo statistics are taken into account.

m_H [GeV]	S/B	Significance for 5 fb^{-1}		$\mathcal{L}_{\text{disc}}$ [fb]	
		no bkg syst	with bak syst	no bkg syst	with bkg syst
150	0.61	6.6	4.0	3.0	8.2
160	1.51	14	7.7	0.58	1.1
165	1.66	15	8.3	0.50	0.90
170	1.19	11	6.3	0.88	1.7
180	0.65	6.7	3.7	2.7	7.3




Figure 10.14: Signal to background ratio (left) and signal significance for a luminosity of 5fb^{-1} (right) as a function of different Higgs masses for the $H \rightarrow WW$ channel.

10.2.2.5 Selections optimization in the region of $M_H=130\text{-}150 \text{ GeV}/c^2$ with $e^+e^-\nu\nu$ final state.

A dedicated optimisation for the $e^+e^-\nu\nu$ final state in the mass range of $130 \leq M_H \leq 150 \text{ GeV}/c^2$ is performed. The largest significance is searched assuming a known M_H . The latest developments in detailed electron reconstruction are used and allow a good rejection of the W +jets background which is characterized by the misidentification of a jet as an electron. New kinematical variables have been designed to reduce the W +jets background as well as the contribution from Drell-Yan events with recoiling jets (Z +jets). For instance, in the signal, the two electrons tend to be close to each other, and the di-electron system is essentially emitted in the central region. On the contrary, in the Z +jets background, the di-electron pair is emitted uniformly η , and the electrons candidates in the W +jets backgrounds are well separated. Other selection criteria relying on the absence of a true source of missing transverse energy in the Z + jets events have been introduced: in the events where the missing transverse energy is mismeasured, it is usually in the same direction as the leading jet. Similarly, the imbalance of the missing energy and the di-lepton system in the transverse plane is exploited.

Both W +jets and Z +jets backgrounds are thus explicitly reduced to a manageable level. Figure 10.15 (left) shows the reconstructed WW transverse mass for the 140 GeV Higgs signal

selection with 10 fb^{-1} . Figure 10.15(right) shows the signal significance as function of the Standard Model Higgs mass for the integrated luminosity of 30 fb^{-1} with and without systematics taken into account. A 3σ observation is possible for Higgs masses from 135 GeV. A 5σ discovery is reached with 60 fb^{-1} .




Figure 10.15: Left plot : the reconstructed WW transverse mass for the 140 GeV Higgs signal selection with 10 fb^{-1} . The dashed lines show the window of events entering in the signal significance calculation. Right plot: the signal significance as function of the Standard Model Higgs mass for an integrated luminosity of 30 fb^{-1} .

10.2.3 The vector boson fusion production with $H \rightarrow \tau\tau$

10.2.3.1 Lepton plus τ jet final state: $H \rightarrow \tau\tau \rightarrow \ell + \tau \text{ jet} + E_T^{\text{miss}}$

Signal, Background Events and Preselections

The signal events were generated using PYTHIA for four different values of the Higgs boson mass: 115, 125, 135 and 145 GeV/c^2 . The Higgs boson was forced to decay to two τ leptons with one τ decaying to leptons and the other τ to hadrons. The TAUOLA package was used to simulate the τ polarization.

For background events, following processes are considered:

QCD $2\tau+2/3j$

The QCD production of $2\tau+2\text{jet}$ and $+3\text{jet}$ events with the invariant mass of two τ leptons, $M_{\tau\tau} > 70 \text{ GeV}/c^2$, was generated using ALPGEN with CTEQ5L PDF. Given the limit of the detector acceptance and requirements in the course of the event reconstruction, all jets were required to satisfy $p_{Tj} > 20 \text{ GeV}$, $|\eta_j| < 5.0$ and $|\Delta R_{jj}| > 0.5$. Further preselections were applied on the two highest p_T jets (j_1 and j_2) reflecting the offline VBF selection cuts: $|\Delta\eta_{j_1j_2}| > 4.0$, $M_{j_1j_2} > 600 \text{ GeV}/c^2$. Then the events $2\tau+2j$ and $2\tau+3j$ were added together with the MLM prescription in PYTHIA to avoid double counting of the jets. The TAUOLA package was used in PYTHIA to force one τ lepton to decay leptonically and the other hadronically.

Electro Weak (EW) production of $2\tau+2j$

The EW production of two τ 's with $M_{\tau\tau} > 70 \text{ GeV}/c^2$ and two jets in the final state was generated using MadGraph with CTEQ5L PDF. Soft preselections were applied during generation with MadGraph on the kinematics of the jets: $p_{Tj} > 20 \text{ GeV}/c$ and $M_{jj} > 500 \text{ GeV}/c^2$.

Further preselection cuts were applied on jets and τ 's given the limit of the detector acceptance and requirements of the event reconstruction: $|\eta_j| < 5.2$, $|\Delta R_{jj}| > 0.5$, $|\Delta R_{\tau\tau}| > 0.4$. The showering and hadronization of the MadGraph parton level events were carried out using PYTHIA where all decay modes of the τ lepton were open.

W+jets

The W+3j and W+4j events with $W \rightarrow \mu\nu$ decays were generated using ALPGEN with CTEQ5M PDF. In addition to the kinematical cuts on jets used for the QCD Z+jets production described above, further preselections were made based on the lepton properties with $|\eta_l| < 3$ and $p_{Tl} > 10 \text{ GeV}/c$. The MLM prescription was applied in PYTHIA.

t \bar{t} \rightarrow WbWb

The $t\bar{t}$ background was generated using PYTHIA, TopRex, ALPGEN, CompHEP and MadGraph. All leptonic W decays were included and no kinematical preselection was applied.

Event Reconstruction and Selection Stream

1. L1 trigger: single isolated e, single μ and combined e- τ triggers.
2. HLT: single isolated e, single μ , combined e- τ and μ - τ triggers.
3. The lepton identification is carried out by the corresponding subsystems in ORCA and the highest E_T candidate is selected from OfflineElectronReco or GlobalMuonReconstructor. For an electron candidate, there are three additional requirements: $E/P > 0.9$, tracker isolation, $(\sum_{0.01 < \Delta R < 0.2}^{trk} P)/E < 0.05$, and E_T of the hottest HCAL tower, $E_T^{\text{Htow}} < 2 \text{ GeV}$. A further cut is applied on the lepton p_T at $15 \text{ GeV}/c$.
The lepton track is used to identify tracks originating from the signal vertex. The tracks are used for the electron isolation, τ tagging and in central jet veto. A track is associated to the signal vertex if its z impact parameter lies within $|\Delta z| < 0.2 \text{ cm}$ from that of the lepton track.
4. The τ -jet identification is carried out offline seeded from the L1/HLT τ candidates. A jet is formed around each candidate which does not coincide with the identified electron, and the jet is passed through a series of τ -tagging criteria. The τ tagging used in HLT (Ref. [68]) has been adapted to offline use with $R_m = 0.1$, $R_s = 0.07$, $R_i = 0.45$, $p_T^{\text{ltr}} = 6 \text{ GeV}/c$ and $p_T^i = 1 \text{ GeV}/c$. The charge of the τ -jet is required to be opposite of the lepton charge, and $E_T^{\text{Htow}} > 2 \text{ GeV}$ is required if the jet coincides with any of the electron candidates. A further cut is applied on the transverse energy of the τ -jet, $E_T > 30 \text{ GeV}$.
5. The jets from the VBF process are identified as the two highest E_T calorimeter jets with $E_{Tj} > 40 \text{ GeV}$, excluding the electron and the τ -jet. The jets are required to satisfy: $|\eta_j| < 4.5$, $\eta_{j1} \times \eta_{j2} < 0$, $\Delta\eta_{j1j2} > 4.5$, $\Delta\phi_{j1j2} < 2.2$, and the invariant mass, $M_{j1j2} > 1 \text{ TeV}$. The jets after these selections will be referred to as tagging jets.
6. A cut is applied on the transverse mass of the lepton- E_T^{miss} system, $M_T(\text{lep}, E_T^{\text{miss}}) < 40 \text{ GeV}$, in order to reject reducible backgrounds.
7. Central jet veto
An event is vetoed if there is an additional jet (j3) with $E_{Tj3}^{\text{raw}} > 10 \text{ GeV}$ in the rapidity gap between the two tagging jets, satisfying the following:
 - $(\eta_{\min} + 0.5) < \eta_{j3} < (\eta_{\max} - 0.5)$
where η_{\min} and η_{\max} correspond to the tagging jets which has smaller and larger value of η respectively.

- $\alpha_{j3} = \sum p_{T\text{trk}}/E_{Tj3}^{\text{raw}} > 0.1$
where $p_{T\text{trk}}$ is the p_T of tracks originating from the signal vertex, which lie within the 0.5 cone around the jet axis, and E_{Tj3}^{raw} is the raw E_T of the jet measured in the calorimeter.

α_{j3} is defined for each additional jet, and the one which satisfies the first criteria and has the highest α_{j3} is considered for the veto.

8. Mass Calculation

The invariant mass of the two reconstructed τ 's is calculated as described in MSSM $H(A) \rightarrow \tau\tau$ analysis (Section 5.2) using collinear approximation of visible part of τ 's and neutrinos. The E_T^{miss} is reconstructed by summing the E_T of the calorimeter towers and the muon candidates, and applying the jet energy corrections (Type 1 E_T^{miss}).

Selection Efficiency

The efficiency of each reconstruction and selection step and the cumulative cross section expected at the LHC are given in Table 10.9. The total selection efficiencies are, 0.32%, 0.34%, 0.42%, 0.39%, for the signal events with the Higgs boson masses, $M_H = 115, 125, 135$ and $145 \text{ GeV}/c^2$ respectively.

For the $W+3/4j$ samples, the efficiencies of some selection cuts have been obtained from factorization of cuts. The triggers and the lepton identification are carried out as other samples, and the remaining steps are carried out in two uncorrelated parallel streams – A: VBF and $M_T(\text{lep}, E_T^{\text{miss}})$ cuts, B: central jet veto, τ tagging and mass calculation – after preselections of forward jets and τ -jet candidates.

Table 10.9: Cumulative cross sections in fb after successive selection cuts. The efficiency (%) of each cut is listed inside the brackets. The entry, “valid mass”, corresponds to the fraction remained after the calculation of the di- τ mass; some events are lost due to the negative reconstructed neutrino energies. For the $W+3/4j$ samples, efficiencies are obtained from factorization of cuts and the τ -jet ID efficiency includes the p_T cut, and the number of events at 30 fb^{-1} (indicated by *) is calculated for all leptonic decay modes of W .

Selection	cross section, σ [fb] (% from previous cut)				
	signal	background			
	$M_H=135$	EW $2\tau+2j$	QCD $\tau\tau+2/3j$	$W+3/4j$	$t\bar{t} \rightarrow WbWb$
Starting σ	82.38	299.	1615.	14.45×10^3	86×10^3
L1	46.50 (56.5)	179.8 (60.1)	543.8 (33.7)	9186. (63.6)	71.39×10^3 (83.0)
L1+HLT	24.60 (52.9)	58.81 (32.7)	201.3 (37.0)	6610. (71.9)	55.42×10^3 (77.6)
lepton ID	23.34 (94.9)	50.67 (86.2)	187.4 (93.1)	6549. (99.1)	54.08×10^3 (97.6)
lepton p_T	23.16 (99.3)	49.13 (97.0)	185.6 (99.0)	6543. (99.9)	53.54×10^3 (99.0)
τ -jet ID	8.276 (35.7)	10.49 (21.3)	39.64 (21.4)	(0.21)	5.056×10^3 (9.4)
τ -jet p_T	6.422 (77.6)	7.360 (70.2)	24.25 (61.2)	-	3.215×10^3 (63.6)
Valid mass	4.461 (69.5)	4.232 (57.5)	14.49 (59.8)	(17.4)	848.6 (26.4)
VBF cuts	0.545 (12.2)	0.391 (9.2)	1.666 (11.5)	(11.0)	2.738 (0.3)
$M_T(\text{lep}, E_T^{\text{miss}})$	0.423 (77.6)	0.322 (82.4)	1.382 (83.0)	(30.5)	0.942 (34.4)
Central Jet Veto	0.344 (81.3)	0.230 (71.4)	0.555 (39.7)	(28.9)	0.224 (23.8)
N events at 30 fb^{-1}	10.3	6.9	16.6	1.5*	6.7

Reconstructed Mass and Fit

The distribution of the invariant mass of two reconstructed τ 's for different samples is shown

in Figure 10.16, where the signal sample with the Higgs boson mass, $M_H = 135 \text{ GeV}/c^2$ is used. A Gaussian function is used to fit the signal distribution, a Breit-Wigner function for the 2τ +jets background from EW and QCD processes, and a second order polynomial for the W +jets and $t\bar{t}$ events. The Higgs boson mass resolution is 9.1%.




Figure 10.16: The invariant mass of two reconstructed τ 's. The number of entries in each histogram is normalized to the expected number of events at an integrated luminosity of 30 fb^{-1} .

Significance

The significance is calculated using a window with a fixed width of $40 \text{ GeV}/c^2$, which slides in $5 \text{ GeV}/c^2$ steps. An optimum window position which maximizes the significance is chosen for each of the four different masses of Higgs boson. The numbers of signal and background events within the window, N_S and N_B , are estimated from the fits to individual samples. The method S_{CP} (Ref. [71]) is used for calculating the significance, including the systematic uncertainty of 7.8% for 30 fb^{-1} and 5.9% for 60 fb^{-1} . The results are summarised in Table 10.10.

It is envisaged that the shapes of the two background distributions will be extracted experimentally from the LHC data in a region unaffected by the signal contribution, using some relaxation of selection cuts. Since the number of background events in the signal region will be estimated using real data, the fitting procedure is the only contribution to the uncertainty in the significance estimate. The fit uncertainty has been evaluated by performing MC trials, randomly generating a mass distribution from the original fit functions and re-fitting the distribution at each trial. With the data, the Higgs boson mass will be estimated by repeating the fitting procedure for different mass hypotheses and finding the value where the χ^2 of the fit is minimized.

10.2.4 Searching for Standard Model Higgs via Vector Boson Fusion in $H \rightarrow W^+W^- \rightarrow \ell^\pm \nu jj$ with m_H from 120 to $250 \text{ GeV}/c^2$

The signal topology of Higgs boson with $H \rightarrow W^+W^- \rightarrow \ell \nu jj$ via vector boson fusion has been shown as a good potential discovery channel for the medium-high mass range ($m_H > 300 \text{ GeV}/c^2$). The final state is characterized as two forward jets, two central jets from W

Table 10.10: The production cross section and significance of the expected number of signal events within the optimum mass window for each of the four different simulated masses of the Higgs boson.

M_H [GeV]	115	125	135	145
Production σ [fb]	4.65×10^3	4.30×10^3	3.98×10^3	3.70×10^3
$\sigma \times \text{BR}(H \rightarrow \tau\tau \rightarrow lj)$ [fb]	157.3	112.9	82.38	45.37
N_S at 30 fb^{-1}	10.5	7.8	7.9	3.6
N_B at 30 fb^{-1}	3.7	2.2	1.8	1.4
Significance at 30 fb^{-1} ($\sigma_B = 7.8\%$)	3.97	3.67	3.94	2.18
Significance at 60 fb^{-1} ($\sigma_B = 5.9\%$)	5.67	5.26	5.64	3.19

hadronic decay, and one high p_T lepton and missing transverse energy (E_T^{miss}) from the W leptonic decay. Extending the use of this channel to the low mass range ($m_H < 300 \text{ GeV}/c^2$) makes valuable physics analysis possible and is complementary to Higgs boson search using $H \rightarrow W^+W^- \rightarrow \ell\nu\ell\nu$. Especially for $160 < m_H < 180 \text{ GeV}/c^2$, the $H \rightarrow ZZ^*$ branching ratio is highly suppressed due to $H \rightarrow W^+W^-$ resonance.

The result of this section shows that in the Higgs mass range between 140 and 200 GeV/c^2 , a significance of $\sim 5 \sigma$ can be achieved with integrated luminosity of 30 fb^{-1} . Major backgrounds include $t\bar{t}$ +jets, $W+t\bar{b}(\bar{t}b)$, W +jets, Z +jets, $WW/WZ/ZZ$ +jets, and QCD events. For WW +jets, the QCD and Electroweak (EW) processes are generated separately.

10.2.4.1 Event Selection Strategy

Major difficulties concerning the low mass Higgs analysis using $\ell\nu jj$ final state include: many background processes of very large cross section have one lepton and multiple jets in the final states; simulating the requisite huge number of background events is both a computing and analysis challenge; deep selection cuts and heavy exploitation of physics signal characteristics are necessary to suppress backgrounds and enhance the statistical significance of the signal, however they can lead to large systematic uncertainties; a relatively low Higgs boson mass limits the application of high jet E_T thresholds that would normally be used to suppress backgrounds, in contrast to the situation at high mass; low E_T^{miss} and low E_T jets affect the resolution of Higgs mass. To meet these challenges, a robust reconstruction and selection strategy is developed.

Low p_T objects are ignored (e.g. leptons with $p_T < 10 \text{ GeV}/c$ and jets with $E_T < 25 \text{ GeV}$). The jet E_T threshold is chosen around 25 GeV where there is a stable signal to background ratio (S/B), so that the systematic uncertainty of jet energy scale is minimized (Fig. 10.17). Due to a number of soft jets in the central detector region, the hadronic W reconstruction looks for a di-jet mass with least error to the true W mass. The extra jet veto after forward jet tagging and hadronic W reconstruction is applied. Two schemes are studied: full extra jet veto ($N_{\text{extra}} < 1$) and loose extra jet veto ($N_{\text{extra}} < 2$). The full extra jet veto is very powerful in reducing the $t\bar{t}$ +jets and W +jets background.

The selection chain is divided into two major step: basic selection (Table 10.11) and optimized selection. This strategy helps optimize the selection cuts and factorize the selection efficiency to evaluate the systematic uncertainty and QCD background efficiency.




Figure 10.17: Multiple jet selection efficiency (requiring at least 4 jets in an event) as a function of jet E_T threshold. The efficiency is normalized to the rate with jet E_T threshold of 16 GeV for each sample. The physics channels include: $t\bar{t}$ + jets (solid square), W + 3jets (open circle), W + 4 jets (solid triangle), and VBF Higgs with $m_H = 170 \text{ GeV}/c^2$ (open square)

Table 10.11: Summary of basic event selection cuts

Selection	Configuration
Lepton selection	calorimeter-based e/μ isolation $30 < p_T < 120 \text{ GeV}/c$ $\Delta R_{\ell,j} > 0.5$
Jet selection	$N_{\text{jet}} \geq 4$ jets with $E_T > 25 \text{ GeV}$ $E_T^{\text{miss}} > 30 \text{ GeV}$
Forward jet tagging	$E_T > 30 \text{ GeV}$ $\eta_1 \cdot \eta_2 < 0$ $ \eta_1 - \eta_2 > 3.8$ $m_{qq} > 800 \text{ GeV}/c^2$
Hadronic-W	$\Delta m_W < 25 \text{ GeV}/c^2$ ($m_H \geq 160 \text{ GeV}/c^2$) $30 < m_W < 90 \text{ GeV}/c^2$ ($m_H < 160 \text{ GeV}/c^2$) select di-jet with the least Δm_W
Leptonic-W	using lepton and E_T^{miss} select Leptonic-W candidates of smaller $\Delta R(\text{Lepton} - W, \text{Hadronic} - W)$

The optimized selection for $m_H \geq 160 \text{ GeV}/c^2$ ($m_H < 160 \text{ GeV}/c^2$) includes 3 steps:

- $E_T^{\text{FH}} > 45$ (40) GeV, $E_T^{\text{FL}} > 35$ (30) GeV, $\Delta\eta > 4.2$, and $m_{qq} > 1000 \text{ GeV}/c^2$. E_T^{FH} (E_T^{FL}) is the high (low) jet E_T threshold for forward jets.
- $E_T^{\text{CH}} > 30$ GeV, $E_T^{\text{CL}} > 25$ GeV, $\Delta m_W < 20 \text{ GeV}/c^2$ ($30 < m_W < 90 \text{ GeV}/c^2$), and $N_{\text{extra}} < 1$. E_T^{CH} (E_T^{CL}) is the high (low) jet E_T threshold for central jets that are used for hadronic-W reconstruction.
- $E_T^{\text{miss}}(qqWW) < 40$ GeV, $\Delta R(\text{lepton}, \text{Hadronic-W}) < 2.0$, and $\Delta R(\text{Leptonic-W}, \text{Hadronic-W}) < 1.0$. $E_T^{\text{miss}}(qqWW)$ is the E_T^{miss} of qqWW system that includes reconstructed Higgs and two forward jets.

The efficiency of basic selection and three steps of optimized selection is summarized in Table 10.12 and 10.13 for $m_H \geq 160 \text{ GeV}/c^2$ and $m_H < 160 \text{ GeV}/c^2$ respectively. Loose extra jet veto with tightening cuts: $m_{qq} > 1200 \text{ GeV}/c^2$ and $\Delta R(\text{lepton}, \text{Hadronic-W}) < 1.6$, gives a conservative result.

Table 10.12: Cross section (fb) of the signal and background in optimized selection with $m_H \geq 160 \text{ GeV}/c^2$ for full extra jet veto

Channels	Basic Selection	Step 1	Step 2	Step 3
VBF Higgs ($m_H=160$)	16.15	9.531	4.580	2.989
VBF Higgs ($m_H=170$)	15.99	9.814	4.828	3.006
VBF Higgs ($m_H=180$)	16.28	9.916	4.711	2.738
VBF Higgs ($m_H=190$)	14.16	9.363	4.294	2.340
VBF Higgs ($m_H=200$)	13.78	8.626	4.341	1.983
VBF Higgs ($m_H=210$)	13.43	8.211	4.080	1.571
VBF Higgs ($m_H=220$)	13.35	8.227	4.128	1.259
VBF Higgs ($m_H=250$)	10.71	6.900	3.426	0.810
$t\bar{t}$ + jets	1494.2	626.5	16.751	1.232
WW + jets (QCD)	9.27	1.265	0.422	< 0.008
WW + jets (EW)	7.88	9.683	4.454	< 0.0277
ZZ + jets	1.00	0.269	0.0245	< 0.001
ZW + jets	7.23	2.335	0.223	< 0.001
W + $t\bar{b}$ ($\bar{t}b$)	92.8	35.21	4.427	< 0.05787
W + 4j ($W \rightarrow e/\mu/\tau + \nu$)	1110.8	583.0	72.066	0.323
Z + 4j ($Z \rightarrow ee/\mu\mu$)	82.3	3.713	0.141	0.0104
Z + 3j ($Z \rightarrow ee/\mu\mu$)	72.4	2.313	0.233	< 0.0067
Sum of Background	3579.7	1492.5	167.38	1.565

Table 10.13: Cross section (fb) of signal and background in optimized selection with $m_H < 160 \text{ GeV}/c^2$ for full extra jet veto

Channels	Basic Selection	Step 1	Step 2	Step 3
VBF Higgs ($m_H=120$)	1.28	0.951	0.363	0.231
VBF Higgs ($m_H=130$)	4.03	3.004	1.125	0.664
VBF Higgs ($m_H=140$)	7.12	5.520	2.369	1.656
VBF Higgs ($m_H=150$)	11.01	8.345	3.505	2.317
$t\bar{t}$ + jets	1483.0	859.5	20.94	0.493
WW + jets (QCD)	9.70	4.215	0.422	< 0.004
WW + jets (EW)	7.94	11.21	5.395	< 0.0277
ZZ + jets	0.96	0.465	0.0979	< 0.001
ZW + jets	7.45	3.781	0.334	< 0.01
W + $t\bar{b}$ ($\bar{t}b$)	101.5	54.37	6.799	< 0.0289
W + 4j ($W \rightarrow e/\mu/\tau + \nu$)	1110.7	778.5	118.9	0.667
Z + 4j ($Z \rightarrow ee/\mu\mu$)	81.3	4.700	0.152	0.00522
Z + 3j ($Z \rightarrow ee/\mu\mu$)	70.0	3.160	0.353	< 0.01333
Sum of Background	3630.6	2066.5	267.2	1.164

10.2.4.2 Discovery Potential

Using optimized selection cuts, the overall S/B and significance at 30 fb^{-1} with respect to various Higgs boson masses is shown in Fig. 10.18.

The reconstructed Higgs mass distribution from signal and background at 60 fb^{-1} is shown in Fig. 10.19.

In the following, a quantitative estimation of the background shape in reconstructed Higgs boson mass distribution as a function of E_T^{miss} in qqWW system is performed by using four regions in the Higgs boson mass distribution:

- Region A: $m_H < 186 \text{ GeV}/c^2$
- Region B: $186 < m_H < 203 \text{ GeV}/c^2$




Figure 10.18: The significance of discovery potential at 30 fb^{-1} verse Higgs boson masses. The high (low) curves of S/B and significance correspond to full (loose) extra jet veto




Figure 10.19: Results of VBF Higgs mass reconstruction based on signal+background (blue) and background (black)

- Region C: $203 < m_H < 228 \text{ GeV}/c^2$
- Region D: $228 < m_H < 262 \text{ GeV}/c^2$

For a given E_T^{miss} cut, total number of events in these regions and fraction of events in each region are calculated for VBF Higgs boson signal and the background respectively Fig. 10.20):

- For VBF Higgs boson signal, the shape of the Higgs boson mass distribution is very stable, because the fraction of events in each region does not change with the E_T^{miss} cut. Region A dominates the mass distribution which accounts for $\sim 2/3$ of the events.
- For the background events, the change mainly occurs for $E_T^{\text{miss}} < 80 \text{ GeV}$. There is a roughly 10% difference in the fraction of events for Region A and C between the low and high E_T^{miss} cut. A much lower bump from the background events is expected, which mainly lies on the tail of the signal's distribution.

10.2.4.3 Estimate of QCD Background

In order to estimate the contamination of QCD background in the final result, the overall selection chain are divided into three groups: leptonic-W chain, hadronic-W chain and optimization chain. The leptonic-W chain and hadronic- W chain are independent, while optimization chain will need to be conducted after Leptonic-W and Hadttonic-W chains. This




Figure 10.20: The fraction of events in different regions for the overall background (left) and VBF Higgs signal (right) as a function of E_T^{miss} cuts. Region A (close square), Region B (open square), and Region C (open circle).

strategy allows to factorize the selection efficiency of QCD events for each selection chain:

- Leptonic-W chain includes: lepton trigger, lepton isolation, leptonic-W selection. The selection efficiency for QCD events is E_L .
- Hadronic-W chain includes: jet topology cuts (≥ 5 jets, E_T^{miss} cut), forward jet tagging, hadronic-W selection. The reason we require 5 jet in the topology cut is because we need save one jet faking an isolated e/μ ($p_T \geq 30$ GeV/c), then rest 4 jets used for two forward jets and central jets of hadronic-W reconstruction and selection. The selection efficiency for QCD events is E_H .
- For QCD events, since there is no way we can get a single event that pass two previous chains, so we directly use W+jets efficiency to estimate the efficiency of QCD events. In another word, we treat those QCD events passing basic selection as leptonic background. The selection efficiency for QCD events is E_O .

The overall selection efficiency for QCD events is: $E = E_L \times E_H \times E_O$. The E_L and E_H is directly measured from QCD samples. The number of events estimated for 60 fb^{-1} for each qcd \hat{p}_T range is summarized in the table. The overall QCD contamination is estimated as 2-5 events for an upper limit, which causes possible 2-4% increase of background, which has almost no change in the significance.

Following is the summary of selection efficiency for QCD events:

Table 10.14: Factorized QCD Event Selection Efficiency

QCD \hat{p}_T	L.W Chain	H.W Chain	Opt Chain	Evt in 60 fb^{-1}
qcd_20_30	0	6.3E-4	5.0E-4	
qcd_30_50	0	1.3E-3	5.0E-4	
qcd_50_80	0	3.4E-3	5.0E-4	
qcd_80_120	$<3.8\text{E-6}$	6.7E-3	5.0E-4	<2.57
qcd_120_170	5.0E-6	1.3E-2	5.0E-4	1.1
qcd_170_230	5.6E-6	2.2E-2	5.0E-4	0.44
qcd_230_300	1.1E-5	3.3E-2	5.0E-4	0.31
qcd_300_380	2.4E-5	3.4E-2	5.0E-4	0.19
qcd_380_470	2.6E-5	3.9E-2	5.0E-4	0.08
qcd_470_600	1.6E-5	4.3E-2	5.0E-4	0.02

10.2.4.4 Experimental Analysis Method

The selection cut based on E_T^{miss} in qqWW system shown in previous section provides a strong suppression of background. The distinct distribution of E_T^{miss} in qqWW system between the VBF Higgs signal and background make it a valuable experimental signature for VBF Higgs boson analysis. In the following, a method of identifying excess of events with the presence of VBF Higgs boson signal is described, which uses the phase space defined by E_T^{miss} of qqWW system. Three regions with respect to E_T^{miss} are defined as:

- Region A: $E_T^{\text{miss}} < 40$ GeV.
- Region B: $40 < E_T^{\text{miss}} < 55$ GeV
- Region C: $55 < E_T^{\text{miss}} < 65$ GeV

The number of events in each region as defined as N_A , N_B and N_C respectively. A loose selection cut based on ΔR between the lepton and hadronic W is applied:

- $\Delta R = 0.6, 0.9, 1.2, 1.5, 1.8, 2.1, 2.4, 2.7, 3.0, 3.5, 4.0, 5.0, \text{ and } 6.0$.
- The other selection cuts except E_T^{miss} and ΔR are the same as those optimized ones.

Using the phase space defined above, we can investigate the correlation of the number of events in each region with respect to various values of ΔR cut. This correlation is mainly affected by whether there is the existence of VBF Higgs signal (called Scenario of Signal + Background) or just the background (called Scenario of Background Only). The selection efficiency will play a less important role in the correlation. It is expected that Region A will be influenced by VBF Higgs signal, while the rest two regions get much less influence.

Fig. 10.21 shows the ratio of the number of events between each of two regions ($R_{AB} = N_A/N_B$, $R_{AC} = N_A/N_C$, and $R_{BC} = N_B/N_C$). It can be seen that R_{AB} and R_{AC} are highly affected by the existence of VBF Higgs between two scenarios, while there is almost no change in R_{BC} .




Figure 10.21: The ratio of number of events as a function of ΔR for Region A to Region B (left), Region A and Region C (middle), and Region B to Region C (right). Two scenarios are used: Signal + Background (open square) and Background Only (solid square) respectively.

The difference between two scenarios provides an experimental way to identify the existence of VBF Higgs in addition to the Higgs boson mass peak. Especially using R_{AC} as shown Fig. 10.21 (b), the value from Background Only Scenario is about 0.8, but the Signal + Background Scenario can give a much larger ratio which can be identified easily.

The scale of the ratios (defined as R^{s+b}/R^b , where R^{s+b} is the ratio of Signal + Background Scenario, R^b is the ratio of Background Only Scenario) between two scenarios are illustrated in Fig. 10.22, which shows the excess in the low E_T^{miss} range (Region A) causes significant increase of the scale as ΔR goes lower. In general, the ratios between different regions defined by E_T^{miss} provide a good probe of VBF Higgs signature.




Figure 10.22: The ratio of Signal + Background Scenario to Background Only Scenario as a function of ΔR for Region A to Region B (open square), Region A to Region C (solid square), and Region B to Region C (open circle)

10.2.4.5 Detector Systematic Uncertainties and Control

Several calorimeter level systematic uncertainties have significant impact on this channel including: jet energy scale and resolution, E_T^{miss} scale and resolution, and calorimeter-based lepton isolation cut. Their impacts on the rate of signal (S), background (B) and S/B are summarized in Table. 10.15. The total detector level systematic uncertainty is about 16% in the absolute rate of background in the final result.

Table 10.15: Systematic Uncertainties due to Jet and E_T^{miss}

Source	S	B	S/B
Jet energy scale	10.6%	14.5%	5.2%
Jet energy resolution	0.1%	2.0%	2.0%
E_T^{miss}	2.5%	1.2%	1.7%
Lepton isolation	1.4%	1.3%	0.5%

The data driven technique is able to significantly reduce the detector level systematic uncertainties. For example, the largest uncertainty comes from the selection efficiency with respect to lowest jet E_T threshold. The event rate of the background near this threshold can be measured from data and used to tune the MC prediction, which leaves much less uncertainty due to the systematic bias of jet energy scale. Ignoring the uncertainty in the rate for from lowest jet E_T threshold, the uncertainty of jet energy scale only causes about 5.5% error in the rest of the selection chain which immediately reduces the total detector level systematic uncertainty down to 10% level.

10.2.5 Vector boson fusion production with $H \rightarrow \gamma\gamma$

Signal and background

Higgs boson signal

Table 10.16: Cross section for vector boson fusion Higgs boson production process

Higgs mass	σ_{VBF}	$\text{Br}(H \rightarrow \gamma\gamma)$	$\sigma_{VBF}\text{BR}$
110 GeV/ c^2	4.7 pb	1.90×10^{-3}	8.93 fb
115 GeV/ c^2	4.3 pb	2.08×10^{-3}	8.94 fb
120 GeV/ c^2	4.1 pb	2.19×10^{-3}	8.98 fb
130 GeV/ c^2	4.0 pb	2.23×10^{-3}	8.92 fb
140 GeV/ c^2	3.7 pb	1.94×10^{-3}	7.18 fb
150 GeV/ c^2	3.4 pb	1.39×10^{-3}	4.72 fb

The cross section for the vector boson fusion Higgs boson production was calculated by the program VV2H is shown in Table 10.16. The most up-to-date next-to-leading order corrections (so-called K factors) were included in the cross section calculation. The branching ratio $\text{Br}(H \rightarrow \gamma\gamma)$ and the product of the VBF cross section and branching ratio are also shown in Table 10.16. The branching ratio was calculated with the program HDECAY.

Different types of background

The search for the $H \rightarrow \gamma\gamma$ signal at LHC is affected by four types of backgrounds:

- The prompt di-photon production from the quark annihilation and gluon fusion diagrams, which provides an intrinsic or ‘irreducible’ background.
- The γ + jets production consisting of two parts: i) prompt photon + the second photon coming from the outgoing quark due to bremsstrahlung and ii) prompt photon + neutral hadron (mostly isolated π^0) in a jet.
- The background from QCD hadronic jets, where an electromagnetic energy deposit results from the decay of neutral hadrons (especially isolated π^0 s) in a jet.
- Drell Yan process with e^-e^+ in a final state which could mimic photons when correspondent electron tracks will not be assigned to the superclusters during the reconstruction.

Table 10.17 shows the cross sections of different types of background, along with the corresponding threshold of the transverse momentum of the parton in the hard interaction process P_T^{hard} that was used for generation.

In the framework of CompHEP approach, the backgrounds are identified on the level of Feynman diagram sets

- The irreducible QCD background $\gamma\gamma+2$ jets from the 34 partonic subprocesses $2 \rightarrow 4$. Main contribution comes from the subprocesses with u -quark and gluon in the initial state $ug \rightarrow \gamma\gamma gu$ and $gu \rightarrow \gamma\gamma gu$. Subleading contributions come from $\bar{u}g \rightarrow \gamma\gamma g\bar{u}$, $g\bar{u} \rightarrow \gamma\gamma g\bar{u}$, $uu \rightarrow \gamma\gamma uu$ and $gg \rightarrow \gamma\gamma u\bar{u}$.
- The irreducible electroweak background $\gamma\gamma+2$ jets from 20 partonic subprocesses $2 \rightarrow 4$. Main contribution comes from the subprocesses with u, d -quarks in the initial state $ud \rightarrow \gamma\gamma ud$ and $du \rightarrow \gamma\gamma ud$.
- The misidentification QCD background in the channel $pp \rightarrow \gamma+3$ jets from 50 partonic subprocesses $2 \rightarrow 4$. Main contribution comes from the subprocesses

Table 10.17: Cross sections of different types of background, calculated at leading order.

Background process	P_T^{hard} (GeV/c)	Cross section (pb)
QCD hadronic jets (eg05_jets_2g_pt50up)	50	$2.8 \cdot 10^7$
γ + jets (eg03_gamjet)	30	$4.93 \cdot 10^4$
Gluon fusion (eg03_twophoton_box)	20	83
Quark annihilation (eg03_twophoton_born)	20	83
Drell Yan (sm05_dy2e)	15	$4.1 \cdot 10^3$
γ + 3 jets, QCD (CompHEP)	–	5970
$\gamma\gamma$ + 2jets, QCD (CompHEP)	–	47.24
$\gamma\gamma$ + 2jets, EW (CompHEP)	–	0.33

with u -quark and gluon in the initial state $ug \rightarrow \gamma ggu$ and $gu \rightarrow \gamma ggu$.

- The misidentification EW background in the channel $pp \rightarrow \gamma + 3$ jets from 24 partonic subprocesses $2 \rightarrow 4$, main contribution is given by $ud \rightarrow \gamma gud$ and $du \rightarrow \gamma gud$. Total cross section 5.15 pb with the same partonic level cuts as defined above.

QCD hadronic jets generator level preselection

Selection based on the generated particles was devised, aimed at selecting events which could produce in the detector two electromagnetic showers consistent with isolated photons. In order to apply cuts on the invariant mass of the two candidates an attempt to estimate lower and upper limits to the energy of the candidates that will be reconstructed after the simulation was done.

The idea of this preselection, is just to pick up events that will give rise to energy depositions in ECAL large enough and isolated enough to be important for this analysis. Preselection algorithm is getting all particles which might deposit electromagnetic energy in ECAL, and looking around each particle in a narrow cone, to find another, may be less energetic particles which will make deposits in ECAL as well, and will potentially be reconstructed as one supercluster. In addition to that, a very loose tracker isolation was applied: three charged particles were required in a cone $\Delta R = 0.2$ around the “supercluster candidate”, described above, per one “supercluster candidate”, and no more than 6 per two first most energetic candidates.

After that some other cuts were applied for the “supercluster candidates” as well - $p_T > 37.5$ GeV/c for most energetic one and $p_T > 22.5$ GeV/c for the second most energetic one. The invariant mass of the first most energetic and second most energetic “supercluster candidates” should be more than $90 \text{ GeV}/c^2$ for the purpose of this analysis.

$\gamma + 3$ jets and $\gamma\gamma + 2$ jets CompHEP produced backgrounds

Partonic level events generation was done with CompHEP version 4.2p1 for $pp \rightarrow \gamma\gamma + 2$ jets subprocess with the preselection cuts:

- $p_T^\gamma > 20 \text{ GeV}$
- $p_T^j > 20 \text{ GeV}$
- $\Delta R_{ij} > 0.4$

34 partonic subprocesses for the QCD background and 20 partonic subprocesses for the elec-

troweak background were taken into account.

For the subprocess $pp \rightarrow \gamma + 3jets$ (50 partonic subprocesses, QCD misidentification background) preselection cuts were:

- $p_T^\gamma > 20$ GeV
- $p_T^j > 20$ GeV
- $\Delta R_{ij} > 0.4$
- two jets in opposite hemispheres with the rapidity gap greater than 3.5

The parameter values $\alpha_e = 1/137$ (i.e. $e = 0.3029$) and $\alpha_s = 0.1298$ ($g_s = \sqrt{4\pi\alpha_s} = 1.2772$), $m_Z = 91.19$ GeV, $\sin\theta_w = 0.481$, $\Gamma_Z = 2.44$ GeV, $\Gamma_W = 2.03$ GeV. PDF set was taken to be CTEQ5L, factorization scale 50 GeV.

Hadronization was done by PYTHIA and the same preselection was applied for both QCD $\gamma + 3jets$ and QCD $\gamma\gamma + 2jets$ CompHEP datasets, as it was described above for QCD hadronic jets in this Section. Rejection factors are 2.5 for $\gamma\gamma + 2jets$ dataset and 7.8 for $\gamma + 3jets$ dataset.

Simulation tools and Monte Carlo data samples

For the Higgs boson background study the following data samples were used:

- Signal:
 - $H \rightarrow \gamma\gamma$ at five different Higgs boson masses ($m_H=115, 120, 130, 140$ and 150 GeV/ c^2):
 - 115 GeV - 16000 events,
 - 120 GeV - 192920 events,
 - 130 GeV - 10700 events,
 - 140 GeV - 20000 events,
 - 150 GeV - 20000 events,
- Background:
 - 4.5M preselected QCD hadronic jet events;
 - 5.5M preselected $\gamma + jets$ events;
 - 1M preselected quark annihilation events;
 - 1M preselected gluon fusion events.
 - 1M preselected Drell Yan events
 - 40k preselected QCD $\gamma + 3jets$ CompHEP events
 - 200k preselected QCD $\gamma\gamma + 2jets$ CompHEP events
 - 41k preselected electroweak $\gamma\gamma + 2jets$ CompHEP events

All events were generated with PYTHIA 6.227 at $\sqrt{s}=14$ TeV with the CTEQ5L parton density structure functions. The simulation of particles propagated through the detector is done using OSCAR version 3.6.5.

The simulated events were digitized using the reconstruction and analysis program ORCA 8.7.3. Pileup events, corresponding to the luminosity of 2×10^{33} /cm²/s were added.

The numbers of generated and fully simulated events are shown in Table 10.18 for different types of background. In the last column the corresponding equivalent integrated luminosity is shown.

Table 10.18: Number of generated and simulated events for different types of background.

Background process	Number of generated events	Rejection rate	Number of simulated events	L_{intg} (fb^{-1})
QCD hadronic jets	31.2×10^9	6048	4.5M	~ 1
γ + jets	1.3×10^8	20.0	5.5M	~ 2
Gluon fusion	2.25×10^6	2	1M	~ 52
Quark annihilation	3.0×10^6	2.7	1M	~ 55
Drell Yan	1.0×10^6	1	1M	0.25
γ + 3jets QCD	0.3×10^6	7.8	40k	0.05
$\gamma\gamma$ + 2jets QCD	0.5×10^6	2.56	200k	6
$\gamma\gamma$ + 2jets EW	41×10^3	1	41k	120

Selection and reconstruction

Photon reconstruction

Energy deposited by electromagnetic showers in the ECAL are constructed into basic clusters using the island clustering algorithm. The minimum value of the transverse energy for the seed crystal of a basic cluster is 0.5 GeV. These basic clusters are in turn clustered into superclusters to recover the energy radiated by electrons or positrons from photon conversions, which fall outside the seed shower cluster. The hybrid algorithm is used to reconstruct the supercluster in the barrel, and the island algorithm is used in the endcap. The reconstructed supercluster is associated with a photon candidate.

The following photon selection was applied, which includes the trigger, kinematical and geometrical acceptance cuts. The 12 GeV Level-1 double-isolated electron trigger was applied to both the signal and background data samples. Both photon candidates had to match Level-1 trigger objects, such that, the distance R ($R = \sqrt{\delta\eta^2 + \delta\phi^2}$) between the photon candidate and trigger object be less than 0.5. The transverse energies of the two photon candidates were required to be greater than 40 GeV and 25 GeV respectively. The fiducial volume in rapidity was restricted to $|\eta| < 1.4442$ in the barrel and $1.566 < |\eta| < 2.5$ in the endcap for both photon candidates. Efficiencies of 12 GeV L1 double-isolated trigger, L1 matching procedure and photon candidates transverse energy cuts with invariant mass in the range 115 - 170 GeV/ c^2 can be found in Table 10.19).

Table 10.19: Efficiencies of 12 GeV L1 double-isolated trigger and L1 matching procedure with invariant mass in the range 115 - 170 GeV/ c^2 .

	signal	gluon fusion	quark annihilation	QCD jets	γ +jets	Drell Yan	γ +3jets QCD	$\gamma\gamma$ +2jets QCD	$\gamma\gamma$ +2jets EW
L1 trigger	92.2%	80.1%	79.9 %	1.7 %	15.2 %	13.2%	20.6%	60%	67.7%
L1 match	92.0 %	78.9 %	78.6 %	1.2%	13.9%	13.0%	20.2%	59.5%	67.5%

Energy correction

The difference between the photon measured energy and the true generated energy, $E_{\text{meas}} - E_{\text{true}}$, was considered as a function of the ratio $S1/S9$, where $S9$ is the energy in a 3×3 crystal matrix, centered around most energetic crystal, $S1$ is the energy of the most energetic

crystal in the matrix. This dependence on $S1/S9$ shows the influence of the photon impact point with respect to the crystal center. The measured energy E_{meas} was corrected for this difference.

Higgs boson vertex reconstruction

Three different algorithms were studied.

- **P_T balance** - the P_T balance for charged particle tracks along the reconstructed Higgs boson direction is defined as $P_T^B = -\sum P_{Ti} \cos \theta_i$, where θ_i is the angle between the Higgs boson and track i direction in the transverse plane
- **Maximal P_T** : the primary vertex is selected as the vertex with the track of highest P_T
- **Number of charged particle tracks** above P_T cutoff in pixel vertex. The primary vertex is selected as the vertex with a largest number of tracks.

To compare different vertex reconstruction algorithms, the number of events reconstructed in a $5 \text{ GeV}/c^2$ mass window are determined. The P_T balance and Maximal P_T algorithms give exactly the same number of events, while track counting algorithm gives a few percent less efficiency. The Higgs boson efficiency in $5 \text{ GeV}/c^2$ mass window is improved by 15%.

Isolation

Isolation based on the tracker

The tracker isolation criteria are based on the number of charged particle tracks with p_T greater than a p_T threshold, p_T^{thresh} , calculated in a cone R ($R = \sqrt{\delta\eta^2 + \delta\phi^2}$) around the photon candidate. The algorithm contains three parameters:

- The size of the cone R around the photon candidate, wherein the number of charged tracks is counted.
- The p_T threshold, p_T^{thresh} . Only charged particle tracks with p_T greater than p_T^{thresh} are considered in isolation calculations.
- The 'number of tracks' threshold N^{thresh} . If the number of charged particle tracks in cone R with p_T greater than the chosen p_T^{thresh} is greater than N^{thresh} , then the photon candidate is considered non-isolated, otherwise isolated.

The jet rejection factor is very sensitive to the 'number of tracks' threshold, N^{thresh} . By increasing N^{thresh} from 0 to 1, the Higgs boson signal efficiency is improved by 6-10%, but the jet rejection factor drops by a factor of ~ 2 . Therefore, the parameter N^{thresh} was fixed to zero. The cone size $R = 0.30$ and $p_T^{\text{thresh}} = 1.5 \text{ GeV}/c$ were used in this study.

Isolation based on the electromagnetic calorimeter

The isolation criteria are based on the sum of transverse energies deposited in basic clusters in some cone R ($R = \sqrt{\delta\eta^2 + \delta\phi^2}$) around a photon candidate. The basic clusters that belong to the photon candidate's supercluster are not counted as part of the sum. The algorithm contains four parameters:

- The size of the cone R around the photon candidate wherein the transverse energies deposited in the basic clusters are summed.
- The transverse energy sum threshold E_T^{thresh} . If the sum of transverse energies is below this threshold, the photon candidate is considered isolated, otherwise

non-isolated.

- The ratio, r , of the transverse energy sum in all surrounded basic clusters to the transverse energy of the most energetic super cluster.
- The ratio, hoe , of the super-cluster energy deposited in HCAL to the energy of this super-cluster, deposited in ECAL.

There is no strong dependence of the QCD hadronic jet rejection factor on the cone size R , though slightly better rejection factors are empirically obtained for a cone size $R = 0.30 - 0.35$. The cone size $R = 0.30$ is used in this study. The transverse energy sum thresholds, E_T^{thresh} , were chosen to be 1.2 GeV in the barrel and 1.6 GeV in the endcap. The ratio, $r = 0.01$, of the transverse energy sum in all surrounded basic clusters to the transverse energy of the most energetic super cluster was used for this study. The HCAL/ECAL ratio of the energies of the super-cluster, deposited in HCAL over the super-cluster energy deposited in ECAL was set to 0.1 .

Forward jet tagging

Jet tagging was done based on the reconstructed jets by E_t recombination scheme based on EcalPlusHcalTowers with 0.5 GeV threshold, ConeCut ΔR variable was set to 0.7 and JetEtCut variable was set to 10 GeV. No additional Monte Carlo calibration was done.

The selection criteria for the forward tagged jet are

$$p_T^{\text{jet}} > 20 \text{ GeV}/c, \quad |\eta_{\text{jet}}| \leq 4.5, \quad \Delta R_{\gamma\text{jet}} \geq 0.5$$

$$\Delta\eta_{\text{jets}} = |\eta_{\text{jet1}} - \eta_{\text{jet2}}| \geq 4.0, \quad \eta_{\text{jet1}} \times \eta_{\text{jet2}} < 0$$

Two additional cuts were applied to the already selected two forward jets in order to reduce the background even more than it was done with forward jet tagging procedure:

- $p_t^{\text{jet1}} > 50 \text{ GeV}$, where p_t^{jet1} is the transverse momentum of the first most energetic forward jet, selected by forward jet tagging procedure, described above.
- $p_t^{\text{jet2}} > 25 \text{ GeV}$, where p_t^{jet2} is the transverse momentum of the second most energetic forward jet, selected by forward jet tagging procedure, described above.
- $m_{j1j2} > 500 \text{ GeV}$, where m_{j1j2} is the invariant mass of the two most energetic forward jets, selected by forward jet tagging procedure, described above.

To be able to reduce the total number of events passed through all cuts from CompHEP background sample $\gamma\gamma + 2\text{jets}$ the cut on the transverse momentum of the second most energetic forward jet, selected by forward jet tagging procedure, was adjusted:

- $p_t^{\text{jet2}} > 35 \text{ GeV}$, where p_t^{jet2} is the transverse momentum of the second most energetic forward jet, selected by forward jet tagging procedure, described above.

To suppress CompHEP background further, additional cut was introduced:

$$- \min[\eta_{\text{jet1}}, \eta_{\text{jet2}}] + 0.7 < \eta_{\gamma1,2} < \max[\eta_{\text{jet1}}, \eta_{\text{jet2}}] - 0.7$$

Results

Only the results, produced by CompHEP, will be presented here.

Usual $2 \rightarrow 2$ subprocesses in PYTHIA are contain part of NLO contributions if ISR and FSR

Table 10.20: Factorization procedure for all CompHEP-produced background samples, no mass window

	$\gamma\gamma+2\text{jets EW}$	$\gamma\gamma+2\text{jets QCD}$	$\gamma+3\text{jets QCD}$
Total Nev	41000	192227	37559
Prep.cuts	4938	7359	1558
Isol cuts	1000	847	16
Jet cuts	2019	617	228

are switched on. But this contribution is adequate only in cases when isolation of emitted photon or gluon from quark or gluon is not very large. If this isolation required to be large enough, PYTHIA underestimates this contribution. For the purposes of this study, this part of NLO contribution in PYTHIA will be named further as ‘‘PYTHIA NLO’’.

CompHEP datasets have $2 \rightarrow 4$ subprocesses with further FSR and ISR switched on in PYTHIA during the hadronization. These CompHEP complete set of diagrams can be considered as NLO for usual PYTHIA $2 \rightarrow 2$ subprocesses with ‘‘PYTHIA NNLO’’ in terms of previous paragraph.

Keeping in mind that CompHEP cannot calculate box-like diagrams, three PYTHIA background samples - quark annihilation, $\gamma + \text{jets}$ and QCD hadronic jets - can be replaced by four CompHEP calculated background samples - $\gamma + 3\text{jets}$ and $\gamma\gamma + 2\text{jets}$ (QCD and electroweak). This replacement will change picture substantially.

$\gamma + 3\text{jets QCD and electroweak backgrounds estimation}$

Due to the limited statistics, rescaling and factorization should be applied to reduce statistical errors.

Factorization procedure was done as follows (see Table 10.20). All analysis was splitted on three different subsets of cuts:

- 1) all preliminary cuts were done to make sure that we have two photon candidates and two jets in event, passed through the requirements : two have L1 electromagnetic isolated trigger, to have more than two photons matched with L1 trigger objects, to have more than two jets and event passed through jet tagging procedure. This subset of cuts will be mentioned as ‘‘prep. cuts’’ in Table 10.20.
- 2) isolation cuts - all cuts including photon E_t cuts, track and ECAL isolation for both photons, $P_T^H > 50 \text{ GeV}/c$, HCAL/ECAL ≤ 0.1 . All these cuts will be mentioned in Table 10.20 as ‘‘isol. cuts’’.
- 3) jet cuts - all cuts including $p_T^{jet1} > 50 \text{ GeV}/c$, $p_T^{jet2} > 35 \text{ GeV}/c$, $M_{j1j2} > 500 \text{ GeV}/c^2$, and geometry cut: $\min[\eta_{jet1}, \eta_{jet2}] + 0.7 < \eta_{\gamma1,2} < \max[\eta_{jet1}, \eta_{jet2}] - 0.7$.

After that splitting, we will pass all events through first subset of cuts, preliminary cuts, and will pass through last two subset separately to increase the statistics of events passed through the cuts.

Based on QCD samples $\gamma+3\text{jets}$ and $\gamma\gamma+2\text{jets}$ we can estimate rejection factors for the photon to be a photon and for jet to be a photon.

From QCD $\gamma\gamma+2\text{jets}$ sample: $(r_{\gamma \rightarrow \gamma}^{QCD})^2 = 847/192227$. From $\gamma+3\text{jets QCD}$ sample: $(r_{\gamma \rightarrow \gamma}^{QCD}) * (r_{jet \rightarrow \gamma}^{QCD}) = 16/37559$.

From these two equations we have got next values for $r_{\gamma \rightarrow \gamma}^{QCD}$ and $r_{jet \rightarrow \gamma}^{QCD}$:

$$r_{\gamma \rightarrow \gamma}^{QCD} = 0.066, \quad r_{jet \rightarrow \gamma}^{QCD} = 6.45 * 10^{-3}$$

Keeping this in mind, we can get $r_{\gamma \rightarrow \gamma}^{EW}$ value from $\gamma\gamma + 2jets$ electroweak CompHEP background sample: $(r_{\gamma \rightarrow \gamma}^{EW})^2 = 1000/41000$, hence $r_{\gamma \rightarrow \gamma}^{EW} = 0.156$.

To be able to estimate electroweak part of $\gamma+3jets$ ComHEP background, we need to know $r_{jet \rightarrow \gamma}^{EW}$.

Main assumption here is $r_{jet \rightarrow \gamma}^{EW} = r_{jet \rightarrow \gamma}^{QCD} = 6.45 * 10^{-3}$.

The rest jet cuts give us additional factor at the level from 0.084 to 0.15. We will use $r_{jet}^{EW} = 0.1$.


Now we can calculate an average number of events in mass window 5 GeV (rescaling probability will be again 1/12 as above in Scenario I) around each invariant mass of Higgs boson. Keeping in mind that total cross section of $\gamma+3jets$ CompHEP background will be 5.15 pb, and normalizing to 60 fb^{-1} , we can count the events:

$$N_{ev} = 5150 \text{ fb} * 60 \text{ fb}^{-1} * r_{jet \rightarrow \gamma}^{EW} * r_{\gamma \rightarrow \gamma}^{EW} * r_{jet}^{EW} * 1/12 = 5000 \text{ fb} * 30 \text{ fb}^{-1} * 0.156 * 6.45 * 10^{-3} * 0.1 * 1/12 = 30/12 \approx 2.5 \text{ ev}$$

So, in a total, we got an upper limit on the number of events in 5 GeV mass window - 2.5 events normalized to 60 fb^{-1} .

Based on this digit, we will estimate the upper limit on the number of events in $\gamma + 3jets$ QCD CompHEP background. Looking at the Table 10.21, we will make **second assumption - the ratio of events between $\gamma\gamma+2jets$ QCD and EW will be the same as $\gamma+3jets$ QCD and EW**. This assumption is a bit more questionable than previous one, but in this case we will again getting larger numbers of events through this estimation than if we would produce enough statistics, because $\gamma+3jets$ QCD background will suppressed remarkably well and our procedure, likely, overestimates the number of events passed through the analysis.

The data themselves will be looked at to estimate and cross-check the total background at very high energy. If Higgs boson is discovered, the number of events in the side-bands of the Higgs boson and their mass dependence will be used to estimate the number of background events under the peak, provided there is enough statistics in the side-bands. In this approach, the uncertainties on the background cross-sections, the PDF and on the luminosity measurement are highly reduced. For Higgs boson production the two regions left and right of the Higgs boson can be used. We estimate a reduction of the signal significance of the order of 10-20%. The dependency of significance vs Higgs boson invariant mass for side-band method of background estimation, normalized to 30 and 60 fb^{-1} , can be found on Figure 10.23.

 This case Higgs boson signal and significance, normalized to 60 fb^{-1} , are shown in Table 10.21.

10.2.6 Associated WH production with $H \rightarrow WW^{(*)} \rightarrow 2\ell 2\nu$

The cross-section for this process exhibits a maximum near the WW resonance, due to the combined behavior of the production cross-section and the Higgs boson branching ratio. We

Table 10.21: Higgs boson signal for different masses and significance, normalized to 60 fb^{-1} (mass window $5.0 \text{ GeV}/c^2$).

	$m_H = 115$ GeV/c^2	$m_H = 120$ GeV/c^2	$m_H = 130$ GeV/c^2	$m_H = 140$ GeV/c^2	$m_H = 150$ GeV/c^2
N_s	20.2	21.1	19.1	15.7	11.2
gluon fusion	0	0	0	0	0
$\gamma+3\text{jets}$ (QCD)	2.7	4.7	3.5	2.0	5.8
$\gamma+3\text{jets}$ (EW)	2.5	2.5	2.5	2.5	2.5
$\gamma\gamma + 2\text{jets}$ (QCD)	11.2	13.2	9.85	8.9	4.6
$\gamma\gamma + 2\text{jets}$ (EW)	10	7.0	7.0	11.0	2.0
Drell Yan	0	0	0	0	0
N_b	26.0	26.2	21.4	28.2	14.9
δN_b	2.8	3.2	2.4	3.0	1.8
S	3.07	3.15	3.21	2.32	2.30




Figure 10.23: Significance vs Higgs boson invariant mass, normalized to 30 and 60 fb^{-1} for instant luminosity $2 * 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$

Table 10.22: Background processes considered into the present analysis. The cross-section includes the decay of W and Z bosons into leptons. The generator and the number of events processed are also shown together with the corresponding weight for a luminosity of $1fb^{-1}$. For comparison, the signal varies, with the same hypotheses, from 5 fb to 20 fb in the considered mass range.

Background	Cross-section	Generator	MC statistic	weight ($1fb^{-1}$)
WWW($3l^{\pm}$)	4.95 fb	CompHep	10000	5.19×10^{-4}
WZ($3l^{\pm}$)	1.71 pb	Pythia	50000	3.46×10^{-2}
ZZ($4l^{\pm}$)	0.17 pb	Pythia	50000	3.67×10^{-3}
$t\bar{t}(1^{+}1^{-}b\bar{b})$	90.9 pb	TopRex	100000	0.93
Wt($1^{+}1^{-}b$)	5.25 pb	TopRex	50000	0.11

will therefore focus on the intermediate mass region between $120\text{ GeV}/c^2$ and $190\text{ GeV}/c^2$, where the cross-section exceeds 300 fb. We decided to investigate events containing three leptons in the final state, concentrating on electrons and muons (including leptonic tau decays).

10.2.6.1 Signal and background generation


Starting at $115\text{ GeV}/c^2$, we considered Higgs boson with masses of 125, 130, 140, 150, 160, 170, 180 and $190\text{ GeV}/c^2$. Events were generated with PYTHIA for each of the nine Higgs boson masses, without any kinematical cut. W bosons are forced to decay leptonically (e, μ, τ).

All Standard-Model processes likely to produce three leptons must be considered as background for this analysis. This includes events where three leptons are actually produced in the hard process but also events with a fake or missed lepton. One particular case is the production of leptons in the semi-leptonic decay of a B meson. In the present analysis, we considered the production of WWW, WZ, ZZ, $t\bar{t}$, and Wt. Most of the processes are simulated with PYTHIA, except for WWW, which is generated with CompHep, and Wt generated with TopRex. In all cases, PYTHIA is used for the hadronization step. Table 10.35 shows the cross-section, the generator used and the number of events produced.


10.2.6.2 Selection streams at Level-1 and HLT

The global (cumulative) trigger efficiency after L1 and HLT is found to reach 72% for a $140\text{ GeV}/c^2$ Higgs boson using the full trigger table. Main contributions come from single and double leptonic (e and μ) triggers (65%). There is a small contribution from the missing transverse energy trigger (E_T^{miss}) and from combined ($e \wedge \tau$) and ($\mu \wedge \tau$) triggers, further reduced by the event selection, which favors multi-leptonic patterns. For this analysis, we restricted ourselves to triggers known to have the highest impact on the total efficiency: single- and double-electron and muon triggers. Figure 10.42(a) shows the efficiency for each (exclusive) trigger pattern, given the above choice of interesting bits.

Details about the efficiency for each type of event (defined from the number of muons, electrons and taus in the event) are given in Figure 10.42(b). Events containing one or more muons are more easily retained (efficiency reaches 85% for events with three muons) while tau events are only marginally selected (efficiency: 12%). Efficiency rises slightly with the Higgs boson mass, from 58% at $115\text{ GeV}/c^2$ to 74% at $190\text{ GeV}/c^2$.



(a)



(b)

Figure 10.24: (a) Trigger efficiency by trigger pattern, for the signal. Efficiency is calculated as N_x/N_{HLT} , where x is one of the 2^4 exclusive trigger classes. “Others” stands for unconsidered trigger patterns; (b) Trigger efficiency for each class of Monte-Carlo events. Results are given after L1 and after HLT. Efficiency is computed as the ratio between the number of triggered events and the total number of generated events.

Table 10.23: Trigger efficiency for each source of background. Efficiency at HLT with the restricted trigger set used in the present analysis is also presented.

Background	L1 efficiency	HLT efficiency	$e, ee, \mu, \mu\mu$ HLT efficiency
WWW($3l^\pm$)	0.87	0.79	0.73
Wt($1^{+1-}b$)	0.88	0.78	0.67
WZ($3l^\pm$)	0.8	0.72	0.65
ZZ($4l^\pm$)	0.78	0.69	0.64
$t\bar{t}(1^{+1-}b\bar{b})$	0.91	0.79	0.65

Table 10.36 shows the trigger efficiency for each source of background. Efficiency, when restricting ourselves to single- and double-electron and muon triggers, varies from 64% to 73%, which is the same magnitude as the trigger efficiency for signal events. This restricted trigger is 15% (for $t\bar{t}$) to 5% (for ZZ) less efficient than the inclusive High-Level trigger.

10.2.6.3 Off-line selection

Electrons and muons are reconstructed using default offline reconstruction algorithms. For electrons, additional quality cuts are applied: the energy measured by ECAL and the momentum obtained by the tracker must agree within 50%, and the ratio of energy measured by HCAL and ECAL must be lower than 0.15. Only leptons with $P_t^l > 14 \text{ GeV}/c$ are retained. A first set of selection criteria is applied to select signal-like topologies, requiring three and only three leptons, for a total charge of either +1 or -1. A cut on the distance in the z direction between the points of closest approach of lepton tracks to the beam is applied to ensure that all of the three leptons are coming from the same interaction. The two closest (in the $\eta - \phi$ plane) opposite-sign leptons are then assigned to the Higgs boson decay. The angle between leptons attributed to the Higgs boson can be used to distinguish signal

and background. The acollinearity between two leptons is defined as the angle between the two leptons, in the space, and their acoplanarity is defined as the same angle projected onto the transverse plane. Both the acollinearity ($\theta_{\text{aco}} < 1.75$ rad) and the acoplanarity ($0.1 \text{ rad} < \phi_{\text{aco}} < 0.75$ rad) between the leptons are used, as they provide complementary information. Leptons required to be isolated in the tracker, i.e. the angle between the lepton's track and the closest track above $3 \text{ GeV}/c$ must be more than 0.2. A jet veto is applied rejecting events with a jet, reconstructed with the iterative cone algorithm ($r=0.7$), above 25 GeV in the central region ($|\eta| < 2.1$). An additional B veto is applied, imposing that no single B-jet is reconstructed by the default *combined B-tag* algorithm. This removes low-energy bjets passing the standard jet veto. A cut on the invariant mass of any pair of leptons compatible with the Z hypothesis (via charge flavor and invariant mass constraints, $M_Z \notin [65, 115]$) is used to reject ZZ and WZ events. Finally, kinematical cuts are used: $E_T^{\text{miss}} > 50 \text{ GeV}/c$, $M_T(W_3) > 40 \text{ GeV}/c^2$ and $Pt(\sum \vec{P}_l) > 40 \text{ GeV}/c$, where $Pt(\sum \vec{P}_l)$ is the transverse momentum of the vector sum of momenta of all three leptons, and $M_T(W_3)$ is the reconstructed transverse mass of the associated W boson:

$$M_T(W_3) = \sqrt{2 * P_T^{l_3} E_T^{\text{miss}} (1 - \cos \Delta\phi_{l_3 p_T})}, \quad (10.3)$$

with $P_T^{l_3}$ being the transverse momentum of the lepton not associated to the Higgs boson, E_T^{miss} the missing transverse momentum, and $\Delta\phi_{l_3 p_T}$ the polar angle between the lepton and the missing transverse momentum. Optimized cuts are summarized in Table 10.37.

The Higgs boson transverse mass is computed from the two chosen leptons and from the missing transverse momentum:

$$M_T(H) = \sqrt{M_T^{ll^2} + 2E_T^{ll} p_T - 2P_T^{ll} p_T \cos \Delta\phi_{ll p_T}}, \quad (10.4)$$

Figure 10.43 shows the distribution of this quantity for the signal, on top of remaining background, after all cuts for a Higgs boson mass of $140 \text{ GeV}/c^2$ and an integrated luminosity of 100 fb^{-1} . The cumulated efficiency (including trigger and event selection) depends on the Higgs boson mass hypothesis. Starting at 0.5% for a mass hypothesis of $115 \text{ GeV}/c^2$, the efficiency rises to a maximum at the WW resonance (1.3%). Beyond the WW production threshold, efficiency drops since W bosons start to be boosted in the Higgs boson frame, which influences the angular distribution of leptons. Efficiency in that region could certainly be improved by optimizing the analysis for a Higgs boson mass of $190 \text{ GeV}/c^2$.

10.2.6.4 Systematic uncertainties

Systematic sources considered in this study are related to the normalization of backgrounds, to the reconstruction, the event selection, the luminosity and the structure functions of protons.

The uncertainty on the background normalization is usually estimated from the scale-dependence of the NLO cross-sections. This scale dependence is not known to us. This uncertainty can be large, since NLO effects are important. For events of interest for this study, the differences between leading-order and next-to-leading-order calculations are large: 45% for $t\bar{t}$, 40% for WZ and 25% for ZZ events. Before tuning them with real data, and thus especially at low

[‡]No energy correction is applied since we are mostly handling fake low-energy jets from pile-up and underlying events.

Table 10.24: Summary of the optimized selection cuts. The cross-section for the signal and backgrounds, for each step in the selection, is given in fb. An upper limit for the Wt and $t\bar{t}$ cross-sections is given when no simulated event remains.

Cut		Signal (fb) 140 GeV/c ²	Background (fb)				
Id.	Type		$t\bar{t}$	Wt	ZW	ZZ	WWW
0	L1 \wedge HLT	12.24	72067	4115.8	1238.4	118.438	3.91
1	$N_{lept} = 3, ch_{\Sigma} = 1$	3.81	16432.7	680.0	339.4	34.65	1.05
2	Lepton cuts	2.67	5629.1	245.3	245.9	23.53	0.70
3	Angular cuts	0.87	400.6	15.0	18.3	2.29	0.11
4	B veto	0.43	3.85	0.42	9.77	1.19	0.06
5	Jet veto	0.27	< 1.93	0.31	7.26	0.58	0.04
6	Z veto	0.21	< 1.93	0.21	0.40	0.08	0.03
7-9	Topological	0.13	< 1.93	< 0.11	0.06	0.01	0.02




Figure 10.25: Reconstructed transverse mass from Equation 10.10 for a 140 GeV/c² Higgs boson and an integrated luminosity of 100 fb⁻¹.




Figure 10.26: Distribution of the acoplanarity for pseudo-experiments, fitted by a signal+background shape, as described in the text.

integrated luminosity, it will be impossible to trust background estimates from Monte-Carlo programs. Hence, background will be normalized to signal-free regions of the phase-space.

By looking at the acoplanarity distribution when the angular cuts are not applied, data can be fitted to a sum of signal and background shapes. For that purpose, the signal is described by a sigmoid distribution, while the background remains constant. The Monte Carlo distribution for Signal and background are first fitted independently, and the shapes obtained that way are used to fit data from pseudo-experiments (Figure 10.44). The uncertainty on the background normalization is then related to the uncertainty on the background level in that fit. The uncertainty on the background level is found to be 15% for an integrated luminosity of 100 fb^{-1} , and rises up to 20% for 30 fb^{-1} . That value will be used in the following.

Reconstruction and selection uncertainties mainly arise from the jet veto, the b veto and lepton reconstruction. Experience from Tevatron tells us that a typical 2% uncertainty on lepton reconstruction efficiency has to be considered, while 5% uncertainty comes from lepton isolation [474]. Since three leptons are present in our analysis, 12% uncertainty from lepton reconstruction and selection has been assumed. The additional uncertainties from the jet veto and the b veto will be assumed to be 5% each, which also seems reasonable in the light of Tevatron data.

To take into account other uncertainties related to the event selection, cuts are varied within the resolution of the associated quantity. The signal efficiency and background rejection are found to be stable with respect to such variations. A conservative value of 3% for the associated uncertainty is considered in the following.

The last considered uncertainty comes from the product of the luminosity and the proton structure functions, known as the parton luminosity. Considering these two quantities separately, 5% uncertainty on the luminosity is generally assumed, while the uncertainty from the proton parton distribution function (PDF) is taken to be 4% [475]. This latter uncertainty is reduced by the process considered, for which the mid-x region (where uncertainties are small) dominates. This gives an uncertainty of 6.4% on the parton luminosity.

The additional source of systematic uncertainties arising from the limited Monte Carlo statis-




Figure 10.27: (a) Luminosity needed to obtain a 5σ significance using the likelihood-ratio method, with systematics only, Monte-Carlo statistical uncertainties only, or with both effects considered; (b) luminosity needed to exclude a Higgs boson at 95% C.L. if no excess is observed, using the same method.

tics is also considered in the following result. With the Likelihood ratio method we are using, this is done bin per bin in the distributions of signal and background, so that a single value cannot be quoted. For the time being, this has a large impact on the results, but this effect will easily be reduced in the future, as more events become available.

10.2.6.5 Signal significance

In order to integrate the effect of systematic uncertainties and to exploit the discriminative power from the transverse mass distribution, the likelihood-ratio method (S_{CL}) is used. Figure 10.45(a) shows the luminosity needed to obtain a 5σ significance using this method, with systematics only, with Monte-Carlo statistical uncertainties, or with both effects considered. Figure 10.45(b) shows the luminosity needed to exclude a Higgs boson at 95% C.L. if no excess is observed, using the same method. Less than 50fb^{-1} are required in most of the mass range, while only 20fb^{-1} are needed at $170 \text{GeV}/c^2$.

One important motivation for studying this channel is also that it is one of the only allowed signatures for a fermiophobic Higgs boson model. If the Higgs boson does not couple to fermions, the usual gluon-fusion diagrams are indeed forbidden, as well as $b\bar{b}$ decays. A fermiophobic Higgs boson will present a large cross-section at low mass, as the branching ratio does not drop down as in the Standard Model. Figure 10.46(a) shows the luminosity needed to obtain a 5σ significance for a fermiophobic Higgs boson. Compared to Figure 10.45, the needed luminosity is found to be similar in the most favorable mass region for the Standard Model (around $170 \text{GeV}/c^2$) and above, but far better results are obtained in the low mass region. After 100fb^{-1} , all masses between the LEP limit and $175 \text{GeV}/c^2$ will be covered by this analysis alone. Figure 10.46(b) shows the luminosity needed to exclude a fermiophobic Higgs boson at 95% C.L. if no excess is observed. In the absence of signal, less than 30fb^{-1} are required to reject any fermiophobic Higgs boson up to $175 \text{GeV}/c^2$.




Figure 10.28: Results obtained using the benchmark fermiophobic model; (a) Luminosity needed to obtain a 5σ significance using the likelihood-ratio method, with systematics only, Monte-Carlo statistical uncertainties only, or with both effects considered; (b) luminosity needed to exclude a Higgs boson at 95% C.L. if no excess is observed, using the same method.

10.2.6.6 Measurement of Γ_W

In the present study, the partial decay width of the Higgs boson into W bosons, Γ_W , is determined with a good accuracy. This is possible using the narrow width approximation and benefiting from the fact that only the g_{HWW} coupling is involved in the Feynman diagram for the signal.

From the number of observed events, and assuming a good description of the background by the Monte-Carlo simulations, the product of the signal cross-section and the branching ratio of the Higgs boson into WW can be computed as:

$$\sigma \times BR = \frac{N_s}{\epsilon \mathcal{L}} = \frac{N - N_b}{\epsilon \mathcal{L}}, \quad (10.5)$$

with ϵ representing the combined trigger and selection efficiency. The uncertainty on this measurement is then found to be

$$\left(\frac{\Delta(\sigma \times BR)}{\sigma \times BR} \right)^2 = \frac{(\Delta N)^2}{N_s^2} + \frac{(\Delta N_b)^2}{N_s^2} + \frac{(\Delta \mathcal{L})^2}{\mathcal{L}^2} + \frac{(\Delta \epsilon)^2}{\epsilon^2}. \quad (10.6)$$

In this expression, ΔN corresponds to the statistical uncertainty on N , and will be assumed to be \sqrt{N} . The other three ΔN_b , $\Delta \mathcal{L}$ and $\Delta \epsilon$ quantities are dominated by systematic error sources described in Section 10.2.9.4.

In the narrow width approximation, the measured product of signal cross-section and branching ratio of the Higgs boson into WW can also be expressed from partial and total widths of the Higgs boson:

$$\sigma \times BR = \frac{\sigma^{SM} \Gamma_W}{\Gamma_W^{SM}} \times \frac{\Gamma_W}{\Gamma_{tot}} \equiv T \Gamma_W^2. \quad (10.7)$$




Figure 10.29: (a) Resolution on the product of cross-section and branching ratio into WW as a function of the Higgs boson mass, for an integrated luminosity of $100fb^{-1}$. (b) Resolution on the Γ_W partial width as a function of the Higgs boson mass, for the same integrated luminosity.

This allows us to write the uncertainty on the Higgs boson width into W bosons as

$$\left(\frac{\Delta\Gamma_W}{\Gamma_W}\right)^2 = \frac{1}{4} \left(\frac{(\Delta\sigma \times BR)^2}{(\sigma \times BR)^2} + \frac{(\Delta T)^2}{T^2} \right). \quad (10.8)$$

In this expression, the uncertainty on the cross-section is the quantity obtained in Equation 10.12, and the uncertainty on the T factor comes from theoretical uncertainties and expected experimental uncertainties on Γ_{tot} . Assuming $100fb^{-1}$ luminosity and the aforementioned systematic sources, the uncertainty on the cross-section is found to be 36% for a $150\text{ GeV}/c^2$ Higgs boson. The corresponding uncertainty on the Γ_W partial width is 21%.



These results depend on the Higgs boson mass, and are presented Figure 10.47.

10.2.7 Associated $t\bar{t}H$ production with $H \rightarrow \gamma\gamma$

10.2.7.1 Introduction

A Standard Model or two-doublet neutral Higgs boson produced in association with a $t\bar{t}$ pair, with $H(h^0) \rightarrow \gamma\gamma$ would share a fully reconstructible mass peak with the inclusive $H(h^0) \rightarrow \gamma\gamma$ signature. But like the WH and ZH channels[476], the signature could contain an isolated high-transverse-momentum charged lepton which can be used both to discriminate against QCD background and reconstruct the primary vertex; the associated production channels could hence be less dependent on photon energy resolution. In particular, the presence of two top quarks would tend to produce high-multiplicity events, which could offer additional discriminating power against light jet QCD background. And in the case of the two-doublet MSSM, the gluon fusion Higgs production channel could in fact be subject to suppression with respect to the associated production channels in the case of top-stop

degeneracy (“maximal mixing”)[477]. Prior generator-level studies for the detection of the SM[478] and MSSM [479] Higgs bosons in CMS via this channel have indicated a signal-to-background ratio of approximately 1. A full simulation study in the ATLAS Physics Technical Design Report [480] has predicted a signal significance of $S\sqrt{B} = 4.3-2.8$ for $m_H = 100-140$ GeV with a signal efficiency of $\sim 30\%$. A more recent, related ATLAS study involving a 2-photon signature accompanied by missing energy [481] has indicated, for 100fb^{-1} , a signal-to-background ratio of ~ 2 for $m_H = 120$ GeV.

10.2.7.2 Signal production cross-sections, event rates and event generation

Production cross-sections for $t\bar{t}H$ have been calculated at next-to-leading order [465, 153, 466]. Taking the branching ratio for $H \rightarrow \gamma\gamma$ from HDECAY [9] and assuming in addition that the decay of exactly one of the top quarks yields a lepton (electron or muon) from $W^\pm \rightarrow l + \nu_l$ (including the possibility of tau lepton decays to muons or electrons), we estimate for several Higgs boson masses the number of signal events for 30 and 100 fb^{-1} (Table 10.25):

Table 10.25: Estimated number of signal events for $t\bar{t}H, H \rightarrow \gamma\gamma$, assuming NLO production cross sections [153], Higgs branching ratios to 2 photons [9], and 1 electron or muon from top decay (including from tau lepton decays).

Higgs Boson Mass (GeV)	After 30 fb^{-1}	After 100 fb^{-1}
115	20.80	69.33
120	19.61	65.36
130	15.96	53.20
140	11.20	37.33

Figure 10.30 shows typical Feynman diagrams of the signal process.




Figure 10.30: Typical diagrams for $t\bar{t}H$ production with $H \rightarrow \gamma\gamma$ and at least one charged lepton from the decay of a top or antitop quark.

Signal events were generated with both the MADGRAPH [73, 482, 483] and ALPGEN [152, 484, 485] LO exact matrix element generators, for each of the Higgs boson masses shown in Table 10.25. Events from both generators were found to yield comparable LO cross-section results and kinematical distributions. The LO cross-sections were also found to agree with those from the program HQQ [8] at the percent level. The samples analyzed were those generated with ALPGEN, containing ~ 4500 simulated and reconstructed events for each of the four Higgs boson mass values.

It should be noted that for the current study all signal events have been generated such that exactly one of the two W bosons from the two top quarks decays leptonically. It can be assumed, however, that the event selection which will be described below will also have some efficiency for events where both W bosons yield leptons, thus potentially increasing the total number of signal events expected to be observed.

10.2.7.3 Background Processes Considered and Event generation

Standard Model processes resulting in both irreducible and reducible backgrounds have been identified. A background is called irreducible if it is capable of giving rise to the same signature on the particle level as that searched for in a signal event, that is to say, a lepton and two photons ($l\gamma\gamma$). Among the irreducible processes, special care has been taken to properly treat the $t\bar{t}\gamma\gamma$ background. Feynman diagrams of three possible types of $t\bar{t}\gamma\gamma$ processes considered are shown in Figure 10.31. In the first case, called “Type 1”, both photons are radiated from either outgoing top quark or incoming parton lines. In the third case, called “Type 3”, both are radiated from top quark decay products. The second case, “Type 2” combines one photon radiated according to “Type 1” with the second radiated according to “Type 3”. (A fourth process arises from both photons being radiated from different decay products of the *same* top quark; for the relevant event selection (see pertinent section below) with $m_{\gamma\gamma} > 70$ GeV we have verified that this contribution is completely negligible). Since at the time of undertaking this study no matrix element generator included either the Types 2 or 3 processes, a collaboration was begun with the authors of ALPGEN to add them. Also added to ALPGEN was the process $W\gamma\gamma + 4$ jets. Where applicable in the ALPGEN samples, top quarks and W bosons are decayed within ALPGEN which assures preservation of spin correlation information which could impact kinematical distributions.




Figure 10.31: A subsample of the relative Feynman graphs illustrating the three types of $t\bar{t}\gamma\gamma$ processes.

Table 10.26 lists the considered irreducible background processes, the generators used to either generate or cross-check event samples, the LO cross-section with statistical errors, the number of events expected for 30 (100) fb^{-1} of integrated luminosity, the number of events generated, simulated, reconstructed and analyzed as well as the equivalent integrated luminosity, which ranges from 400 to over 6000 fb^{-1} . The cross-sections reflect preselection criteria imposed at generator-level which are described in the next section. In the processes involving real top quarks as well as in the $W\gamma\gamma + 4j$ process, one top quark/the W boson was forced to decay leptonically, and the stated cross-section therefore implicitly includes the relevant branching ratio. It can be seen that the effect of the inclusion of background Types 2 and 3 is to augment the total initial contribution (before selection) from $t\bar{t}\gamma\gamma$ by approximately one order of magnitude.

A background is called reducible if at least one element of the final-state signature is mistakenly identified due to incomplete detector coverage or other instrumental effects. This could arise if one or more electrons or jets are misidentified as photons, or a jet as an electron or a muon. It has been heretofore possible to evaluate only the irreducible backgrounds discussed

Table 10.26: Cross-sections at leading order (statistical errors in parentheses), number of events generated, simulated and reconstructed, and equivalent integrated luminosity for the irreducible backgrounds considered.

Process	$\sigma \times \text{BR}$ [fb] (1 $W \rightarrow l\nu$)	Ngen	N 30 fb ⁻¹	N 100 fb ⁻¹	Generator	N simul./ reconstr.	N Anal.	Anal. Eq. Int. Lumi. [fb ⁻¹]
tt $\gamma\gamma$ 1	1.6 ($\leq 1/\text{mil}$)	52202	48	160	AL, MG	10000	4695	6250
tt $\gamma\gamma$ 2	6.1 ($\leq 1\%$)	6238	183	610	AL	6000	5109	1000
tt $\gamma\gamma$ 3	4.9 ($\leq 1\%$)	2967	147	490	AL	2500	2250	510
W $\gamma\gamma$ 4j	11.5 (1.2%)	4587	345	1150	AL	4500	3957	400

above with acceptable statistics, so only these will be presented here. Low-statistics tests on most of the reducible background processes have been performed, and strong requirements have been implemented in the following selection in order to veto them.

10.2.7.4 Event simulation and reconstruction

All generated signal and background events were fragmented and hadronized with PYTHIA [238, 62] version 6.227, using the CTEQ5L [29] PDFs. They were then simulated, digitized and reconstructed using the standard CMS tools. All samples were digitized with high-luminosity ($10^{34} \text{cm}^{-2} \text{s}^{-1}$) pileup.

10.2.7.5 Description of Kine preselections

No generator-level preselections were made on signal events. For the irreducible background events, the following preselection was made:

- $m_{\gamma\gamma} \geq 80$ GeV for all four processes;
- $p_{T\gamma} \geq 20$ GeV, $|\eta_{\gamma}| \leq 2.5$ (MADGRAPH) or $p_{T\gamma} \geq 15$ GeV, $|\eta_{\gamma}| \leq 2.7$ (ALPGEN) for all four processes;
- $p_{Tl} \geq 15$ GeV for all processes except tt $\gamma\gamma$ 1;
- $p_{Tj} \geq 15$ GeV, $|\eta_{j,l}| \leq 2.7$, $\Delta R(l,j \text{ or } j,j \text{ or } \gamma,j \text{ or } \gamma,\gamma) \geq 0.3$ for the process W $\gamma\gamma$ 4j, where 'j' refers to one of the four additional light quark jets;

where p_T refers to the transverse momentum of the particle, η its rapidity and $\Delta R = \sqrt{(\Delta\eta)^2 + \Delta\phi^2}$ where ϕ is the azimuthal angle.

The intersection (most restrictive set) of the above generator-level criteria except that pertaining to the additional light quark jets was then imposed on all signal and background event samples at the particle level.

10.2.7.6 Description of selection streams at Level-1 and HLT

The events are passed through the online Level-1 and High-Level-Triggers (HLT) simulation, configured for high luminosity ($10^{34} \text{cm}^{-2} \text{s}^{-1}$). Table 10.27 shows the response of the CMS Level-1 and global HLT triggers to each signal and background sample, normalized to the number of events having both true Higgs photons (at the particle level) within the CMS electromagnetic calorimeter acceptance ($|\eta| < 2.5$). Also shown is the percentage of events triggered incrementally by each of the several HLT subtriggers, normalized to the global HLT response, which demonstrates that between 98 and 99% of within-acceptance signal events

should be triggered by the photonic HLT subtriggers alone[§], and over 99% if the single muon subtrigger is added.

Table 10.27: The response of the CMS Level-1 and global HLT triggers to each signal and background sample, normalized to the number of events having both true Higgs photons within the CMS electromagnetic calorimeter acceptance ($|\eta| < 2.5$). Also shown is the percentage of events triggered incrementally by each of the several HLT subtriggers, normalized to the global HLT response.

trigger	115	120	130	140	ttgg	ttgg2	ttgg3	W2 γ 4j
$ \eta < 2.5$	88.16	87.47	88.33	88.04	100	100	100	100
L1 + HLT	97.15	97.42	97.92	98.11	94.09	91.89	83.29	92.55
gg	95.58	96.00	96.56	96.65	93.78	92.32	93.86	92.74
+ g	98.57	98.59	99.03	99.14	97.19	97.35	97.12	96.04
+ μ	99.71	99.57	99.76	99.78	99.16	99.07	99.09	98.80
+ e	99.85	99.78	99.85	99.88	99.45	99.37	99.79	99.26

10.2.7.7 Description of off-line selection

In some prior studies of associated Higgs production [476], the two photons with the highest values of p_T have been assumed to have come from the Higgs boson. However we have observed, first at particle-level and now in the study of reconstructed events (see the pertinent curve in Figure 10.33(left)) that the use of this criterion results in considerable sidebands in the two-photon invariant mass distribution, at the level of approximately 10% at particle-level and 30% for reconstructed $t\bar{t}H$ events with $H \rightarrow \gamma\gamma$. Investigation has shown that this faulty combinatorial choice concerns overwhelmingly the photon with the second-highest p_T , since the photon with the highest p_T is not a Higgs photon only at the level of approximately 1/mil. The origin of these ‘fake’ second Higgs photons is approximately 80% from π^0 s, 10% from η s, a few percent from ω s, and the remainder from other particles. Fully 80% of these fake Higgs photon ‘mother’ particles appear to come from parton showers whose origin is one of the two final-state top quarks, and as such are peculiar to the $t\bar{t}H$ process. The other 20% come from showering from the initial-state protons and hence are common to all the associated production channels. In this way we can estimate that the level of the effect on the WH and ZH processes would be approximately 2-3%.

We have investigated possible methods of improving the Higgs photon selection so as to improve this situation. In particular, we have evaluated the performance on our signal samples of the photon isolation variables investigated and used by the $H \rightarrow \gamma\gamma$ inclusive analysis [?]. For photons in the ECAL barrel, we obtain the best performance from the sum of E_T of ECAL basic clusters within a cone of $\Delta R < 0.35$, after removing the E_T of those basic clusters constructed with the Island algorithm included in the supercluster matched ($\Delta R < 0.2$) with the offline photon itself (‘ECALiso’). For photons in the ECAL endcap, we obtain the best performance with the sum of E_T of HCAL calorimeter towers within $\Delta R < 0.4$ of the photon candidate (‘HCALiso’). We then studied the effect of varying the isolation cone radii for the two variables (see for example Figure 10.32 (left)) and found that better performance is possible against combinatorial photons with smaller cone sizes than those used in the $H \rightarrow \gamma\gamma$ inclusive analysis. Finally, performance can be further improved by considering linear combinations of the isolation variables (see for example Figure 10.32 (right)).

[§]The threshold for the single photon HLT trigger is $E_T > 100$ GeV, and those of the double photon trigger $E_{T1} > 35$ GeV and $E_{T2} > 20$ GeV, at an instantaneous luminosity of $L=10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ [?]




Figure 10.32: Left: Efficiency for true Higgs photons vs. that for combinatorial photons in the ECAL barrel for the ECAL isolation variable, for different isolation cone sizes. Right: Efficiency for true Higgs photons vs. that for combinatorial photons in the Ecal endcap, for several linear combinations of isolation variables.

For this study, the two highest p_T Offline Photons satisfying the following requirement on the isolation energy $\text{Iso} = \text{HcalIso} + (2 \cdot \text{EcalIso})$ were retained as Higgs photon candidates:

- For photons in the barrel: $\text{Iso} < 25 \text{ GeV}$
- For photons in the endcap: $\text{Iso} < 22 \text{ GeV}$

with $\Delta R < 0.25$ instead of 0.35 for EcalIso and $\Delta R < 0.3$ instead of 0.4 for HcalIso . These values yield approximately 95% efficiency for true Higgs photons and less than 40% for combinatorial photons. This strategy successfully restores approximately one-half of the true Higgs photon pairs previously lost to misidentification when selection based on only photon p_T is used, as is demonstrated by Figure 10.33.




Figure 10.33: Left: Invariant mass of the two Higgs photon candidates selected according to p_T alone (dark grey curve), p_T and isolation (light grey curve), and where both candidates are geometrically matched to particle-level Higgs photons (medium grey curve). Right: Distribution of the p_T of jets from pile-up events. Jets not matched to generated particle jets from the signal are considered to be pile-up jets.

A similar technique can be employed for the selection of candidate leptons from top quark decays (via a W boson). For OfflineElectrons, once again the best performance against combinatorial non-W-from-top electrons is obtained with the previously-defined ECALiso vari-

able. For GlobalMuons, the transverse momentum of tracks in a cone of radius $\Delta R < 0.25$ ('IsoByTkPt025') performs significantly better than the transverse calorimetric energy in a cone of radius $\Delta R < 0.3$ ('IsoByCaloEt03') for 'true' W leptons[¶] in our signal samples. For this study, we therefore retained as the W-decay (top) lepton candidate the highest E_T OfflineElectron or highest p_T GlobalMuon satisfying the following requirement:

- For electrons, IsoEcal < 5 GeV,
- For muons, IsoByTkPt02 < 9 GeV

These values yield $\sim 92\%$ efficiency for 'true' W leptons and approximately 35% for combinatorial leptons. In the selection criteria involving photons described below, as well as those involving leptons described thereafter, the pertinent distributions are constructed using the Higgs photon and W lepton candidates selected according to the procedure combining p_T and isolation described above.

Five variables involving the Higgs photon candidates have demonstrated effective performance against the irreducible background events: The p_T of each of the two OfflinePhoton candidates ($'p_{T\gamma_1}', 'p_{T\gamma_2}'$), the sum of their p_T ($'p_{T\gamma_1} + p_{T\gamma_2}'$), the angular distance between them ($'\Delta R_{\gamma_1, \gamma_2}'$), and $\cos \theta^*$, where $\tan \theta^* = \frac{|\vec{p}_i| \sin \theta_i}{\gamma(|\vec{p}_i| \cos \theta_i - \beta E_i)}$, and E_i and θ_i refer respectively to the energy of and the 3-space angle between either of the two Higgs photon candidate directions and the direction of their joint 4-vector, in the laboratory frame (the scalar nature of the Higgs boson should assure a flat distribution of this variable for signal events, and one peaked in the forward and backward directions for background events). With the goal of conserving a signal efficiency of between 85 and 95% per selection criterion, we have established the following eventwise selection involving the two Higgs photon candidates:

- $p_{T\gamma_1, \gamma_2} \geq 50, 18$ GeV
- $p_{T\gamma_1} + p_{T\gamma_2} \geq 85$ GeV
- $\Delta R_{\gamma_1, \gamma_2} \leq 3.2$
- $\cos \theta^* \leq 0.85$

Three variables involving the W lepton candidates have demonstrated effective performance against the irreducible background events: The E_T (OfflineElectron) or p_T (OfflineMuon) of the candidate, and the angular distances between the candidate and each of the two Higgs photon candidates ($'\Delta R'_{\gamma_1, lepton}, '\Delta R'_{\gamma_2, lepton}$). With the goal of conserving a signal efficiency of between 85 and 95% per selection criterion, we have established the following eventwise selection involving the W lepton candidate:

- $p_{Tlepton} \geq 15$ GeV
- $\Delta R_{\gamma_1, lepton}, \Delta R_{\gamma_2, lepton} \geq 0.3, 1.0$

In order to remove part of the backgrounds studied here and also other backgrounds, in particular from QCD processes, we take advantage of the high jet multiplicity of our signal events as well as the presence of two real top quarks yielding b-quark jets as decay products. Jets including those possibly corresponding to b-quarks are constructed with the Iterative Cone [?] algorithm with a cone radius of $\Delta R = 0.5$ and then calibrated with $\gamma + \text{jet}$ Monte Carlo events. A discriminant ('BtagDisc') is then calculated for each candidate b-quark jet

[¶]as for Higgs photons, considered to be those OfflineElectrons or GlobalMuons lying within a cone of radius $\Delta R < 0.1$ of a particle-level electron or muon from a W boson which itself is a decay product of one of the final-state top quarks

with the Combined BTag[?] algorithm. We require the presence of a minimum number of jets having a value of p_T greater than 60 GeV, which permits the removal of jets from pile-up from consideration (we consider a reconstructed jet to be from pile-up if it is not geometrically matched with a particle-level jet, which has been constructed using the same algorithm and parameters as the reconstruction-level jets). Figure 10.33(right) shows the p_T distribution of the pile-up jets thus identified in a signal sample with $m_H = 115$ GeV. We require more than four jets with $p_T > 60$ GeV.

To specifically target the $W+2\gamma+$ jets background (and eventually other non-b-quark reducible backgrounds), we make limited use of tagging of b-quark jets. We require that at least one candidate jet having $p_T > 60$ GeV have $B_{tagDisc} > 0.8$.

10.2.7.8 Performance of Off-line Selection

Tables 10.28 and 10.29 show the progression of the signal ($m_H = 115$ GeV) and background samples through the selection. Prior to checking for the L1 trigger decision, we apply the preselection at particle-level described in Section

10.2.7.9 Description of Kine preselections

to all signal and background samples. The number of surviving events is expressed as an effective cross-section in fb. The final results are also expressed as numbers of surviving signal and total background events with statistical errors, for both 30 and 100 fb^{-1} .

Table 10.28: Progression of the signal ($m_H = 115$ GeV) and background samples through the trigger portion of the selection, expressed as an effective cross-section in fb. Efficiencies with respect to the previous sequential requirement are expressed as percentages.

Requirement	MH=115	ttgg1	ttgg2	ttgg3	W2 γ 4j
Before selection	0.693 (100.0)	1.59 (100.0)	6.12 (100.0)	4.95 (100.0)	11.4 (100.0)
Preselection	0.533 (76.8)	1.4 (87.9)	5.05 (82.5)	3.94 (79.6)	11.3 (98.9)
L1 + HLT Accept	0.517 (97.0)	1.34 (95.4)	4.71 (93.4)	3.36 (85.7)	10.5 (93.0)
HLT $\gamma\gamma, \gamma$ accept	0.508 (98.3)	1.30 (96.9)	4.57 (96.9)	3.25 (96.6)	10.0 (96.0)

10.2.7.10 Uncertainties, Systematic Errors, and Strategy for Background Measurement from Data

To estimate the systematic error on the surviving signal cross-section, we have considered the following global source of error which is applied directly to the estimated number of signal events:

- Luminosity $< 3\%$

The error due to the inclusion/non-inclusion of initial and final-state photon radiation at the fragmentation/hadronisation level as well as that due to the matrix element generator used (ALPGEN or MADGRAPH) were found to be insignificant.

We have also considered the following sources of error relevant to the detector:

- Electron/Photon/Muon identification: 1% per identified object
- Efficiency to tag jets containing b quarks: 5% per identified b-quark jet

Table 10.29: Progression of the signal ($m_H = 115$ GeV) and background samples through the offline portion of the selection, expressed as an effective cross-section in fb. Efficiencies with respect to the previous sequential requirement are expressed as percentages.

Requirement	Criterion	MH=115	ttgg1	ttgg2	ttgg3	W2 γ 4j
2 γ (w/o pix. seed)	-	0.506 (100.0)	1.29 (100.0)	4.56 (100.0)	3.24 (100.0)	10.0 (100.0)
2 is. γ (w/o pix. seed)	-	0.482 (95.2)	1.22 (94.0)	3.96 (86.8)	2.53 (78.2)	9.58 (95.7)
pT γ_1 (GeV)	> 50	0.432 (90.0)	1.04 (85.3)	3.14 (79.4)	1.48 (58.5)	7.90 (82.5)
pT γ_2 (GeV)	> 18	0.386 (89.2)	0.88 (84.7)	2.25 (71.6)	1.03 (69.7)	6.72 (85.0)
pT γ_1 +pT γ_2 (GeV)	> 85	0.379 (98.2)	0.847 (96.3)	2.17 (96.5)	0.926 (89.8)	6.40 (95.3)
$\Delta R(\gamma_1 \gamma_2)$ (GeV)	< 3.2	0.364 (96.4)	0.738 (87.2)	1.86 (85.9)	0.719 (77.7)	5.30 (82.8)
$\cos \theta^*$	< 0.85	0.332 (91.4)	0.589 (79.8)	1.48 (79.5)	0.583 (81.0)	4.36 (82.3)
pT $_{lep}$ isolated (GeV)	>15	0.238 (72.2)	0.443 (75.2)	0.984 (66.4)	0.387(66.4)	3.15 (72.3)
$\Delta R(\gamma_1 l)$	>0.3	0.236 (99.0)	0.441 (99.5)	0.925 (94.0)	0.321 (83.0)	3.14 (99.6)
$\Delta R(\gamma_2 l)$	>1.0	0.208 (87.4)	0.389 (88.2)	0.607 (65.7)	0.163 (50.7)	2.34 (74.6)
N jets w/ $p_T > 60$ GeV	≥ 4	0.179 (86.2)	0.338 (87.0)	0.455 (74.9)	0.110 (67.6)	1.79 (76.6)
Btag Disc of 1 jet >0.8	>0.8	0.110 (61.6)	0.217 (64.0)	0.276 (60.7)	0.051 (46.0)	0.294 (16.4)
$M_H \pm 1.5$ GeV		0.074 (67.1)	0.005 (2.51)	0.011 (3.86)	<0.002 (3.92)	<0.003 (1.02)
N_{evts} 30 fb $^{-1}$		2.22+/-0.10	0.483 +/-0.158			
N_{evts} 100 fb $^{-1}$		7.42+/-0.334	1.61+/-0.53			

- Uncertainty on jet energy: 3%

Only the effect of the uncertainty on the jet energy scale is evaluated by propagation through the selection, and yields a net uncertainty of +1.6/-3.9% for a Higgs boson mass of 115 GeV. All the above contributions are summed in quadrature and a systematic error of +6.3/-7.2% is obtained for the number of signal events for a Higgs boson mass of 115 GeV. The uncertainty on the number of surviving background events, calculated below and amounting to an average of $\pm 15\%$, is finally added to the above quadratic sum to represent the observation of a signal peak atop background, finally yielding an error of +16.3/-16.6% on the number of events in a peak containing both signal and background events, corresponding to the case of a signal cross-section measurement.

The background spectrum can be fit from the sidebands surrounding the positions of the putative Higgs boson masses. The fit to a decreasing exponential function is shown by the grey curve in Figure 10.34 (left). For this purpose, the bin width has been chosen to be large enough (20 GeV) to have a sufficient number of events for a luminosity of 100 fb $^{-1}$. The bin centered around the generated Higgs boson mass is not used for the fitting procedure. To evaluate the error on the obtained number of background events, we fit test distributions obtained with the average of the bin contents according to a Poisson distribution. These gedanken experiments correspond to possible real data sets in the future. One such fit is shown in the black curve in figure 10.34 (left). The distribution of the number of background events thus obtained is shown in figure 10.34(right). The width of the gaussian fit of the distribution yields an estimate of the systematic error of the estimation of the number of background events. The mean and the error thus obtained are used in Table 10.31 to compute the signal significance.

10.2.7.11 Results for the Standard Model Higgs Boson

Tables 10.30 and 10.31 show, for each of the four Standard Model Higgs boson mass values considered, the signal selection efficiency, the total number of surviving signal and background events with statistical errors (from the number of generated events), and the signal significance calculated using the $S_{c_{12}}$ (for a number of background events inferior to 1) and




Figure 10.34: Left: Background estimation from the fit of the sidebands: Example of a gedanken experiment giving a possible set of real data points consistent with the Poisson distribution of the simulated number of surviving background events. The fit through these points is superimposed on the original fit. Right: Distribution of the estimated number of background events from the fit of the gedanken experiments.

ScP estimators[486], after the imposition of our selection, after 30 and 100 fb⁻¹, respectively. Also reported is the signal significance calculated taking into account the systematic errors discussed in the above section, for 100 fb⁻¹ of integrated luminosity.

Table 10.30: Signal selection efficiency, total number of surviving signal and background events with statistical errors (from the number of generated events), and the signal significance calculated using the Sc₁₂ estimator, without systematic errors taken into account, for each of the four Standard Model Higgs boson mass values considered after 30 fb⁻¹ of integrated luminosity.

Higgs Boson Mass (GeV)	115	120	130	140
Sig. Selection Eff. (%)	10.7	11.2	11.3	11.3
Number Signal	2.22 +/- 0.10	2.20 +/- 0.10	1.79 +/- 0.08	1.26 +/- 0.06
Total Number Bckgd	0.483 +/- 0.158	0.838 +/- 0.185	0.595 +/- 0.199	0.330 +/- 0.154
Signal Significance (Sc ₁₂)	1.898	1.655	1.546	1.373

Table 10.31: Signal selection efficiency, total number of surviving signal and background events, and the signal significance calculated using the ScP estimator, with and without systematic errors taken into account, for each of the four Standard Model Higgs boson mass values considered after 100 fb⁻¹ of integrated luminosity. In the case without systematic errors, the errors shown are statistical and come strictly from the number of generated events. In the case with systematic errors, the errors correspond to the fitted sigma of the distribution in Figure 10.34(right).

Higgs Boson Mass (GeV)	115	120	130	140
Sig. Selection Eff. (%)	10.7	11.2	11.3	11.3
Number Signal NS	7.42 +/- 0.33	7.33 +/- 0.33	5.96 +/- 0.27	4.21 +/- 0.19
Total Number Bcgkd	1.61 +/- 0.53	2.79 +/- 0.62	1.98 +/- 0.66	1.10 +/- 0.51
Total Number Bcgkd from fit w. syst.	2.23 +/- 0.34	1.94 +/- 0.32	1.60 +/- 0.22	1.39 +/- 0.22
Signal Significance (ScP)	3.541	3.662	3.257	2.510
Signal Significance (ScP) w. syst.	3.414	3.523	3.184	2.453

Figure 10.35 shows, for Higgs boson mass values of 115 and 140 GeV, the invariant mass spectrum of the two Higgs photon candidates after all selection criteria have been applied.




Figure 10.35: Invariant mass of the two Higgs photon candidates added to that of the sum of the surviving backgrounds after imposition of all selection requirements, for $m_H = 115$ GeV (left) and 140 GeV (right).

10.2.7.12 Conclusion

A full-reconstruction-level sequential study has been performed for the channel $t\bar{t}H(h^0)$, with $H(h^0) \rightarrow \gamma\gamma$, taking into account irreducible backgrounds not previously studied. The ratio of signal to background events is approximately 4:1 representing a factor of 2 improvement over prior CMS and ATLAS studies. Signal observability in excess of 3σ is indicated for masses up to 130 GeV with full simulation and reconstruction and with estimated systematic errors taken into account for 100 fb^{-1} of integrated luminosity.

10.2.8 Associated WH,ZH production with $H \rightarrow \gamma\gamma$

Compared to the gluon-gluon fusion channel $gg \rightarrow H \rightarrow \gamma\gamma$, the associated production channels WH/ZH [487, 488] suffer from a much lower production cross section. Several advantages, however, make these channels attractive when the decay of the gauge boson results in a charged lepton: requiring an additional relatively high transverse-momentum lepton greatly reduces the significant QCD background in the $\gamma\gamma$ topology and improves the primary vertex reconstruction [489]. In the context of supersymmetric models, maximal mixing in the stop sector could result in a strong suppression of the $gg \rightarrow h$ signal, which the associated production channels would not be subject to [490]. The searched-for final state is therefore comprised of 2 isolated photons and at least one isolated electron or muon. Prior generator-level or fast simulation studies [478, 479, 491, 492] conclude to the possibility of a discovery at the LHC in this channel.

10.2.8.1 Event generation and reconstruction

All the process considered in this study have been simulated at the leading order. Signal events were generated by the matrix element generator COMPHEP [34] for Higgs boson masses ranging from 90 to 150 GeV/c^2 , in steps of 5 GeV/c^2 . Total cross-sections have been rescaled accordingly to the NLO calculation [8]. K-factors from 1.15 to 1.16 are obtained on the whole mass range. Branching ratios for $H \rightarrow \gamma\gamma$ were taken from HDECAY program [9]. The irreducible backgrounds from the processes $W\gamma\gamma$ et $Z\gamma\gamma$ were also generated with COMPHEP, with the same K-factors applied as those pertinent to the signal. Fragmentation and hadronization was performed by PYTHIA [238].

The reducible background processes $\gamma\gamma$, γ -jet, $W\gamma$, $b\bar{b}$, and $t\bar{t}$, retained due to their capacity to mimic the $l\gamma\gamma$ signal, have been generated with PYTHIA and leading order cross-sections were considered, except for the $t\bar{t}$ production where a NLO cross section of 840 pb is used [266].

To ensure an efficient generation and preserve sufficient statistics of the most signal like events, a preselection is applied at generator level. Three electromagnetic candidates or two electromagnetic candidates and one muon candidate with $E_T > 20$ GeV and $|\eta| < 2.7$ are required. An electromagnetic candidate is obtained by clustering electrons and photons in a $\Delta\eta = 0.09$, $\Delta\phi = 0.09$ window. Muon candidates are either μ , τ , π , or K particles.

The generated events were passed through the Geant-3 simulation of CMS [13]. The events were then digitized and reconstructed with the standard CMS software [15] with the addition of pile-up event corresponding to the high luminosity phase ($\mathcal{L} = 10^{34} \text{ cm}^{-2}\text{s}^{-1}$).

10.2.8.2 Trigger selection

Events are required to pass the global Level 1 trigger [493] and the double photon High Level Trigger (HLT) [?] configured for the high luminosity phase. The trigger efficiencies for the preselected signal events are higher than 95% on the whole Higgs boson mass range (90-150 GeV/ c^2) as shown in Table 10.32 and 10.33.

10.2.8.3 Offline event selection

To suppress the reducible backgrounds, four discriminant combined variables are first constructed using a likelihood ratio method to estimate the isolation of the photons, the quality of the lepton reconstruction, the isolation of the lepton and the QCD / multi-jets nature of the event. The reference histograms for the four likelihoods are all produced on independent simulated event samples after a very loose preselection requiring two offline photons and one electron or muon reconstructed by the standard algorithms. Photon candidates with a matching seed in the pixel detector are rejected. The two photons with the highest transverse energy are assigned to the Higgs boson decay. Several isolation variables [494] were tested in the likelihood and the best performance is obtained with the sum of the transverse energy of the basic clusters within a cone $\Delta R < 0.3$ around the photon, excluding the basic clusters belonging to the photon supercluster and the sum of the transverse energy of the HCAL towers within a cone $\Delta R < 0.3$ around the photon.

Then the offline lepton with the highest E_T is selected. The reconstruction quality of the electron is carefully checked. The four variables yielding the most significant discriminating power are the ratio E/p between the electron energy as measured in the calorimeter and its momentum measured by the tracker, the hadronic energy fraction E_{had}/E , the distance $\Delta\eta$ between the track and the associated supercluster and the ratio R_9 between the sum of the energies of 9 crystals (3x3 matrix centred on the maximum-energy crystal) and the energy of the corresponding supercluster. In the case of muons the purity obtained by the standard CMS reconstruction algorithms has already proven sufficient; therefore, no additional criteria are applied.

For the lepton isolation, similar calorimeter variables as for photons are used. In addition, the number of pixel lines within a cone $\Delta R < 0.3$ around the lepton improves the discriminative power of the likelihood.

Finally a global discriminant variable against multijet background is constructed. The rejection of π^0 faking signal photons, effective against QCD backgrounds, has been accomplished by a neural network procedure exploiting the information on the lateral profile of the electromagnetic shower. Variables involving the multiplicity of reconstructed objects in the electromagnetic calorimeter improve the discriminating power.

Table 10.32: Expected rates (in fb) after each stage of the event selection for signals ($m_H = 120 \text{ GeV}/c^2$) and irreducible backgrounds. Errors are statistical only.

	WH	ZH	$W\gamma\gamma$	$Z\gamma\gamma$
$\sigma \cdot \text{BR}$	0.810	0.137	-	-
Preselection: $\sigma \cdot \text{BR} \cdot \epsilon$	0.460	0.0440	13.58	18.92
Double photons HLT	0.439 ± 0.005	0.0423 ± 0.0004	8.80 ± 0.04	12.13 ± 0.07
2 isolated photons	0.387 ± 0.005	0.0370 ± 0.0004	7.14 ± 0.04	6.51 ± 0.04
1 good quality lepton	0.331 ± 0.004	0.0350 ± 0.0003	5.56 ± 0.04	4.58 ± 0.03
Lepton isolation	0.299 ± 0.004	0.0318 ± 0.0003	4.83 ± 0.04	4.11 ± 0.03
QCD rejection	0.281 ± 0.004	0.0273 ± 0.0003	4.50 ± 0.04	3.53 ± 0.03
$80 < m_{\gamma\gamma} < 160$	0.271 ± 0.004	0.0259 ± 0.0003	2.04 ± 0.02	1.42 ± 0.02

Table 10.33: Expected rates (in fb) after each stage of the event selection for reducible backgrounds. Contributions of the different \hat{p}_T bins are summed. Errors are statistical only.

	$\gamma\gamma$	$W\gamma$	$b\bar{b}$	$t\bar{t}$	γ -jet (jet)
$\sigma \cdot \text{BR}$	1.1×10^5	5.79×10^3	1.78×10^9	86.2×10^3	1.21×10^8
Preselection: $\sigma \cdot \text{BR} \cdot \epsilon$	270.1	26.5	2.96×10^5	6.00×10^3	7.16×10^4
Double photons HLT	197.7 ± 1.0	16.8 ± 0.1	77120 ± 764	1948 ± 17	35045 ± 256
2 isolated photons	161.6 ± 0.8	9.97 ± 0.07	682 ± 72	31.2 ± 2.2	7235 ± 115
1 good quality lepton	27.3 ± 0.3	7.98 ± 0.07	311 ± 49	23.5 ± 1.9	2552 ± 68
Lepton isolation	9.8 ± 0.2	6.59 ± 0.06	(0.87)	14.2 ± 1.5	209 ± 20
QCD rejection	7.6 ± 0.2	5.74 ± 0.06	(0.003)	(0.35)	(6.6)
$80 < m_{\gamma\gamma} < 160$	3.2 ± 0.1	2.40 ± 0.04	(0.001)	(0.26)	(3.7)

The results of the selection applied on the four combined variables are presented in Tables 10.32 and 10.33. The multijet backgrounds are entirely suppressed. To obtain a more precise estimation of these backgrounds, the cut factorization method has been applied and the result is given between parantheses in Table 10.33. After rejecting events outside the 80 - $160 \text{ GeV}/c^2$ di-photon mass window, the expected rate of events is $0.297 \pm 0.003 \text{ fb}$ for signal and $13.1 \pm 2.6 \text{ fb}$ for background. Some simple kinematical variables are then used to form a final likelihood. The best discrimination was obtained with the transverse energy of the photons and of the lepton, the ΔR distances between lepton and each photon, the missing transverse energy, and the $\Delta\Phi$ angle between the directions of the missing transverse energy and of the highest E_T photon. The distribution of the resulting combined variable y is shown in Figure 10.36 for a Higgs boson mass of $120 \text{ GeV}/c^2$.

10.2.8.4 Statistical method and optimization

The statistical methods developed by the LEP Higgs working group [495, 496] are used in this analysis to optimize the selection criteria and determine the statistical significance of the final results. To form the test-statistic, the three obvious variables to be used are the counting rates, the $\gamma\gamma$ invariant mass and the kinematic likelihood variable y . The limited statistics of the MC events prohibit however the use of a two-dimensional method for the determination of the Higgs boson discovery potential. So, only the counting rates and shape of the reconstructed $\gamma\gamma$ mass distribution will be used along with a cut on the combined




Figure 10.36: Distribution of the final combined variable for the signal ($m_H = 120 \text{ GeV}/c^2$) and for the background. The optimal working point is obtained with a cut $\log(y) > 0.35$.

likelihood variable y . The optimal working point is the y cut value which maximizes the discovery potential as shown in Figure 10.37.




Figure 10.37: Left: Reconstructed $\gamma\gamma$ mass for different selection values on the final combined variable y for an integrated luminosity of 100 fb^{-1} . Right: Statistical significance as a function of the cut on the combined variable $\log(y)$, for $m_H = 120 \text{ GeV}/c^2$ and an integrated luminosity of 100 fb^{-1} . The highest significance is obtained with a cut $\log(y) > 0.35$.

The list of the optimal working points obtained for the different Higgs boson mass hypotheses is given in Table 10.34. The significance and the expected number of signal and background events are given for a luminosity of 100 fb^{-1} . For the γ -jet, $t\bar{t}$ and $b\bar{b}$ backgrounds, the rates are estimated by the method of cut factorization.

10.2.8.5 Use of real data in sidebands - Systematic uncertainties

The signal is characterized by a strongly peaked di-photon mass and the $m_{\gamma\gamma}$ distribution of the background is quite smooth at the considered working points. Therefore, when real data will be available, the data taken in $m_{\gamma\gamma}$ sidebands will be used to optimize the likelihood

Table 10.34: Optimal working points for different Higgs boson mass hypotheses. The significance and the expected number of signal and background events are given for an integrated luminosity of 100 fb^{-1} .

m_H (GeV/ c^2)	working point $\log(y) >$	significance	WH	ZH	$W\gamma\gamma$	$Z\gamma\gamma$	$W\gamma$	$\gamma\gamma$	γ -jet	$t\bar{t}$	$b\bar{b}$
115	0.41	4.30σ	22.1	1.8	49.3	30.9	33.0	10.2	1.7	0.16	10×10^{-5}
120	0.35	4.09σ	20.7	1.6	51.2	36.2	34.5	12.4	1.9	0.15	10×10^{-5}
130	0.68	3.64σ	14.6	1.3	30.7	16.9	18.7	6.0	1.4	0.10	4×10^{-5}
140	0.99	3.35σ	11.4	1.0	18.9	10.3	10.6	3.7	1.0	0.04	1×10^{-5}
150	0.83	2.87σ	10.4	0.9	20.2	11.7	12.3	5.4	1.1	0.03	3×10^{-5}

analysis and to estimate the background.

Likelihood optimization with sideband events - No kinematic observables were used to construct the four primary likelihoods aimed at rejecting multijet events to avoid correlations with the di-photon mass. If the shapes of the distributions of the variables used in the likelihoods are sufficiently similar for different di-photon mass regions, then data taken outside the signal region can be used to optimize the likelihoods. To test the feasibility of the method, a sample of “fake real data” (the number of MC events for each background is equal to the expected number of events for a given luminosity) taken in the $20 < m_{\gamma\gamma} < 80 \text{ GeV}/c^2$ sideband is used to produce the reference S/B histograms of the likelihoods. The equivalent luminosity of the sample is limited to 132 pb^{-1} by the available statistics and is composed of 4682 $b\bar{b}$, 465 γ -jet, 222 $t\bar{t}$, 2 $\gamma\gamma$, 1 $W\gamma$ and 1 $Z\gamma\gamma$ events. The performance obtained with the likelihood on the events in the $80 - 160 \text{ GeV}/c^2$ band is compared to the results obtained by the standard analysis optimized with the full MC statistics available. For the four global discriminant variables, up to 20% loss of efficiency is observed for the same rejection power. The degradation of the performance is mainly due to the insufficient statistics of γ -jet and $t\bar{t}$ events in 132 pb^{-1} of data. To increase the size of the “fake data sample”, gedanken experiments were generated using the fitted shapes of the variables used in the likelihoods (correlations between the variables are neglected). The results are presented in Figure 10.38 for the photon isolation likelihood. An integrated luminosity of 5 fb^{-1} will be sufficient to optimize the four primary likelihoods with the real data taken in the $m_{\gamma\gamma}$ sideband and to reproduce the results obtained when using the full MC statistics.

Background measurement from Data - The $m_{\gamma\gamma}$ distribution of the background is smooth enough to be easily fit in the sideband to estimate the background in the signal region. To optimize the method (size and position of the window, bin width, choice of the fit function) and to estimate the uncertainty on background, 10000 signal+background pseudo-experiments were generated for each Higgs boson mass and luminosity hypothesis, as illustrated in Figure 10.39. For a luminosity of 100 fb^{-1} and a Higgs boson mass of $120 \text{ GeV}/c^2$, the background is measured with a precision of 11%, and with a precision of 6.6% for 300 fb^{-1} .

Systematic uncertainties for signal and cross-section measurement - The theoretical cross-section error due to the scale variation are estimated to $\pm 3\%$ for WH and ZH production for all considered Higgs boson masses [8]. The uncertainty on the parton density function of the CTEQ collaboration [29] is almost constant for the associated production $q\bar{q} \rightarrow VH$ at the LHC and of the order of 4% over a Higgs boson mass range between 100 and $200 \text{ GeV}/c^2$ [497]. The error on the measured luminosity is expected to be 3% for luminosity above 30 fb^{-1} . The error on the lepton or photon reconstruction and identification has been estimated to 1%




Figure 10.38: Comparison of the performance obtained when optimizing the photon isolation likelihood with a sample of 132 fb^{-1} of “fake real data” taken in the $20 < m_{\gamma\gamma} < 80 \text{ GeV}/c^2$ sideband (dash-dotted line) with the performance obtained by the standard analysis using the full MC statistics (solid line). To increase the available statistics in the sideband, gedanken experiments were generated for an equivalent luminosity of 5 fb^{-1} . The results of the optimization on these sideband events is represented by the dotted line.




Figure 10.39: Left: Background measurement in the signal region with a fit on the $m_{\gamma\gamma}$ sideband. The fit of the full MC statistics is represented by the solid light gray line. The fit of the fake data (dark grey) is performed on the sidebands (*i.e.* after the exclusion of the signal window represented by the dotted line). Two gedanken experiments are represented for an integrated luminosity of 300 fb^{-1} and a $120 \text{ GeV}/c^2$ Higgs boson mass hypothesis. Right: Relative uncertainty on the background estimation by the sideband fit method as a function of the integrated luminosity with LHC running at high luminosity for a Higgs boson mass of $120 \text{ GeV}/c^2$.

for each photon and lepton. An error of 5% on the missing transverse energy [498] propagated in the final likelihood gives a $-1.08\% +0.49\%$ variation of the final signal rate for $m_H = 120 \text{ GeV}/c^2$. The quadratic sum of all these errors gives a 6% total error on the expected signal rate.

In the case of a Higgs boson discovery, this channel will be used to measure the cross-section

times the branching ratio:

$$\sigma_s \times BR = \frac{N_s}{\epsilon_{sel} \mathcal{L}} = \frac{N - N_b^{fit}}{\epsilon_{sel} \mathcal{L}}$$

where N_s is the number of signal events given by the difference between the total number N of observed events and the number N_b^{fit} of background events measured by the sideband fit. The total uncertainty on the measure is given by:

$$\left(\frac{\Delta(\sigma_s \times BR)}{\sigma_s \times BR} \right)^2 = \left(\frac{\Delta N}{N - N_b^{fit}} \right)^2 + \left(\frac{\Delta N_b^{fit}}{N - N_b^{fit}} \right)^2 + \left(\frac{\Delta \mathcal{L}}{\mathcal{L}} \right)^2 + \left(\frac{\Delta \epsilon_{sel}}{\epsilon_{sel}} \right)^2$$

The expected precision on the $\sigma \times BR$ measurement is represented as a function of the integrated luminosity in Figure 10.40. For a 120 GeV/c² Higgs boson, the product of the cross-section and branching ratio will be measured with a precision of 35% after one year of LHC running at high luminosity, and with a precision of 19% after three years of high luminosity running.




Figure 10.40: Left: Precision on the measurement of the product of cross-section and branching ratio as a function of the integrated luminosity with LHC running at high luminosity for a 120 GeV/c² Higgs boson. Right: Statistical significance for different Higgs boson mass hypotheses as a function of the integrated luminosity with LHC running at high luminosity. The 1σ systematic uncertainty is represented by the grey band.




Figure 10.41: Statistical significance (left) and luminosity needed for a 5 σ or 3 σ observation (right) as a function of m_H . The 1σ systematic uncertainty is represented by the grey band.

Table 10.35: Background processes considered into the present analysis. The cross-section includes the decay of W and Z bosons into leptons. The generator and the number of events processed are also shown together with the corresponding weight for a luminosity of $1fb^{-1}$. For comparison, the signal varies, with the same hypotheses, from 5 fb to 20 fb in the considered mass range.

Background	Cross-section	Generator	MC statistic	weight ($1fb^{-1}$)
WWW($3l^{\pm}$)	4.95 fb	CompHep	10000	5.19×10^{-4}
WZ($3l^{\pm}$)	1.71 pb	Pythia	50000	3.46×10^{-2}
ZZ($4l^{\pm}$)	0.17 pb	Pythia	50000	3.67×10^{-3}
$t\bar{t}(1^{+}1^{-}b\bar{b})$	90.9 pb	TopRex	100000	0.93
Wt($1^{+}1^{-}b$)	5.25 pb	TopRex	50000	0.11

10.2.8.6 Results for the Standard Model Higgs Boson

The statistical significance is represented as a function of the luminosity in Figure 10.40 for different m_H hypothesis. The statistical significance and the luminosity needed for a 5σ or 3σ observation are represented as a function of m_H in Figure 10.41. One year of high luminosity running allows the observation at 3σ of the SM Higgs boson up to $m_H = 150 \text{ GeV}/c^2$, and five years of running at high luminosity are required to reach a 5σ discovery.

10.2.9 Associated WH production with $H \rightarrow WW^{(*)} \rightarrow 2\ell 2\nu$

The cross-section for this process exhibits a maximum near the WW resonance, due to the combined behavior of the production cross-section and the Higgs boson branching ratio. We will therefore focus on the intermediate mass region between $120 \text{ GeV}/c^2$ and $190 \text{ GeV}/c^2$, where the cross-section exceeds 300 fb. We decided to investigate events containing three leptons in the final state, concentrating on electrons and muons (including leptonic tau decays).


10.2.9.1 Signal and background generation

Starting at $115 \text{ GeV}/c^2$, we considered Higgs boson with masses of 125, 130, 140, 150, 160, 170, 180 and $190 \text{ GeV}/c^2$. Events were generated with PYTHIA for each of the nine Higgs boson masses, without any kinematical cut. W bosons are forced to decay leptonically (e, μ, τ).


All Standard-Model processes likely to produce three leptons must be considered as background for this analysis. This includes events where three leptons are actually produced in the hard process but also events with a fake or missed lepton. One particular case is the production of leptons in the semi-leptonic decay of a B meson. In the present analysis, we considered the production of WWW, WZ, ZZ, $t\bar{t}$, and Wt. Most of the processes are simulated with PYTHIA, except for WWW, which is generated with CompHep, and Wt generated with TopRex. In all cases, PYTHIA is used for the hadronization step. Table 10.35 shows the cross-section, the generator used and the number of events produced.

10.2.9.2 Selection streams at Level-1 and HLT

The global (cumulative) trigger efficiency after L1 and HLT is found to reach 72% for a $140 \text{ GeV}/c^2$ Higgs boson using the full trigger table. Main contributions come from single



(a)



(b)

Figure 10.42: (a) Trigger efficiency by trigger pattern, for the signal. Efficiency is calculated as N_x/N_{HLT} , where x is one of the 2^4 exclusive trigger classes. “Others” stands for unconsidered trigger patterns; (b) Trigger efficiency for each class of Monte-Carlo events. Results are given after L1 and after HLT. Efficiency is computed as the ratio between the number of triggered events and the total number of generated events.

Table 10.36: Trigger efficiency for each source of background. Efficiency at HLT with the restricted trigger set used in the present analysis is also presented.

Background	L1 efficiency	HLT efficiency	$e, ee, \mu, \mu\mu$ HLT efficiency
WWW($3l^\pm$)	0.87	0.79	0.73
Wt($1^{+1-}b$)	0.88	0.78	0.67
WZ($3l^\pm$)	0.8	0.72	0.65
ZZ($4l^\pm$)	0.78	0.69	0.64
$t\bar{t}(1^{+1-}b\bar{b})$	0.91	0.79	0.65

and double leptonic (e and μ) triggers (65%). There is a small contribution from the missing transverse energy trigger (E_T^{miss}) and from combined ($e \wedge \tau$) and ($\mu \wedge \tau$) triggers, further reduced by the event selection, which favors multi-leptonic patterns. For this analysis, we restricted ourselves to triggers known to have the highest impact on the total efficiency: single- and double-electron and muon triggers. Figure 10.42(a) shows the efficiency for each (exclusive) trigger pattern, given the above choice of interesting bits.

Details about the efficiency for each type of event (defined from the number of muons, electrons and taus in the event) are given in Figure 10.42(b). Events containing one or more muons are more easily retained (efficiency reaches 85% for events with three muons) while tau events are only marginally selected (efficiency: 12%). Efficiency rises slightly with the Higgs boson mass, from 58% at $115 \text{ GeV}/c^2$ to 74% at $190 \text{ GeV}/c^2$.

Table 10.36 shows the trigger efficiency for each source of background. Efficiency, when restricting ourselves to single- and double-electron and muon triggers, varies from 64% to 73%, which is the same magnitude as the trigger efficiency for signal events. This restricted trigger is 15% (for $t\bar{t}$) to 5% (for ZZ) less efficient than the inclusive High-Level trigger.

10.2.9.3 Off-line selection

Electrons and muons are reconstructed using default offline reconstruction algorithms. For electrons, additional quality cuts are applied: the energy measured by ECAL and the momentum obtained by the tracker must agree within 50%, and the ratio of energy measured by HCAL and ECAL must be lower than 0.15. Only leptons with $P_t^l > 14 \text{ GeV}/c$ are retained. A first set of selection criteria is applied to select signal-like topologies, requiring three and only three leptons, for a total charge of either +1 or -1. A cut on the distance in the z direction between the points of closest approach of lepton tracks to the beam is applied to ensure that all of the three leptons are coming from the same interaction. The two closest (in the $\eta - \phi$ plane) opposite-sign leptons are then assigned to the Higgs boson decay. The angle between leptons attributed to the Higgs boson can be used to distinguish signal and background. The acollinearity between two leptons is defined as the angle between the two leptons, in the space, and their acoplanarity is defined as the same angle projected onto the transverse plane. Both the acollinearity ($\theta_{\text{aco}} < 1.75 \text{ rad}$) and the acoplanarity ($0.1 \text{ rad} < \phi_{\text{aco}} < 0.75 \text{ rad}$) between the leptons are used, as they provide complementary information. Leptons required to be isolated in the tracker, i.e. the angle between the lepton's track and the closest track above $3 \text{ GeV}/c$ must be more than 0.2. A jet veto is applied rejecting events with a jet, reconstructed with the iterative cone algorithm ($r=0.7$), above 25 GeV in the central region ($|\eta| < 2.1$). An additional B veto is applied, imposing that no single B-jet is reconstructed by the default *combined B-tag* algorithm. This removes low-energy bjets passing the standard jet veto. A cut on the invariant mass of any pair of leptons compatible with the Z hypothesis (via charge flavor and invariant mass constraints, $M_Z \notin [65, 115]$) is used to reject ZZ and WZ events. Finally, kinematical cuts are used: $E_T^{\text{miss}} > 50 \text{ GeV}/c$, $M_T(W_3) > 40 \text{ GeV}/c^2$ and $Pt(\sum \vec{P}_l) > 40 \text{ GeV}/c$, where $Pt(\sum \vec{P}_l)$ is the transverse momentum of the vector sum of momenta of all three leptons, and $M_T(W_3)$ is the reconstructed transverse mass of the associated W boson:

$$M_T(W_3) = \sqrt{2 * P_T^{l_3} E_T^{\text{miss}} (1 - \cos \Delta\phi_{l_3 p_T})}, \quad (10.9)$$

with $P_T^{l_3}$ being the transverse momentum of the lepton not associated to the Higgs boson, E_T^{miss} the missing transverse momentum, and $\Delta\phi_{l_3 p_T}$ the polar angle between the lepton and the missing transverse momentum. Optimized cuts are summarized in Table 10.37.

The Higgs boson transverse mass is computed from the two chosen leptons and from the missing transverse momentum:

$$M_T(H) = \sqrt{M_T^{ll^2} + 2E_T^{ll} p_T - 2P_T^{ll} p_T \cos \Delta\phi_{ll p_T}}, \quad (10.10)$$

Figure 10.43 shows the distribution of this quantity for the signal, on top of remaining background, after all cuts for a Higgs boson mass of $140 \text{ GeV}/c^2$ and an integrated luminosity of 100 fb^{-1} . The cumulated efficiency (including trigger and event selection) depends on the Higgs boson mass hypothesis. Starting at 0.5% for a mass hypothesis of $115 \text{ GeV}/c^2$, the efficiency rises to a maximum at the WW resonance (1.3%). Beyond the WW production threshold, efficiency drops since W bosons start to be boosted in the Higgs boson frame, which influences the angular distribution of leptons. Efficiency in that region could certainly be improved by optimizing the analysis for a Higgs boson mass of $190 \text{ GeV}/c^2$.

^{||}No energy correction is applied since we are mostly handling fake low-energy jets from pile-up and underlying events.

Table 10.37: Summary of the optimized selection cuts. The cross-section for the signal and backgrounds, for each step in the selection, is given in fb. An upper limit for the Wt and $t\bar{t}$ cross-sections is given when no simulated event remains.

Cut		Signal (fb) 140 GeV/c ²	Background (fb)				
Id.	Type		$t\bar{t}$	Wt	ZW	ZZ	WWW
0	L1 \wedge HLT	12.24	72067	4115.8	1238.4	118.438	3.91
1	$N_{lept} = 3, ch_{\Sigma} = 1$	3.81	16432.7	680.0	339.4	34.65	1.05
2	Lepton cuts	2.67	5629.1	245.3	245.9	23.53	0.70
3	Angular cuts	0.87	400.6	15.0	18.3	2.29	0.11
4	B veto	0.43	3.85	0.42	9.77	1.19	0.06
5	Jet veto	0.27	< 1.93	0.31	7.26	0.58	0.04
6	Z veto	0.21	< 1.93	0.21	0.40	0.08	0.03
7-9	Topological	0.13	< 1.93	< 0.11	0.06	0.01	0.02




Figure 10.43: Reconstructed transverse mass from Equation 10.10 for a 140 GeV/c² Higgs boson and an integrated lumisonity of 100 fb⁻¹.




Figure 10.44: Distribution of the acoplanarity for pseudo-experiments, fitted by a signal+background shape, as described in the text.

10.2.9.4 Systematic uncertainties

Systematic sources considered in this study are related to the normalization of backgrounds, to the reconstruction, the event selection, the luminosity and the structure functions of protons.

The uncertainty on the background normalization is usually estimated from the scale-dependence of the NLO cross-sections. This scale dependence is not known to us. This uncertainty can be large, since NLO effects are important. For events of interest for this study, the differences between leading-order and next-to-leading-order calculations are large: 45% for $t\bar{t}$, 40% for WZ and 25% for ZZ events. Before tuning them with real data, and thus especially at low integrated luminosity, it will be impossible to trust background estimates from Monte-Carlo programs. Hence, background will be normalized to signal-free regions of the phase-space.

By looking at the acoplanarity distribution when the angular cuts are not applied, data can be fitted to a sum of signal and background shapes. For that purpose, the signal is described by a sigmoid distribution, while the background remains constant. The Monte Carlo distribution for Signal and background are first fitted independently, and the shapes obtained that way are used to fit data from pseudo-experiments (Figure 10.44). The uncertainty on the background normalization is then related to the uncertainty on the background level in that fit. The uncertainty on the background level is found to be 15% for an integrated luminosity of 100 fb^{-1} , and rises up to 20% for 30 fb^{-1} . That value will be used in the following.

Reconstruction and selection uncertainties mainly arise from the jet veto, the b veto and lepton reconstruction. Experience from Tevatron tells us that a typical 2% uncertainty on lepton reconstruction efficiency has to be considered, while 5% uncertainty comes from lepton isolation [474] Since three leptons are present in our analysis, 12% uncertainty from lepton reconstruction and selection has been assumed. The additional uncertainties from the jet veto and the b veto will be assumed to be 5% each, which also seems reasonable in the light of Tevatron data.

To take into account other uncertainties related to the event selection, cuts are varied within the resolution of the associated quantity. The signal efficiency and background rejection




Figure 10.45: (a) Luminosity needed to obtain a 5σ significance using the likelihood-ratio method, with systematics only, Monte-Carlo statistical uncertainties only, or with both effects considered; (b) luminosity needed to exclude a Higgs boson at 95% C.L. if no excess is observed, using the same method.

are found to be stable with respect to such variations. A conservative value of 3% for the associated uncertainty is considered in the following.

The last considered uncertainty comes from the product of the luminosity and the proton structure functions, known as the parton luminosity. Considering these two quantities separately, 5% uncertainty on the luminosity is generally assumed, while the uncertainty from the proton parton distribution function (PDF) is taken to be 4% [475]. This latter uncertainty is reduced by the process considered, for which the mid- x region (where uncertainties are small) dominates. This gives an uncertainty of 6.4% on the parton luminosity.

The additional source of systematic uncertainties arising from the limited Monte Carlo statistics is also considered in the following result. With the Likelihood ratio method we are using, this is done bin per bin in the distributions of signal and background, so that a single value cannot be quoted. For the time being, this has a large impact on the results, but this effect will easily be reduced in the future, as more events become available.

10.2.9.5 Signal significance

In order to integrate the effect of systematic uncertainties and to exploit the discriminative power from the transverse mass distribution, the likelihood-ratio method (S_{CL}) is used. Figure 10.45(a) shows the luminosity needed to obtain a 5σ significance using this method, with systematics only, with Monte-Carlo statistical uncertainties, or with both effects considered. Figure 10.45(b) shows the luminosity needed to exclude a Higgs boson at 95% C.L. if no excess is observed, using the same method. Less than $50 fb^{-1}$ are required in most of the mass range, while only $20 fb^{-1}$ are needed at $170 GeV/c^2$.

One important motivation for studying this channel is also that it is one of the only allowed signatures for a fermiophobic Higgs boson model. If the Higgs boson does not couple to fermions, the usual gluon-fusion diagrams are indeed forbidden, as well as $b\bar{b}$ decays. A




Figure 10.46: Results obtained using the benchmark fermiophobic model; (a) Luminosity needed to obtain a 5σ significance using the likelihood-ratio method, with systematics only, Monte-Carlo statistical uncertainties only, or with both effects considered; (b) luminosity needed to exclude a Higgs boson at 95% C.L. if no excess is observed, using the same method.

fermiophobic Higgs boson will present a large cross-section at low mass, as the branching ratio does not drop down as in the Standard Model. Figure 10.46(a) shows the luminosity needed to obtain a 5σ significance for a fermiophobic Higgs boson. Compared to Figure 10.45, the needed luminosity is found to be similar in the most favorable mass region for the Standard Model (around $170 \text{ GeV}/c^2$) and above, but far better results are obtained in the low mass region. After 100 fb^{-1} , all masses between the LEP limit and $175 \text{ GeV}/c^2$ will be covered by this analysis alone. Figure 10.46(b) shows the luminosity needed to exclude a fermiophobic Higgs boson at 95% C.L. if no excess is observed. In the absence of signal, less than 30 fb^{-1} are required to reject any fermiophobic Higgs boson up to $175 \text{ GeV}/c^2$.

10.2.9.6 Measurement of Γ_W

In the present study, the partial decay width of the Higgs boson into W bosons, Γ_W , is determined with a good accuracy. This is possible using the narrow width approximation and benefiting from the fact that only the g_{HWW} coupling is involved in the Feynman diagram for the signal.

From the number of observed events, and assuming a good description of the background by the Monte-Carlo simulations, the product of the signal cross-section and the branching ratio of the Higgs boson into WW can be computed as:

$$\sigma \times BR = \frac{N_s}{\epsilon \mathcal{L}} = \frac{N - N_b}{\epsilon \mathcal{L}}, \quad (10.11)$$

with ϵ representing the combined trigger and selection efficiency. The uncertainty on this measurement is then found to be

$$\left(\frac{\Delta(\sigma \times BR)}{\sigma \times BR} \right)^2 = \frac{(\Delta N)^2}{N_s^2} + \frac{(\Delta N_b)^2}{N_s^2} + \frac{(\Delta \mathcal{L})^2}{\mathcal{L}^2} + \frac{(\Delta \epsilon)^2}{\epsilon^2}. \quad (10.12)$$




Figure 10.47: (a) Resolution on the product of cross-section and branching ratio into WW as a function of the Higgs boson mass, for an integrated luminosity of $100fb^{-1}$. (b) Resolution on the Γ_W partial width as a function of the Higgs boson mass, for the same integrated luminosity.

In this expression, ΔN corresponds to the statistical uncertainty on N , and will be assumed to be \sqrt{N} . The other three ΔN_b , $\Delta \mathcal{L}$ and $\Delta \epsilon$ quantities are dominated by systematic error sources described in Section 10.2.9.4.

In the narrow width approximation, the measured product of signal cross-section and branching ratio of the Higgs boson into WW can also be expressed from partial and total widths of the Higgs boson:

$$\sigma \times BR = \frac{\sigma^{SM} \Gamma_W}{\Gamma_W^{SM}} \times \frac{\Gamma_W}{\Gamma_{tot}} \equiv T \Gamma_W^2. \quad (10.13)$$

This allows us to write the uncertainty on the Higgs boson width into W bosons as

$$\left(\frac{\Delta \Gamma_W}{\Gamma_W} \right)^2 = \frac{1}{4} \left(\frac{(\Delta \sigma \times BR)^2}{(\sigma \times BR)^2} + \frac{(\Delta T)^2}{T^2} \right). \quad (10.14)$$

In this expression, the uncertainty on the cross-section is the quantity obtained in Equation 10.12, and the uncertainty on the T factor comes from theoretical uncertainties and expected experimental uncertainties on Γ_{tot} . Assuming $100fb^{-1}$ luminosity and the aforementioned systematic sources, the uncertainty on the cross-section is found to be 36% for a $150 GeV/c^2$ Higgs boson. The corresponding uncertainty on the Γ_W partial width is 21%. These results depend on the Higgs boson mass, and are presented Figure 10.47.



10.3.1 Accuracy of the Higgs boson mass measurement

FIXME: preliminary

Figure 10.48 shows the statistical precision of the Higgs boson mass measurement for the 30 fb^{-1} using inclusive Higgs boson production $pp \rightarrow H + X$ and the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ \rightarrow 4\ell$ decay modes.




Figure 10.48: The statistical precision of the Higgs boson mass measurement for the 30 fb^{-1} using inclusive Higgs boson production $pp \rightarrow H + X$ and the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ \rightarrow 4\ell$ decay modes.

10.3.2 Discovery reach for the Standard Model Higgs boson

This section summarizes the discovery reach for the Standard Model Higgs boson. The cross sections and branching ratios for the Higgs boson were calculated at next-to-leading order using the programs HDECAY [32], HIGLU [31], VV2H, V2HV and HQQ [8].

Figure 10.49 shows the integrated luminosity needed for the 5σ discovery of the inclusive Higgs boson production $pp \rightarrow H + X$ with the Higgs boson decay modes $H \rightarrow \gamma\gamma$, $H \rightarrow ZZ \rightarrow 4\ell$, and $H \rightarrow WW \rightarrow 2\ell 2\nu$.

Figure 10.50 shows the discovery significance as a function of the Higgs boson mass for 30 fb^{-1} of the integrated luminosity for the different Higgs boson production and decay channels.




Figure 10.49: The integrated luminosity needed for the 5σ discovery of the inclusive Higgs boson production $pp \rightarrow H + X$ with the Higgs boson decay modes $H \rightarrow \gamma\gamma$, $H \rightarrow ZZ \rightarrow 4\ell$, and $H \rightarrow WW \rightarrow 2\ell 2\nu$.




Figure 10.50: The discovery significance as a function of the Higgs boson mass for 30 fb^{-1} of the integrated luminosity for the different Higgs boson production and decay channels




Figure 10.51: Definitions of the angles in the $\Phi \rightarrow ZZ \rightarrow 2e2\mu$ process.

10.3.3 Study of CP properties of the Higgs boson using angle correlation in the $\Phi \rightarrow ZZ \rightarrow e^+e^-\mu^+\mu^-$ process

The most general ΦVV coupling ($V = W^\pm, Z^0$) for spin-0 Higgs boson Φ^{**} looks as follows [499, 500, 501, 502]:

$$\mathcal{C}_{\Phi VV}^{J=0} = \kappa \cdot g^{\mu\nu} + \frac{\zeta}{m_V^2} \cdot p^\mu p^\nu + \frac{\eta}{m_V^2} \cdot \epsilon^{\mu\nu\rho\sigma} k_{1\rho} k_{2\sigma}, \quad (10.15)$$

where k_1, k_2 are four-momenta of vector bosons V and $p \equiv k_1 + k_2$ is four-momentum of the Higgs boson. In the present analysis a simplified version of above ΦVV coupling (Eq. 10.15) is studied with a Standard-Model-like scalar and a pseudoscalar contributions (i.e. $\kappa, \eta \neq 0$ and $\zeta = 0$). To study deviations from the Standard Model ΦZZ coupling we take $\kappa = 1^{\dagger\dagger}$. The decay width for the $\Phi \rightarrow ZZ \rightarrow (\ell_1 \bar{\ell}_1)(\ell_2 \bar{\ell}_2)$ process consists now of three terms: a scalar one (denoted by H), a pseudoscalar one $\sim \eta^2$ (denoted by A) and the interference term violating CP $\sim \eta$ (denoted by I):

$$d\Gamma(\eta) \sim H + \eta I + \eta^2 A. \quad (10.16)$$

This way the Standard-Model scalar ($\eta = 0$) and the pseudoscalar (in the limit $|\eta| \rightarrow \infty$) contributions could be recovered. It is convenient to introduce a new parameter ξ , defined by $\tan \xi \equiv \eta$, which is finite and has values between $-\pi/2$ and $\pi/2$. Expressions for H, A and I can be found in article [500].

In study of the CP-parity of the Higgs boson we use two angular distributions. The first one is a distribution of the angle φ (called plane or azimuthal angle) between the planes of two decaying Z 's in the Higgs boson rest frame. The negatively charged leptons were used to fix planes' orientations. The second one is a distribution of the polar angle θ , in the Z rest frame, between negatively charged lepton and the direction of motion of Z boson in the Higgs boson rest frame (Fig. 10.51).

10.3.3.1 MC samples

The Higgs boson signal samples were generated using PYTHIA [62] for three masses of the Higgs boson ($m_\Phi = 200, 300, 400$ GeV). The analysis was performed for scalar, pseudoscalar and CP-violating states, the latter for $\tan \xi = \pm 0.1, \pm 0.4, \pm 1, \pm 4$. Samples for scalar, pseudoscalar and $\tan \xi = \pm 1$ states contain 10 000 events, while each of remaining samples contains 5 000 events. The predicted production cross-sections $\sigma_\Phi, \sigma_\Phi \cdot BR(\Phi \rightarrow 4\ell)$ and

**In the present section symbol Φ means the Higgs particle with unspecified CP-parity, while H (h) and A mean the scalar and pseudoscalar Higgs particles, respectively.

^{††}The ΦVV coupling with $\kappa = 1$ and arbitrary η is implemented in the PYTHIA generator.

Table 10.38: Production cross-sections of the signal. Errors are statistical only.

mass (GeV)	σ_Φ (fb)	$\sigma_\Phi \cdot BR(\Phi \rightarrow 4\ell)$ (fb)	$\sigma_\Phi \cdot \epsilon \cdot BR(\Phi \rightarrow 4\ell)$ (fb)
200	$17.86 \cdot 10^3$	38.75	7.65 ± 0.09
300	$9.41 \cdot 10^3$	24.03	5.08 ± 0.06
400	$8.71 \cdot 10^3$	20.15	4.45 ± 0.05

Table 10.39: Production cross-sections and number of used events for background processes. Errors are statistical only.

process	σ_{bkg} (fb)	$\sigma_{bkg} \cdot BR$ (fb)	$\sigma_{bkg} \cdot \epsilon \cdot BR$ (fb)	# events
ZZ/γ^*	$28.9 \cdot 10^3$	730.27	39.75 ± 0.34	20k
$t\bar{t}$	$840 \cdot 10^3$	$87.2 \cdot 10^3$	775.08 ± 4.84	48k
$Zb\bar{b}$	$525 \cdot 10^3$	$9.49 \cdot 10^3$	116.38 ± 3.22	5k

$\sigma_\Phi \cdot \epsilon \cdot BR(\Phi \rightarrow 4\ell)$, where ϵ is the preselection efficiency, are summarized in Table 10.38. We assume the Standard Model cross-section [8] and the Standard Model branching ratio [9] for each value of the ξ parameter (independently on the CP-parity of the Higgs boson). A dependence of the analysis' results on the assumed cross-section is discussed in Section 10.3.3.4. Following background processes were considered:

1. $ZZ/\gamma^* \rightarrow e^+e^-\mu^+\mu^-$ (irreducible background; generated with PYTHIA, with $m_{\gamma^*} > 5$ GeV).
2. $t\bar{t} \rightarrow W^+W^-b\bar{b} \rightarrow e^+e^-\mu^+\mu^- X$ (generated with PYTHIA).
3. $Zb\bar{b} \rightarrow e^+e^-\mu^+\mu^- X$ (generated with CompHEP, with $p_t^b > 1$ GeV, $|\eta^b| < 2.5$ and $81 < m_{Z^*} < 101$ GeV).

The information about the generated background samples are summarized in Table 10.39. The minimum-bias pile-up events for the low LHC luminosity were added to each signal and background sample.

10.3.3.2 Selection

We use selection criteria (for four isolated leptons) developed in the Standard-Model Higgs boson searches for the $H \rightarrow ZZ \rightarrow e^+e^-\mu^+\mu^-$ process (see Section 10.2.1). The reconstructed angular distributions after the final selection for the signal (with mass $m_\Phi=300$ GeV) for various values of the parameter ξ , and for the background are shown in Fig. 10.52.

10.3.3.3 Determination of the parameter ξ

The parameter ξ was determined by maximization of the likelihood function $\mathcal{L}(\xi, R)$, which was constructed from angular distributions and invariant mass distribution of four leptons, for the signal and the background. The function depends on two parameters: ξ describing CP of the signal, and R describing fraction of the signal in the data sample. The function has the following form:

$$\mathcal{L}(\xi, R) \equiv 2 \sum_{x_i \in \text{data}} \log Q(\xi, R; x_i), \text{ where } Q(\xi, R; x_i) \equiv R \cdot \mathcal{PDF}_S(\xi; x_i) + (1 - R) \cdot \mathcal{PDF}_B(x_i). \quad (10.17)$$




Figure 10.52: The φ -distributions (left) and the θ -distributions (right) for various values of the parameter ξ after final selection (normalized to 60 fb^{-1}). The signal for $m_\phi=300 \text{ GeV}$ and $\xi = 0$ (scalar), $\xi = -\pi/4$, $\xi = +\pi/4$ and $|\xi| = \pi/2$ (pseudoscalar), respectively (empty histograms). The ZZ background - filled-up histogram.

$\mathcal{PDF}_B(x_i)$ and $\mathcal{PDF}_S(\xi; x_i)$ are probability density functions for background and signal respectively; $\{x_i\}$ are values of the measured quantities (angles and invariant mass) in the data event i . They are products of probability densities \mathcal{P}^M , \mathcal{P}^φ , $\mathcal{P}^{\cos\theta_{1,2}}$ of four leptons invariant mass and angles φ and $\cos\theta_{1,2}$: $\mathcal{PDF} \equiv \mathcal{P}^M \cdot \mathcal{P}^\varphi \cdot \mathcal{P}^{\cos\theta_1} \cdot \mathcal{P}^{\cos\theta_2}$ for the signal and background. \mathcal{P}^M , \mathcal{P}^φ , $\mathcal{P}^{\cos\theta_{1,2}}$ are obtained by the Monte Carlo technique, using normalized histograms of given quantities after the final selection.

A part of the function \mathcal{Q} which describes angular distributions of signal depends on the parameter ξ , namely from Eq. (10.16) we obtain:

$$\mathcal{P}(\xi) \equiv (\mathcal{P}_S^\varphi \cdot \mathcal{P}_S^{\cos\theta_1} \cdot \mathcal{P}_S^{\cos\theta_2})(\xi) \equiv (\mathcal{H} + \tan\xi \cdot \mathcal{I} + \tan^2\xi \cdot a^2\mathcal{A}) / (1 + a^2 \tan^2\xi), \quad (10.18)$$

where: $\mathcal{H} \equiv \mathcal{P}_H^\varphi \cdot \mathcal{P}_H^{\cos\theta_1} \cdot \mathcal{P}_H^{\cos\theta_2}$ and $\mathcal{A} \equiv \mathcal{P}_A^\varphi \cdot \mathcal{P}_A^{\cos\theta_1} \cdot \mathcal{P}_A^{\cos\theta_2}$ are probability densities obtained by the Monte Carlo technique for the scalar (H) and the pseudoscalar (A), respectively. The parameter a^2 is a (mass dependent) relative strength of the pseudoscalar and scalar couplings. For example $a^2=0.51, 1.65, 1.79$ for $m_\phi=200, 300, 400 \text{ GeV}$, respectively. \mathcal{I} is a normalized product of angular distributions for the CP-violating term. \mathcal{I} is not positive, and its integral is equal to zero, so it is not possible to simulate it separately. The \mathcal{I} contribution can be obtained indirectly from the combined probability density for the signal with non-zero value of the parameter ξ . For example by introducing $\mathcal{P}_+ \equiv \mathcal{P}(\pi/4) = (\mathcal{H} + \mathcal{I} + a^2\mathcal{A}) / (1 + a^2)$ and $\mathcal{P}_- \equiv \mathcal{P}(-\pi/4) = (\mathcal{H} - \mathcal{I} + a^2\mathcal{A}) / (1 + a^2)$ we have $\mathcal{I} = (1 + a^2) / 2 \cdot (\mathcal{P}_+ - \mathcal{P}_-)$. Finally we obtain:

$$\mathcal{P}(\xi) \equiv [\mathcal{H} + \tan\xi \cdot \frac{1 + a^2}{2} \cdot (\mathcal{P}_+ - \mathcal{P}_-) + \tan^2\xi \cdot a^2\mathcal{A}] / (1 + a^2 \tan^2\xi). \quad (10.19)$$

10.3.3.4 Results

After selection all background contribution but $ZZ/\gamma^* \rightarrow e^+e^-\mu^+\mu^-$ are negligible, therefore only these events were used to construct probability density function for the background. The ZZ/γ^* sample contains 15 000 events at the generator level. Signal probability density functions were constructed using samples of scalar Higgs boson (H), pseudoscalar (A) and \mathcal{P}_+ , \mathcal{P}_- samples, each containing 8 000 events at the generator level. Likelihood functions were constructed independently for three masses of the Higgs boson ($m_\phi=200, 300, 400 \text{ GeV}$).




Figure 10.53: Reconstructed value of the parameter ξ as function of the generated value of the parameter ξ , for $\mathcal{L}=60 \text{ fb}^{-1}$, for Higgs boson mass $m_\Phi=200, 300, 400 \text{ GeV}$. Uncertainties correspond to one standard deviation.

For each value of the parameter ξ and for each Higgs-boson mass we made 200 pseudoexperiments for the integrated luminosity $\mathcal{L}=60 \text{ fb}^{-1}$. For each pseudoexperiment we randomly selected events from the signal and background samples to form a test sample. The number of selected events was given by a Poisson probability distribution with mean defined by the process cross-section and the examined luminosity. Then we performed a maximization of the likelihood function $\mathcal{L}(\xi, \mathcal{R})$ for the test sample to obtain a value of the parameter ξ . The expected and reconstructed values of the parameter ξ with its uncertainty, obtained for three masses of the Higgs boson are shown in Fig. 10.53.

In our analysis the Standard-Model signal cross-section and branching ratio were used as a reference. However both of them may change for other Higgs models. An influence of possible suppression (enhancement) factor C^2 of the signal (in respect to the Standard Model expectation) on the reconstructed ξ were studied. We found that value of ξ slightly depends on size of suppression (enhancement). On the other hand, the uncertainty of ξ is approximately proportional to $1/C$ (i.e. it depends on square-root of number of events, what one can expect):

$$\Delta\xi(\xi, C^2) \equiv \frac{\sigma_0(\xi)}{\sqrt{C^2}}. \quad (10.20)$$

A value of $\sigma_0(\xi)$ (a precision factor) can be determined from the fit. Taking this into account we parametrize a relative error of distinction between an arbitrary value of ξ and a reference value ξ_0 (e.g. $\xi_0=0$, scalar) as follows:

$$\frac{\sigma_{|\xi-\xi_0|}(C^2)}{|\xi-\xi_0|} = \frac{\Delta\xi(\xi, C^2)}{|\xi-\xi_0|}. \quad (10.21)$$

The requirement of exclusion of $\xi \neq \xi_0$ at the level of “ N sigmas” could be written as $\sigma_{|\xi-\xi_0|}(C^2) = |\xi-\xi_0|/N$:

$$C^2(N) = N^2 \frac{\sigma_0^2(\xi)}{(\xi-\xi_0)^2}. \quad (10.22)$$

The exclusion contours for $N=1, 3$ and for $\xi_0=0$ (scalar) are shown in Fig. 10.54.




Figure 10.54: Exclusion contours for scalar Higgs boson as a function of the enhancement factor C^2 for the Higgs boson masses $m_\Phi=200, 300, 400$ GeV (from left to right). The excluded area lies above curves. Results were obtained for 60 fb^{-1} .

Chapter 11

MSSM Higgs bosons



11.1 Introduction

Supersymmetric extensions of the SM [503, 504, 505, 506, 317] are strongly motivated by the idea of providing a solution of the hierarchy problem in the Higgs sector. They allow for a light Higgs particle in the context of GUTs [507], in contrast with the SM, where the extrapolation requires an unsatisfactory fine-tuning of the SM parameters. Supersymmetry is a symmetry between fermionic and bosonic degrees of freedom and thus the most general symmetry of the S -matrix. The minimal supersymmetric extension of the SM (MSSM) yields a prediction of the Weinberg angle in agreement with present experimental measurements if embedded in a SUSY-GUT [508, 509]. Moreover, it does not exhibit any quadratic divergences, in contrast with the SM Higgs sector. Owing to the large top quark mass SUSY-GUTs develop electroweak symmetry breaking at the electroweak scale dynamically [510, 511, 512, 513]. The lightest supersymmetric particle offers a proper candidate for the Cold Dark Matter content of the universe, if R -parity is conserved. Finally, local supersymmetry enforces gravitational interactions.


In the MSSM two isospin Higgs doublets have to be introduced in order to preserve supersymmetry [514, 515, 511]. After the electroweak symmetry-breaking mechanism, three of the eight degrees of freedom are absorbed by the Z and W gauge bosons, leading to the existence of five elementary Higgs particles. These consist of two CP-even neutral (scalar) particles h, H , one CP-odd neutral (pseudoscalar) particle A , and two charged particles H^\pm . In order to describe the MSSM Higgs sector one has to introduce four masses M_h, M_H, M_A and M_{H^\pm} and two additional parameters, which define the properties of the scalar particles and their interactions with gauge bosons and fermions: the mixing angle β , related to the ratio of the two vacuum expectation values, $\tan \beta = v_2/v_1$, and the mixing angle α in the neutral CP-even sector. Due to supersymmetry there are several relations among these parameters, and only two of them are independent at leading order. In the absence of CP-violation they are usually chosen as M_A and $\tan \beta$. The other Higgs-boson masses and mixing angles are calculable in terms of the other MSSM parameters. Measuring the masses and angles will constitute an important consistency check of the MSSM.

At tree-level the following mass hierarchies hold: $M_h < M_Z, M_A < M_H$ and $M_W < M_{H^\pm}$. The tree-level bound on M_h receives large corrections from SUSY-breaking effects in the Yukawa sector of the theory. The leading one-loop correction is proportional to m_t^4 . The leading logarithmic one-loop term (for vanishing mixing between the scalar top quarks)

reads [516, 517, 518, 519, 520, 521, 522]

$$\Delta M_h^2 = \frac{3G_\mu m_t^4}{\sqrt{2}\pi^2 \sin^2 \beta} \ln \left(\frac{m_{\tilde{t}_1} m_{\tilde{t}_2}}{m_t^2} \right), \quad (11.1)$$

where G_μ is the Fermi constant, and $m_{\tilde{t}_{1,2}}$ are the two stop masses. Corrections of this kind have drastic effects on the predicted value of M_h and many other observables in the MSSM Higgs sector. The higher-order contributions can shift M_h by 50–100% [523, 524, 525, 526, 527, 137, 528, 529, 530, 531, 532, 533, 138, 534]. The corrections to the MSSM Higgs boson sector have been evaluated in several approaches. The status of the available calculations can be summarized as follows. For the one-loop part, the complete result within the MSSM is known [518, 516, 517, 522, 535, 536, 537, 538]. The by far dominant one-loop contribution is the $\mathcal{O}(\alpha_t)$ term due to top and stop loops ($\alpha_t \equiv h_t^2/(4\pi)$, h_t being the top-quark Yukawa coupling). Concerning the two-loop effects, their computation is quite advanced and has now reached a stage such that all the presumably dominant contributions are known [524, 525, 526, 527, 137, 539, 528, 529, 540, 541, 542, 531, 532, 533, 534, 543, 544, 545, 546, 547, 548, 549]. They include (evaluated for vanishing external momenta) the strong corrections, $\mathcal{O}(\alpha_t \alpha_s)$, and Yukawa corrections, $\mathcal{O}(\alpha_t^2)$, to the dominant one-loop $\mathcal{O}(\alpha_t)$ term, as well as the strong corrections to the bottom/sbottom one-loop $\mathcal{O}(\alpha_b)$ term ($\alpha_b \equiv h_b^2/(4\pi)$), i.e. the $\mathcal{O}(\alpha_b \alpha_s)$ contribution. The latter can be relevant for large values of $\tan \beta$. For the (s)bottom corrections the all-order resummation of the $\tan \beta$ -enhanced terms, $\mathcal{O}(\alpha_b (\alpha_s \tan \beta)^n)$, has also been computed. Finally, the $\mathcal{O}(\alpha_t \alpha_b)$ and $\mathcal{O}(\alpha_b^2)$ corrections have been obtained. The higher-order corrections shift the upper bound of M_h to $M_h \lesssim 135$ GeV [137, 138]. The remaining theoretical uncertainty on M_h has been estimated to be below ~ 3 GeV [138, 550]. Besides the masses of the Higgs bosons, also their couplings are affected by large higher-order corrections (see below).

An important feature of the MSSM Higgs sector is that for large pseudoscalar masses M_A the light scalar Higgs mass reaches its upper bound and becomes SM-like. Moreover, for large values of $\tan \beta$ the down(up)-type Yukawa couplings are strongly enhanced (suppressed) apart from the region, where the light (heavy) scalar is at its upper (lower) mass bound. The atively corrected Higgs masses are depicted in Fig. 11.1.




Figure 11.1: The CP-even and charged MSSM Higgs boson masses as a function of M_A for $\tan \beta = 3$ and 30, including radiative corrections [551].

The LEP experiments have searched for the MSSM Higgs bosons via the Higgs-strahlung process $e^+e^- \rightarrow Z + h/H$ and the associated production $e^+e^- \rightarrow A + h/H$ for the neutral Higgs particles and $e^+e^- \rightarrow H^+H^-$ for the charged Higgs bosons. Neutral Higgs masses $M_A \lesssim 91.9$ GeV and $M_{h/H} \lesssim 91$ GeV are excluded [552] as well as charged Higgs masses $M_{H^\pm} \lesssim 78.6$ GeV [553].

The lightest Higgs boson h will mainly decay into $b\bar{b}$ and $\tau^+\tau^-$ pairs, since its mass is below ~ 135 GeV, see Fig. 11.2a. Close to its upper bound in mass all decay modes as for the SM Higgs boson open up rapidly. For large values of $\tan\beta$ the heavy scalar and pseudoscalar Higgs particles H, A will decay dominantly into $b\bar{b}, \tau^+\tau^-$ pairs, too, due to the enhanced Yukawa couplings for down-type fermions. The branching ratios for the decays into $b\bar{b}$ and $\tau^+\tau^-$ are about 90% and 10% respectively. Other heavy scalar Higgs decay modes as $H \rightarrow t\bar{t}, W^+W^-, ZZ, hh, AA$ develop sizeable branching ratios only for small values of $\tan\beta$ (see Fig. 11.2b) and analogously the pseudoscalar Higgs decays $A \rightarrow t\bar{t}, gg, Zh$ (see Fig. 11.2c). The charged Higgs bosons decay mainly into $\tau\nu_\tau$ pairs for $M_{H^\pm} \lesssim 180$ GeV and into tb final states above (see Fig. 11.2d). All other decay modes do not acquire branching ratios larger than a few per cent. The (SUSY-)QCD [386, 387, 388, 389, 390, 391, 392, 554, 535, 548] and (SUSY-)electroweak corrections [393, 394, 395, 396, 554, 555] to the fermionic decay modes are sizeable. In addition to the usual large QCD corrections, significant corrections arise from virtual sbottom/stop and gluino/gaugino exchange contributions in the $h, H, A \rightarrow b\bar{b}$ and $H^\pm \rightarrow tb$ decay modes [554, 535, 548, 555]. The dominant part of the latter corrections can be absorbed in improved bottom Yukawa couplings. In this way these contributions can also be resummed up to all orders thus yielding reliable perturbative results [549, 546]. The rare photonic decay modes $h, H, A \rightarrow \gamma\gamma$ are mediated by W, t, b loops as in the SM Higgs case and additional contributions from charged Higgs bosons, charginos and sfermions, if these virtual particles are light enough [8, 370, 371]. The QCD corrections to these decay modes can reach a few per cent in the relevant mass regions [397, 398, 399, 400, 401, 402, 403]. If decays into supersymmetric particles, i.e. gauginos and sfermions, are possible, they acquire significant branching ratios and can even be the dominant decay modes [556, 557, 8, 370, 371]. In contrast to the SM the total widths of the MSSM Higgs bosons do not exceed several tens of GeV, so that the MSSM Higgs particles appear as narrow resonances.

The dominant neutral MSSM Higgs production mechanisms for small and moderate values of $\tan\beta$ are the gluon fusion processes

$$gg \rightarrow h, H, A$$

which are mediated by top and bottom loops as in the SM case, but in addition by stop and sbottom loops for the scalar Higgs bosons h, H , if the squark masses are below about 400 GeV [558]. The NLO QCD corrections to the quark loops are known in the heavy quark limit as well as including the full quark mass dependence [410, 411, 412, 414, 415, 416, 417]. They increase the cross sections by up about 100% for smaller $\tan\beta$ and up to about 40% for very large $\tan\beta$, where the bottom loop contributions become dominant due to the strongly enhanced bottom Yukawa couplings. The limit of heavy quarks is only applicable for $\tan\beta \lesssim 5$ within about 20–25%, if full mass dependence of the LO terms is taken into account [8, 370, 371, 413]. Thus the available NNLO QCD corrections in the heavy quark limit [418, 419, 420, 421] can only be used for small and moderate $\tan\beta$, while for large $\tan\beta$ one has to rely on the fully massive NLO results [410, 411, 412]. The QCD corrections to the squark loops are only known in the heavy squark limit [558] and the full SUSY-QCD corrections in the limit of heavy squarks and gluinos [559, 560, 561, 562]. The pure QCD corrections are of about the




Fig. 11.2a




Fig. 11.2b

Figure 11.2: Branching ratios of the MSSM Higgs bosons h, H, A, H^\pm for non-SUSY decay modes as a function of the masses for two values of $\tan\beta = 3, 30$ and maximal mixing. The common squark mass has been chosen as $M_S = 1$ TeV. The other SUSY-parameters have been chosen as $M_2 = m_{\tilde{g}} = \mu = 1$ TeV and $A_{t,b} = 2783$ (2483) TeV for $\tan\beta = 3$ (30).




Fig. 11.2c




Fig. 11.2d

Figure 11.2: :Continued.

same size as those to the quark loops thus rendering the total K factor of similar size as for the quark loops alone with a maximal deviation of about 10% [558]. The pure SUSY-QCD corrections are small [559, 560, 561, 562]. The NNLL resummation of the SM Higgs cross section [422] can also be applied to the scalar MSSM Higgs cross sections in the regions, where the heavy quark and squark limits are valid. The same is also true for the NLO QCD corrections to the p_T distributions [429, 430, 431, 432, 433] and the NNLL resummation of soft gluon effects [434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444], i.e. for small values of $\tan\beta$, M_H and p_T only. However, for large values of $\tan\beta$ the p_T distributions are only known at LO, since the bottom loops are dominant and the heavy top limit is not valid. An important consequence is that the p_T distributions of the neutral Higgs bosons are softer than for small values of $\tan\beta$ [563].

The vector-boson fusion processes [450, 452]

$$pp \rightarrow qq \rightarrow qq + WW/ZZ \rightarrow qq + h/H$$

play an important role for the light scalar Higgs boson h close to its upper mass bound, where it becomes SM-like, and for the heavy scalar Higgs particle H at its lower mass bound. In the other regions the cross sections are suppressed by the additional SUSY-factors of the Higgs couplings. The NLO QCD corrections to the total cross section and the distributions can be taken from the SM Higgs case and are of the same size [453, 454]. The SUSY-QCD corrections mediated by virtual gluino and squark exchange at the vertices turned out to be small [564].

Higgs-strahlung off W, Z gauge bosons [455, 456]

$$pp \rightarrow q\bar{q} \rightarrow Z^*/W^* \rightarrow H + Z/W$$

does not play a major role for the neutral MSSM Higgs bosons at the LHC. The NLO [457] and NNLO [458] QCD corrections are the same as in the SM case, and the SUSY-QCD corrections are small [564]. The SUSY-electroweak corrections are unknown.

Higgs radiation off top quarks [460, 461, 462, 463, 464]

$$pp \rightarrow q\bar{q}/gg \rightarrow h/H/A + t\bar{t}$$

plays a significant role at the LHC for the light scalar Higgs particle only. The NLO QCD corrections are the same as for the SM Higgs boson with modified top and bottom Yukawa couplings and are thus of moderate size [465, 153, 466]. The SUSY-QCD corrections have been computed recently for the light scalar case [565]. They are of moderate size.

For large values of $\tan\beta$ Higgs radiation off bottom quarks [460, 461, 462, 463, 464]

$$pp \rightarrow q\bar{q}/gg \rightarrow h/H/A + b\bar{b}$$

constitutes the dominant Higgs production process. The NLO QCD corrections can be taken from the analogous calculation involving top quarks. However, they turn out to be very large [566, 567, ?]. The main reason is that the integration over the transverse momenta of the final state bottom quarks generates large logarithmic contributions. The resummation of the latter requires the introduction of bottom quark densities in the proton, since the large logarithms are related to the DGLAP-evolution of these densities. Their DGLAP-evolution resums them. This leads to an approximate approach starting from the process [568, ?] (see Fig. 11.3a)




Figure 11.3: Typical diagrams for all Higgs boson production mechanisms related to Higgs radiation off bottom quarks at leading order: (a) $b\bar{b} \rightarrow h/H/A$, (b) $gb \rightarrow b + h/H/A$, (c) $gg \rightarrow b\bar{b} + h/H/A$.

$$pp \rightarrow b\bar{b} \rightarrow h/H/A$$

at LO, where the transverse momenta of the incoming bottom quarks, their masses and their off-shellness are neglected. The NLO [569, 570] and NNLO [571] QCD corrections to this bottom-initiated process are known and of moderate size, if the running bottom Yukawa coupling at the scale of the Higgs mass is introduced. At NNLO the full process $gg \rightarrow h/H/A + b\bar{b}$ (see Fig. 11.3c) contributes for the first time. At this order a proper matching to the fully massive result for this process can be performed [572, 573] so that the final expression provides an improved result, which takes into account the resummation of the large logarithms and mass effects. The fully exclusive $gg \rightarrow h/H/A + b\bar{b}$ process, calculated with four active parton flavors in a fixed flavour number scheme (FFNS), and this improved resummed result, calculated with 5 active parton flavours in the variable flavour number scheme (VFNS), will converge against the same value at higher perturbative orders. The best agreement between the NLO FFNS and NNLO VFNS is achieved, if the factorization scale of the bottom quark densities is chosen as about a quarter of the Higgs mass [574, 575]. If only one of the final state bottom jets accompanying the Higgs particle is tagged, the LO bottom-initiated process is $gb \rightarrow b + h/H/A$ (see Fig. 11.3b), the NLO QCD corrections of which have been calculated [?]. They turn out to reach $\mathcal{O}(40 - 50\%)$. The situation concerning the comparison with the FFNS at NLO is analogous to the total cross section. Agreement within the respective theoretical uncertainties is found for a factorization scale of about a quarter of the Higgs mass [574]. If both bottom jets accompanying the Higgs boson in the final state are tagged, one has to rely on the fully exclusive calculation for $gg \rightarrow b\bar{b} + h/H/A$.

All neutral MSSM Higgs production cross sections including the NLO QCD corrections are shown in Fig. 11.4.

The dominant charged Higgs production process is the associated production with heavy quarks [576, 577, 578] (see Fig. 11.5a)

$$pp \rightarrow q\bar{q}, gg \rightarrow H^- + t\bar{b} \quad \text{and c.c.}$$

The NLO QCD and SUSY-QCD corrections have very recently been computed [579, ?]. They are of significant size due to the large logarithms arising from the transverse-momentum integration of the bottom quark in the final state and the large SUSY-QCD corrections to the bottom Yukawa coupling. The large logarithms can be resummed by the introduction of bottom quark densities in the proton in complete analogy to the neutral Higgs case. In this approach the LO process is $gb \rightarrow H^- t$ and the charge conjugate. The NLO SUSY-QCD corrections have been derived in [580, 581, 582, 583] and found to be of significant size. This process, however, relies on the same approximations as all bottom-initiated processes. A quantitative comparison of the processes $gb \rightarrow H^- t$ and $gg \rightarrow H^- + t\bar{b}$ at NLO is missing so far.




Fig. 11.4a




Fig. 11.4b

Figure 11.4: Neutral MSSM Higgs production cross sections at the LHC for gluon fusion $gg \rightarrow \Phi$, vector-boson fusion $qq \rightarrow qqVV \rightarrow qqh/qqH$, Higgs-strahlung $q\bar{q} \rightarrow V^* \rightarrow hV/HV$ and the associated production $gg, q\bar{q} \rightarrow b\bar{b}\Phi/t\bar{t}\Phi$, including all known QCD corrections. (a) h, H production for $\tan\beta = 3$, (b) h, H production for $\tan\beta = 30$, (c) A production for $\tan\beta = 3$, (d) A production for $\tan\beta = 30$. The same parameters as in Fig. 11.2 have been adopted.






Fig. 11.4c




Fig. 11.4d

Figure 11.4: :Continued.




Figure 11.5: Typical diagrams for charged Higgs boson production mechanisms at leading order: (a) $gg \rightarrow H^- t\bar{b}$, (b) $q\bar{q} \rightarrow H^+ H^-$, (c) $gg \rightarrow H^+ H^-$, (d) $gg \rightarrow W^+ H^-$, (e) $b\bar{b} \rightarrow H^+ H^-$, (f) $b\bar{b} \rightarrow W^+ H^-$.

The second important charged Higgs production process is charged Higgs pair production in a Drell–Yan type process (see Fig. 11.5b)

$$pp \rightarrow q\bar{q} \rightarrow H^+ H^-$$

which is mediated by s -channel photon and Z -boson exchange. The NLO QCD corrections can be taken from the Drell–Yan process and are of moderate size as in the case of the neutral Higgs-strahlung process discussed before. The genuine SUSY–QCD corrections, mediated by virtual gluino and squark exchange in the initial state, are small [564].

Charged Higgs pairs can also be produced from gg initial states by the loop-mediated process [584, 585, 586, 587, 588] (see Fig. 11.5c)

$$pp \rightarrow gg \rightarrow H^+ H^-$$

where the dominant contributions emerge from top and bottom quark loops as well as stop and sbottom loops, if the squark masses are light enough. The NLO corrections to this process are unknown. This cross section is of similar size as the bottom-initiated process [588] (see Fig. 11.5e)

$$pp \rightarrow b\bar{b} \rightarrow H^+ H^-$$

which relies on the approximations required by the introduction of the bottom densities as discussed before and is known at NLO [589]. The SUSY–QCD corrections are of significant size. The pure QCD corrections and the genuine SUSY–QCD corrections can be of opposite sign.

Finally, charged Higgs bosons can be produced in association with a W boson [590, 591, 592] (see Fig. 11.5d)

$$pp \rightarrow gg \rightarrow H^+ W^- \quad \text{and c.c.}$$

which is generated by top-bottom quark loops and stop-sbottom loops, if the squark masses are small enough. This process is known at LO only. The same final state also arises from the process [590, 591, 593] (see Fig. 11.5f)

$$pp \rightarrow b\bar{b} \rightarrow H^+W^- \quad \text{and c.c.}$$

which is based on the approximations of the VFNS. The QCD corrections have been calculated and turn out to be of moderate size [594, 595].

2 Higgs boson channels

11.2.1 Associated $b\bar{b}H$ production with $H \rightarrow \tau\tau \rightarrow e^\pm\mu^\mp + E_T^{\text{miss}}$

Compared to the hadronic and semileptonic final states described in Section 5.2, the fully leptonic final states are suppressed by relatively small branching ratio $\text{BR}(\tau \rightarrow \mu\nu\nu) \sim 0.174$ and $\text{BR}(\tau \rightarrow e\nu\nu) \sim 0.178$, but the signal is clean and easy to trigger.

The signal consists of events in which the Higgs boson decays into two tau leptons which in turn decay leptonically. Two possibilities exist, either to select any-two-lepton final states, which have larger signal rate, or electron+muon final states for which the background is easier to suppress. Here the electron+muon final state is chosen.

The main backgrounds for $H/A \rightarrow \tau\tau$ with $e\mu$ final state are the Drell-Yan $\tau\tau$ production, the $t\bar{t}$ and the Wt production where the W boson coming from top quark decay decays leptonically, the $\tau\tau b\bar{b}$ production, and the $b\bar{b}$ background with b quarks decaying semileptonically. Other backgrounds are pairs of vector bosons WW or WZ decaying into leptonic final states, but their contribution is small. The $\tau\tau c\bar{c}$ background is also found negligible. The most biggest background arises from those $t\bar{t}$ and Drell-Yan events which involve genuine τ 's and b jets and produce events very similar to the signal. No SUSY particle background is assumed.

11.2.1.1 Event generation

The Higgs boson signal is generated with PYTHIA [238]. The signal cross sections and branching ratios are calculated with FeynHiggs [136]. TAUOLA package [146] is used for leptonic τ decays in the signal events.

The Drell-Yan $\tau\tau$ production, $b\bar{b}$, WW , WZ and ZZ backgrounds are generated with PYTHIA. The Drell-Yan $\tau\tau$ next-to-leading order cross section of 1891 pb calculated with the program MCFM [48] for $M_{\tau\tau} > 80\text{GeV}/c^2$ is used. The $\tau\tau b\bar{b}$ background is generated with CompHEP [34] with no p_T and η cuts applied on b quarks and the leading order cross section calculated with CompHEP are used. The Z/γ^* generation is split into two bins of generated $\tau\tau$ mass $m_{\tau\tau}$: $80 - 100\text{GeV}/c^2$ and $> 100\text{GeV}/c^2$, and the $\tau\tau b\bar{b}$ is generated in the $\tau\tau$ mass bins of $60-100\text{GeV}/c^2$ and $> 100\text{GeV}/c^2$.

The $t\bar{t}$ background is generated with TopREX [35] and PYTHIA and the single top (Wt) events are generated with TopREX. A cross section of 840 and 60 pb is used for $t\bar{t}$ and Wt events, respectively.

11.2.1.2 Level 1 and HLT selections

The events are triggered with the single and the double electron and muon triggers. The p_T threshold for single muons is 19 GeV/c, for single electrons 26 GeV/c, for double muons 7 GeV/c and for double electrons 14.5 GeV/c. The Level 1 trigger efficiency for the signal of $M_A=200$ GeV/ c^2 is 0.96, and the overall trigger efficiency including the HLT is 0.82. The corresponding trigger efficiencies for the Drell-Yan $\tau\tau$, the $\tau\tau b\bar{b}$, the $t\bar{t}$ and the Wt backgrounds are 0.18, 0.29, 0.68 and 0.68, respectively.

In the future also a combined e+mu trigger with symmetric thresholds of 10 GeV/c for the electron and muon will be included. No large gain is expected since events passing e+mu trigger are most probably already triggered by the single muon trigger.

11.2.1.3 Offline selections

The basic event selection is a requirement of two isolated leptons (one e and one μ) with $p_T > 20$ GeV/c in the central detector acceptance region $|\eta| < 2.5$ coming from a reconstructed primary vertex (PV). The electron candidates are required to pass electron identification cuts described in [147]. The efficiency for the electron identification is about 90% for electrons passing the trigger. The leptons are defined isolated when there are no other tracks from the primary vertex with $p_T > 1$ GeV/c within a cone $\Delta R = \sqrt{\Delta\varphi^2 + \Delta\eta^2} \leq 0.4$ around the lepton. The p_T cut and the isolation reduce efficiently the backgrounds with soft leptons ($b\bar{b}, c\bar{c}, \dots$).

The b jets associated with the Higgs boson provide a powerful tool to separate the $b\bar{b}H/A$ events from the Drell-Yan background. The Drell-Yan background in which Z/γ^* decay into a tau pair has a large cross section compared to the Higgs production. However, these events are mostly produced with no associated jets, and if they have associated jets they are mostly light quark and gluon jets. Therefore the Drell-Yan background can be suppressed by requiring a reconstructed jets present in the event, and even further by requiring that the associated jets are identified as b jets. The b jets associated with the Higgs bosons are generally very soft, which makes their tagging a challenging task. For low jet E_T values the track multiplicity and momenta tend to be low, and many jets do not have enough significant tracks to be identified as a b jet. As a consequence the b tagging efficiency is not very high. The b tagging efficiency of 43% per jet for the signal events with 2% of the mistagging rate is found.

The $t\bar{t}$ background cannot be suppressed with b tagging due the presence of two energetic genuine b jets in the event. In fact, the jet reconstruction and the b-tagging efficiencies are higher for b jets in $t\bar{t}$ events than for those associated with the signal. This can be exploited using a central jet veto: if more than one jet is found, the event is rejected. The threshold of 20 GeV is set on the calibrated E_T for the jets within the tracker acceptance region, $|\eta| < 2.5$. A suppression factor of 8 is obtained against the $t\bar{t}$ background with an efficiency of 60% for the signal.

A missing energy measurement is needed for estimating the fraction of the energy carried away by neutrinos. This information is used in the Higgs boson mass reconstruction. The amount of missing transverse energy is small and close to the detector resolution.

The τ 's from the Higgs boson with $M_A = 200$ GeV/ c^2 travel on average about 5 mm before they decay. Therefore the leptons coming from τ decays are displaced relative to the primary vertex [596]. The track impact parameter measurements in the transverse plane for the two

leptons are combined quadratically into one variable $\sigma_{ip} = \sigma_{ip}(\tau_1) \oplus \sigma_{ip}(\tau_2)$, where $\sigma_{ip}(\tau_1, \tau_2)$ are significances of the lepton impact parameters. The leptons in $t\bar{t}$ background come mostly from W decays. The $t\bar{t}$ events with two intermediate τ 's cannot be suppressed by using impact parameter.

The neutrinos-charged lepton collinear approximation method for the mass reconstruction in $H/A \rightarrow \tau\tau$ is described in section 5.2.5. The mass reconstruction is possible when the two leptons are not in a back-to-back configuration. The back-to-back events are removed with a cut on the angle between the two leptons in the transverse plane $\Delta\varphi(e, \mu) < 175^\circ$. Uncertainties of the missing transverse energy measurement can lead to negative neutrino energies. For the signal $\sim 40\%$ of events are lost when the positive neutrino energies are required. This requirement, however, yields a further suppression of the $t\bar{t}$ and Wt backgrounds, since for these backgrounds the neutrinos are generally not emitted along the lepton directions. The efficiencies of $E_{\nu 1, \nu 2} > 0$ cut for these backgrounds are about 17% and 15%, respectively. The reconstructed $\tau\tau$ mass with 30 fb^{-1} after all selections, but the mass window, is shown in Figures 11.6 and 11.7. In the Figures the signal of $M_A=140, \tan\beta=20$ and $200 \text{ GeV}/c^2, \tan\beta=25$ in the m_h^{max} scenario and the backgrounds are presented.




Figure 11.6: The $\tau\tau$ reconstructed mass with 30 fb^{-1} after all selections, but the mass window. The signal of $M_A = 140 \text{ GeV}/c^2$ and $\tan\beta = 20$ in the m_h^{max} scenario and the backgrounds are shown.




Figure 11.7: The $\tau\tau$ reconstructed mass with 30 fb^{-1} after all selections, but the mass window. The signal of $M_A = 200 \text{ GeV}/c^2$ and $\tan\beta = 25$ in the m_h^{max} scenario and the backgrounds are shown.

11.2.1.4 Expected number of events

Table 11.1 shows the cross section times branching ratio for the backgrounds for each step of the selections. The signal cross sections for $M_A = 140, 200$ and $250 \text{ GeV}/c^2$ and $\tan\beta=20$ in the m_h^{max} scenario are shown in Table 11.2. The expected number of events with 30 fb^{-1} after all cuts, but mass window, is also shown in Tables 11.1 and 11.2. The expected number of events after all cuts including the mass window is shown for the signal and the total background in Table 11.3.

Table 11.1: The background cross section times branching ratio (in pb) for each step of the selections. The expected number of events at 30 fb^{-1} is also shown.

	Z, γ^*	bbZ, γ^*	$t\bar{t}$	tW	bb	VV
$\sigma \times BR$	233.1	3.422	86.2	6.16	36170	7.88
Level 1	83.9	1.85	72.2	5.37	811	5.16
HLT	42.6	0.981	53.7	4.17	78.0	4.10
reconstructed PV	40.8	0.952	53.3	4.11	78.1	3.92
isol $e+\mu, p_T$ cut	1.10	0.0270	5.65	0.452	0.0378	0.288
$Q_e + Q_\mu = 0$	1.09	0.0268	5.62	0.451	0.0374	0.248
$\sigma_{ip}(e) \oplus \sigma_{ip}(\mu)$	0.296	0.00745	0.791	0.0550	0.0254	0.0255
$N_{jets} > 0$	0.0127	0.00527	0.778	0.0509	0.00654	0.0115
b tagging	0.00457	0.00289	0.608	0.0341	0.00312	0.000547
jet veto	0.00344	0.00124	0.0745	0.0166	0.000179	0.000265
$\Delta\varphi(e, \mu)$	0.00295	0.00116	0.0696	0.0159	0.000142	0.000259
$E_{\nu 1, \nu 2} > 0$	0.00124	0.000486	0.0119	0.00246	0.0000661	0.0000546
N_{ev} at 30 fb^{-1}	37.1	14.6	355.8	73.7	2.0	1.6

Table 11.2: The signal cross section times branching ratio (in pb) for $M_A=140, 200$ and $250 \text{ GeV}/c^2$ and $\tan\beta=20$ in the m_h^{\max} scenario for each step of the selections. The expected number of events at 30 fb^{-1} is also shown.

m_A	140	200	250
$\sigma \times BR$ (pb)	3.468	1.123	0.493
L1	3.238	1.079	0.479
HLT	2.585	0.923	0.419
reconstructed PV	2.434	0.866	0.395
isol $e+\mu, p_T$ cut	0.258	0.116	0.0613
$Q_e + Q_\mu = 0$	0.256	0.116	0.0612
$\sigma_{ip}(e) \oplus \sigma_{ip}(\mu)$	0.0859	0.044	0.0260
$N_{jets} > 0$	0.0375	0.0216	0.0130
b tagging	0.0177	0.0104	0.00649
jet veto	0.0115	0.00619	0.00390
$\Delta\varphi(e, \mu)$	0.0106	0.00554	0.00351
$E_{\nu 1, \nu 2} > 0$	0.00601	0.00340	0.00222
N_{ev} at 30 fb^{-1}	180	102	67

11.2.1.5 Systematic uncertainties and the discovery reach.

The uncertainty of the event selection efficiency is related to the uncertainty of the lepton identification efficiency, the jet energy and the missing energy scale and the b tagging efficiency. The jet energy and the missing energy scale uncertainty gives the uncertainty of 7.3% on the $t\bar{t}$ background, which is the dominant background. The uncertainty of the lepton identification efficiency of 2% is used for both electrons and muons. The uncertainty of the b tagging efficiency, 5%, can be estimated from $t\bar{t}$ events as in Ref.[75]. The 5% uncertainty of the mistagging efficiency is assumed [597]. The 5.8% uncertainty of the theoretical prediction of the $t\bar{t}$ cross section is taken. The total systematic uncertainty including the luminosity uncertainty 3% yields a 12% uncertainty for the total background.

The signal significance S with 30 fb^{-1} for the signal of $M_A = 140, 200$ and $250 \text{ GeV}/c^2$ and $\tan\beta=20$ in the m_h^{\max} scenario is shown in Table 11.2 without and with the background systematic uncertainty taken into account. Figure 11.8 shows the discovery reach in the M_A - $\tan\beta$ plane in the m_h^{\max} scenario with 30 fb^{-1} . The lower (upper) curve corresponds to the $S=3$ when the background systematic uncertainty is not taken (taken) into account.

Table 11.3: The expected number of the signal plus background and the background events in a given mass windows for 30 fb^{-1} and the signal significance S without and with the background systematic uncertainty taken into account.

	$\Delta m_{\tau\tau}$	N_{S+N_B}	N_B	$S_{\text{no syst.}}$	$S_{\text{sys.}}$
$m_A = 140 \text{ GeV}/c^2, \tan\beta = 20$	100 - 200 GeV/c^2	225	107	9.9	7.3
$m_A = 200 \text{ GeV}/c^2, \tan\beta = 20$	140 - 250 GeV/c^2	163	109	4.8	3.1
$m_A = 250 \text{ GeV}/c^2, \tan\beta = 20$	160 - 380 GeV/c^2	244	204	2.7	1.4




Figure 11.8: The discovery region for $gg \rightarrow H/A, H/A \rightarrow \tau\tau \rightarrow e\mu + X$ channel in M_A - $\tan\beta$ in the m_h^{max} scenario with 30 fb^{-1} .

11.2.2 Associated $b\bar{b}H$ production with $H \rightarrow \mu^+\mu^-$

In the process $h/A/Hb\bar{b} \rightarrow \mu^+\mu^-b\bar{b}$ the MSSM Higgs bosons are produced in association with a $b\bar{b}$ quark pair, and decay in a $\mu^+\mu^-$ lepton pair. The muon pair final state, although not favorite in terms of Higgs decay branching ratio, allows a full reconstruction of the Higgs final state, and can be used to directly measure the particle mass and width. Varying the A Higgs boson mass, it is possible to distinguish three different regimes: the decoupling regime, the low M_A regime and the intense coupling regime. The main background contribution comes from the Drell-Yan, the top pair production and $Zb\bar{b}$ events. Any other background process, with at least two muons in the final state, was found to be negligible.

11.2.2.1 Online Selection

For the online selection, the event must be triggered as a two muon event both by the L1 and the HLT.

11.2.2.2 Offline Selection

Muon identification

The signal is characterized by two well reconstructed, isolated muons. Therefore the event is accepted if there are at least two muons, with opposite charge, both of them satisfying the

following conditions:

- the number of hits in the Muon detector $N_{hits} > 8$;
- the transverse momentum $p_T > 20 \text{ GeV}/c$;
- a cone of $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.35$ is defined around the reconstructed muon track. Then the variable E_{iso} is obtained as the sum of the energies measured by all the detectors inside this cone. The muon is defined *isolated* if $E_{iso} < 10 \text{ GeV}$.

Top rejection

The rejection of $t\bar{t}$ events is based on two selection cuts and exploits the presence of the neutrino in the top decay chain and of two well reconstructed energetic jets.

The event is accepted if the following conditions are satisfied:

- the missing transverse energy is less than 40 GeV;
- the jets, reconstructed with the Iterative Cone Algorithm [304], must have the transverse energy less than 45 GeV and $|\eta| < 5.0$.

B-tagging

The presence of the b jets is the feature which is exploited to suppress Drell-Yan background, which otherwise will be dominant, especially for di-muon invariant masses below 200 GeV/c².

The b quarks in signal events are mainly produced in the forward region, with lower p_T with respect to b quarks coming from $t\bar{t}$ background.

Two different strategies, based on two distinct cuts, have been developed for the b-tagging:

1. The event must contain at least one b-tagged jet using the *CombinedBTagging* algorithm [598]. This algorithm has been designed to tag mainly central b jets of high transverse energy: thus it is not optimized for the b jets of the signal (in the following this cut will be defined *hard b-tag*).
2. The tracks in the event are classified as *good tracks* if they satisfy:
 - at least 6 hits in the Tracker detector of which at least one belonging to the pixel detectors;
 - the transverse momentum $p_T > 2.4 \text{ GeV}/c$;
 - the pseudorapidity $|\eta| < 2.4$;
 - the transverse impact parameter $IP < 0.5 \text{ cm}$;
 - a fit to the track model with $X^2/ndf < 5$.

The event must contain at least two good tracks with transverse impact parameter (IP) in the range $0.01 < IP < 0.1 \text{ cm}$ (only one track if $0.02 < IP < 0.075 \text{ cm}$).

The first strategy consists on applying the cut 1) only. The second strategy is the logical OR between cuts 1) and 2) (in the following this cut will be defined *soft b-tag*).

Results have been calculated for both selections and the one with the best significance has been considered.

11.2.2.3 Fitting procedure

Figure 11.9 shows the distribution of reconstructed di-muon invariant mass, including the main background and, as an example, the signal with $M_A = 150 \text{ GeV}/c^2$ and $\tan\beta = 40$. The plot has been obtained assuming an integrated luminosity of 30 fb^{-1} and the hard b-tag cut. The signal is visible as a peak over a background that exponentially decreases with increasing $M_{\mu\mu}$.




Figure 11.9: Fitting procedure applied to the di-muon reconstruction mass for the main background and for the signal sample with $M_A = 150 \text{ GeV}/c^2$ and $\tan\beta = 40$.

The background is estimated by fitting the di-muon mass distribution in the off-peak regions, where the signal is not present. To identify this region, the *TSpectrum* class in root is used: this class allows to find a signal peak over a background distribution.

The function used in this analysis to parametrize the background has three free parameters:

$$f_B(M_{\mu\mu}; P_0, P_1, P_2) = P_0 \times \frac{\Gamma_Z}{2\pi \left((M_{\mu\mu} - M_Z)^2 + \left(\frac{\Gamma_Z}{2}\right)^2 \right)} + P_1 + P_2 \times M_{\mu\mu} \quad (11.2)$$

After that the background parametrization function is determined by fitting the background in the off-peak region, a binned likelihood fit method, with three free parameters, is applied over the whole $M_{\mu\mu}$ range using the function:

$$f_{tot}(M_{\mu\mu}; M_A, \sigma_{\mu\mu}, \Gamma_A, N_S) = (N_{TOT} - N_S) \times pdf_B(M_{\mu\mu}) + N_S \times V(M_{\mu\mu}; M_A, \sigma_{\mu\mu}, \Gamma_A) \quad (11.3)$$

where $pdf_B(M_{\mu\mu})$ is the probability distribution function for the background with fixed parameters, and the second is the Voigt function, i.e. the convolution function between a Gaussian and a Breit-Wigner. The three free parameters are the number of signal events (N_S), the MSSM Higgs boson mass (M_A) and width (Γ_A). The quantity $\sigma_{\mu\mu}$ is the CMS resolution for $M_{\mu\mu}$ and its value is found from real data fitting the Z peak in the Drell-Yan distribution.

To estimate the significance for the potential discovery of the Higgs boson, the likelihood fit is performed in the *signal + background* hypothesis (L_{S+B}) and in the *background* hypothesis (L_B). The significance is defined [96] as:

$$S_L = \sqrt{2(\ln L_{S+B} - \ln L_B)} \quad (11.4)$$

11.2.2.4 Results

Table 11.4 summarizes the selection cut efficiency for background and signal. The first set of cuts, down to the Jet Veto cut, is always applied. After that two different b-tags are considered.

Table 11.4: Effect of the selection cuts on the background and signal cross section (all values in pb). Efficiency w.r.t. previous cut in % in brackets. The *no cut* value for the top pair background refers to the inclusive $t\bar{t}$ production. Other background sources not listed turned out to be negligible.

	top pairs	Drell-Yan $M_{\mu\mu} > 115 \text{ GeV}/c^2$	Zbb $M_{\mu\mu} > 100 \text{ GeV}/c^2$	signal (130,30)
No cuts	840	27.8	1.05	0.309
preselection cut	20.9 (2.5)	13.0 (46.8)	0.778 (74.1)	0.245 (79.2)
L1	19.8 (94.7)	11.9 (91.3)	0.720 (92.5)	0.226 (92.2)
HLT	17.1 (86.1)	11.8 (99.3)	0.712 (98.9)	0.223 (98.7)
Muon Id	5.23 (30.7)	10.4 (87.9)	0.569 (79.9)	0.183 (81.8)
Missing Et	1.20 (23.)	9.51 (91.7)	0.503 (88.4)	0.163 (89.2)
Jet Veto	0.317 (26.4)	8.37 (88.1)	0.418 (83.1)	0.138 (84.5)
Soft b-tag	0.238 (75.2)	0.916 (10.9)	0.146 (35.0)	0.0424 (30.9)
Events after 30 fb^{-1}	7140	27480	4380	1272
Hard b-tag	0.173 (54.7)	0.0697 (0.83)	0.0616 (14.7)	0.0154 (11.2)
Events after 30 fb^{-1}	5190	2091	1848	462

There will be no use of Montecarlo and theoretical input to compute the significance of the observed signal. Some systematic effects may be introduced by the experimental technique to fit the background. To estimate such effects, the fitting procedure has been repeated fixing one of the parameters to the measured value increased by its error. Thus a new significance is obtained and the difference with the previous one is taken as the corresponding systematic uncertainty (ΔS). From this the final significance is obtained as $S - \Delta S$, differing by from 5 to 40% with respect to the one calculated without systematic uncertainty.

Decoupling regime

Table 11.5 shows the significance as a function of $\tan \beta$, for an Higgs mass of 150 and 200 GeV/c^2 : in general, where the fitting procedure works properly, the significance is greater than five. Best results are obtained for low values of M_A (as the cross section increases with decreasing Higgs mass) and for high values of $\tan \beta$ (the cross section is proportional to $\tan^2 \beta$).

Low M_A regime

In the low M_A regime the background is particularly large due to the presence of the Z^0 peak. Therefore the signal peak is hidden for the integrated luminosity considered in this study. Better results could be obtained in the LHC high luminosity phase.

Table 11.5: Significance for the decoupling regimes.

Luminosity (fb^{-1})	$\tan \beta = 15$	$\tan \beta = 30$	$\tan \beta = 40$	$\tan \beta = 50$
$M_A = 150 \text{ GeV}/c^2$ - soft b-tag				
10	-	-	6.5	7.9
20	-	7.2	10.3	12.1
30	-	9.7	13.0	15.4
$M_A = 150 \text{ GeV}/c^2$ - hard b-tag				
10	-	3.8	5.7	6.7
20	-	6.2	7.3	9.8
30	-	8.8	9.8	13.1
$M_A = 200 \text{ GeV}/c^2$ - soft b-tag				
20	-	-	3.1	5.2
30	-	-	4.7	5.7

Intensive coupling regime

The intensive coupling regime is particularly interesting because all the three neutral Higgs bosons contribute to the signal peak. Each boson has rather small intrinsic width (less than $3 \text{ GeV}/c^2$ for $\tan \beta = 30$) which, at the generation level, is smaller than the mass difference. However, once that the mass resolution is taken into account, it becomes impossible to separate the three peaks.

The significance, on the other hand, is quite good despite the vicinity of the Z^0 peak, because the signal cross section is large: thus the discovery can be already done with an integrated luminosity of 20 fb^{-1} . Table 11.6 summarizes the significance obtained for the three signal samples as a function of the integrated luminosity.

Table 11.6: Significance for the intensive coupling regime as a function of the integrated luminosity, for different M_A values.

Luminosity (fb^{-1})	$M_A = 125 \text{ GeV}/c^2$	$M_A = 130 \text{ GeV}/c^2$	$M_A = 135 \text{ GeV}/c^2$
20	7.1	5.4	5.1
30	9.8	7.6	7.1

Figure 11.10 shows the discovery contour plot in the plane $(M_A, \tan \beta)$ obtained with this analysis.

11.2.2.5 $\tan \beta$ measurement

The peculiar feature of the two muon channel is the full reconstruction of the Higgs final state, which allows a direct measurement of the quantity Γ_H . Therefore the measurement of $\tan \beta$ exploiting its relation with Γ_H will be discussed. As shown in section 11.2.2, the measurement can be done only if A is in the range $150 < M_A < 200 \text{ GeV}/c^2$.

Figure 11.11 compares the intrinsic Higgs boson width with the measured one for $M_A = 150 \text{ GeV}/c^2$. Fitting the mass distribution with a Voigt function, the contribution to the Higgs peak from the muon invariant mass resolution is subtracted. However, another effect must be taken in account: the degeneration of the two neutral Higgs bosons, A and H, is not perfect. $M_A - M_H$ is plotted as a function of $\tan \beta$ (blue circles): this effect is particularly evident for $M_A = 150 \text{ GeV}/c^2$ and for low $\tan \beta$ (where the mass difference is greater than the intrinsic width). Thus the measured effective width is not the intrinsic one (magenta tri-




Figure 11.10: Discovery contour plot for the MSSM neutral Higgs in two muon analysis: signal points inside the grey area have a significance > 5 with an integrated luminosity of 30 fb^{-1} . The structure of the contour plot near the minimum is due to the features of the signal in the intense coupling regime. The dashed line refers to the analysis without systematic uncertainties. It must be pointed out that the contour plot does not correspond to a significance equal to 5 for any values of M_A . In particular, for $M_A < 180 \text{ GeV}/c^2$, the contour plot is actually determined by the possibility to perform a successful fit to the data, due to the low statistics. For such mass values, the contour plot corresponds to a significance which is actually slightly larger than 5. Only for $M_A > 180 \text{ GeV}/c^2$ the contour plot also corresponds to a significance equal to 5. This explains why the effect of the inclusion of the systematic uncertainty is visible only in this mass range. For $M_A < 180 \text{ GeV}/c^2$, the fit fails even if systematic uncertainties are not included in the analysis, and the contour plot does not change.

angles), but it is the sum of the intrinsic width and of Higgs mass difference (green triangles): $\Gamma_A + (M_H - M_A)$.

Figure 11.12 shows the uncertainty on the $\tan \beta$ measurement that can be obtained if the MSSM relation between the Higgs boson width and $\tan \beta$ is exploited. To calculate these values, a theoretical uncertainty of 15% [546] is included.

11.2.3 Associated $b\bar{b}H$ production with $H \rightarrow b\bar{b}$

In the high $\tan \beta$ region ($\tan \beta > 10$), the dominant channel for observing the heavier scalar and the pseudoscalar Higgs bosons is the 4 b-jet final state signature where two b-jets come from the Higgs decay, and the other b-jets are associated to the Higgs production. Nevertheless, the challenge of this topology is driven by the huge background expected for the final signature of 2 soft b-jets + 2 hard b-jets, the latter coming from the Higgs decay.

In this analysis **NAME:** [cmsnote for AN2006/023] a study of the observability of this channel using the fast simulation framework of CMS [287] is performed. Signal is also studied with full GEANT4 [599] CMS detector simulation [14] which allows to validate the fast simulation samples.

This channel can be considered as a cross-check for the discovery once it is known which Higgs boson masses (observed for instance in $b\bar{b}H/A \rightarrow b\bar{b}\tau^+\tau^-$ channel) must be looked at. It constitutes the only channel to measure the $b\bar{b}$ decay for the heavier A/H Higgs bosons. It can be also used in combination with the $\tau\tau$ mode to calculate the relative $A(H) \rightarrow b\bar{b}/A(H) \rightarrow \tau\tau$ branching ratio.




Figure 11.11: Compare between the expected effective Higgs boson width and the measured one as a function of $\tan\beta$ for $M_A = 150 \text{ GeV}$




Figure 11.12: Uncertainty on the $\tan\beta$ measurement obtained from the Higgs boson width measurement with an integrated luminosity of 30 fb^{-1} .

11.2.3.0.1 Event Generation

Signal events $b\bar{b}H, H \rightarrow b\bar{b}$ were produced using PYTHIA ?? for 4 values of m_A : 200, 500, 600 and $800 \text{ GeV}/c^2$. The signal cross sections and branching ratios were calculated with FeynHiggs [136]. The $\tan\beta$ value chosen for generation was 50. In the considered $(m_A, \tan\beta)$ region, A and H Higgs bosons have almost the same mass and the two masses can not be distinguished.

Among the SM processes, backgrounds for this channel come mainly from QCD multi-jet production; events with four real b-jets can be considered almost an irreducible background. Backgrounds have been generated with PYTHIA QCD jet production processes, additional jets were produced from gluon splitting and from Initial and Final State Radiation.

The generation of backgrounds has been weighted in order to get a similar statistics in the whole relevant p_t range, the transverse momentum in the centre of frame of the main process. A minimum $p_t > 50 \text{ GeV}/c$ cut is required by PYTHIA to avoid divergence at low p_t . Production was split in p_t bins of 50 GeV/c from 50 to 1000 GeV/c .

11.2.3.0.2 Event preselection

About 800 million Monte-Carlo events were generated and passed to a pre-selection, requiring a final state containing at least three heavy (b or c) quarks (in any jet), four PYCELL ?? jets reconstructed with the standard iterative cone algorithm using an opening angle of $\Delta R = 0.5$ in the $|\eta| < 4.5$ region, $p_{T_2} > 50 \text{ GeV}/c$ and $p_{T_4} > 10 \text{ GeV}/c$.

This pre-selection greatly reduces the number of events that need to pass through the detector simulation, conserving a proper background shape estimation in the mass region of interest, while the signal is not much affected since it contains always 4 heavy quarks. After pre-selection, around 30 million events were passed to the detector simulation. The QQ + light quark background (with Q=b,c) was found to be less than 1% of total background at final level of selection cuts.

11.2.3.0.3 Online Selection

This channel can be triggered at L1 by the jet trigger, mainly by the single jet trigger, which has no effect on the significance for $m_A > 400 \text{ GeV}/c^2$. Nevertheless, L1 thresholds are superseded by HLT requirements.

At HLT, inclusive b-jets trigger [600] has been considered, which is much more critical due to the L2 E_T thresholds on the leading jets. For masses below $\sim 600 \text{ GeV}/c^2$ those thresholds cut off the mass distribution peak.

Due to the high content in b-jets expected for the signal analyzed, lower E_T thresholds and two b-tagged jets among the two highest- E_T jets are expected to improve the signal acceptance and will allow to improve the significance of this channel.

11.2.3.0.4 Off-line Selection

Analysis has been performed in fast simulated signal and background samples where pile-up process was not taken into account, once it was checked with full simulation on signal events that its effect was not present after requiring $p_T > 30 \text{ GeV}/c$ jets.

The jets in the final state are reconstructed with the standard iterative cone algorithm [304] using an opening angle of $\Delta R = 0.5$. Seeds for the cones were selected from all EcalPlusHcalUETowers (projective Towers that sum the energy deposits in 5x5 crystals in the electromagnetic and one cell in the hadronic calorimeter, using a sliding energy cut on tower), which minimizes the fake jet rate, and MC calibration method [601] was applied. A first simple selection was applied on the event requiring at least four jets in calorimeter η acceptance, followed by kinematical cuts on 1st, 2nd and 4th calibrated jets (p_T ordered) depending upon the m_A point considered according to table 11.7, independently of the $\tan \beta$ value. The cut on the 4th p_T jet is motivated by reliability of the analysis simulation w.r.t. PU effect and has no signal selection motivation.

Table 11.7: Off-line p_T (in GeV/ c) selection cuts for different Higgs mass values considered.

m_A	200	500	600	800
p_{T_1}	90	200	220	260
p_{T_2}	80	180	200	240
p_{T_4}	30			

Subsequently pseudo-rapidities in the range of the tracker, $|\eta| < 2.4$, are required. Combined b-tag as described in [598] has been used. At least three b-tagged jets (discriminant variable > 2), among the 4 highest p_T jets, are requested in the analysis; two of them must be the two highest p_T jets. It would also have been possible to be less restrictive and accept events where only 3 of the 4 jets are in the tracker acceptance and as b-quarks, with the other outside the tracker acceptance, but this option will not be considered in this analysis.

Finally, a variable of shape in the phase space is used. The centrality, defined as:

$$\text{Centrality} = \frac{\sum E_T}{\sqrt{(\sum E)^2 + (\sum P_z)^2}} \quad (11.5)$$

using the four p_T highest jets in the event, is used to discriminate between signal and background, given its independency from signal mass. The analysis uses the discrimination power of this variable to reject background events with values lower than 0.7.


Table 11.8 summarizes the selection cut efficiencies for background and signal.

Table 11.8: Signal selection cumulative acceptances for $m_A = 600 \text{ GeV}/c^2$ and background cumulative efficiencies. $\tan \beta$ value chosen is 50. The samples used to calculate numbers given in this table are statistically independent from the ones used to optimize the cuts.

Selection	Signal efficiency	Background efficiency	S/B (full mass range)
None	1	1	1.85E-07
Preselection	5.14E-01	5.94E-03	1.60E-05
At least 4 jets	5.01E-01	5.85E-03	1.58E-05
$P_T^{(jet1)}$	3.10E-01	1.57E-04	3.66E-04
$P_T^{(jet2)}$	1.86E-01	4.76E-05	7.21E-04
$P_T^{(jet4)}$	1.02E-01	3.24E-05	5.82E-04
Jets in $ \eta \leq 2.4$	8.25E-02	2.26E-05	6.73E-04
b-tagging of 1 jet	3.61E-02	2.44E-06	2.73E-03
b-tagging of 2 jets	1.69E-02	2.81E-07	1.11E-02
b-tagging of 3 jets	8.57E-03	5.62E-08	2.82E-02
centrality > 0.7	7.05E-03	3.69E-08	3.52E-02

11.2.3.0.5 Results

The criterion for the presence of signal is based on the distribution of the reconstructed Higgs boson mass, considering as mass estimator the invariant mass distribution of the two

leading p_T jets. The signal significance, S/\sqrt{B} is calculated in the mass window which maximizes this ratio. Figure 11.13 shows the reconstructed Higgs boson mass distribution  signal and background after the off-line and on-line selection as expected in 60 fb^{-1} .




Figure 11.13: Higgs boson mass distributions after off-line and on-line selections for signal $m_A = 600 \text{ GeV}/c^2$ (black in foreground), background (solid line) and signal plus background (dashed line) in 60 fb^{-1} .

Table 11.9: Significance S/\sqrt{B} in optimized mass window after selection cuts in the HLT Trigger and no-Trigger scenario for the different Higgs mass values considered at $\tan \beta = 50$. Last line shows the low limit of $\tan \beta$ where the 5σ discovery is possible with 60 fb^{-1} in the absence of systematics.

m_A	200	500	600	800
Only L1 Trigger, No HLT trigger	30.9	10.4	7.7	2.3
HLT inclusive b-jet Trigger	2.9	6.4	5.6	3.4
Low limit $\tan \beta$ reachable with 5σ	71	44	47	62

The signal significances in the optimized mass window after all the cuts applied excluding and including the HLT in the analysis chain, can be found on table 11.9. HLT decreases the significance up to a factor 10 for low masses ($m_A = 200 \text{ GeV}/c^2$). For higher masses, $\sim m_A = 500 \text{ GeV}/c^2$, this factor is reduced to less than 2. $\tan \beta$ values greater than 44 for $m_A = 500 \text{ GeV}/c^2$ can be seen with 5σ significance in the absence of systematics.

11.2.3.0.6 Background estimation from data

Given the low S/B ratio and the similarities of the signal and background distributions, a careful evaluation of the background has to be performed. The best source of background events will come from real data samples, when available, as it is being done at the Tevatron experiments [602]. The QCD multi-jet background will be determined from data by

normalizing distributions outside of the signal region, once the mass of the Higgs is known from other channels for example. Data will be also used to extract the background shape with possibly the help of Monte-Carlo. The overall background normalization will be therefore determined by fitting the leading two jets invariant mass spectrum in triply b-tagged events outside of the hypothesized signal region to the estimated shape for triply b-tagged background.




Figure 11.14: Two-sigma significance contour including some systematic error in the background calculation from Monte-Carlo shape. Statistics are for 60 fb^{-1} .

Figure 11.14 shows the effect of different systematic error values in the expected number of background events from the Monte-Carlo shape fits. The significance used for this plot is defined by: $s = \frac{S}{\sqrt{B + (\varepsilon B)^2}}$, with S the number of signal events in the mass window, B the number of background events in the same window and ε the relative background count systematics. Due to the huge number of background events expected, for errors greater than $\sim 2.0\%$, the systematic uncertainty is larger than the number of the expected signal events even for the maximal cross section times branching ratios point.

The potential of this channel is limited by the low signal-to-background ratio and the similarity of the signal and background distribution shapes. So far it is not known how to control the background uncertainty in order to be able to observe an MSSM Higgs in the four-b final state.

11.2.4 Charged Higgs boson of $M_H < m_t$ in $t\bar{t} \rightarrow H^\pm W^\mp b\bar{b}$ production with $H^\pm \rightarrow \tau^\pm \nu$, $\tau \rightarrow \text{hadrons } \nu$ and $W^\mp \rightarrow \ell^\mp \nu$

11.2.4.1 Event Generation and Cross Sections of Signal and Background Events

The charged Higgs boson of the MSSM can be produced in top quark decays, $t \rightarrow H^\pm b$, if $m_{H^\pm} < m_t - m_b$. The branching ratio of top decay to charged Higgs depends on both m_{H^\pm} and $\tan \beta$ as shown in Fig. 11.15. The corresponding top decay to $W^\pm b$ decreases with

increasing $\tan\beta$ so as to keep the sum of branching ratios almost at unity. While the top decay to H^\pm or W^\pm depends on $\tan\beta$, the light charged Higgs decay to $\tau\nu$ is almost independent of $\tan\beta$ (for $\tan\beta > 10$) and is $\sim 98\%$ for all $\tan\beta > 10$ and $m_{H^\pm} < m_t$ as shown in Fig. 11.16.




Figure 11.15: Branching ratio of top decay to H^\pm vs $\tan\beta$




Figure 11.16: Branching ratio of charged Higgs decay to different final states for $\tan\beta = 20$.

There are two different final states for $t\bar{t} \rightarrow H^\pm W^\mp b\bar{b}$ events depending on W^\pm decay to leptons or jets. In this section the leptonic decay of W^\pm boson is chosen and signal events are triggered by the single lepton trigger (e, μ). The τ lepton is forced to decay to hadrons. Table 11.10 shows the cross section times branching ratio of $t\bar{t} \rightarrow H^\pm W^\mp b\bar{b} \rightarrow \tau\nu_\tau \ell\nu_\ell b\bar{b}, \tau \rightarrow \text{hadrons}$ for $\tan\beta = 20$. In

Table 11.10: Cross section times branching ratio of $t\bar{t} \rightarrow H^\pm W^\mp b\bar{b} \rightarrow \tau\nu_\tau \ell\nu_\ell b\bar{b}, \tau \rightarrow \text{hadrons}$ for $\tan\beta = 20$

$m_{H^\pm} (\text{GeV}/c^2)$	140	150	160	170
Cross section [pb]	10.70	5.06	1.83	0.16

this analysis for $m_{H^\pm} = 170 \text{ GeV}/c^2$ both $t\bar{t} + g\bar{b}$ and $g\bar{g} \rightarrow t\bar{b}H^\pm$ were used for comparison. The NLO cross section times branching ratio of signal events with $m_{H^\pm} \simeq m_t$ is listed in Table 11.11.

The background channels consist of $t\bar{t}$ events with at least a single lepton (e or μ) and τ -jets or jets which could fake τ -jets, $W^\pm + 3 \text{ jet}$ events and also single top events which have a small contribution. The cross section of main background channels are shown in Table 11.12.

Table 11.11: Cross section times branching ratio of signal events for $m_{H^\pm} \simeq m_t$ according to NLO calculations in [582] for $\tan\beta = 20$.

Channel	$g\bar{b} \rightarrow tH^\pm \rightarrow \ell\nu_\ell b\tau\nu_\tau$ ($\tau \rightarrow \text{hadrons}$) $m_{H^\pm} = 170 \text{ GeV}/c^2$	$g\bar{g} \rightarrow t\bar{b}H^\pm \rightarrow \ell\nu_\ell b\bar{b}\tau\nu_\tau$ ($\tau \rightarrow \text{hadrons}$) $m_{H^\pm} = 170 \text{ GeV}/c^2$
Cross section [pb]	0.14	0.30

Table 11.12: Cross section times branching ratio of background events

Channel	$t\bar{t} \rightarrow W^+W^-bb$ $\rightarrow \ell\nu_\ell\tau\nu_\tau b\bar{b}$ ($\tau \rightarrow$ hadrons)	$t\bar{t} \rightarrow W^+W^-b\bar{b}$ $\rightarrow \ell\nu_\ell\ell'\nu_{\ell'}b\bar{b}$ $\ell, \ell' = e$ or μ	$t\bar{t} \rightarrow W^+W^-b\bar{b}$ $\rightarrow \ell\nu_\ell jjb\bar{b}$	$W^\pm + 3$ jets $W^\pm \rightarrow e$ or μ
Cross section [pb]	25.8	39.7	245.6	840

11.2.4.2 Online Event Selection and Offline Reconstruction

Event Trigger

First events are triggered by the Level 1 and HLT package using the single lepton trigger.

Jet Reconstruction, Correction

Then more than or equal to 3 jets are required in the signal and background events to suppress $W^\pm + n$ jets with $n < 3$. The jet reconstruction is performed using Iterative Cone algorithm with GammaJet correction applied. A jet is accepted if $E_T > 40$ GeV after GammaJet correction is applied.

b tagging

Only 1 b-jet is required in this analysis. This requirement will be justified later in the text when the selection efficiencies for signal and background events are presented.

Offline τ tagging

Since events are triggered by the existing lepton (electron or μ), τ jets are identified using an offline τ tagging algorithm which uses Level 1 τ objects for jet reconstruction. The first jet satisfying the conditions of $E_T > 20$ GeV and Hottest Hcal Tower $E_T > 2$ GeV is used as a τ candidate. A matching cone with $\Delta R = 0.1$, isolation cone with $\Delta R = 0.4$ and a signal cone with $\Delta R = 0.07$ is defined for checking isolation requirements in the tracker. The ECal isolation requirement is defined as

$$P_{\text{isol.}} = \sum_{\text{crystals}, \Delta R_{\text{crystal}, \tau\text{-jet}} < 0.4} E_{T\text{crystal}} - \sum_{\text{crystals}, \Delta R_{\text{crystal}, \tau\text{-jet}} < 0.13} E_{T\text{crystal}} < 5.6 \text{ GeV} \quad (11.6)$$

When the tracker and ECAL isolation cuts are applied, the τ -jet E_T is required to be more than 40 GeV and the leading track of τ jet is required to carry at least 80% of the τ energy and finally the charges of the τ lepton and the lepton in the event should satisfy the requirement $Q(\ell) + Q(\tau) = 0$.

Missing E_T

The missing E_T is reconstructed with the energy corrections applied to jets (type I correction) and a cut on the reconstructed missing E_T ($E_T^{\text{miss}} > 70$ GeV) is applied as a rejection tool against background events especially $W^\pm + 3$ jets.

11.2.4.3 Selection Efficiencies

Tables 11.13, 11.14, 11.15 show the selection cuts and their efficiencies for signal and background samples. Other background events such as Wbb , Zbb with $W \rightarrow \ell\nu$ ($\ell = e, \mu$) and $Z \rightarrow ee$, or $\tau\tau$ turned out to be negligible. Single top background contribution is also small but was considered in the analysis for signal significance calculations.

Table 11.13: List of selection cuts and their efficiencies for signal events with $m_{H^\pm} < 170 \text{ GeV}/c^2$ for $\tan\beta = 20$. Numbers in each row show the remaining cross section after applying the corresponding cut. Numbers in parentheses are relative efficiencies in percent.

	$t\bar{t} \rightarrow H^\pm W^\mp b\bar{b}$ $\rightarrow \ell\nu_\ell\tau\nu_\tau b\bar{b}$ $m_{H^\pm} = 140 \text{ GeV}/c^2$	$t\bar{t} \rightarrow H^\pm W^\mp b\bar{b}$ $\rightarrow \ell\nu_\ell\tau\nu_\tau b\bar{b}$ $m_{H^\pm} = 150 \text{ GeV}/c^2$	$t\bar{t} \rightarrow H^\pm W^\mp b\bar{b}$ $\rightarrow \ell\nu_\ell\tau\nu_\tau b\bar{b}$ $m_{H^\pm} = 160 \text{ GeV}/c^2$
$\sigma \times \text{BR}[\text{fb}]$	10.7×10^3	5060.	1830.
L1 + HLT	5170.5(48.3)	2456.3(48.5)	888.9(48.6)
≥ 3 jets	1889.7(36.5)	795.0(32.4)	264.3(29.7)
≥ 1 b jet	1103.5(58.4)	427.4(53.8)	131.4(49.7)
< 2 b jets	883.0(80.0)	358.7(83.9)	119.2(90.7)
Having L1 τ	878.4(99.5)	357.4(99.6)	119.0(99.8)
τ -jet reconstruction	875.0(99.6)	356.5(99.7)	118.8(99.8)
Hottest HCAL tower $E_T > 2. \text{ GeV}$	778.0(88.9)	316.1(88.6)	105.9(89.1)
Tracker isolation	378.2(48.6)	163.5(51.7)	52.7(49.8)
Ecal isolation	292.9(77.4)	134.2(82.1)	43.1(81.8)
$\tau E_T > 40. \text{ GeV}$	244.3(83.4)	113.0(84.2)	36.5(84.7)
Pleading track/ $E_\tau > 0.8$	102.3(41.9)	50.7(44.8)	16.8(45.9)
$Q(\ell) + Q(\tau) = 0$	88.0(86.0)	42.4(83.6)	14.6(87.0)
$\text{MET} > 70 \text{ GeV}$	51.0(58.0)	25.4(59.9)	9.2(63.3)
Expected Number of events after 10 fb^{-1}	510	254	92

Table 11.14: List of selection cuts and their efficiencies for signal events with $m_{H^\pm} = 170 \text{ GeV}/c^2$ for $\tan\beta = 20$. Numbers in each row show the remaining cross section after applying the corresponding cut. Numbers in parentheses are relative efficiencies in percent.

	$t\bar{t} \rightarrow H^\pm W^\mp b\bar{b}$ $\rightarrow \ell\nu_\ell\tau\nu_\tau b\bar{b}$ $m_{H^\pm} = 170 \text{ GeV}/c^2$	$g b \rightarrow t H^\pm$ $\rightarrow \ell\nu_\ell\tau\nu_\tau b$ $m_{H^\pm} = 170 \text{ GeV}/c^2$	$g g \rightarrow t b H^\pm$ $\rightarrow \ell\nu_\ell\tau\nu_\tau b\bar{b}$ $m_{H^\pm} = 170 \text{ GeV}/c^2$
$\sigma \times \text{BR}[\text{fb}]$	157.	140.	297.
L1 + HLT	78.0(49.7)	70.5(50.4)	145.4(48.9)
≥ 3 jets	23.2(29.7)	21.7(30.7)	55.3(38.0)
≥ 1 b jet	11.5(49.4)	11.7(54.1)	31.9(57.7)
< 2 b jets	10.9(94.8)	10.0(85.5)	25.8(80.9)
Having L1 τ	10.8(99.8)	10.0(99.6)	25.7(99.4)
τ -jet reconstruction	10.8(99.9)	10.0(99.9)	25.5(99.1)
Hottest HCAL tower $E_T > 2. \text{ GeV}$	9.6(88.4)	8.9(88.8)	22.6(88.9)
Tracker isolation	4.9(51.3)	5.1(57.2)	11.4(50.5)
Ecal isolation	4.2(84.9)	4.3(84.5)	9.6(84.4)
$\tau E_T > 40. \text{ GeV}$	3.8(90.9)	3.9(90.6)	8.6(89.2)
Pleading track/ $E_\tau > 0.8$	1.6(41.7)	1.8(45.9)	3.4(39.6)
$Q(\ell) + Q(\tau) = 0$	1.3(84.4)	1.6(87.2)	2.8(82.6)
$\text{MET} > 70 \text{ GeV}$	0.8(61.7)	1.0(65.2)	1.6(55.3)
Expected Number of events after 10 fb^{-1}	8	10	16

Table 11.15: List of selection cuts and their efficiencies for background events. Numbers in each row show the remaining cross section after applying the corresponding cut. Numbers in parentheses are relative efficiencies in percent.

	$t\bar{t} \rightarrow W^+W^-bb \rightarrow \ell\nu_\ell\tau\nu_\tau b\bar{b}$	$t\bar{t} \rightarrow W^+W^-bb \rightarrow \ell\nu_\ell\ell'\nu_{\ell'}b\bar{b}$	$t\bar{t} \rightarrow W^+W^-bb \rightarrow \ell\nu_\ell jjb\bar{b}$	$W^\pm + 3 \text{ jets}$ $W^\pm \rightarrow \ell\nu_\ell$
$\sigma \times \text{BR}[\text{fb}]$	25.8×10^3	39.8×10^3	245.6×10^3	$840. \times 10^3$
L1 + HLT	12101.2(46.9)	28429.1(71.4)	99506.6(40.5)	287280(34.2)
≥ 3 jets	5105.2(42.2)	11306.6(39.8)	66038.6(66.4)	114050(39.7)
≥ 1 b jet	3428.3(67.1)	7622.0(67.4)	43433.0(65.8)	24292.7(21.3)
< 2 b jets	2325.7(67.8)	5262.7(69.0)	29003.4(66.8)	21207.5(87.3)
Having L1 τ	2310.7(99.3)	5233.7(99.4)	28698.8(98.9)	20613.7(97.2)
τ -jet reconstruction	2303.6(99.7)	5224.4(99.8)	28465.0(99.2)	19438.7(94.3)
Hottest HCAL tower $E_T > 2. \text{ GeV}$	2034.1(88.3)	3850.6(73.7)	26635.1(93.6)	17125.5(88.1)
Tracker isolation	798.7(39.3)	1120.6(29.1)	6653.3(25.0)	5411.7(31.6)
Ecal isolation	545.6(68.3)	519.5(46.3)	2952.8(44.4)	2554.3(47.2)
$\tau E_T > 40. \text{ GeV}$	405.8(74.4)	341.8(65.8)	1946.8(65.9)	1312.9(51.4)
Pleading track/ $E_\tau > 0.8$	123.5(30.4)	131.9(38.6)	377.9(19.4)	224.5(17.1)
$Q(\ell) + Q(\tau) = 0$	95.7(77.5)	56.7(43.0)	78.8(20.9)	27.1(12.1)
$\text{MET} > 70 \text{ GeV}$	51.6(53.9)	29.3(51.8)	36.6(46.4)	10.7(39.3)
Expected Number of events after 10 fb^{-1}	516	293	366	107

11.2.4.4 Systematic Uncertainties

The systematic uncertainties in the signal statistical significance calculation include the experimental selection uncertainty of the background events and the theoretical cross section calculation uncertainty of the $t\bar{t}$ and single top background. All systematic uncertainties are added quadratically to the background statistical fluctuation in the significance calculation. The $t\bar{t}$ background uncertainty is taken into account as in Eq. 11.7.

$$\Delta_{\text{sys.}}^{t\bar{t}} = \Delta_{\text{lepton reconstruction}} \oplus \Delta_{\geq 3 \text{ jet selection}} \oplus \Delta_{1 \text{ b-jet tagging}} \oplus \Delta_{1 \tau \text{ tagging}} \oplus \Delta_{\text{lumi.}} \oplus \Delta_{\text{theo.}}^{t\bar{t}} \quad (11.7)$$

The $W^\pm + 3 \text{ jets}$ background is assumed to be measured from the real data. The uncertainty of the measurement is estimated by propagating the contribution of events counted in the background area to the signal area and cancelling the common selection cuts efficiencies and their uncertainties. Eq. 11.8 describes how systematic uncertainties are taken into account in $W^\pm + 3 \text{ jets}$ cross section measurement.

$$\Delta_{\text{sys.}}^{W^\pm+3 \text{ jets}} = \Delta_{\text{stat.}} \oplus \frac{\Delta N_B^{t\bar{t}}}{N_B^{W^\pm+3 \text{ jets}}} \oplus \Delta_{3 \text{ non-b-jet}} \oplus \Delta_{\text{b-jet mistagging}} \oplus \Delta_{\tau \text{ mistagging}} \quad (11.8)$$

Table 11.16 lists different sources of systematic uncertainties and their used values corresponding to 30 fb^{-1} in this analysis.

11.2.4.5 5σ Discovery Contour

Figures 11.17 and 11.18 show the light charged Higgs discovery contour in the $(m_{H^+}, \tan\beta)$ and $(m_{A^0}, \tan\beta)$ planes including the systematic uncertainties. It should be noted that this analysis is systematics dominated and there could be alternative approaches where the systematic uncertainties cancel down to a reasonable level.

Table 11.16: The values of different selection uncertainties for $t\bar{t}$ and $W^\pm + 3$ jets background events at 30 fb^{-1} .

Scale uncertainty of $t\bar{t}$ cross section	5%
PDF uncertainty of $t\bar{t}$ cross section	2.5%
b-tagging	5%
τ -tagging	4%
Lepton identification	2%
Jet energy scale	3%
Mistagging a non-b-jet as a b-jet	5%
Mistagging a jet as a τ -jet	2%
Non-b-jet identification (anti-b-tagging)	5%
Luminosity uncertainty	5%




Figure 11.17: The 5σ contour in the $(m_{H^\pm}, \tan\beta)$ plane for light charged Higgs discovery including the effect of systematic uncertainties.




Figure 11.18: The 5σ contour in the $(m_{A^0}, \tan\beta)$ plane for light charged Higgs discovery including the effect of systematic uncertainties.

11.2.5 Charged Higgs boson of $M_H > m_t$ in $gg \rightarrow t\bar{t}H^\pm$ production with $H^\pm \rightarrow \tau^\pm\nu, \tau \rightarrow \text{hadrons } \nu$ and $W^\mp \rightarrow jj$

The $H^\pm \rightarrow \tau^\pm\nu_\tau$ decay mode with fully hadronic final state of the charged Higgs bosons in the associated production with a top quark has been shown to lead to a clean and almost background-free signature at large $\tan\beta$ with several particle level [603] and fast simulation [385],[604], [605],[384] studies. The advantages of this decay mode in association of top are the large missing transverse energy from H^\pm , the possibility to disentangle the hadronic τ decay from the hadronic jets, the possibility to reconstruct the top mass to suppress the multi-jet backgrounds, and, in particular, τ helicity correlations favouring the $H^\pm \rightarrow \tau^\pm\nu_\tau$ decay over the $W^\pm \rightarrow \tau^\pm\nu_\tau$ decay. The main backgrounds are due to genuine τ 's in multi-jet events from $t\bar{t}$ with $t_1 \rightarrow b\tau\nu_\tau$, $t_2 \rightarrow bqq$, Wt with $W_1 \rightarrow \tau\nu_\tau$, $W_2 \rightarrow qq'$ and $W+3$ jets with $W \rightarrow \tau\nu_\tau$. The hadronic QCD multi-jet events can lead to a background through fake τ 's or through the uncertainty of E_T^{miss} measurement.

11.2.5.1 Helicity correlations

The polarization states for the τ^+ from $H^+ \rightarrow \tau^+\nu_\tau$ and from $W^+ \rightarrow \tau^+\nu_\tau$ are opposite due to the spin-parity properties of the decaying particle. The angular distribution of a pion from the τ^+ in the CM frame has the form $(1+P_\tau\cos\theta)$, which leads to more energetic pions in the laboratory frame for the signal ($P_\tau = 1$) than for the background ($P_\tau = -1$) [606],[603]. The $\tau^\pm \rightarrow \pi^\pm\nu_\tau$ decay channel presents 12.5% of the hadronic decay modes. Similarly, the signal pions are more energetic in the τ decays to vector mesons and subsequent decays to one charged pion in the longitudinal polarization states of the vector meson, $\tau^\pm \rightarrow \rho_L^\pm\nu_\tau \rightarrow \pi^\pm\pi^0\nu_\tau$ (26%) and $\tau^\pm \rightarrow a_{1L}^\pm\nu_\tau \rightarrow \pi^\pm\pi^0\pi^0\nu_\tau$ (7.5%). For the transverse polarization states of the vector meson the situation is opposite with more energetic pions from the background. The small contributions from K^* and K in the τ decays lead to similar effects. The helicity correlations can be expressed as a function of the τ -jet momentum fraction carried by the charged pion $R_\tau = p_\pi/p_{\tau\text{ jet}}$. As is shown in Refs.[606],[603] the $\tau^\pm \rightarrow \pi^\pm\nu_\tau$ decay leads to a δ -function at $R_\tau = 1$, the $\rho_L^\pm\nu_\tau \rightarrow \pi^\pm\pi^0\nu_\tau$ has contributions at $R_\tau \sim 1$ and $R_\tau \sim 0$, $\rho_R^\pm\nu_\tau \rightarrow \pi^\pm\pi^0\nu_\tau$ and $a_{1R}^\pm\nu_\tau \rightarrow \pi^\pm\pi^0\pi^0\nu_\tau$ have largest contributions around $R_\tau \sim 0.5$ while $a_{1L}^\pm\nu_\tau \rightarrow \pi^\pm\pi^0\pi^0\nu_\tau$ peaks at $R_\tau \sim 0$.

11.2.5.2 Event generation and simulation

The $gb \rightarrow tH^\pm$ and $gg \rightarrow tbH^\pm$ processes contribute to the production of a heavy single charged Higgs boson in association of top quarks. In the $gb \rightarrow tH^\pm$ process the b quark is considered as a massless parton of the incoming proton. Logarithmic factors of the form $\log(p_T^b/m_b)$, due to the collinear b quarks, can be resummed to give a well defined cross section. The $gg \rightarrow tbH^\pm$ process, where the bottom quarks from the incoming gluons are considered massive, is of the order α_s^2 and is part of the next-to-leading order (NLO) corrections to the leading order (LO) process $gb \rightarrow tH^\pm$. These processes lead to somewhat different dynamics of the final state objects, visible in particular as a more energetic associated b quark in the $gg \rightarrow tbH^\pm$ process [607]. Near the top threshold, $m_{H^\pm} \sim m_t$, only the exclusive process $gg \rightarrow tbH^\pm$ can lead to a correct event description. As the correct description of merging these two processes is not possible in the full simulation, signal events were generated with the $gg \rightarrow tbH^\pm$ process over the full mass range with PYTHIA [62]. The cross sections were normalized to the NLO results of Refs. [608],[582]. The mass of the charged Higgs boson and the $H^\pm \rightarrow \tau\nu_\tau$ branching fraction were calculated with FeynHiggs [136]. The $t\bar{t}$ background was generated with PYTHIA, the Wt background with TOPREX [35], the $W+3\text{jet}$ background with MadGraph [73] and the QCD multi-jet background with PYTHIA. The production cross sections for the background processes were normalized to the NLO cross sections. Preselections at the particle level, requiring at least one jet with $E_T > 80$ GeV reconstructed with PYTHIA in a cone of 0.5 and containing at least one charged hadron with $p_T > 60$ GeV/ c , were applied to the $t\bar{t}$ and Wt backgrounds. The τ decays were performed with TAUOLA [146] for the signal and backgrounds.

The analysis is based on pre-digitised event samples for the signal and backgrounds. For these samples, the response of the CMS detector has been simulated with the CMSIM [13] and OSCAR [14] packages including the pile-up corresponding to the luminosity of $2 \times 10^{33} \text{cm}^{-2} \text{s}^{-1}$. For the reconstruction the version ORCA874 of the CMS OO reconstruction package [15] was used. Tracks, jets and missing transverse energy were reconstructed with the dedicated packages of the CMS OO software. The primary vertex was reconstructed summing the transverse momenta of the associated tracks and selecting the vertex giving the largest sum.

The hadronic jets were reconstructed with an iterative cone algorithm taking a cone size of 0.5 [156]. The jet energies were corrected with factors obtained with Monte-Carlo simulation method.

11.2.5.3 Event selection

Due to an energetic τ jet from H^\pm the $gg \rightarrow tbH^\pm$, $H^\pm \rightarrow \tau^\pm\nu$, $\tau \rightarrow \text{hadrons } \nu$, $W^\mp \rightarrow jj$ events can be triggered at the Level-1 with a single τ -jet trigger [68],[268]. The E_T threshold needed to suppress the hadronic jets at Level-1 for the given bandwidth of 2 kHz is 93 GeV at low luminosity [68]. Energy corrections, derived from Monte Carlo data, were applied on the reconstructed Level-1 τ jet. At the HLT, the cut $E_T^{\text{miss}} > 67$ GeV was used in the missing transverse energy calculated summing the cells of the Electromagnetic calorimeter (ECAL) and the corresponding Hadron calorimeter (HCAL) cells (trigger towers) without energy corrections. For the single τ trigger, the tracks around the jet direction are reconstructed in the full silicon tracker, limiting, however, the number of hits to six. The track with maximum p_T was searched for in a cone of 0.1 around the jet direction. A small cone with $\Delta R = 0.065$ around the leading track was taken as the signal cone requiring this signal cone to be isolated in a larger cone of $\Delta R = 0.4$ counting the tracks with $p_T > 1$ GeV/ c . The cut $p_T > 25$ GeV/ c , was applied on the leading track. Efficiencies of the Level1 and HLT trigger are shown in Tables 11.17 and 11.18, respectively. Purity of the τ trigger for the signal events is higher than 80%. A small fraction ($< 1\%$) of the $\tau \rightarrow e\nu_\tau\nu_\tau$ decays passes the trigger for the signal events while for the backgrounds a significant fraction of the events are triggered with an electron from τ . The larger contamination in the background events is due to the boosted W 's required for the τ jet to pass the trigger threshold. Due to this boost the E_T^{miss} from $W^\pm \rightarrow \tau\nu_\tau$ is enhanced by the two neutrinos from the $\tau \rightarrow e\nu_e\nu_\tau$ decay.

In the off-line reconstruction the transverse mass from the τ jet and missing transverse energy requires a fully hadronic event, where E_T^{miss} originates mainly from the H^\pm . Other sources of E_T^{miss} in the signal events are the leptonic W decays and the semileptonic b quark decays. The events with leptonic W decays can be removed with a veto on isolated leptons. The reconstructed electrons and muons were first required to be isolated in the tracker demanding that no track with $p_T > 1$ GeV/ c was found in a cone of $\Delta R = 0.4$ around the lepton direction. The fraction of events containing at least one reconstructed muon with $p_T > 15$ GeV/ c is 24.1%. An isolated muon is found in 8.9% of the signal events and about 84% of these muons originate from $W \rightarrow \mu\nu_\mu$. The fraction of events containing at least one reconstructed electron candidate with $p_T > 15$ GeV/ c is 72.4% and an isolated electron candidate 41.7%. The hadronic contamination can be removed with matching requirements between the track and the calorimeter deposition and with calorimeter profile cuts. The ratio of the energy summed in the ECAL cells to the energy summed in the HCAL cells behind the ECAL cells (required to be < 0.1) as well the track momentum-ECAL cluster matching ($E_{\text{cluster}}/p_{\text{track}} > 0.6$), described in Ref. [147], were found to be the most powerful variables against the hadronic contamination. The fraction of events removed with a veto on the identified electrons is 7.9%, from which 93.3% are due to genuine electrons from $W \rightarrow e\nu_e$.

The missing transverse energy (E_T^{miss}) was reconstructed from the full calorimeter response summing the ECAL-plus-HCAL towers [142]. For this measurement the hadronic jets with $E_T > 20$ GeV were calibrated with correction factors obtained from a simulation of QCD photon+jet events. The τ jet was reconstructed in the calorimeter around the L1 τ -jet direction in a cone of 0.4 applying energy corrections evaluated for one- and three-prong τ decays.

Table 11.17: Cross section times branching fraction, efficiency for the selection cuts and final number of events for $m_T(\tau \text{ jet}, E_T^{\text{miss}}) > 100 \text{ GeV}/c^2$ and for $\Delta\phi(\tau \text{ jet}, E_T^{\text{miss}}) > 60^\circ$ with an integrated luminosity of 30 fb^{-1} for the signal events with $m_{H^\pm} = 170, 180, 200$ and $400 \text{ GeV}/c^2$ and $\tan\beta = 30$.

$m_{H^\pm} \text{ (GeV}/c^2)$	171.6	180.4	201.0	400.4
$\sigma(\text{NLO}) \times \text{BR} \text{ (fb)}$	2091	1904	1193	58
Level-1 trigger	1122.9 (53.7%)	1058.6 (55.6%)	694.3 (58.2%)	43.8 (75.6%)
HLT trigger	186.1 (16.6%)	197.8 (18.7%)	147.5 (21.2%)	18.6 (42.4%)
Primary vertex	184.4 (99.1%)	196.2 (99.2%)	146.3 (99.2%)	18.5 (99.2%)
Isolated lepton veto	145.3 (78.8%)	160.3 (81.7%)	120.3 (82.2%)	15.6 (85.0%)
$E_T^{\text{miss}} > 100 \text{ GeV}$	102.6 (70.6%)	107.7 (67.2%)	82.0 (68.2%)	12.6 (80.7%)
$E_T^{\tau \text{ jet}} > 100 \text{ GeV}$	51.8 (50.5%)	56.4 (52.4%)	42.7 (52.1%)	10.3 (81.8%)
$R_\tau > 0.8$	17.3 (33.4%)	17.8 (31.5%)	14.6 (34.2%)	3.52 (34.2%)
1 or 3 signal tracks	16.5 (95.3%)	17.3 (97.1%)	14.0 (95.9%)	3.41 (97.0%)
Tracker isolation	15.4 (93.2%)	16.2 (94.0%)	13.3 (94.9%)	3.20(93.7%)
ECAL isolation	14.5 (94.4%)	15.4 (95.0%)	12.7 (95.7%)	3.06 (95.8%)
$E_T^{\text{max(HCAL cell)}} > 2 \text{ GeV}$	14.0 (96.5%)	14.4 (93.3%)	12.2 (95.5%)	3.03 (98.7%)
$\text{IP}_T^{\text{leading track}} < 0.3 \text{ mm}$	13.8 (97.8%)	14.1 (98.2%)	12.0 (99.0%)	3.00 (99.3%)
$N_{\text{hits}}^{\text{leading track}} \geq 10$	13.3 (95.9%)	13.6 (96.5%)	11.4 (94.6%)	3.00 (96.5%)
$\geq 3 \text{ jets}, E_T > 20 \text{ GeV}$	9.8 (74.4%)	11.0 (80.9%)	8.8 (77.4%)	2.15 (71.9%)
$140 < m_{\text{top}} < 210 \text{ GeV}/c^2$	7.1 (72.6%)	7.4 (67.2%)	5.6 (63.7%)	1.43 (66.6%)
Discriminator > 1.5	3.1 (43.7%)	3.0 (39.9%)	2.4 (42.7%)	0.57 (40.3%)
$E_T^{\text{b jet}} > 30 \text{ GeV}$	2.9 (93.2%)	2.8 (95.2%)	2.2 (91.6%)	0.50 (88.2%)
Jet veto, $E_T^{\text{jet}} > 25 \text{ GeV}$	1.0 (35.2%)	0.97 (34.6%)	0.8 (36.4%)	0.21 (40.9%)
$E_T^{\text{Higgs}} > 50 \text{ GeV}$	0.94 (91.9%)	0.97 (100%)	0.8 (100%)	0.20 (95.1%)
$m_T > 100 \text{ GeV}/c^2$	0.73 (77.3%)	0.76 (78.4%)	0.60 (74.9%)	0.19 (94.8%)
$N^{\text{ev}}, m_T > 100 \text{ GeV}/c^2$	21.8 ± 5.3	22.8 ± 4.9	17.9 ± 3.6	5.5 ± 0.7
$\Delta\phi(\tau, E_T^{\text{miss}}) > 60^\circ$	0.30 (31.9%)	0.28 (28.5%)	0.42 (53.1%)	0.18 (93.1%)
$N^{\text{ev}}, \Delta\phi(\tau, E_T^{\text{miss}}) > 60^\circ$	9.0 ± 3.4	9.3 ± 3.1 (28.5%)	12.7 ± 3.0	5.4 ± 0.7

The offline E_T cut on the τ jet was taken to be $E_T^{\text{jet}} > 100 \text{ GeV}$, close to the Level-1 threshold of 93 GeV . The tracks were reconstructed inside the jet reconstruction cone. The leading track was searched for in a cone of $\Delta R = 0.1$ around the τ -jet direction. For an efficient isolation against the hadronic jets a small signal cone of $\Delta R = 0.04$ was selected. The fraction of signal events with $m_{H^\pm} = 200 \text{ GeV}/c^2$, where the one-prong (three-prong) τ decays lead to one (three) track(s) with $p_T > 1 \text{ GeV}/c$ in the signal cone, was found to be in 92.3% (64%). Accidental track reconstruction problems, like shared hits, can lead to a fake large- p_T tracks in the hadronic jets [268],[7]. These fake leading tracks are possible in the hadronic multi-jet events but can appear also in the $t\bar{t}$, Wt and $W + 3 \text{ jet}$ backgrounds if the E_T of the τ jet is below the trigger threshold and the event is triggered with a τ -like hadronic jet. The fake tracks can be suppressed with an upper bound in the transverse impact parameter of the leading track, $\text{IP}_T^{\text{leading track}} < 0.3 \text{ mm}$, and requiring at least 10 hits in the full tracker. The fraction of $\tau \rightarrow e\nu\nu$ events passing the full τ selection was found to be 3% for the $t\bar{t}$ background. This contamination can be efficiently suppressed requiring at least one energetic HCAL cell with $E_T > 2 \text{ GeV}$ within $\Delta R < 0.15$ around the τ -jet direction [268].

The τ helicity correlations are best exploited requiring the leading track to carry at least

Table 11.18: Cross section times branching fraction, efficiency for the selection cuts and final number of events for $m_T(\tau \text{ jet}, E_T^{\text{miss}}) > 100 \text{ GeV}/c^2$ and for $\Delta\phi(\tau \text{ jet}, E_T^{\text{miss}}) > 60^\circ$ with an integrated luminosity of 30 fb^{-1} for the $t\bar{t}$, Wt , $W^\pm + 3\text{jets}$ and QCD multi-jet backgrounds background.

	$t\bar{t}$	Wt	$W^\pm + 3\text{jets}$
$\sigma(\text{NLO}) \times \text{BR} \text{ (fb)}$	123820	9140	4.19×10^5
Preselection	6440 (5.2%)	237.6 (2.6%)	
Level-1 trigger	4730 (73.4%)	185.6 (78.1%)	1.25×10^5 (29.8%)
HLT trigger	320 (6.9%)	20.5 (11.1%)	4.19×10^3 (3.4%)
Primary vertex	319 (99.8%)	20.4 (99.7%)	4190 (100%)
Isolated lepton veto	314 (89.4%)	18.4 (89.9%)	3456 (82.5%)
$E_T^{\text{miss}} > 100 \text{ GeV}$	267.4 (85.1%)	15.9 (86.6%)	2674 (77.1%)
$E_T^{\tau \text{ jet}} > 100 \text{ GeV}$	167.4 (62.6%)	10.7 (67.2%)	1280 (69.2%)
$R_\tau > 0.8$	35.5 (21.2%)	2.53 (23.7%)	175.4 (13.7%)
1 or 3 signal tracks	31.2 (88.0%)	2.37 (93.7%)	149.3 (85.1%)
Tracker isolation	27.8 (89.1%)	2.18 (91.9%)	132.9 (89.2%)
ECAL isolation	26.1 (93.7%)	2.07 (94.9%)	125.1 (94.1%)
$E_T^{\text{max(HCAL cell)}} > 2 \text{ GeV}$	24.1 (92.4%)	1.95 (94.2%)	105.1 (84.0%)
$IP_T^{\text{leading track}} < 0.3 \text{ mm}$	21.4 (88.8%)	1.92 (98.3%)	88.4 (84.1%)
$N_{\text{hits}}^{\text{leading track}} \geq 10$	19.9 (92.9%)	1.81 (94.4%)	84.6 (95.7%)
$\geq 3 \text{ jets}, E_T > 20 \text{ GeV}$	17.3 (87.0%)	1.04 (57.6%)	67.5 (79.8%)
$140 < m_{\text{top}} < 210 \text{ GeV}/c^2$	12.2 (70.4%)	0.71 (67.7%)	26.6 (39.4%)
Discriminator > 1.5	5.81 (47.7%)	0.34 (48.1%)	1.09 (4.1%)
$E_T^{\text{b jet}} > 30 \text{ GeV}$	5.27 (90.6%)	0.30 (89.2%)	0.82 (75.1%)
Jet veto, $E_T^{\text{jet}} > 25 \text{ GeV}$	1.48 (28.1%)	0.24 (78.0%)	0.14 (17.2%)
$E_T^{\text{Higgs}} > 50 \text{ GeV}$	1.44 (97.1%)	0.23 (98.6%)	0.14 (98.3%)
$m_T(\tau \text{ jet}, E_T^{\text{miss}}) > 100 \text{ GeV}/c^2$	0.03 (2.0%)	0.003 (1.3%)	0.02 (10.3%)
Events for $m_T > 100 \text{ GeV}/c^2$	0.86 ± 0.33	0.09 ± 0.04	0.60 ± 0.60
$\Delta\phi(\tau \text{ jet}, E_T^{\text{miss}}) > 60^\circ$	0.01 (1.0%)	9.2×10^{-4} (0.4%)	0.013 (6.7%)
Events for $\Delta\phi(\tau \text{ jet}, E_T^{\text{miss}}) > 60^\circ$	0.43 ± 0.25	0.03 ± 0.02	0.39 ± 0.39

80% of the τ jet energy. The efficiencies for the $t\bar{t}$ and Wt events, shown in Tables 11.17 and 11.18, are affected with the preselection cuts and do not show the expected background suppression for $R_\tau > 0.8$. This cut suppresses the three-prong τ decays leaving as the fraction of three-prong τ decays after all selection cuts 3.1% for the signal events with $m_{H^\pm} = 200 \text{ GeV}/c^2$. To suppress hadronic jet backgrounds isolation was performed counting tracks with $p_T > 1 \text{ GeV}/c$ in the region of $0.04 < \Delta R < 0.4$. As for the single τ trigger isolation in the ECAL is not exploited at the trigger level, the ECAL cells were counted in the cone of $0.1 < \Delta R < 0.4$ around the leading track to enhance the isolation power.

Due to limited MC statistics, the trigger simulation was not used in the estimation of the hadronic multi-jet background. Events with at least one jet with $E_T > 100 \text{ GeV}$ containing a track with $p_T > 80 \text{ GeV}/c$ were saved for further study. Efficiency for this selection was found to be 5.55×10^{-3} in the interval $170 < \hat{p}_T < 380 \text{ GeV}/c$. The τ selection cuts, except the E_T^{jet} threshold, are not correlated with the E_T^{miss} cut. Therefore the selection was factorized to E_T^{miss} selection and to τ selection. The efficiency of the τ selection cuts on the preselected events was found to be 1.65%. Combined to the preselection the full τ -selection efficiency

for hadronic multi-jet events in this interval is 9.2×10^{-5} .

The $gg \rightarrow tbH^\pm$ events contain two b jets, one from the decay of the top quark and one associated b jet from the production process. The associated b quark is preferentially emitted in the forward directions and is distributed to smaller p_T values than the b quark from top decay. In about 20% of the signal events events, however, this b quark is more energetic than the b quark from top decay thus contaminating the spectrum of the identified b jet for top reconstruction. The event reconstruction was continued for the events where at least three hadronic jets with $E_T^{\text{jet}} > 20$ GeV were found. A probabilistic secondary vertex algorithm with a discriminator cut was used [148]. The fraction of events where the best tagged jet is the b from $t \rightarrow bW$ was found to be 61%. The fractions of the associated b and the q from $W \rightarrow qq$ were found to be $\sim 26\%$ and $\sim 8\%$, respectively.

The top-quark mass was reconstructed with minimization of a χ^2 distribution made from the reconstructed and nominal top and W masses by $\chi^2 = ((m_{jj} - m_W)/\sigma_W)^2 + ((m_{jjj} - m_{\text{top}})/\sigma_{\text{top}})^2$, where m_{jj} and m_{jjj} are the invariant masses of all two- and three-jet combinations in the event and σ_W and σ_{top} are the gaussian widths of the reconstructed true W and top mass distributions. The jet assigned to the top but not to the W presents the b jet from top. For a better reconstruction efficiency, in the presence of a significant contamination from the associated b quark any of the three jets assigned to the top was tagged requiring discriminator > 1.5 and $E_T > 30$ GeV. A mass resolution of $\sim 11\%$ and a mean reconstructed mass of ~ 176 GeV/ c^2 were obtained, with a fraction of about 40% of correct jet assignments. For a further suppression of $t\bar{t}$ background, the additional jets after top reconstruction were searched for within $|\eta| < 2.5$. The E_T threshold for the veto jets was set to 25 GeV. The efficiency of this method has decreased from the fast simulation results [604] mainly due to more energetic associated b jets in $gg \rightarrow btH^\pm$ with respect to the $gb \rightarrow tH^\pm$ events.

For the $t\bar{t}$, Wt and W+3jet backgrounds the configuration of large E_T^{miss} and large $E_T^{\tau\text{jet}}$ can be reached only for strongly boosted W. Therefore to suppress background from events triggered with a fake τ from a hadronic jet recoiling against the genuine τ jet, a lower bound ($E_T^H > 50$ GeV) was set on the p_T reconstructed from the τ jet and the missing transverse energy.

The large E_T thresholds lead to an almost two-body situation between the τ jet and missing transverse energy. Therefore an upper edge can be expected in the transverse mass $m_T = \sqrt{(2 \times E_T^{\tau\text{jet}} \times E_T^{\text{miss}} \times (1 - \Delta\phi(\tau\text{ jet}, E_T^{\text{miss}})))}$ at m_{H^\pm} for the signal and at m_W for the $t\bar{t}$, Wt and W+3jet backgrounds. The boost required for the $t\bar{t}$, Wt and W+3jet backgrounds to pass the E_T thresholds, leads to small opening angles $\Delta\phi(\tau\text{ jet}, E_T^{\text{miss}})$ in the transverse plane. Requiring $\Delta\phi > 60^\circ$ removes most of the background for $m_T < 100$ GeV/ c^2 . The m_T distributions for the signal and total background are shown in Figs. 11.19 and 11.20 for $m_{H^\pm} = 170$ and 400 GeV/ c^2 and $\tan\beta = 30$, without cut in $\Delta\phi(\tau\text{ jet}, E_T^{\text{miss}})$.

Tables 11.17 and 11.18 show the cross sections and efficiency for the selection cuts for the signal events with $m_{H^\pm} = 170, 180, 200$ and 400 GeV/ c^2 and $\tan\beta = 30$. The trigger efficiency and the efficiency of the primary vertex reconstruction are also shown. Table 11.18 shows the same for the $t\bar{t}$, Wt and W+3jet backgrounds. For the QCD multi-jet background the number of events where at least three jets are found after E_T^{miss} and τ selection was estimated without the τ selection cuts. At this level of selection the QCD multi-jet events can be assumed to be similar to the W+3jet events at the same selection level. Therefore the efficiency of the remaining selection cuts was taken from the W+3jet events yielding an estimate of 0.1 ± 0.1 events for $m_T(\tau\text{ jet}, E_T^{\text{miss}}) > 100$ GeV/ c^2 .




Figure 11.19: Transverse mass reconstructed from the τ jet and missing transverse energy for the $gg \rightarrow tbH^\pm, t \rightarrow bW, W^\mp \rightarrow jj$ signal (dark histogram) with $m_{H^\pm} = 170 \text{ GeV}/c^2$, $\tan\beta=30$ and for the total background (light histogram) for an integrated luminosity of 30 fb^{-1} .




Figure 11.20: Transverse mass reconstructed from the τ jet and missing transverse energy for the $gg \rightarrow tbH^\pm, t \rightarrow bW, W^\mp \rightarrow jj$ signal (dark histogram) with $m_{H^\pm} = 400 \text{ GeV}/c^2$, $\tan\beta=30$ and for the total background (light histogram) for an integrated luminosity of 30 fb^{-1} .

11.2.5.4 Systematic uncertainties on background determination

Background in the signal area $m_T(\tau \text{ jet}, E_T^{\text{miss}}) > 100 \text{ GeV}/c^2$ may arise from two main sources, the tail due to measurement uncertainties in the backgrounds with $W \rightarrow \tau\nu$ decays, and the possibility of fake τ jets mainly in the $W+3\text{jet}$ and QCD multi-jet events. The level of the backgrounds with $W \rightarrow \tau\nu$ decays can be measured from data exploiting the precise muon momentum measurement in $W+3\text{jets}$, $W \rightarrow \mu\nu$, selecting events in the tail of the transverse mass distribution. The probability of a hadronic jet faking the τ jets can be measured exploiting the precise photon energy measurement in the photon+jet events, as proposed in Ref. [268]. For this work a Monte-Carlo method was chosen assuming that the probability of the background events to migrate to the signal area depends mainly on the precision of the jet energy and E_T^{miss} measurements. The systematic uncertainty due to the energy scale was estimated varying the jet energy and the E_T^{miss} values with the expected energy scale uncertainties yielding the average values of 3% and 2% for the uncertainties of the efficiency of the E_T^{miss} cut and the efficiency of the selection of three hadronic jets for top reconstruction, respectively. The uncertainty of the τ identification has been estimated to be 8% for the τ jets in the E_T scales of $Z \rightarrow \tau\tau$ decays [?]. For the b-tagging uncertainty a conservative estimate of 5% was taken. The theoretical uncertainty on the $t\bar{t}$ cross section due to scale variation and the uncertainty choice of PDF has been estimated to be 5.6% [150]. These values yield 11% for the total systematic uncertainty for the $t\bar{t}$ background. For the $W+3\text{jet}$ and QCD multi-jet backgrounds the uncertainties due to present MC statistics strongly dominate the measurement uncertainties and therefore the statistical uncertainties are used. The total number of background events in the signal area $m_T(\tau \text{ jet}, E_T^{\text{miss}}) > 100 \text{ GeV}$ is 1.7 ± 1.0 events, including the systematic and MC uncertainties

Table 11.19: Value of $\tan\beta$, cross section times branching fraction, number of events and the statistical significance for a total background of 1.7 ± 1.0 events with and without background uncertainty, for the signal with $m_{H^\pm} = 170$ to 600 GeV/c^2 ($m_A = 150$ to 600 GeV/c^2) and for an integrated luminosity of 30 fb^{-1}

m_{H^\pm} (GeV/c^2)	171.6	180.4	201.0	300.9	400.7	600.8
$\tan\beta$	20	20	20	30	30	30
$\sigma(\text{NLO}) \times \text{BR}$ (pb)	0.968	0.866	0.543	0.182	0.058	0.0086
Events for 30 fb^{-1}	10.9 ± 2.4	10.4 ± 2.3	8.2 ± 1.6	12.8 ± 1.9	5.5 ± 0.7	1.1 ± 0.1
Significance, Syst.= 0	5.36	5.16	4.3	6.05	3.13	0.77
Significance	4.10	3.94	3.25	4.65	2.35	0.48

11.2.5.5 Discovery potential

Table 11.19 shows the number of signal events for $m_{H^\pm} = 170$ to 200 GeV/c^2 with $\tan\beta = 20$ and for $m_{H^\pm} = 300$ to 600 GeV/c^2 with $\tan\beta = 30$ and the statistical significance calculated according to poisson statistics [486] with and without background uncertainty for the total background of 1.7 ± 1.0 events. The results are shown for an integrated luminosity of 30 fb^{-1} . For the $t\bar{t}$ background the estimated systematic uncertainty of 11% is included. Figure 11.21 shows the 5σ -discovery potential for the heavy charged Higgs bosons as a function of m_A and $\tan\beta$ in the maximal mixing scenario with $\mu = 200$ GeV/c^2 with and without systematic uncertainties.




Figure 11.21: The 5σ -discovery region in the m_A , $\tan\beta$ plane for $gg \rightarrow tbH^\pm$, $H^\pm \rightarrow \tau\nu_\tau$ with an integrated luminosity of 30 fb^{-1} . The discovery regions with and without systematic uncertainties are shown. The regions excluded by the LEP and Tevatron searches are also shown in the figure.

11.2.6 Charged Higgs boson of $M_H > m_t$ in $gg \rightarrow tbH^\pm$ production with $H^\pm \rightarrow tb$

The branching fractions for the decay channels of the charged Higgs boson depend strongly on its mass (see Figure 11.2). For masses above $m_t + m_b$, the channel $H^\pm \rightarrow tb$ opens up. Two production channels and corresponding final states were considered in the search for

charged Higgs bosons in the $H^\pm \rightarrow tb$ decay channel [609]:

$$gb \rightarrow tH^\pm \rightarrow ttb \rightarrow W^+W^-bbb \rightarrow qq'\mu\nu_\mu bbb, \quad (11.9)$$

$$gg \rightarrow tH^\pm b \rightarrow ttbb \rightarrow W^+W^-bbbb \rightarrow qq'\mu\nu_\mu bbbb. \quad (11.10)$$

These final states are the most interesting ones from the experimental point of view because an isolated muon is present to trigger on and the branching fraction into this decay is high ($\sim 30\%$).

The inclusive final state (11.9) is studied using triple b tagging within the parametrized simulation framework of CMS [287]. The final state (11.10), where a fourth b jet is resolved in the detector, is studied with full `GEANT4` [599] CMS detector simulation [14]. Production of the H^\pm bosons through heavy sparticle cascades is not taken into account. In addition, supersymmetric particles are supposed to be heavy enough, such that supersymmetric decays of the H^\pm can be neglected.

11.2.6.1 Signal and background simulation

Events from the process (11.9) are modelled by considering the initial b quark as a massless parton from the corresponding parton density in the proton. On the other hand, events from the process (11.10) are described with massive spectator b quarks.

The calculation of the total signal cross section was performed at NLO [608], starting from the process (11.9). When calculating the cross section for both processes (11.9) and (11.10) to all orders, however, one expects to obtain the same result, as they both describe the same physics. Therefore, for both processes, the cross section was rescaled to the NLO result for the $pp \rightarrow tH^\pm X$ channel.

The signal cross section is sensitive to the two parameters $\tan\beta$ and m_{H^\pm} (Figures 11.22 and 11.23). The cross section is enhanced at small and large values of $\tan\beta$, with a minimum at $\tan\beta = \sqrt{m_t/m_b} \approx 6$. Furthermore, the cross section decreases rapidly with rising m_{H^\pm} . The generation of both processes (11.9) and (11.10) was performed with `PYTHIA` [62], forcing the decay $H^\pm \rightarrow tb$ of the charged Higgs boson. The branching fraction $\text{BR}(H^\pm \rightarrow tb)$ for this decay process was calculated with `HDECAY 3.0` [32].

The main background to charged Higgs boson production and decay through $pp \rightarrow tH^\pm(b) \rightarrow ttb(b)$ is by far the Standard Model top-quark pair production with additional jets. Other potential multi-jet backgrounds are much smaller and neglected.

In the case of process (11.9), the leading order background comes from SM $pp \rightarrow t\bar{t}b$ and $pp \rightarrow t\bar{t} + \text{jet}$ production, where in the latter the extra jet is misidentified as a b jet. The event simulation was performed using the matrix element generator `MadGraph/MadEvent` [73], interfaced to `PYTHIA` for parton shower, fragmentation and hadronization, with a cut $p_T > 10 \text{ GeV}/c$ on the transverse momentum and $|\eta| < 2.5$ on the pseudorapidity of the extra jet. This resulted in a cross section of 678 pb.

The background for process (11.10) consists of the irreducible $pp \rightarrow t\bar{t}b\bar{b}$ and the reducible $pp \rightarrow t\bar{t}jj$ process, where in the latter two jets are misidentified as b jets. Both these backgrounds were simulated using the `CompHEP` generator [34]. The generator level cuts $p_T > 15 \text{ GeV}$ and $|\eta| < 3$ were applied on the partons produced in association with the $t\bar{t}$ pair. A separation cut $\Delta R > 0.3$ was also imposed. This resulted in a cross section of 3.285 pb for the $pp \rightarrow t\bar{t}b\bar{b}$ process and 507.8 pb for $pp \rightarrow t\bar{t}jj$ production. Care was taken to avoid double




Figure 11.22: NLO cross section for $pp \rightarrow tH^\pm X$ as a function of m_{H^\pm} .




Figure 11.23: NLO cross section for $pp \rightarrow tH^\pm X$ as a function of $\tan\beta$.

counting between the $pp \rightarrow t\bar{t}b\bar{b}$ and $pp \rightarrow t\bar{t}jj$ processes and the cross section for $pp \rightarrow t\bar{t}jj$ was scaled to the result from a similar ALPGEN generation, where a jet matching technique was applied to more rigorously handle the transition between the hard interaction and the parton shower.

11.2.6.2 Event selection and reconstruction

On the final states (11.9) and (11.10) a basic event selection is applied on the reconstructed objects (Tables 11.20 and 11.21). Events passing the single muon HLT trigger are required to have at least one muon with $p_T > 20$ GeV and $|\eta| < 2.5$, at least respectively five or six calibrated jets with $p_T > 25$ GeV and $|\eta| < 2.5$ and at least respectively three or four of these jets tagged as b jet with a secondary vertex-based algorithm [148].

Table 11.20: Event selection yield for $\tan\beta = 30$ and an integrated luminosity of 30 fb^{-1} .

30 fb^{-1}	$t\bar{t}b/\bar{t}j$	$gb \rightarrow tH^\pm (\tan\beta = 30)$					
$m_{H^\pm} (\text{GeV}/c^2)$		263	311	359	408	457	506
cross section \times BR (pb)	678	0.850	0.570	0.377	0.251	0.169	0.116
# events before cuts	20.3M	25 489	17 088	11 319	7 529	5 063	3 472
single muon HLT	17%	16%	16%	16%	16%	16%	16%
1 muon	95%	95%	95%	95%	96%	96%	96%
5 jets	18%	35%	42%	44%	46%	49%	51%
3 b-tagged jets	6%	27%	29%	30%	32%	31%	29%
# remaining events	32 880	364	314	230	171	116	80

In both final states (11.9) and (11.10) the best jet association is selected with a likelihood ratio technique, which combines information from kinematical properties of the extra jets, b-tagging of all jets and the result of a kinematic fit on the $t\bar{t}$ system, imposing both W^\pm and t mass constraints. Starting from the chosen jet association the Higgs boson mass was reconstructed. An ambiguity remains, as it is not possible to know which top quark candidate the additional b jet should be combined with. In Figure 11.24 the reconstructed charged Higgs

Table 11.21: Event selection yield for $\tan\beta = 30$ and an integrated luminosity of 30 fb^{-1} .

30 fb^{-1}	$t\bar{t}b\bar{b}$	$t\bar{t}jj$	$gg \rightarrow tbH^\pm (\tan\beta = 30)$					
$m_{H^\pm} (\text{GeV}/c^2)$			263	311	359	408	457	506
cross section \times BR (pb)	2.386	235.8	0.850	0.570	0.377	0.251	0.169	0.116
# events before cuts	71 580	7.07M	25 489	17 088	11 319	7 529	5 063	3 472
single muon HLT	19%	19%	13%	13%	13%	13%	13%	13%
1 muon	96%	97%	96%	95%	97%	97%	97%	97%
6 jets	19%	23%	19%	23%	25%	26%	28%	31%
4 b-tagged jets	7%	0.55%	6%	5%	7%	7%	5%	6%
# remaining events	179	1 623	37	24	25	18	9	8

boson mass with hadronically decaying top is shown for correct and wrong jet pairings in the case of three tagged b jets and for $m_{H^\pm} = 311 \text{ GeV}/c^2$. Due to the large combinatorial background, the mass information is of limited use for the separation between signal and background, and is therefore not used further on in the analysis.

11.2.6.3 Background suppression

To suppress the large $t\bar{t}$ +jets background, observables were identified that have different properties for signal and background events. These observables were combined into an overall discriminator. In the case of process (11.9) the b-tagging information for the extra jet was used, together with the p_T of the softest jet from the W^\pm decay and the ratio of the E_T of the sixth jet and the fifth. For the process (11.10) only the b-tagging information for the two extra jets was used. In Figures 11.25 and 11.26 the resulting discriminator distributions are shown for the process (11.9) and (11.10) respectively.




Figure 11.24: Reconstructed H^\pm mass with hadronically decaying top for the chosen jet association ($m_{H^\pm} = 311 \text{ GeV}/c^2$).




Figure 11.25: Distribution of the discriminator used to distinguish between signal 11.9 and background.




Figure 11.26: Distribution of the discriminator used to distinguish between signal 11.10 and background.

11.2.6.4 Discovery reach and systematics

A cut on the discriminating variables of Figures 11.25 and ?? was optimized to obtain the maximal statistical significance for an integrated luminosity of 30 fb^{-1} . The signal cross section required for a significance of 5, corresponding to a discovery, was derived and translated into a minimal value of $\tan\beta$ needed for a discovery for a given value of m_A . Performing this analysis and optimization at different values of m_A a discovery contour was obtained in the MSSM $(\tan\beta, m_A)$ plane.

The background is large in both final states and therefore the effect of systematic uncertainties on the knowledge of the background is important. A possible way to estimate the background level from data is to require one b-tagged jet less. After such a selection it is possible to calculate the expected number of background events plus its uncertainty, when tagging a third or fourth b jet. Optimistically the uncertainty on the mistag rate can be taken as 5%. Possible large theoretical uncertainties related to this method, like the ratio of events with real extra b jets and events with only light extra jets, should still be accounted for.

Depending on the expected systematic uncertainty on the background level the maximal significance was searched. In Figures 11.27 and 11.28 the discovery contours are plotted for the final states (11.9) and (11.10) respectively, when supposing perfect knowledge of the background cross section ($\epsilon = 0$), a 1% uncertainty ($\epsilon = 0.01$), and a 3% uncertainty ($\epsilon = 0.03$). From the above estimate of the systematic uncertainty on the number of background events, the conclusion is drawn that, neglecting SUSY cascade decays, no visibility for this channel is obtained in the MSSM parameter space during the low luminosity phase of LHC.




Figure 11.27: Discovery contour for the charged Higgs boson in the $H^\pm \rightarrow tb$ decay, applying 3 b tags, for 30 fb^{-1} ; systematic uncertainties on the background of $\epsilon = 0\%$, 1% and $\epsilon = 3\%$ are taken into account.




Figure 11.28: Discovery contour for the charged Higgs boson in the $H^\pm \rightarrow tb$ decay, applying 4 b tags, for 30 fb^{-1} ; systematic uncertainties on the background of $\epsilon = 0\%$, $\epsilon = 1\%$ and $\epsilon = 3\%$ are taken into account.

11.2.7 Search for the $A \rightarrow Zh$ decay with $Z \rightarrow \ell^+\ell^-$, $h \rightarrow b\bar{b}$

The observation of the CP-odd pseudo-scalar Higgs (A) via its decay into a Z boson and the lighter CP-even scalar Higgs (h) followed by $Z \rightarrow e^+e^-$, $\mu^+\mu^-$ and $h \rightarrow b\bar{b}$ decays provides an interesting way to detect A and h simultaneously. The largest branching ratio of the $A \rightarrow Zh$ appears for low $\tan\beta$ and $m_Z + m_h \leq m_A \leq 2m_{top}$ mass region. The main production




Figure 11.29: The cross-section time branching ratios as a function of M_A for $\tan\beta=1$ in the m_h^{\max} scenario with $\mu=M_2=200$ GeV/c^2 (circles) and $\mu=M_2=600$ GeV/c^2 (triangles)

mechanism for A at low $\tan\beta$ is via $gg,qq \rightarrow A$.

The decays of the A into charginos and neutralinos ($A \rightarrow \chi\chi$), however, can dominate at the certain values of μ and M_2 , Higgs-higgsino and $SU(2)$ gaugino mass parameters, since the masses of charginos and neutralinos as well as their couplings to the Higgs bosons depend on μ and M_2 (in addition to $\tan\beta$ and M_A). The large values of μ and M_2 are the favorite ones for the observation of the $A \rightarrow Zh$ channel.

In Figure 11.29 and 11.30 the production cross section times the appropriate branching ratios (including $Z \rightarrow e^+e^-$, $\mu^+\mu^-$ and $h \rightarrow b\bar{b}$ decays) is shown as a function of M_A in the m_h^{\max} scenario with $\mu=M_2=200$ GeV/c^2 and $\mu=M_2=600$ GeV/c^2 for two values of the $\tan\beta$, 1 and 5. One can see that the difference in the total cross sections for the two choices of the μ and M_2 parameters can be as large as the one order of magnitude. The $A \rightarrow Zh$ analysis and the discovery reach presented below was evaluated in the m_h^{\max} scenario with $\mu=M_2=600$ GeV/c^2 .

11.2.7.1 Event Generation , Simulation and Reconstruction

The Higgs boson production processes, $gg \rightarrow A$ and $pp \rightarrow A b\bar{b}$, were generated using PYTHIA 6.225 [62] for three values of M_A (250, 300, 350 GeV/c^2) and two values of $\tan\beta$ (1.0, 5.0). No preselection at generation level was applied. The Standard Model backgrounds considered are: the $Zb\bar{b}$ generated with CompHEP [47]) and ZZ , ZW , Z +jets, W +jets and $t\bar{t}$ generated with PYTHIA 6.215. Events were fully simulated and digitized with pile-up corresponding to a luminosity of $2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. Offline reconstruction of electrons and muons and b tagging were performed using standard algorithms.




Figure 11.30: The cross-section time branching ratios as a function of M_A for $\tan\beta=5$ in the m_h^{\max} scenario with $\mu=M_2=200$ GeV/c^2 (circles) and $\mu=M_2=600$ GeV/c^2 (triangles)

11.2.7.2 Online Selection

The events are required to pass the global Level-1 (L1) and High Level Trigger (HLT) di-muon or di-electron selections since there will always be a real Z in the event decaying into two high p_T electrons or muons. The inclusion of the single muon and electron triggers was found does not improve the discovery reach in the M_A - $\tan\beta$ plane.

11.2.7.3 Off-line Event Selection

The baseline selection requires two opposite sign high p_T isolated leptons (e or μ) and two high E_T tagged b-jets separated from the leptons with $\Delta R(\ell, j) \geq 0.7$. Muons must have $|\eta| < 2.4$ and electrons should be in the ECAL fiducial region ($|\eta| < 2.5$ with $1.444 < |\eta| < 1.566$ region excluded). The event is required to have small missing E_T and reconstructed invariant mass of the leptons close to the Z mass in order to reject a significant fraction of the $t\bar{t}$ background.

Table 11.22 summarizes the basic selection variables and the thresholds. The variation of the signal significance with the change of the p_T thresholds on the electrons, muons and b-jets, and the thresholds on the b-tagging discriminant for the two tagged jets has been checked. No significant variation was found with the small changes of the cut values presented in Table 11.22.

Table 11.22: Selection variables and thresholds

Selection Variable	Threshold
most energetic electron/muon p_T	$> 30 \text{ GeV}/c$
second-most energetic electron/muon p_T	$> 15 \text{ GeV}/c$
most energetic b-jet E_T	$> 25 \text{ GeV}$
second-most energetic b-jet E_T	$> 20 \text{ GeV}$
missing E_T	$< 60 \text{ GeV}$
most energetic b-jet discriminator	> 1.5
second-most energetic b-jet discriminator	> 0.5
Z mass cut	$84 \text{ GeV}/c^2 < M_Z < 96 \text{ GeV}/c^2$
Z p_T	$> 30.0 \text{ GeV}/c$

11.2.7.4 Results

The selection efficiencies for the signal vary from 5% to 12% depending on the M_A and $\tan\beta$ values as well as the production mechanism. The details can be found in [610]. The next-to-leading order (NLO) background cross sections before and after selections are shown in Table 11.23.

The signal and the background distributions of $M_{b\bar{b}}$ and $M_{\ell+\ell-b\bar{b}}$ after selections are shown in Figure 11.31 and Figure 11.32 respectively for 30 fb^{-1} of integrated luminosity.

11.2.7.5 Systematic uncertainties

The method to evaluate the background from the real data measuring the background in the signal free (normalization) region is proposed. The background uncertainty then consists of

Table 11.23: Background Efficiencies

	NLO cross sections (fb)	
	before selection	after selection
$Zb\bar{b}, Z \rightarrow ee, \mu\mu, \tau\tau$	112830	415.26
$t\bar{t}, W \rightarrow e\nu, \mu\nu, \tau\nu$	88500	70.8
$Z+\text{jets}, Z \rightarrow ee, \mu\mu, \tau\tau$	5300000	83.05
$W+\text{jets}, W \rightarrow e\nu, \mu\nu, \tau\nu$	47900000	0.0
ZZ (inclusive)	14985	7.34
ZW (inclusive)	49422	1.98




Figure 11.31: Distribution of $M_{b\bar{b}}$ for signal and background after event selection for 30 fb^{-1} of integrated luminosity. Red (dark gray), yellow (light gray) and green (medium gray) distributions represent the $Zb\bar{b}$, $t\bar{t}$ and $Z+\text{jets}$ backgrounds. Blue (black) distribution is the signal ($M_A=300, \tan\beta=2$) and black dots the data (sum of the signal and the background).




Figure 11.32: Distribution of $M_{\ell^+\ell^-b\bar{b}}$ for signal and background after event selection for 30 fb^{-1} of integrated luminosity. Red (dark gray), yellow (light gray) and green (medium gray) distributions represent the $Zb\bar{b}$, $t\bar{t}$ and $Z+\text{jets}$ backgrounds. Blue (black) distribution is the signal ($M_A=300, \tan\beta=2$) and black dots the data (sum of the signal and the background).

the statistical uncertainty of the background measurement in the normalization region and systematic uncertainty of the ratio of the background in the signal and the normalization region.

The normalization region for the $t\bar{t}$ background is defined by same selections as for the signal search, except the missing E_T which is required to be bigger than 120 GeV. With a such selections 544 events were found for 30 fb^{-1} with high purity (93.4%), thus giving the statistical uncertainty of 4.4%. The distribution of $M_{\ell^+\ell^-b\bar{b}}$ in the $t\bar{t}$ normalization region can be seen in Figure 11.33. The contamination comes mainly from $Zb\bar{b}$ events (6%). The 5% missing E_T scale uncertainty gives 18.5% uncertainty on the number of the $t\bar{t}$ events in the signal region. Therefore the overall uncertainty in the estimation of the $t\bar{t}$ background is 19.0%.

For the irreducible $Zb\bar{b}$ background a similar idea can be used. In order to suppress the $t\bar{t}$ contribution as much as possible, the missing $E_T < 40 \text{ GeV}$ was used. Applying a lower cut in the $M_{\ell^+\ell^-b\bar{b}}$ distribution of $500 \text{ GeV}/c^2$, 920 $Zb\bar{b}$ events were found with a purity




Figure 11.33: Distribution of $M_{\ell+\ell-b\bar{b}}$ in the $t\bar{t}$ background normalization region. Color code is as in Fig. 11.32




Figure 11.34: Distribution of $M_{\ell+\ell-b\bar{b}}$ used in the $Zb\bar{b}$ background estimation. Color code is as in Fig. 11.32

around 95% for 30fb^{-1} . Contamination comes mainly from $t\bar{t}$ events. The accuracy of measuring the $Zb\bar{b}$ background is around 3.4% taking into account only statistics. The distribution of $M_{\ell+\ell-b\bar{b}}$ for those events can be seen in Figure 11.34 before the application of the $M_{\ell+\ell-b\bar{b}} > 500 \text{ GeV}/c^2$ cut. The uncertainty 5% of the missing E_T scale and the uncertainty 3% of the jet energy scale lead to correspondingly 3.6% and 2.5% of the uncertainty of the $Zb\bar{b}$ background estimate in the signal region. So the overall uncertainty in the estimation of the $Zb\bar{b}$ background is 5.6%.

11.2.7.6 Discovery reach in M_A - $\tan\beta$ plane.

Figure 11.35 shows the 5σ discovery contours in the $(M_A, \tan\beta)$ plane for 30 and 60 fb^{-1} of integrated luminosity in the m_h^{max} scenario with $\mu = M_2 = 600 \text{ GeV}/c^2$. For the calculation of the signal significance the signal and background events were counted in mass windows of $\pm\sigma$ around the reconstructed mass of M_h and M_A . Since only three different M_A masses and two $\tan\beta$ values are available, the estimations for the rest of $M_A, \tan\beta$ parameter space was done using extra/inter-polations of the signal efficiencies from the available parameter points. The statistical significance for 5,10% (dashed lines) as well as the estimated (line) uncertainty for the background is also shown for 30 fb^{-1} of integrated luminosity.

11.2.8 Search for $A^0/H^0 \rightarrow \chi_2^0\chi_2^0 \rightarrow 4l + E_T^{\text{miss}}$ channel in mSUGRA

11.2.8.1 Introduction

In some regions of the SUSY parameter space, heavy neutral Higgs bosons can be searched for using their decay modes to supersymmetric particles. This is the case in particular in the difficult low and intermediate $\tan\beta$ region of the parameter space which is not accessible through the $A^0/H^0 \rightarrow \tau\tau$ decay channel as the coupling of the Higgs boson to taus is not sufficiently enhanced.

One of the most promising channel is the A^0/H^0 decay into a pair of next-to-lightest neutralinos, χ_2^0 , followed by the decay $\chi_2^0 \rightarrow l^+l^-\chi_1^0$ (with $l = e, \mu$). This process results in a




Figure 11.35: The 5σ discovery contours for 30 and 60 fb^{-1} integrated luminosity with and without systematic uncertainties taken into account. The effect of underestimation or overestimation of the background systematic uncertainty can be seen in the curve of 30 fb^{-1} .

clean four leptons plus missing transverse energy final state:

$$A^0/H^0 \rightarrow \chi_2^0 \chi_2^0 \rightarrow 4l + E_T^{miss} .$$

There are two main categories of backgrounds to such process: SUSY and Standard Model backgrounds. In the SUSY category the dominant source of background is the production of leptons from the decays of squarks and gluinos which cascade to charginos and neutralinos. Unlike the neutralinos from the Higgs boson decay, the leptons in this case are produced in association with quarks and gluons. Therefore, the associated large hadronic activity can be used to suppress this type of background. An additional but smaller source of backgrounds come from the direct production of slepton or gaugino pairs via the Drell-Yan processes and the direct production of χ_2^0 pairs. The rejection of these backgrounds is more difficult, as the hadronic activity in these events is very small. In the Standard Model category, three processes which yield to the same signature of 4 leptons in the final state contribute as backgrounds: ZZ^*/γ^* , $Zb\bar{b}$ and $t\bar{t}$.

11.2.8.2 Analysis

The study is performed in the minimal Super Gravity constrained version of the MSSM (mSUGRA). To determine the regions where the signal has a sizeable branching ratio times cross section, a scan of the parameters space $(m_0, m_{1/2})$ for $\tan\beta = 5, 10$, $sign(\mu) = +$ and $A_0 = 0$ is performed. Three benchmark points are defined for the evaluation of CMS sensitivity. The corresponding mSUGRA parameters are presented in Table 11.24.

The signal and SUSY background datasets are generated using ISASUGRA and PYTHIA. A preselection at generator level is applied, asking for $e^+e^-\mu^+\mu^-$ final state with $e(\mu) p_T > 7(5)$ GeV and $|\eta| < 2.5$. The detector simulation is carried out using FAMOS_1.4.0.

Point	m_0 (GeV)	$m_{1/2}$ (GeV)	A_0 (GeV)	$\tan\beta$	$sign(\mu)$
A	60	175	0	10	+
B	80	200	0	5	+
C	50	150	0	5	+

Table 11.24: Chosen benchmark points.

The online selection of the events is a logical or of the di-electron and di-muon triggers. The offline reconstruction of electrons and muons is performed using FAMOS standard algorithms. Events are then analyzed as follow:

- $e^+e^-\mu^+\mu^-$ final state is selected;
- the four leptons are required to be isolated;
- a jet veto is applied, requiring no jets with $p_T > 25$ GeV;
- events must have E_T^{miss} and $p_{T(III)}$ less than 80 GeV;
- a Z veto is imposed, i.e. events with a dilepton pair with invariant mass in the range $m_Z \pm 10$ GeV are rejected;
- further optimizations are performed by introducing an upper limit to the dilepton invariant masses and by applying a cut on the four lepton invariant mass.

The signal acceptances w.r.t the production cross section times branching ratio are 6.3%, 5.1% and 2.5% respectively for point A, B and C, whereas the acceptances for SUSY backgrounds are $1.5 \cdot 10^{-4}\%$, $3.6 \cdot 10^{-4}\%$ and $2.6 \cdot 10^{-4}\%$ respectively w.r.t the total the SUSY production cross section.

11.2.8.3 Results

Figure 11.36 presents the invariant mass distribution of the four leptons for the 3 benchmark points. Results are given for an integrated luminosity of 30 fb^{-1} .




Figure 11.36: Four lepton invariant mass distributions for the 3 benchmark points. Distributions are shown for the signal + backgrounds (points) and for the contribution of each process (histograms).

Figure 11.37 shows the extrapolated 5σ -discovery contours in the $(m_0, m_{1/2})$ plane, for an

integrated luminosity of 30 fb^{-1} . The values of the other mSUGRA parameters are $A_0 = 0$, $\text{sign}(\mu) = +$ and $\tan\beta = 5, 10$. The complex structure of the high significance region is mainly determined by the effective cross section of $A^0/H^0 \rightarrow \chi_2^0 \chi_2^0 \rightarrow 4l + E_T^{\text{miss}}$. The A^0/H^0 could therefore be discovered through their decays to neutralino pairs in the region $150 < m_{1/2} < 250$ and $m_0 < 120$ for $\tan\beta = 10$ and in the region $150 < m_{1/2} < 250$ and $30 < m_0 < 120$ for $\tan\beta = 5$.




Figure 11.37: For integrated luminosity of 30 fb^{-1} 5σ -discovery contours for $A^0/H^0 \rightarrow \chi_2^0 \chi_2^0 \rightarrow 4l + E_T^{\text{miss}}$ channel in the $(m_0, m_{1/2})$ plane for fixed $A_0 = 0$, $\text{sign}(\mu) = +$ and $\tan\beta = 5, 10$.

3 Discovery reach and measurement of MSSM parameters

11.3.1 Benchmark scenarios for MSSM Higgs boson searches

11.3.1.1 Why benchmarks — which benchmarks?

The tree-level values for the CP-even Higgs bosons of the MSSM, M_h and M_H , are determined by $\tan\beta$, the CP-odd Higgs-boson mass M_A , and the Z boson mass M_Z . The mass of the charged Higgs boson, M_{H^\pm} , is given in terms of M_A and the W boson mass, M_W . Beyond the tree-level, the main correction to the Higgs boson masses stems from the t/\tilde{t} sector, and for large values of $\tan\beta$ also from the b/\tilde{b} sector, see Section 11.1. Subleading corrections come from all other sectors of the MSSM. In this way the Higgs sector phenomenology is connected to the full spectrum of the MSSM via radiative corrections.

In the unconstrained version of the MSSM no particular SUSY breaking mechanism is assumed, but rather a parameterization of all possible soft SUSY breaking terms is used. This leads to more than a hundred parameters (masses, mixing angles, phases) in this model in addition to the ones of the Standard Model. While a detailed scanning over the more-than-hundred-dimensional parameter space of the MSSM is clearly not practicable, even a sampling of three- or four-dimensional parameter space of certain SUSY-breaking models (such as mSUGRA, GMSB or AMSB) is beyond the present capabilities for phenomenological studies, in particular when it comes to simulating experimental signatures within the detectors. For this reason one often resorts to specific benchmark scenarios, i.e. one studies only specific parameter points [611, 612] or samples of one- or two-dimensional parameter

space [613, 614, 251], which exhibit specific characteristics of the MSSM parameter space. Benchmark scenarios of this kind are often used, for instance, for studying the performance of different experiments at the same collider. Similarly, detailed experimental simulations of MSSM particle production with identical parameters in the framework of different colliders can be very helpful for developing strategies for combining pieces of information obtained at different machines [615].

The question of which parameter choices are useful as benchmark scenarios depends on the purpose of the actual investigation. If one is interested, for instance, in setting exclusion limits on the SUSY parameter space from the non-observation of SUSY signals at the experiments performed up to now, it is useful to use a benchmark scenario which gives rise to “conservative” exclusion bounds. An example of a benchmark scenario of this kind is the m_h^{\max} -scenario [614] used for the Higgs search at LEP [552]. It gives rise to maximal values of the lightest CP-even Higgs-boson mass (for fixed values of the top-quark mass and the SUSY scale) and thus allows one to set conservative bounds on $\tan\beta$ and M_A [530]. Another application of benchmark scenarios is to study “typical” experimental signatures of SUSY models and to investigate the experimental sensitivities and the achievable experimental precisions for these cases. For this purpose it seems reasonable to choose “typical” (a notion which is of course difficult to define) and theoretically well motivated parameters of certain SUSY-breaking scenarios. Examples of this kind are the benchmark scenarios used so far for investigating SUSY searches at the LHC [611, 612] and at the ILC [616]. As a further possible goal of benchmark scenarios, one can choose them so that they account for a wide variety of SUSY phenomenology. For this purpose, it can also be useful to consider “pathological” regions of parameter space or “worst-case” scenarios. Examples for this are the “small α_{eff} scenario” [614] for the Higgs search at LEP, for which the decay $h \rightarrow b\bar{b}$ or $h \rightarrow \tau^+\tau^-$ can be significantly suppressed.

A related issue concerning the definition of appropriate benchmarks is whether a benchmark scenario chosen for investigating physics at a certain experiment or for testing a certain sector of the theory should be compatible with additional information from other experiments (or concerning other sectors of the theory). This refers in particular to constraints from cosmology (by demanding that SUSY should give rise to an acceptable dark matter density [617, 618, 619, 620]) and low-energy measurements such as the rate for $b \rightarrow s\gamma$ [621, 622] and the anomalous magnetic moment of the muon, $(g-2)_\mu$ [623, 624]. On the one hand, applying constraints of this kind gives rise to “more realistic” benchmark scenarios. On the other hand, one relies in this way on further assumptions (and has to take account of experimental and theoretical uncertainties related to these additional constraints), and it could eventually turn out that one has inappropriately narrowed down the range of possibilities by applying these constraints. This applies in particular if slight modifications of the model under consideration are possible that have a minor impact on collider phenomenology but could significantly alter the bounds from cosmology and low-energy experiments. For instance, the presence of small flavor mixing terms in the SUSY Lagrangian could severely affect the prediction for $\text{BR}(b \rightarrow s\gamma)$, while allowing a small amount of R-parity violation in the model would strongly affect the constraints from dark matter relic abundance while leaving collider phenomenology essentially unchanged. The extent to which additional constraints of this kind should be applied to possible benchmark scenarios is related to the actual purpose of the benchmark scenario. For setting exclusion bounds in a particular sector (e.g. the Higgs sector) it seems preferable to apply constraints from this sector only.

11.3.1.2 The relevant MSSM parameters

Beyond the tree-level, the main correction to the Higgs boson masses and couplings comes from the t/\tilde{t} sector, and for large values of $\tan\beta$ also from the b/\tilde{b} sector. In order to fix our notations, we list the conventions for the inputs from the scalar top and scalar bottom sector of the MSSM: the mass matrices in the basis of the current eigenstates \tilde{t}_L, \tilde{t}_R and \tilde{b}_L, \tilde{b}_R are given by

$$\mathcal{M}_{\tilde{t}}^2 = \begin{pmatrix} M_{\tilde{t}_L}^2 + m_t^2 + \cos 2\beta(\frac{1}{2} - \frac{2}{3}s_W^2)M_Z^2 & m_t X_t \\ m_t X_t & M_{\tilde{t}_R}^2 + m_t^2 + \frac{2}{3}\cos 2\beta s_W^2 M_Z^2 \end{pmatrix}, \quad (11.11)$$

$$\mathcal{M}_{\tilde{b}}^2 = \begin{pmatrix} M_{\tilde{b}_L}^2 + m_b^2 + \cos 2\beta(-\frac{1}{2} + \frac{1}{3}s_W^2)M_Z^2 & m_b X_b \\ m_b X_b & M_{\tilde{b}_R}^2 + m_b^2 - \frac{1}{3}\cos 2\beta s_W^2 M_Z^2 \end{pmatrix} \quad (11.12)$$

where

$$m_t X_t = m_t(A_t - \mu \cot\beta), \quad m_b X_b = m_b(A_b - \mu \tan\beta). \quad (11.13)$$

Here A_t denotes the trilinear Higgs–stop coupling, A_b denotes the Higgs–sbottom coupling, and μ is the higgsino mass parameter.

SU(2) gauge invariance leads to the relation

$$M_{\tilde{t}_L} = M_{\tilde{b}_L}. \quad (11.14)$$

For the numerical evaluation, a convenient choice is

$$M_{\tilde{t}_L} = M_{\tilde{b}_L} = M_{\tilde{t}_R} = M_{\tilde{b}_R} =: M_{\text{SUSY}}. \quad (11.15)$$

We furthermore use the short-hand notation

$$M_{\tilde{S}}^2 := M_{\text{SUSY}}^2 + m_t^2. \quad (11.16)$$

Accordingly, the most important parameters for the corrections in the Higgs sector are m_t , M_{SUSY} , X_t and X_b (or equivalently A_t and A_b), μ and $\tan\beta$. The Higgs sector observables furthermore depend on the SU(2) gaugino mass parameter, M_2 . The other gaugino mass parameter, M_1 , is usually fixed via the GUT relation

$$M_1 = \frac{5}{3} \frac{s_W^2}{c_W^2} M_2. \quad (11.17)$$

At the loop level also the gluino mass, $m_{\tilde{g}}$, enters the predictions for the Higgs-boson phenomenology.

It should be noted in this context that the results for Higgs boson sector observables have been obtained in different schemes. Most commonly these are the on-shell (OS) renormalization scheme (in the Feynman-diagrammatic (FD) approach), and $\overline{\text{MS}}$ scheme (for the renormalization group (RG) approach) [625]. Owing to the different schemes used in the FD and the RG approach for the renormalization in the scalar top sector, the parameters X_t and M_{SUSY} are also scheme-dependent in the two approaches. This difference between the corresponding parameters has to be taken into account when defining the benchmark scenarios.

In a simple approximation the relation between the parameters in the different schemes is at $\mathcal{O}(\alpha_s)$ given by [625]

$$M_S^{2,\overline{\text{MS}}} \approx M_S^{2,\text{OS}} - \frac{8}{3} \frac{\alpha_s}{\pi} M_S^2, \quad (11.18)$$

$$X_t^{\overline{\text{MS}}} \approx X_t^{\text{OS}} + \frac{\alpha_s}{3\pi} M_S \left(8 + 4 \frac{X_t}{M_S} - 3 \frac{X_t}{M_S} \log \left(\frac{m_t^2}{M_S^2} \right) \right). \quad (11.19)$$

At large $\tan \beta$ and large $|\mu|$ the corrections from the b/\bar{b} sector can become especially important. The leading effects are included in the effective Lagrangian formalism [549]. Numerically this is by far the dominant part of the contributions from the sbottom sector (see also Refs. [533, 534]). The effective Lagrangian is given by

$$\begin{aligned} \mathcal{L} = \frac{g}{2M_W} \frac{\bar{m}_b}{1 + \Delta_b} \left[\right. & \tan \beta A i \bar{b} \gamma_5 b + \sqrt{2} V_{tb} \tan \beta H^+ \bar{t}_L b_R \\ & + \left(\frac{\sin \alpha}{\cos \beta} - \Delta_b \frac{\cos \alpha}{\sin \beta} \right) h \bar{b}_L b_R \\ & \left. - \left(\frac{\cos \alpha}{\cos \beta} + \Delta_b \frac{\sin \alpha}{\sin \beta} \right) H \bar{b}_L b_R \right] + \text{h.c.} . \quad (11.20) \end{aligned}$$

Here \bar{m}_b denotes the running bottom quark mass including SM QCD corrections. The prefactor $1/(1 + \Delta_b)$ in Eq. 11.20 arises from the resummation of the leading corrections to all orders. The function Δ_b consists of two main contributions, an $\mathcal{O}(\alpha_s)$ correction from a sbottom–gluino loop and an $\mathcal{O}(\alpha_t)$ correction from a stop–higgsino loop. The explicit form of Δ_b in the limit of $M_S \gg m_t$ and $\tan \beta \gg 1$ reads [549]

$$\Delta_b = \frac{2\alpha_s}{3\pi} m_{\bar{g}} \mu \tan \beta \times I(m_{\bar{b}_1}, m_{\bar{b}_2}, m_{\bar{g}}) + \frac{\alpha_t}{4\pi} A_t \mu \tan \beta \times I(m_{\bar{t}_1}, m_{\bar{t}_2}, \mu). \quad (11.21)$$

The function I is given by

$$\begin{aligned} I(a, b, c) &= \frac{1}{(a^2 - b^2)(b^2 - c^2)(a^2 - c^2)} \left(a^2 b^2 \log \frac{a^2}{b^2} + b^2 c^2 \log \frac{b^2}{c^2} + c^2 a^2 \log \frac{c^2}{a^2} \right) \\ &\sim \frac{1}{\max(a^2, b^2, c^2)}. \end{aligned} \quad (11.22)$$

It becomes obvious that the size and the sign of μ is especially relevant for this type of corrections.

11.3.1.3 The benchmark scenarios

Since at the tree-level the Higgs sector of the MSSM is governed by two parameters (in addition to M_Z and the SM gauge couplings), it seems reasonable to define benchmarks in which all SUSY parameters are fixed and only the two tree-level parameters, M_A and $\tan \beta$ are varied. For the search of the heavy MSSM Higgs bosons corrections from the b/\bar{b} sector can be especially relevant. In this case it is also appropriate to vary μ . We review the definition of the benchmark scenarios as defined in Refs. [614, 251]. Another very important parameter

is the top-quark mass. For sake of simplicity and to make different analyses readily comparable to each other a fixed value of $m_t = 175$ GeV can be used. Alternatively the current experimental value can be used as input.

The m_h^{\max} scenario

This scenario was designed to obtain conservative $\tan \beta$ exclusion bounds [530] at LEP [552]. The parameters are chosen such that the maximum possible Higgs-boson mass as a function of $\tan \beta$ is obtained (for fixed M_{SUSY} , and M_A set to its maximal value, $M_A = 1$ TeV). The parameters are*:

$$\begin{aligned} m_t &= 175 \text{ GeV}, & M_{\text{SUSY}} &= 1 \text{ TeV}, & \mu &= 200 \text{ GeV}, & M_2 &= 200 \text{ GeV}, \\ X_t^{\text{OS}} &= 2 M_{\text{SUSY}} \text{ (FD calculation)}, & X_t^{\overline{\text{MS}}} &= \sqrt{6} M_{\text{SUSY}} \text{ (RG calculation)} \\ A_b &= A_t, & m_{\tilde{g}} &= 0.8 M_{\text{SUSY}}. \end{aligned} \quad (11.23)$$

The no-mixing scenario

This benchmark scenario is the same as the m_h^{\max} scenario, but with vanishing mixing in the \tilde{t} sector and with a higher SUSY mass scale to avoid the LEP Higgs bounds [552, 54],

$$\begin{aligned} m_t &= 175 \text{ GeV}, & M_{\text{SUSY}} &= 2 \text{ TeV}, & \mu &= 200 \text{ GeV}, & M_2 &= 200 \text{ GeV}, \\ X_t &= 0 \text{ (FD/RG calculation)}, & A_b &= A_t, & m_{\tilde{g}} &= 0.8 M_{\text{SUSY}}. \end{aligned} \quad (11.24)$$

The gluophobic Higgs scenario

In this scenario the main production cross section for the light Higgs boson at the LHC, $gg \rightarrow h$, is strongly suppressed. This can happen due to a cancellation between the top quark and the stop quark loops in the production vertex (see Ref. [490]). This cancellation is more effective for small \tilde{t} masses and hence for relatively large values of the \tilde{t} mixing parameter, X_t . The partial width of the most relevant decay mode, $\Gamma(h \rightarrow \gamma\gamma)$, is affected much less, since it is dominated by the W boson loop. The parameters are:

$$\begin{aligned} m_t &= 175 \text{ GeV}, & M_{\text{SUSY}} &= 350 \text{ GeV}, & \mu &= 300 \text{ GeV}, & M_2 &= 300 \text{ GeV}, \\ X_t^{\text{OS}} &= -750 \text{ GeV (FD calculation)}, & X_t^{\overline{\text{MS}}} &= -770 \text{ GeV (RG calculation)} \\ A_b &= A_t, & m_{\tilde{g}} &= 500 \text{ GeV}. \end{aligned} \quad (11.25)$$

In the left plot of Fig. 11.38 we show $[\sigma \times \text{BR}]_{\text{MSSM}}/[\sigma \times \text{BR}]_{\text{SM}}$ for the channel $gg \rightarrow h \rightarrow \gamma\gamma$ in the $M_A - \tan \beta$ -plane. This channel can be strongly suppressed over the whole parameter plane, rendering this detection channel difficult.

The small α_{eff} scenario

Besides the channel $gg \rightarrow h \rightarrow \gamma\gamma$ at the LHC, other channels for light Higgs searches at the Tevatron and at the LHC rely on the decays $h \rightarrow \bar{b}b$ and $h \rightarrow \tau^+\tau^-$. If α_{eff} is small, these two decay channels can be heavily suppressed in the MSSM due to the additional factor

* Better agreement with $\text{BR}(b \rightarrow s\gamma)$ constraints is obtained for the other sign of X_t (called the “constrained m_h^{\max} ” scenario). However, this lowers the maximum M_h values by ~ 5 GeV.




Figure 11.38: $[\sigma \times \text{BR}]_{\text{MSSM}}/[\sigma \times \text{BR}]_{\text{SM}}$ is shown for the channels $gg \rightarrow h \rightarrow \gamma\gamma$ in the gluophobic Higgs scenario (left plot) and $WW \rightarrow h \rightarrow \tau^+\tau^-$ in the small α_{eff} scenarios (right plot) in the $M_A - \tan\beta$ -plane. The white-dotted area is excluded by LEP Higgs searches.

– $\sin\alpha_{\text{eff}}/\cos\beta$ compared to the SM coupling. Such a suppression occurs for large $\tan\beta$ and not too large M_A for the following parameters:

$$\begin{aligned}
 m_t &= 175 \text{ GeV}, & M_{\text{SUSY}} &= 800 \text{ GeV}, & \mu &= 2.5 M_{\text{SUSY}}, & M_2 &= 500 \text{ GeV}, \\
 X_t^{\text{OS}} &= -1100 \text{ GeV (FD calculation)}, & X_t^{\overline{\text{MS}}} &= -1200 \text{ GeV (RG calculation)} \\
 A_b &= A_t, & m_{\tilde{g}} &= 500 \text{ GeV}.
 \end{aligned} \tag{11.26}$$

In the right plot of Fig. 11.38 we show $[\sigma \times \text{BR}]_{\text{MSSM}}/[\sigma \times \text{BR}]_{\text{SM}}$ for the channel $WW \rightarrow h \rightarrow \tau^+\tau^-$ in the $M_A - \tan\beta$ -plane. Significant suppression occurs for large $\tan\beta$, $\tan\beta > 20$, and small to moderate M_A , $M_A < 400$ GeV. Thus, Higgs boson search via the WW fusion channel will be difficult in these parts of the parameter space.

11.3.1.4 Variation of μ

The most sensitive channels for detecting heavy MSSM Higgs bosons at the LHC are the channel $pp \rightarrow H/A + X$, $H/A \rightarrow \tau^+\tau^-$ (making use of different decay modes of the two τ leptons) and the channel $tH^\pm, H^\pm \rightarrow \tau\nu_\tau$ (for $M_{H^\pm} > m_t$). These channels show good prospects for $M_A \gg M_Z$ and large $\tan\beta$.

As discussed above, in this part of the parameter space the corrections from the b/\tilde{b} sector can be very important and thus the size and the sign of μ can play a dominant role. This leads to the definition of an extension of the m_h^{max} and the no-mixing scenario by the following values of μ [251]

$$\mu = \pm 200, \pm 500, \pm 1000 \text{ GeV}, \tag{11.27}$$

allowing both an enhancement and a suppression of the bottom Yukawa coupling and taking into account the limits from direct searches for charginos at LEP. It should be noted that the values $\mu = -500, -1000$ GeV can lead to such a large enhancement of the bottom Yukawa coupling that a perturbative treatment is no longer possible in the region of very large values of $\tan\beta$. Some care is therefore necessary to assess up to which values of μ reliable results can be obtained.

A further variation of the discovery reach is caused by the decays of the heavy Higgs bosons into supersymmetric particles. For a given value of μ , the rates of these decay modes are strongly dependent on the particular values of the weak gaugino mass parameters M_2 and M_1 . Since the Higgs couplings to neutralinos and charginos depend strongly on the admixture between higgsino and gaugino states, the rate of these processes is strongly suppressed for large values of $|\mu| > 500$ GeV. In general, the effects of the decays $H/A \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^0, \tilde{\chi}_k^\pm \tilde{\chi}_l^\mp$ only play a role for $M_A > |\mu| + M_1$. Outside this range the dependence of the rates on μ is relatively weak.

11.3.2 Discovery reach in the M_A - $\tan\beta$ plane

This section summarizes the discovery reach in the M_A - $\tan\beta$ plane for the charged and the neutral MSSM Higgs bosons in the m_h^{\max} scenario. The cross sections and branching ratios for the neutral Higgs boson and the branching ratios for the charged Higgs boson were calculated with FeynHiggs2.3.2 [136, 137, 138]. The next-to-leading order cross section for the charged Higgs production was taken from Refs. [608],[582].

Figure 11.39 shows the 5σ discovery regions for the charged Higgs boson produced in the $pp \rightarrow tbH^\pm$ process with the $H^\pm \rightarrow \tau^\pm \nu_\tau$ decay mode.




Figure 11.39: The 5σ discovery regions for the charged Higgs boson with the $\tau\nu$ decay mode in the m_h^{\max} scenario.

Figure 11.40 shows the 5σ discovery regions for the neutral Higgs boson ϕ ($\phi=h, H, A$) produced in the association with b quarks $pp \rightarrow b\bar{b}\phi$ with the $\phi \rightarrow \mu\mu$ and $\phi \rightarrow \tau\tau$ decay modes. In both Figures the discovery reach was evaluated in the m_h^{\max} scenario with $\mu=200$ GeV/ c^2 (See Section 11.3.1). The discovery reach was evaluated also in the extended m_h^{\max}

scenario (see Section 11.3.1.3 and [251]) with the values of $\mu = -200$ and ± 500 GeV/c^2 . The Figure 11.41 presents the variation of the 5σ discovery potential for the neutral Higgs bosons with the $\tau\tau \rightarrow \mu + \text{jet}$ decay mode. The combination of the effects from supersymmetric radiative corrections and decay modes into supersymmetric particles gives rise to a rather complicated dependence of the discovery contour on μ . This can result in a variation of the discovery region, especially for large M_A and large $\tan\beta$. For the positive values of μ the inclusion of the supersymmetric radiative corrections leads to a shift of the discovery region toward higher values of $\tan\beta$.




Figure 11.41: Variation of the 5σ discovery potential for the neutral Higgs bosons with the $\tau\tau \rightarrow \mu + \text{jet}$ decay mode in the m_h^{max} scenario.

Figure 11.42 shows the 5σ discovery regions for the light, neutral Higgs boson h from the inclusive $pp \rightarrow h + X$ production with with the $h \rightarrow \gamma\gamma$ decay and for the light and heavy scalar Higgs bosons, h and H , produced in the vector boson fusion $qq \rightarrow qqh(H)$ with the $h(H) \rightarrow \tau\tau \rightarrow \ell + \text{jet}$ decay.




Figure 11.42: The 5σ discovery regions for the light, neutral Higgs boson h from the inclusive $pp \rightarrow h+X$ production with the $h \rightarrow \gamma\gamma$ decay and for the light and heavy scalar Higgs bosons, h and H , produced in the vector boson fusion $qq \rightarrow qqh(H)$ with the $h(H) \rightarrow \tau\tau \rightarrow \ell + \text{jet}$ decay.

Chapter 12

Search for Higgs boson in non SUSY models



12.1 Scalar sector of 5D Randal-Sundrum model

The Randall Sundrum model (RS) [87, 626] has recently received much attention because it could provide a solution to the hierarchy problem [551], by means of an exponential factor in a five dimensional nonfactorizable metric. In the simplest version the RS model is based on a five dimensional universe with two four-dimensional hypersurfaces (branes), located at the boundary of the fifth coordinate. By placing all the Standard Model fields on the visible brane all the mass terms, which are of the order of the Planck mass, are rescaled by the exponential factor, to a scale of the order of a TeV. The fluctuations in the metric in the fifth dimension are described in terms of a scalar field, the radion, which in general mixes with the Higgs boson. This scalar sector of the RS model is parametrized in terms of a dimensionless Higgs boson radion mixing parameter ξ , of the Higgs boson and radion masses m_h , m_ϕ and the vacuum expectation value of the radion field Λ_ϕ .

The phenomenology of Higgs boson and radion at LHC has been subject to several studies [627, 628, 629, 630, 631, 632] concentrating mainly on Higgs and radion processes. The Higgs boson and radion detection is not guaranteed in all the parameter space region. The presence in the Higgs radion sector of trilinear terms opens the possibility of $\phi \rightarrow hh$ and $h \rightarrow \phi\phi$ decays. For example, for $m_h = 120 \text{ GeV}/c^2$, $\Lambda_\phi = 5 \text{ TeV}$ and $m_\phi \sim 250\text{-}350 \text{ GeV}/c^2$ the $\text{BR}(\phi \rightarrow hh)$ ranges between 20 and 30%.

The CMS discovery potential is estimated for the decay of the radion in a pair of Higgs bosons, with $\gamma\gamma\bar{b}b$, $\tau\tau\bar{b}b$ and $b\bar{b}b\bar{b}$ final states and for an integrated luminosity of 30 fb^{-1} . The study has been carried out for the radion mass of $300 \text{ GeV}/c^2$ and the Higgs boson mass of $125 \text{ GeV}/c^2$. The sensitivity was evaluated in the (ξ, Λ_ϕ) plane, with systematic uncertainties included.

A detailed description of the analysis can be found in [633]. A brief summary of the analysis and the results is presented below.

12.1.1 The $\phi \rightarrow hh$ analysis with the $\gamma\gamma\bar{b}b$ and $\tau\tau\bar{b}b$ final states.

Signal events $gg \rightarrow \phi \rightarrow hh$ were generated with PYTHIA. The cross sections and branching ratios were evaluated using rescaled NLO cross sections for the SM Higgs boson and a modified HDECAY program. For the radion and a Higgs boson mass points considered ($m_h = 125 \text{ GeV}/c^2$, $m_\phi = 300 \text{ GeV}/c^2$) and for $\Lambda_\phi = 1 \text{ TeV}$ the maximal cross section times branching ratio is 71 fb for $\gamma\gamma\bar{b}b$ final state. For the $\tau\tau\bar{b}b$ final state with the topology considered in the

analysis, one τ lepton decaying leptonically and the other τ lepton decaying hadronically (producing a τ jet), the maximal cross section times branching ratio is 960 fb. This maximal cross section is reached for the radion mixing parameter $\xi = -0.35$.

For the $\gamma\gamma b\bar{b}$ final state the irreducible backgrounds $\gamma\gamma jj$ ($j=u,d,s,g$) (generated with CompHEP) and the $\gamma\gamma c\bar{c}$ and $\gamma\gamma b\bar{b}$ (generated with MadGraph) were studied. The reducible background from γ +three jets and four-jet processes was not evaluated directly, but assumed to be the same as in for the inclusive $h \rightarrow \gamma\gamma$ analysis [6], namely 40 % of the total background after all selection. For the $\tau\tau b\bar{b}$ final state, the $t\bar{t}$, Z+jets, W+jets backgrounds (generated with PYTHIA) and the $b\bar{b}Z$ background (generated with CompHEP) were studied.

The $\gamma\gamma b\bar{b}$ events were required to pass the Level-1 and HLT di-photon trigger. In the off-line analysis two photon candidates with $E_T^{\gamma 1, \gamma 2} > 40, 25$ GeV were required to pass tracker cuts and calorimeter isolation cuts. Events with only two calorimeter jets of $E_T > 30$ GeV and within $|\eta| < 2.4$ were selected. At least one of these jets must be tagged as a b-jet. Finally, the di-photon mass, $M_{\gamma\gamma}$, was required to be in a window of ± 2 GeV/ c^2 , the di-jet mass, $M_{j\bar{j}}$, in a window of ± 30 GeV/ c and the di-photon-di-jet mass, $M_{\gamma\gamma b\bar{b}}$, in a window ± 50 GeV/ c^2 around the Higgs and Radion mass. Figure 12.1 shows the di-jet (left plot) and the di-photon (right plot) mass distribution for the background (open histogram) and the signal of $\phi \rightarrow hh \rightarrow \gamma\gamma b\bar{b}$ (full, black histogram) after all selections except the mass window cuts, and for 30 fb $^{-1}$. The signal is shown for the maximal cross section times branching ratios




Figure 12.1: The di-jet (left plot) and the di-photon (right plot) mass distribution for the background (open histogram) and the signal of $\phi \rightarrow hh \rightarrow \gamma\gamma b\bar{b}$ (full black histogram) after all selections except the mass window cuts with 30 fb $^{-1}$. The signal is shown for the maximal cross section times branching ratios point in $(\xi-\Lambda_\phi)$ plane.

(dashed histogram) and for the signal of $\phi \rightarrow hh \rightarrow \gamma\gamma b\bar{b}$ plus background (solid histogram) after all selections, and for 30 fb $^{-1}$.

The $\tau\tau b\bar{b}$ events were selected by the single electron and muon triggers and by the combined e-plus- τ -jet and the μ -plus- τ -jet triggers. In the off-line analysis a lepton and τ -jet identification was performed. The requirements on the jets were similar to the ones used in the $\gamma\gamma b\bar{b}$ analysis. In addition a cut of the transverse mass of the lepton and missing trans-

verse momentum, $M_T^{l\nu} < 35 \text{ GeV}/c^2$ was applied to suppress the $t\bar{t}$ and W +jets backgrounds. The di- τ -lepton mass was reconstructed using the missing transverse energy as described in Section 5.2.5. The significance of the discovery was calculated using expected number of the signal and background events after the mass window selections: $100 < M_{bj} < 150 \text{ GeV}/c^2$, $100 < M_{\tau\tau} < 160 \text{ GeV}/c^2$ and $280 < M_{\tau\tau} < 330 \text{ GeV}/c^2$. Figure 12.3 (right plot) shows the $M_{\tau\tau b_j}$ distribution for the background (full, gray histogram) and for the signal of $\phi \rightarrow hh \rightarrow \tau\tau b\bar{b}$ plus background (black points with the error bars) after all selections, for 30 fb^{-1} . Fitted curves for the background and the signal plus background are superimposed.




Figure 12.2: Left plot: the $M_{\gamma\gamma b_j}$ distribution for the background (dashed histogram) and for the signal of $\phi \rightarrow hh \rightarrow \gamma\gamma b\bar{b}$ plus background (solid histogram) after all selections for 30 fb^{-1} . Right plot: the $M_{\tau\tau b_j}$ distribution for the background (full gray histogram) and for the signal of $\phi \rightarrow hh \rightarrow \tau\tau b\bar{b}$ plus background (black points with the error bars) after all selections for 30 fb^{-1} . The fitted curves for the background and signal plus background are superimposed. On both plots the signal is shown for the maximal cross section times branching ratios point in $(\xi-\Lambda_\phi)$

The four b-jet final state yields the highest rate for the signal. The maximal cross section times branching ratio at $\Lambda_\phi = 1 \text{ TeV}$ is 10.3 pb , which results in about 3.1×10^5 signal events for 30 fb^{-1} . The effective triggering and selection in the off-line analysis of these events is, however a big challenge due to the huge multi-jet background rate. In fact the remaining background is a few orders of magnitude larger than the signal in the relevant mass range. Techniques can be envisaged to normalize the background directly from a signal free region and predict the number of background events in the signal region. In order to make a 3σ discovery, such extrapolation needs to be performed with a precision of about 0.1%, making this channel essentially hopeless.

The background contribution to the $\gamma\gamma b\bar{b}$ final state can be determined directly from the $\gamma\gamma$ -plus-two-jets data obtained after all selections, except the final mass window cuts on the $M_{\gamma\gamma}$, $M_{j\bar{j}}$ and $M_{\gamma\gamma b\bar{b}}$. The signal-to-background ratio is always less than 10% before the mass cuts are applied. The final cuts on the $M_{\gamma\gamma}$, $M_{j\bar{j}}$ and $M_{\gamma\gamma b\bar{b}}$ introduce a systematic uncertainty on the number of the background events expected after these cuts. This uncertainty is determined by the following factors: the energy scale uncertainty for the photons and jets, and the theoretical uncertainty of the shape of the mass distributions due to the scale and PDF

uncertainties. Figure 12.3 (left plot) shows the 5σ discovery contours for the $\phi \rightarrow hh \rightarrow \gamma\gamma b\bar{b}$ channel for 30 fb^{-1} . The solid (dashed) contour shows the discovery region without (with) the effects of the systematic uncertainties.

For the $\tau\tau b\bar{b}$ final state the background uncertainty due to the experimental selections was estimated to be between 5% and 10% [?]. Figure 12.3 (right plot) shows the 5σ discovery contours for the $\phi \rightarrow hh \rightarrow \tau\tau b\bar{b}$ channel for 30 fb^{-1} . The two contours corresponds to the variation of the background NLO cross sections due to the scale uncertainty. The 5% experimental systematics on the background is taken into account.




Figure 12.3: Left plot: the 5σ discovery contours for the $\phi \rightarrow hh \rightarrow \gamma\gamma b\bar{b}$ channel for 30 fb^{-1} . The solid (dashed) contour shows the discovery region without (with) the effects of the systematic uncertainties (find more explanations in the text). Right plot: the 5σ discovery contours for the $\phi \rightarrow hh \rightarrow \tau\tau b\bar{b}$ channel for 30 fb^{-1} . The two contours corresponds to the variation of the background NLO cross sections due to the scale uncertainty. The 5% experimental systematics on the background is taken into account (see text).

12.2 Search for pair production of doubly charged Higgs boson in the Littlest Higgs model

The main motivation of the Large Hadron Collider (LHC) experiments is to reveal the secrets of electroweak symmetry breaking. If the standard model (SM) Higgs boson will be discovered, the question arises what stabilizes its mass against the Planck scale quadratically divergent radiative corrections. The canonical answer to this question is supersymmetry which implies very rich phenomenology of predicted sparticles in the future collider experiments.

More recently another possibility of formulating the physics of electroweak symmetry breaking, called the little Higgs, was proposed [634, 635, 636]. In those models the SM Higgs boson is a pseudo Goldstone mode of a broken global symmetry and remains light, much lighter than the other new modes of the model which have masses of order the symmetry breaking scale $\mathcal{O}(1) \text{ TeV}$. In order to cancel one-loop quadratic divergences to the SM Higgs mass a new set of heavy gauge bosons W' , Z' with the SM quantum numbers identical to W Z , and a vectorlike heavy quark pair T , \bar{T} with charge $2/3$ must be introduced. Notice that those

12.2. Search for pair production of doubly charged Higgs boson in the Littlest Higgs model 401

fields are put in by hand in order to construct a model with the required properties. However, the minimal model based on the $SU(5)/SO(5)$ global symmetry, the so-called littlest Higgs model [637], has a firm prediction from the symmetry breaking pattern alone: the existence of another $\mathcal{O}(1)$ TeV pseudo Goldstone boson Δ with the $SU(2)_L \times U(1)_Y$ quantum numbers $\Delta \sim (3, 2)$.

Interestingly, the existence of triplet Higgs Δ might also be required to generate Majorana masses to the left-handed neutrinos [638]. Non-zero neutrino masses and mixing is presently the only experimentally verified signal of new physics beyond the SM. In the triplet neutrino mass mechanism [639] the neutrino mass matrix is generated via

$$(m_\nu)_{ij} = (Y_\Delta)_{ij} v_\Delta, \quad (12.1)$$

where $(Y_\Delta)_{ij}$ are the Majorana Yukawa couplings of the triplet to the lepton generations $i, j = e, \mu, \tau$ which are described by the Lagrangian

$$L = i\bar{\ell}_L^c \tau_2 Y_\Delta^{ij} (\tau \cdot \Delta) \ell_{Lj} + h.c., \quad (12.2)$$

and v_Δ is the effective vacuum expectation value of the neutral component of the triplet induced via the explicit coupling of Δ to the SM Higgs doublet H as $\mu \Delta^0 H^0 H^0$. Here μ has a dimension of mass. In the concept of seesaw $\mu \sim M_\Delta$, and the smallness of neutrino masses is attributed to the very high scale of triplet mass M_Δ via the smallness of $v_\Delta = \mu v^2 / M_\Delta^2$, where $v = 174$ GeV.

However, in the littlest Higgs model the triplet mass scale is $\mathcal{O}(1)$ TeV which alone cannot suppress v_Δ . Therefore in this model $\mu \ll M_\Delta$, which can be achieved, for example, via shining from extra dimensions as shown in ref. [640, 641] or if the triplet is related to the Dark Energy of the Universe [642]. In that case $v_\Delta \sim \mathcal{O}(0.1)$ eV while the Yukawa couplings Y_Δ can be large. For the normally hierarchical light neutrino masses neutrino data implies very small Δ decay branching fractions to electrons and $BR(\Delta^{++} \rightarrow \mu^+ \mu^+) \approx BR(\Delta^{++} \rightarrow \tau^+ \tau^+) \approx BR(\Delta^{++} \rightarrow \mu^+ \tau^+) \approx 1/3$. We remind also that v_Δ contributes to the SM oblique corrections, and the precision data fit $\hat{T} < 2 \cdot 10^{-4}$ [643] sets an upper bound $v_\Delta \leq 1.2$ GeV on that parameter.

At LHC Δ^{++} can be produced singly and in pairs. The cross section of the single Δ^{++} production via the WW fusion process [644] $qq \rightarrow q'q' \Delta^{++}$ scales as $\sim v_\Delta^2$. In the context of the littlest Higgs model this process, followed by the decays $\Delta^{++} \rightarrow W^+ W^+$, was studied in ref. [84, 645, 646]. The detailed ATLAS simulation of this channel shows [646] that in order to observe 1 TeV Δ^{++} , one must have $v_\Delta > 29$ GeV. This is in conflict with the precision physics bound $v_\Delta \leq 1.2$ GeV as well as with the neutrino data. Therefore the WW fusion channel is not experimentally promising for the discovery of very heavy doubly charged Higgs.

On the other hand, the Drell-Yan pair production process [647, 644] $pp \rightarrow \Delta^{++} \Delta^{--}$ is not suppressed by any small coupling and its cross section is known up to next to leading order [648] (possible additional contributions from new physics such as Z' are strongly suppressed for any practical purposes). Followed by the lepton number violating decays $\Delta^{\pm\pm} \rightarrow \ell^\pm \ell^\pm$, this process allows to reconstruct $\Delta^{\pm\pm}$ invariant mass from the same charged leptons rendering the SM background to be very small in the signal region. If one also assumes that neutrino masses come from the triplet Higgs interactions, one fixes the $\Delta^{\pm\pm}$ leptonic branching ratios. This allows to test neutrino mass models at LHC.

12.2.1 Search for the final state with four muons

12.2.1.1 Introduction

Here we investigate doubly charged Higgs bosons $\Delta^{\pm\pm}$ pair-produced via the Drell-Yan process, and assume a Branching Ratio of 100% into muons. This provides an almost background free channel.

12.2.1.2 Event Generation

The signal samples are generated using PYTHIA, with doubly charged Higgs bosons pair-produced through the Drell-Yan process. The Higgs bosons are forced to decay into muons. Datasets are produced for several values of the doubly charged Higgs boson mass, ranging from 100 to 800 GeV.

The LO and NLO cross-sections are shown for the signal as a function of the doubly charged Higgs boson mass in Figure 12.4.




Figure 12.4: NLO Cross-section, for $H^{++}H^{--} \rightarrow 4\mu$.

Important backgrounds for this channel with a four muon final state are:

- $t\bar{t} \rightarrow W^+W^-b\bar{b} \rightarrow 2\mu + 2\mu$ (generated with PYTHIA).
- $Zb\bar{b} \rightarrow 2\mu + 2\mu$ (generated with CompHEP [47])
- $Z\bar{Z} \rightarrow 2\mu + 2\mu$ (generated with CompHEP)
- $Z\bar{Z} \rightarrow 2\tau + 2\mu$ (generated with CompHEP)

The contribution of background from $b\bar{b}$ production has also been investigated. $b\bar{b}$ background is the QCD background which yields the highest probability to fake events with multiple muons. It has been found that the $b\bar{b}$ QCD background can be neglected after the online selection and a cut which requires 4 well-reconstructed muons with pseudorapidity $|\eta| < 2.1$ and transverse momentum $p_T > 8$ GeV. The W bosons in the $t\bar{t}$ data sample are forced to decay into electrons, muons and taus. The tau leptons are forced to decay into electrons and

12.2. Search for pair production of doubly charged Higgs boson in the Littlest Higgs model

muons. The Z boson in the $Z\bar{b}b$ sample is generated with $m_{Z/\gamma^*} > 5\text{ GeV}$ and is forced to decay into muons. The Z bosons in the $Z\bar{Z}$ samples are forced to decay into muons with $p_T > 3$ and $|\eta| < 2.5$. The taus in the $Z\bar{Z} \rightarrow 2\tau + 2\mu$ sample decay freely.

On all samples preselection cuts are applied with the following requirements:

- Final state contains two positive and two negative muons
- Transverse momentum $p_T(\mu) > 3\text{ GeV}$ and pseudorapidity $|\eta(\mu)| < 2.4$ for all muons

In the $Z\bar{b}b$ and the $Z\bar{Z}$ sample it is also required that $5\text{ GeV} < m(\mu^+ + \mu^-) < 150\text{ GeV}$ for two of the possible combinations of the muons.

12.2.1.3 Level 1 and HLT selections

The events must be triggered as di-muon events both by the L1 Trigger and the HLT. The p_T threshold for the di-muon HLT is 7 GeV. The L1+HLT efficiency for the signal is $> 99\%$ within uncertainties.

12.2.1.4 Offline reconstruction and selection cuts

The muons are reconstructed by the Global Muon Reconstructor.

The invariant mass of the doubly charged Higgs is reconstructed, by calculating the invariant mass of the two same charge muons with the highest p_T , after all cuts.

Cut on the number of muons

At least 4 muons, with a $p_T > 8\text{ GeV}$ and $|\eta| \leq 2.1$, are required.

Vertex cut

An event, where two or three muons are generated in one collision, and one or two in another, has also to be considered as background to our four muon signal. To suppress this background a vertex cut has been applied.

For each muon in an event the impact point is determined. The impact point is the point of closest approach of the extrapolated muon trajectory to the nominal interaction point. The longitudinal distances Δz_{IPS} between the impact point states of all muons in one event are calculated. The biggest calculated Δz_{IPS} is required to be smaller than 0.05 cm. This is much smaller than the longitudinal size of the luminous region of the LHC beam of about 5 cm. So this cut rejects events with muons from different collision vertices with a probability of roughly 99%.

12.2.1.5 Results

Table 12.1 and table 12.2 show the NLO production cross-section without any forced decay, the cross-section times branching ratio times preselection efficiency and the cross-section times branching ratio times efficiency after each stage of the online and offline event selection. Table 12.1 shows these values for a signal sample and for each of the background samples. Table 12.2 show these values for signal samples with doubly charged Higgs masses 300, 600 and 800 GeV.

Figure 12.5 shows the invariant mass spectrum of the reconstructed $\Delta^{\pm\pm}$ before and after the offline cuts, for $m(\Delta^{\pm\pm}) = 300\text{ GeV}$ and for $m(\Delta^{\pm\pm}) = 600\text{ GeV}$.




Figure 12.5: Invariant reconstructed $\Delta^{\pm\pm}$ mass, (top) after preselection and trigger selection, (bottom) after offline cuts

Table 12.1: Production cross-section (NLO), cross-section times branching ratio times preselection efficiency and cross-section times branching ratio times efficiency after each stage of the online and offline event selection for background events. All values in fb. Errors are statistical only

	$t\bar{t}$	$Zb\bar{b}$	$(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*) \rightarrow 4\mu$	$(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*) \rightarrow 2\mu + 2\tau$
Production cross-section (NLO) [fb]	$840 \cdot 10^3$	$278 \cdot 10^3$	153	212
Preselection: $\sigma \cdot \text{BR} \cdot \epsilon$ [fb]	232 ± 1	289.8 ± 0.9	87.4 ± 0.3	1.63 ± 0.02
Level-1 trigger [fb]	232 ± 1	289 ± 1	87.3 ± 0.3	1.63 ± 0.02
High Level Trigger [fb]	149 ± 1	195 ± 1	69.7 ± 0.3	1.10 ± 0.01
4μ reconstructed ($p_T > 8 \text{ GeV } \eta < 2.1$)[fb]	45.1 ± 0.4	25.1 ± 0.3	18.5 ± 0.1	0.25 ± 0.01
Vertex cut [fb]	22.8 ± 0.3	13.1 ± 0.2	16.9 ± 0.1	0.22 ± 0.01

12.2.1.6 Statistical Interpretation

To interpret the results, the CL_s method [495] is applied, which is based on log-likelihood ratios, calculated for all bins of the invariant mass distribution. CL_s is defined as ratio of the confidence levels for the signal and background hypotheses $CL_s = CL_{s+b}/CL_b$. CL_s can be understood as the probability of excluding an existing signal. $1 - CL_b$ can be understood as the probability for the background distribution to fake a signal.

For high doubly charged higgs boson masses the amount of simulated background events goes to zero. Nevertheless, zero simulated background events do not necessarily mean zero background events in reality.

12.2. Search for pair production of doubly charged Higgs boson in the Littlest Higgs model

Table 12.2: Production cross-section (NLO), cross-section times branching ratio times preselection efficiency and cross-section times branching ratio times efficiency after each stage of the online and offline event selection, for $m_{\Delta^{\pm\pm}} = 300, 600, 800$ GeV All values in fb. Errors are statistical only.

	$m_{\Delta^{\pm\pm}} = 300$ GeV	$m_{\Delta^{\pm\pm}} = 600$ GeV	$m_{\Delta^{\pm\pm}} = 800$ GeV
Production cross-section (NLO) [fb]	19.6	0.909	0.20
Preselection: $\sigma \cdot \text{BR} \cdot \epsilon$ [fb]	17.4 ± 0.3	0.85 ± 0.02	0.190 ± 0.004
Level-1 trigger [fb]	17.3 ± 0.3	0.85 ± 0.02	0.190 ± 0.004
High Level Trigger [fb]	17.1 ± 0.3	0.83 ± 0.02	0.188 ± 0.004
4 μ reconstructed ($p_T > 8$ GeV $\eta < 2.1$) [fb]	13.0 ± 0.2	0.70 ± 0.02	0.158 ± 0.003
Vertex cut [fb]	12.5 ± 0.2	0.67 ± 0.02	0.153 ± 0.003




Figure 12.6: $1 - CL_b$ and CL_s as defined in the Log Likelihood Ratio Method after all selection cuts for an integrated luminosity of 10 fb^{-1}

To estimate the amount of background in this region, empty bins are filled for each background with upper limits to Poisson statistic. Zero background events are compatible with maximal three generated events. Therefore empty bins get filled for each background with three events times the scale factor for an integrated luminosity of 10 fb^{-1} .

The left plot in figure 12.6 shows the $1 - CL_b$ values for different doubly charged Higgs boson masses. For a doubly charged Higgs Boson mass smaller than 650 GeV the signal plus background expectation will exceed the background only expectation by more than 5σ . To claim a discovery, at least three signal events need to be discovered. For a mass of 650 GeV four detectable events remain after all cuts.

The right plot in figure 12.6 shows the CL_s values for different doubly charged Higgs boson masses. If no signal can be detected for an integrated luminosity of 10 fb^{-1} the existence of a doubly charged Higgs Boson in this decay channel can be excluded with 95% confidence up to a mass of 760 GeV.

The ± 1 and ± 2 -sigma bands in figure 12.6 are only for statistical errors.

12.2.1.7 Systematical Uncertainties

The uncertainties on the exclusion limit resulting from systematical errors have yet to be studied in detail, once the detector is running.

The considered backgrounds are also backgrounds to the Standard Model $H \rightarrow ZZ \rightarrow 4\mu$ process. As this process is one of the benchmark processes of the future CMS detector, this backgrounds are studied in detail. The obtained total uncertainty on the background cross section is 1% to 6%. The uncertainty on signal cross section σ_{signal} is 10% to 15%. The uncertainty on the luminosity \mathcal{L} is $\sim 5\%$ for an integrated luminosity of 10 fb^{-1} .

Using a background cross section uncertainty of 6%, a signal cross section uncertainty of 10% and a luminosity uncertainty of 5% the approximated uncertainties on the exclusion mass limit and on the discovery mass limit are:

$$\text{Exclusion Limit} = (760^{+0.5}_{-2}(\sigma_{background}) \pm 10(\sigma_{signal}) \pm 4(\mathcal{L})) \text{ GeV} \quad (12.3)$$

$$\text{Discovery Limit} = (650^{+0.4}_{-0.3}(\sigma_{background})^{+3}_{-0.4}(\sigma_{signal}) \pm 0.2(\mathcal{L})) \text{ GeV} \quad (12.4)$$



12.2.2 Search for the final states with τ leptons

12.2.2.1 Introduction

In this section we discuss the doubly charged Higgs boson pair-production via Drell-Yan process and investigate decays which involve taus and muons. The branching ratios are assumed to be 1/3 for the following three channels: $\Delta^{\pm\pm} \rightarrow 2\mu^{\pm}$, $\Delta^{\pm\pm} \rightarrow \mu^{\pm}\tau^{\pm}$ and $\Delta^{\pm\pm} \rightarrow 2\tau^{\pm}$. The reasoning comes from recent neutrino mixing measurements. As the neutrino mixing matrix and doubly charged Higgs boson decays are directly related then the appropriate branchings can be determined. The above mentioned distribution is the most likely distribution.

12.2.2.2 Event Generation

Signal samples are generated using PYTHIA. Only pair-production through Drell-Yan is assumed (no single production through WW fusion). Datasets are produced from 200 GeV to 600 GeV Higgs mass with 100 GeV increments. In the events produced, the taus can decay both leptonically and hadronically while in analysis we only consider hadronic decays.

Both signal datasets and ZZ datasets have been generated with the aid of CMKIN 5.1.1. Full detector simulation has been performed using OSCAR 3.6.5 in winter mode followed by digitization with ORCA 8.7.4. The full analysis is performed in ORCA 8.9.4. Pileup is not used as it has no effect on muons and tau identification is not affected by presence of low luminosity pile up as shown in [649]. Table 12.3 shows the amount of events generated for the different signal datasets. Table 12.4 shows the amount of events analyzed for background datasets. NLO cross sections used for the background samples can be seen in Table 12.5.

Table 12.3: Number of events generated for signal datasets

$m(\Delta^{\pm\pm})$ in GeV	200	300	400	500	600
Number of events	2762	700	700	900	900

Table 12.4: Number of events analyzed for background datasets.

background	$t\bar{t} \rightarrow 4l$	$Zb\bar{b}$	ZZ	$t\bar{t}Z$
Number of events	1 170 600	666 000	8489	119542

12.2. Search for pair production of doubly charged Higgs boson in the Littlest Higgs model 407

Table 12.5: Background processes cross sections used (in fb)

background	$t\bar{t} \rightarrow 4l$	$Zb\bar{b}$	ZZ	$t\bar{t}Z$
Cross section times BR	$88.4 \cdot 10^3$	$26.2 \cdot 10^3$	212	650

12.2.2.3 Background processes

The important backgrounds which have to be considered for this analysis are as follows:

- $t\bar{t} \rightarrow W^+W^-b\bar{b}$ for this dataset we have used generated samples with PYTHIA, CompHep, AlpGen, TopRex and MadGraph. The W bosons have been
- $t\bar{t}Z \rightarrow W^+W^-Z^0b\bar{b}$ the W and Z bosons are allowed to decay arbitrarily. For this dataset we have used generated samples with CompHep with total integrated luminosity of $184fb^{-1}$.
- $Zb\bar{b}$ where the Z decays to muons or taus. The total integrated luminosity of this dataset is $8fb^{-1}$.
- $(Z^{(*)}/\gamma^*)(Z^{(*)}/\gamma^*)$ where the Z-s are forced to decay leptonically (e, μ, τ). The total integrated luminosity for this dataset is $127fb^{-1}$.

Considering the amount of total available background we use the assumption of total integrated luminosity of $10fb^{-1}$.

12.2.2.4 Event selection

The events must be triggered in HLT (High Level Trigger) with at least one isolated muon present. We do not use L1 trigger explicitly as HLT is only triggered when L1 trigger for a single muon is passed.

In addition to HLT trigger the event will only be used if it is possible to reconstruct the event primary vertex. The vertex is reconstructed using the highest p_{\perp} tracks. If the primary vertex fails to be reconstructed the event is rejected.

The event selection efficiency is dependant on the doubly charged Higgs mass as can be seen in Table 12.6 which in turn is dependent on lepton selection efficiency shown in Table 12.7. Background efficiencies are shown in Table 12.8.

12.2.2.5 Offline reconstruction and selection cuts

The muons are reconstructed using IsolatedGlobalMuonReconstructor. The tau leptons are reconstructed using tau jet candidates and missing transverse energy after selection cuts. The doubly charged Higgs boson invariant mass is reconstructed from the same charge lepton pairs after all selection cuts.

The selection cuts used on muons:

- the transverse momentum must be higher than 50 GeV. For background events 80% of muons have p_{\perp} less than 50 GeV. For 200 GeV signal it is 27% and for higher signal masses it reduces to around 10%. The distribution can be seen on Figure 12.7.
- the distance to primary vertex in z-direction must not exceed 0.03 cm. This does not cut away any muons but limits analysis to leptons coming from the same

primary vertex. At low luminosity it does not cut anything and is just a quality cut for the future use when a single event can have multiple collisions.

The selection cuts used on tau jets:

- for tau jets we consider tau decays which involve 1 or 3 charged tracks. We use ConcreteTauJetWithTracks reconstruction object which implements the tau jet filtering algorithms described in Physics TDR volume 1. Two scenarios are considered. Either 1 or 3 charged tracks in signal cone and no charged tracks in isolation cone or 2 tracks in signal cone and exactly one charged track in the isolation cone.
- the maximum distance to primary vertex in z-direction of any charged track in the tau jet must not exceed 0.2 cm. At low luminosity it does not cut anything and is just a quality cut for the future use when a single event can have multiple collisions.
- the hottest tau jet HCAL tower transverse energy must be higher than 2 GeV. This eliminates high energy electrons which pass through ECAL and confuse the algorithm into considering the electron as a tau jet. This cut eliminates 86% of all electrons considered as tau candidates and only removes 7.5% of real tau jets. The distribution can be seen on Figure 12.8.
- the transverse energy of the tau jet candidate must exceed 50 GeV. It has been chosen to be the same as the cut used on muons.
- no muon track should be in a cone with $\Delta R = 0.3$ constructed around the tau jet candidate. If there is then the candidate is dropped. This eliminates false tau jet candidates which are generated when a charged muon track passes the same region as photons or hadrons. With this cut only a few real tau jets are discarded however most of the false tau jets coming from this misidentification are discarded.

Missing transverse energy (E_T^{miss}) is reconstructed in three steps. As a first step the raw E_T^{miss} is reconstructed from the calorimeters. As step two Type 1 corrections (using the difference of raw reconstructed jets and calibrated jets) are added to the E_T^{miss} . As a final step the muon transverse components are subtracted to get the real missing transverse energy.

Only events with at least four muons or tau jets total are accepted. If the event is accepted a decision is made whether it can be reconstructed. The possible final states are:

- $\Delta^{++}\Delta^{--} \rightarrow 4\mu$ - this channel is investigated in previous subsection.
- $\Delta^{++}\Delta^{--} \rightarrow 3\mu 1\tau$ - this channel is easily reconstructible as there is only one neutrino and it goes the direction of the tau jet.
- $\Delta^{++}\Delta^{--} \rightarrow 2\mu 2\tau$ - this channel can also be reconstructed using the assumption that the neutrinos go the same direction as the tau jets.
- $\Delta^{++}\Delta^{--} \rightarrow 1\mu 3\tau$ - this channel can be reconstructed only with very good E_T^{miss} resolution as it requires an additional assumption that the masses of the two reconstructed Higgs bosons are the same. However the reconstruction is very sensitive to E_T^{miss} accuracy and often the event has to be dropped due to absurd τ -lepton energies. These events are dropped with the invariant mass cut described below.
- $\Delta^{++}\Delta^{--} \rightarrow 4\tau$ - this channel can not be reconstructed and in many cases will not trigger as no muons are found in the event. This also contributes to a lower efficiency at HLT as these events do not pass the single muon HLT trigger.

12.2. Search for pair production of doubly charged Higgs boson in the Littlest Higgs model

Once the event leptons are reconstructed, some additional tasks are performed:

- Z^0 boson reconstruction - if the odd sign pairing gives an invariant mass of 91 ± 5 GeV then these leptons are removed from further use.
- Same charge lepton pairs are reconstructed and only those reconstructed Higgs candidate pairs whose invariant mass difference is within 20% of each other are considered. The difference in background distribution and signal distribution can be seen on Figure 12.8.

12.2.2.6 Efficiencies

Assuming the above mentioned branching ratios the expected theoretical limits on the different final state discovery channels can be investigated. The branching ratio of tau leptons to tau jets is 65%, this gives:

- $\Delta^{++}\Delta^{--} \rightarrow 3\mu 1\tau = 2/9 \text{events} * 0.65 = 14.4\%$.
- $\Delta^{++}\Delta^{--} \rightarrow 2\mu 2\tau = 3/9 \text{events} * 0.65^2 = 14.1\%$.
- $\Delta^{++}\Delta^{--} \rightarrow 1\mu 3\tau = 2/9 \text{events} * 0.65^3 = 6.1\%$.

So the total theoretical limit of discovery when searching with at least one tau jet in the final state gives 34.6% efficiency. Using datasets which have been generated with the above mentioned branching ratios taken into account the efficiencies of the different cuts at different doubly charged Higgs boson masses can be seen in Table 12.6. Taking into account the signal cross section depending on the Higgs boson mass the total number of observable candidates at an integrated luminosity of 10fb^{-1} is given in Table 12.9.

Table 12.6: Trigger and cut efficiencies for signal as a function of the $\Delta^{\pm\pm}$ mass. Total efficiency is the product of the single efficiencies.

$m_{\Delta^{\pm\pm}}$ (GeV)	200	300	400	500	600
HLT trigger	83.7%	86.0%	86.7%	85.8%	88.3%
Primary Vertex	96.9%	98.5%	97.0%	97.5%	98.0%
4 leptons in final state	10.1%	17.2 %	23.6%	24.7%	26.7%
two pairs and at least one τ	44.9%	46.1%	41.7%	53.2%	52.9%
Mass difference	62.5%	77.2%	80.4%	74.3%	63.6%
Total signal efficiency	2.3%	5.1%	6.6%	8.1%	7.7%

Table 12.7: Single muon and tau selection efficiencies and purity.

$m_{\Delta^{\pm\pm}}$ (GeV)	200	300	400	500	600
Single μ selection efficiency	70.7%	82.0%	86.1%	87.2%	89.2%
1 - purity of accepted muons:	0.1%	0.4%	0.8%	0.7%	1.0%
Single τ selection efficiency	36.6%	42.3%	50.6%	53.3%	53.3%
1 - purity of accepted tau jets:	2.2%	2.2%	4.2%	3.6%	3.2%

Table 12.8: Trigger and cut efficiencies for background. Total efficiency is the product of the single efficiencies.

Process	t \bar{t}	t $\bar{t}Z$	ZZ	Zbb
HLT trigger	40.7%	20.3%	40.0%	42.1%
Primary Vertex	99.3%	99.8%	96.7%	98.2%
4 leptons in final state	0.0015%	0.04 %	3.0%	0.0005%
two pairs and at least one τ	-	0.1%	-	-
Mass difference	-	100%	-	-
Total signal efficiency	-	0.0008%	-	-

12.2.2.7 Systematic errors

At the integrated luminosity of $10fb^{-1}$ the cuts implemented above result in almost background free signal. For datasets with integrated luminosity above $10fb^{-1}$ giving zero events ($t\bar{t}, ZZ^*$) we assume the background to be zero events for the purpose of this study. For $t\bar{t}Z$ where we have one event passing all cuts in the dataset we get 0.05 events when scaled with cross section and luminosity. For Zbb where we have background statistics for $8fb^{-1}$ with zero events passing the cuts. We have run the same analysis with p_{\perp} cuts 40 GeV, 30 GeV and 20 GeV with also zero events passing the cuts which confirms our assumption that leptons coming from Zbb are too soft to produce a background for our process. Considering the smallness of the background we assume no background at $10fb^{-1}$ for the following analysis.

The systematic errors used for signal is as follows:

- muon misidentification ($\Delta\mu$) - 1% per muon.
- muon isolation ($\Delta\mu_{isol}$)- 2% per event.
- tau jets ($\Delta\tau$)- 9% per tau jet.
- luminosity ($\Delta\mathcal{L}$) - 5%.
- PDF and scale ($\Delta\sigma$)- 10% (theoretical, not used when we have no background and go to cross section measurement).

As the events are a mixture of different decay modes the total selection efficiency error ($\Delta\epsilon_S$) is calculated per decay channel and then added together using different weights to account for the non-uniform distribution within the dataset

$$\Delta_{3\mu 1\tau} = \sqrt{3\Delta\mu^2 + \Delta\tau^2} = 8.2\%,$$

$$\Delta_{2\mu 2\tau} = \sqrt{2\Delta\mu^2 + 2\Delta\tau^2} = 11.4\%,$$

$$\Delta_{1\mu 3\tau} = \sqrt{\Delta\mu^2 + 3\Delta\tau^2} = 13.9\%,$$

giving

$$\Delta\epsilon_S = \frac{144\Delta_{3\mu 1\tau} + 141\Delta_{2\mu 2\tau} + 61\Delta_{1\mu 3\tau}}{346} = 10.5\%$$

The total systematic error for cross section measurement is then

$$\frac{\Delta\sigma}{\sigma} = \sqrt{\Delta\mu_{isol}^2 + \Delta\mathcal{L}^2 + \Delta\epsilon_S^2} = 13\%.$$

The statistical errors were evaluated constructing the shortest Bayesian confidence interval for the confidence level of 67% [650].

12.2.2.8 Results

The expected number of events and the NLO cross section with expected measurement errors (both statistical and systematic) are given in Table 12.9.

The invariant mass plots for 200, 300, 400, 500 GeV doubly charged Higgs boson searches are shown on Figures 12.9 and 12.10.

12.2. Search for pair production of doubly charged Higgs boson in the Littlest Higgs model 1

Table 12.9: Expected number of events, NLO cross section and expected measurement errors

$m_{\Delta^{\pm\pm}}$ (GeV)	200	300	400	500	600
Expected nr of events at $10fb^{-1}$	26	10	4	2	1
NLO cross section and expected measurement errors (in fb)	$93.9^{+19.3}_{-17.5} \pm 12.2$	$19.6^{+6.6}_{-5.6} \pm 2.5$	$5.9^{+3.4}_{-2.5} \pm 0.8$	$2.2^{+1.9}_{-1.3} \pm 0.3$	$0.9^{+1.3}_{-0.7} \pm 0.1$
Statistical significance	5.37	3.50	2.36	1.69	1.28

The statistical significance and cross section distributions can be seen on Figure 12.11. As no events of background are expected at $10fb^{-1}$ we can detect the doubly charged Higgs boson with a mass of up to ≈ 210 GeV. Reasonable evidence at 3σ level can be seen up to 340 GeV and 95% CL exclusion limit is at 480 GeV.




Figure 12.7: Distribution of μ transverse momentum for background (left) and signal (right).

12.2. Search for pair production of doubly charged Higgs boson in the Littlest Higgs model 13




Figure 12.8: Left: Distribution of hottest HCAL tower p_{\perp} for τ -jet candidates and electrons. Right: Distribution of $m(\Delta_1)/m(\Delta_2)$ for background and signal.




Figure 12.9: Invariant mass plots for $M(\Delta^{\pm\pm})=200$ GeV and 300 GeV.




Figure 12.10: Invariant mass plots for $M(\Delta^{\pm\pm})=400$ GeV and 500 GeV.




Figure 12.11: Left: statistical significance. Right: cross section with statistical errors

12.2. Search for pair production of doubly charged Higgs boson in the Littlest Higgs model




Figure 12.12: Mass reach in dependence to luminosity. 5σ , 3σ levels for discovery are given as well as 95% CL for exclusion if nothing is found by that time.

Chapter 13

Supersymmetry



13.1 Introduction

This chapter presents the results of analyses by which evidence for supersymmetry could be obtained in CMS during the “low luminosity” period of the LHC. After a brief reminder of the main phenomenological features of SUSY in Section 13.2, Section 13.3 is devoted to the outline of the scope of present searches. The emphasis was not on a complete study of a specific point in the parameter space, but rather on covering all relevant signatures by which SUSY might be discovered. For this purpose, a set of test points have been defined, for which a full simulation of the CMS detector was performed, to serve as basis for the analyses. An algorithm allowing the separation of the sparticle decay chains, used in several analyses, is presented in Section 13.4. Sections ?? to 13.12 summarize the searches for SUSY and the reach as a function of luminosity, demonstrating that low mass supersymmetry can be discovered at the LHC with fairly low integrated luminosity for all these signatures in inclusive searches and show the projected reach at the end of the low luminosity run. They are followed by some exclusive studies, mass reconstruction in di-tau final states (Section 13.13), trilepton final states from direct chargino/neutralino production (Section 13.14) and slepton pair production (Section 13.15). A possible violation of lepton number in $\tilde{\chi}_2^0$ decay is studied in Section 13.16. Section 13.17 contains some considerations on the robustness of the considered signatures in scenarios beyond MSUGRA, like for non-universal Higgs masses, and shows that the same signatures would still allow the discovery of supersymmetry. The chapter ends with our conclusion on the CMS reach.

13.2 Summary of Supersymmetry

13.2.1 The MSSM

The Minimal Supersymmetry Model (MSSM) contains the minimal extension of the Standard Model (SM) particle content. Its gauge sector is fully determined by Supersymmetry. But the unknown mechanism for breaking Supersymmetry introduces a large number of free parameters [651] and makes this general model intractable. Therefore, several more constrained models have appeared in the literature. Below, we will focus on a version derived from Supergravity with minimal superpotential and Kähler potential, called MSUGRA, which guarantees universality of gaugino and scalar masses and of trilinear couplings at a high scale. Other SUSY breaking models, like Gauge Mediated Supersymmetry Breaking (GMSB)

or Anomaly Mediated Supersymmetry Breaking (AMSB) have not been included here. R-parity breaking in SUSY is also not considered.

An earlier summary of the potentialities of the CMS experiment at LHC for the discovery of Supersymmetry has been published in 1998 [652]. The potential of the ATLAS experiment for the discovery of supersymmetry was analyzed in [480].

13.2.2 MSUGRA parameters and spectrum

The MSUGRA model of supersymmetry is determined by 5 free parameters defined at the Grand Unification (GUT) scale. If it is assumed that the spontaneous gauge symmetry breaking is induced by radiative corrections, the absolute value of μ is determined from the Z^0 mass. The free parameters are then:

$$m_0, m_{1/2}, A_0, \tan\beta, \text{sign}(\mu) \quad (13.1)$$

They are run down to the electroweak scale by Renormalization Group Equations (RGE) from which the sparticle spectrum, decay branching ratios and production cross sections can be derived.

The gaugino mass parameters M_a at the electroweak scale are approximately :

$$\begin{aligned} M_3 \equiv M_{\tilde{g}} &\simeq 2.7m_{1/2} \\ M_2(M_Z) &\simeq 0.8m_{1/2} \\ M_1(M_Z) &\simeq 0.4m_{1/2} \end{aligned} \quad (13.2)$$

The parameter M_3 determines the gluino mass (after QCD corrections). The masses of neutralinos $\tilde{\chi}_i^0$ ($i = 1 - 4$) and charginos $\tilde{\chi}_i^\pm$ ($i = 1, 2$) are obtained after diagonalizing their mass matrices which are a function of M_1 , M_2 and μ . In the MSUGRA framework, the lightest chargino and the two lightest neutralinos are dominantly gaugino-like with masses close to M_1 and M_2 .

The sfermions of the first two generations have masses given approximately by:

$$\begin{aligned} m_{\tilde{u}_L}^2 &\simeq m_0^2 + 5.0m_{1/2}^2 + 0.35\cos 2\beta M_Z^2 \\ m_{\tilde{d}_L}^2 &\simeq m_0^2 + 5.0m_{1/2}^2 - 0.42\cos 2\beta M_Z^2 \\ m_{\tilde{u}_R}^2 &\simeq m_0^2 + 4.5m_{1/2}^2 + 0.15\cos 2\beta M_Z^2 \\ m_{\tilde{d}_R}^2 &\simeq m_0^2 + 4.4m_{1/2}^2 - 0.07\cos 2\beta M_Z^2 \\ m_{\tilde{e}_L}^2 &\simeq m_0^2 + .49m_{1/2}^2 - 0.27\cos 2\beta M_Z^2 \\ m_{\tilde{\nu}}^2 &\simeq m_0^2 + .49m_{1/2}^2 + 0.50\cos 2\beta M_Z^2 \\ m_{\tilde{e}_R}^2 &\simeq m_0^2 + .15m_{1/2}^2 - 0.23\cos 2\beta M_Z^2 \end{aligned} \quad (13.3)$$

By comparing with the gluino mass, they show that the latter cannot be much larger than the squark mass:

$$M_{\tilde{g}} \lesssim 1.2m_{\tilde{q}} \quad (13.4)$$

This relation (obtained for $m_0 = 0$) is not restricted to the MSUGRA case, as it depends primarily on the α_S contributions to the running down of the mass parameters from the GUT scale.

The masses of the third family scalars are more complicated as the contributions from Yukawa couplings can no longer be neglected and non-negligible off-diagonal elements between left and right states appear (they are proportional to the fermion masses).

13.3 Scope of present searches

13.3.1 Sparticle production and cascade decays

If we assume that Supersymmetry is discovered at the LHC, most likely from fully inclusive studies based on large missing energy and jets, it will be very important to investigate all the typical SUSY signatures to help pin down the underlying model.

If the squarks and/or gluinos are kinematically accessible at the LHC, they are expected to have large production rates. The cross sections for the production of a squark (excluding gluinos) or a gluino at the LHC are displayed in Figure 13.1. The nearly diagonal lines delimit




Figure 13.1: Regions of the m_0 versus $m_{1/2}$ plane showing the production cross-sections and with main squark and gluino decays.

three regions:

- Region 1: in this region, the gluinos are heavier than any of the squarks. The decay chains of the produced sparticles are expected to be

$$\tilde{g} \rightarrow \tilde{q}\bar{q}, \tilde{q} \rightarrow q\chi \tag{13.5}$$

- Region 2: in this region some squarks are heavier, other are lighter than the gluino. Hence, rather complicated decay chains are possible, for instance

$$\tilde{q}_L \rightarrow \tilde{g}q, \tilde{g} \rightarrow \tilde{b}\bar{b}, \tilde{b} \rightarrow b\chi \tag{13.6}$$

as the \tilde{q}_L of the first two generations are expected to be among the heaviest squarks and the \tilde{b}_1 (and \tilde{t}_1) among the lightest.

- Region 3: in this region, the gluinos are lighter than any of the squarks. A typical decay chain is then

$$\tilde{q} \rightarrow \tilde{g}q, \tilde{g} \rightarrow q\bar{q}\chi \quad (13.7)$$

where the gluino gives rise to a three-body decay mediated by a virtual squark.

They will cascade down to the LSP (that we suppose here to be stable). In MSUGRA, the lightest two neutralinos are $\tilde{\chi}_1^0$, which is dominantly Bino-like, and $\tilde{\chi}_2^0$, which is dominantly Wino-like. The \tilde{q}_R then decays almost exclusively directly into $q\tilde{\chi}_1^0$. But the \tilde{q}_L have usually a non-negligible branching ratio to decay via the $\tilde{\chi}_2^0$ or $\tilde{\chi}_1^\pm$. The decay of the $\tilde{\chi}_2^0$ will then provide an excellent signature for the events which can be observed in inclusive searches.

The main decay modes of the $\tilde{\chi}_2^0$, and hence the signatures, are

$$\tilde{\chi}_2^0 \rightarrow \tilde{l}l, \quad (13.8)$$

$$\tilde{\chi}_2^0 \rightarrow \tilde{\nu}\nu, \quad (13.9)$$

$$\tilde{\chi}_2^0 \rightarrow h^0\tilde{\chi}_1^0, \quad (13.10)$$

$$\tilde{\chi}_2^0 \rightarrow Z^0\tilde{\chi}_1^0, \quad (13.11)$$

$$\tilde{\chi}_2^0 \rightarrow l^+l^-\tilde{\chi}_1^0 \quad (13.12)$$

where the last decay is mediated by the exchange of an off-shell Z^0 or \tilde{l} . The first decay corresponds to a gauge interaction coupling a Wino to a slepton-lepton pair and dominates if it is kinematically allowed. When this decay is kinematically forbidden and $m_{1/2}$ is large enough, so that $m(\tilde{\chi}_2^0) - m(\tilde{\chi}_1^0) > m(h^0)$, the next preferred decay is to h^0 . This corresponds to a gaugino-higgsino transition and thus requires a non-zero higgsino component in at least one of the two neutralinos. If also this decay is kinematically forbidden and the neutralino mass difference is sufficient, the $\tilde{\chi}_2^0$ decays to a Z^0 which is suppressed compared to the h^0 decay because it couples to the higgsino component of both neutralinos. When also this decay is kinematically forbidden, direct three-body decays take place. The corresponding regions in the m_0 versus $m_{1/2}$ plane are illustrated for a MSUGRA case in Figure 13.2 (left). The exact boundaries of the areas depend on the assumptions (MSUGRA) and on the value of $\tan\beta$ and the parameter A , but their existence is rather generic. It should be emphasized that the existence of these decay modes is a direct consequence of the gauge structure of the theory and is therefore independent of the model details. Their relative importance at a given SUSY point is, however, model dependent.

In addition to the decays via a $\tilde{\chi}_2^0$, a large fraction of squark decays will proceed via a $\tilde{\chi}_1^\pm$ decay, which may lead to

$$\tilde{\chi}_1^\pm \rightarrow \tilde{l}l, \quad (13.13)$$

$$\tilde{\chi}_1^\pm \rightarrow \tilde{\nu}l, \quad (13.14)$$

$$\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0, \quad (13.15)$$

$$\tilde{\chi}_1^\pm \rightarrow H^\pm\tilde{\chi}_1^0, \quad (13.16)$$

$$\tilde{\chi}_1^\pm \rightarrow l^\pm\nu\tilde{\chi}_1^0, \quad (13.17)$$

where the last decay is mediated by the exchange of an off-shell W , $\tilde{\nu}$ or \tilde{l} . The localization of the chargino decay modes in the $(m_0, m_{1/2})$ plane is illustrated for a MSUGRA case in Figure 13.2 (right).




Figure 13.2: Regions of the m_0 versus $m_{1/2}$ plane with main χ_2^0 decays (left) and main decays of $\tilde{\chi}_1^\pm$ (right).

Further constraints beyond the MSUGRA ones can be imposed, for example the compatibility with the measured relic density. These limit very severely the available parameter space. However, the lack of knowledge of the SUSY breaking mechanism encourages the future experiments to prepare themselves to cope with the broadest possible spectrum of situations. Rather than restricting oneself to a very constrained model, it will be important to understand how to detect departures from the SM in a large variety of topologies and to investigate how to reconstruct the sparticle masses and other SUSY parameters. Of course, there is more information available in the events than just the end points, e.g. momentum asymmetries of the decay leptons, branching ratios and total cross section measurements. These additional informations have so far not been used to a large extent.

13.3.2 Test points for MSUGRA

To cover significantly different experimental signatures, a set of MSUGRA test points have been defined and will be used in the subsequent analyses. First, low mass (LM1 to LM9) test points were chosen to evaluate the sensitivity to SUSY signals in the early period of the LHC but above the Tevatron reach. Then, some high mass test points (HM1 to HM4) near the ultimate reach of the LHC were included.

Their parameters are defined in Table 13.1 and their position in the $(m_0, m_{1/2})$ plane is shown in Figure 13.3. Points LM1, LM2 and LM6 are compatible with WMAP Cold Dark Matter limits in a strict MSUGRA scenario. The other points are not, but can be made compatible with CDM if universality of the Higgs mass parameters is abandoned (NUHM). Quoted branching ratios are from ISASUGRA7.69 [653] (lepton is e or μ). The post-WMAP benchmark points are found in [612], the NUHM points in [654] and the CMS DAQ TDR points in [68].

- **Point LM1 :**
 - Same as post-WMAP benchmark point B' and near DAQ TDR point 4.
 - $m(\tilde{g}) \geq m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is dominant
 - $B(\tilde{\chi}_2^0 \rightarrow \tilde{l}_R l) = 11.2\%$, $B(\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1 \tau) = 46\%$, $B(\tilde{\chi}_1^\pm \rightarrow \tilde{\nu}_l l) = 36\%$
- **Point LM2 :**

Table 13.1: MSUGRA parameter values for the test points. Masses are given in units of GeV/c^2 .

Point	m_0	$m_{1/2}$	$\tan\beta$	$\text{sgn}(\mu)$	A_0
LM1	60	250	10	+	0
LM2	185	350	35	+	0
LM3	330	240	20	+	0
LM4	210	285	10	+	0
LM5	230	360	10	+	0
LM6	85	400	10	+	0
LM7	3000	230	10	+	0
LM8	500	300	10	+	-300
LM9	1450	175	50	+	0
LM10	3000	500	10	+	0
HM1	180	850	10	+	0
HM2	350	800	35	+	0
HM3	700	800	10	+	0
HM4	1350	600	10	+	0




Figure 13.3: Position of the test points in the m_0 versus $m_{1/2}$ plane. The lines in this plane correspond to the assumptions that $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$. The shaded regions are excluded because either the $\tilde{\tau}_1$ would be the LSP or because there is not radiative electroweak symmetry breaking. The regions excluded by the LEP limit on the h^0 or the $\tilde{\chi}_1^\pm$ masses are delineated by dashed lines. The test CMS points are indicated by stars (LM7 and LM10 are outside the boundaries) and the points used in the CMS DAQ TDR by triangles. Also shown are the regions of interest for the decay of the $\tilde{\chi}_2^0$.

- Almost identical to post-WMAP benchmark point I'.
- $m(\tilde{g}) \geq m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is dominant ($\tilde{b}_1 b$ is 25%)
- $B(\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1 \tau) = 96\%$ $B(\tilde{\chi}_1^\pm \rightarrow \tilde{\tau} \nu) = 95\%$

- **Point LM3 :**
 - Same as NUHM point γ and near DAQ TDR point 6.
 - $m(\tilde{g}) < m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is forbidden except $B(\tilde{g} \rightarrow \tilde{b}_{1,2}b) = 85\%$
 - $B(\tilde{\chi}_2^0 \rightarrow ll\tilde{\chi}_1^0) = 3.3\%$, $B(\tilde{\chi}_2^0 \rightarrow \tau\tau\tilde{\chi}_1^0) = 2.2\%$, $B(\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0) = 100\%$
- **Point LM4 :**
 - Near NUHM point α in the on-shell Z^0 decay region
 - $m(\tilde{g}) \geq m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is dominant with $\tilde{g} \rightarrow \tilde{b}_1b = 24\%$
 - $B(\tilde{\chi}_2^0 \rightarrow Z^0\tilde{\chi}_1^0) = 97\%$, $B(\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0) = 100\%$
- **Point LM5 :**
 - In the h^0 decay region, same as NUHM point β .
 - $m(\tilde{g}) \geq m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is dominant with $B(\tilde{g} \rightarrow \tilde{b}_1b) = 19.7\%$ and $B(\tilde{g} \rightarrow \tilde{t}_1t) = 23.4\%$
 - $B(\tilde{\chi}_2^0 \rightarrow h^0\tilde{\chi}_1^0) = 85\%$, $B(\tilde{\chi}_2^0 \rightarrow Z^0\tilde{\chi}_1^0) = 11.5\%$, $B(\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0) = 97\%$
- **Point LM6 :**
 - Same as post-WMAP benchmark point C'.
 - $m(\tilde{g}) \geq m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is dominant
 - $B(\tilde{\chi}_2^0 \rightarrow \tilde{l}_Ll) = 10.8\%$, $B(\tilde{\chi}_2^0 \rightarrow \tilde{l}_Rl) = 1.9\%$, $B(\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1\tau) = 14\%$,
 $B(\tilde{\chi}_1^\pm \rightarrow \tilde{\nu}l) = 44\%$
- **Point LM7 :**
 - Very heavy squarks, outside reach, but light gluino.
 - $m(\tilde{g}) = 678$ GeV, hence $\tilde{g} \rightarrow 3$ -body is dominant
 - $B(\tilde{\chi}_2^0 \rightarrow ll\tilde{\chi}_1^0) = 10\%$, $B(\tilde{\chi}_1^\pm \rightarrow \nu l\tilde{\chi}_1^0) = 33\%$
 - EW chargino-neutralino production cross-section is about 73% of total.
- **Point LM8 :**
 - Gluino lighter than squarks, except \tilde{b}_1 and \tilde{t}_1
 - $m(\tilde{g}) = 745$ GeV, $M(\tilde{t}_1) = 548$ GeV, $\tilde{g} \rightarrow \tilde{t}_1t$ is dominant
 - $B(\tilde{g} \rightarrow \tilde{t}_1t) = 81\%$, $B(\tilde{g} \rightarrow \tilde{b}_1b) = 14\%$, $B(\tilde{q}_L \rightarrow q\tilde{\chi}_2^0) = 26 - 27\%$,
 - $B(\tilde{\chi}_2^0 \rightarrow Z^0\tilde{\chi}_1^0) = 100\%$, $B(\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0) = 100\%$
- **Point LM9 :**
 - Heavy squarks, light gluino. Consistent with EGRET data on diffuse gamma ray spectrum, WMAP results on CDM and MSUGRA [655]. Similar to LM7.
 - $m(\tilde{g}) = 507$ GeV, hence $\tilde{g} \rightarrow 3$ -body is dominant
 - $B(\tilde{\chi}_2^0 \rightarrow ll\tilde{\chi}_1^0) = 6.5\%$, $B(\tilde{\chi}_1^\pm \rightarrow \nu l\tilde{\chi}_1^0) = 22\%$
- **Point LM10 :**
 - Similar to LM7, but heavier gauginos.
 - Very heavy squarks, outside reach, but light gluino.
 - $m(\tilde{g}) = 1295$ GeV, hence $\tilde{g} \rightarrow 3$ -body is dominant
 - $B(\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_4^0) = 11\%$, $B(\tilde{g} \rightarrow tb\tilde{\chi}_2^\pm) = 27\%$
- **Point HM1 :**
 - $m(\tilde{g}) \geq m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is dominant

- $B(\tilde{g} \rightarrow \tilde{t}_1 t) = 25\%$, $B(\tilde{q}_L \rightarrow q\tilde{\chi}_2^0) = 32\%$,
but $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_2^0) = 6\%$, $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_3^0) = 18\%$, $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_4^0) = 9\%$,
- $B(\tilde{\chi}_2^0 \rightarrow \tilde{l}_L l) = 27\%$, $B(\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1 \tau) = 14\%$, $B(\tilde{\chi}_1^\pm \rightarrow \tilde{\nu}_l l) = 37\%$
- **Point HM2 :**
 - $m(\tilde{g}) \geq m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is dominant
 - $B(\tilde{g} \rightarrow \tilde{t}_1 t) = 25\%$, $B(\tilde{q}_L \rightarrow q\tilde{\chi}_2^0) = 32\%$,
but $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_2^0) = 6\%$, $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_3^0) = 20\%$, $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_4^0) = 9\%$,
 - $B(\tilde{\chi}_2^0 \rightarrow \tilde{\tau}_1 \tau) = 78\%$, $B(\tilde{\chi}_1^\pm \rightarrow \tilde{\nu} \tau + \tilde{\tau}_1 \nu) = 13 + 76\%$
- **Point HM3 :**
 - $m(\tilde{g}) \geq m(\tilde{q})$, hence $\tilde{g} \rightarrow \tilde{q}q$ is dominant
 - $B(\tilde{g} \rightarrow \tilde{t}_1 t) = 52\%$, $B(\tilde{q}_L \rightarrow q\tilde{\chi}_2^0) = 32\%$,
but $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_2^0) = 5\%$, $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_3^0) = 20\%$, $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_4^0) = 11\%$,
 - $B(\tilde{\chi}_2^0 \rightarrow h^0 \tilde{\chi}_1^0) = 94\%$, $B(\tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0) = 100\%$
- **Point HM4 :**
 - $m(\tilde{g}) < m(\tilde{q})$, hence $\tilde{q} \rightarrow \tilde{g}q$ is important
 - $B(\tilde{q}_L \rightarrow \tilde{g}q) = 43\%$, $B(\tilde{q}_R \rightarrow \tilde{g}q) = 77 - 93\%$, $B(\tilde{g} \rightarrow \tilde{t}_1 t) = 82\%$,
 - $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_2^0) = 3\%$, $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_3^0) = 22\%$, $B(\tilde{t}_1 \rightarrow t\tilde{\chi}_4^0) = 16\%$,
 - $B(\tilde{\chi}_2^0 \rightarrow h^0 \tilde{\chi}_1^0) = 94\%$, $B(\tilde{\chi}_4^0 \rightarrow h^0 \tilde{\chi}_2^0) = 30\%$, $B(\tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0) = 100\%$



cross sections for the test points are given at NLO and LO from PROSPINO1 in Table

13.4 Hemisphere algorithm for separation of decay chains

13.4.1 Basic idea and goal

In the MSSM, the primary SUSY particles are heavy and tend to be produced with a large Q^2 , whereas the transverse momentum of their decay products with respect to their initial direction is limited by the magnitude of their mass. Moreover, ignoring R_p violation, they are produced in pairs. It may, therefore, be possible to separate the two decay chains by reconstructing the two production directions (in 3D) and collecting the jets and leptons in two clusters according to their "closeness" to these axes. This procedure is inspired by the reconstruction of the thrust or sphericity axis in e^+e^- collisions, except that in hadron collisions two separate axes need to be introduced per event. Note that the back-to-back orientation of the sparticles in the transverse plane cannot be used, as the invisible LSP disturbs significantly the direction of the observable particles.

In hadron colliders like the LHC, the large multiplicity of jets and leptons often lead to a large combinatorial background when trying to reconstruct peaks or to determine end points in effective mass distributions (to reconstruct sparticle masses). Provided the hemisphere algorithm has a large probability to assign correctly the jets to their parents, a reduction of a factor 2 to 4 can be expected in the combinatorial background, which is a significant gain.

The proposed algorithm consists of a recursive method going through the following steps:

- starting off by computing two initial axes (called "seeds" below)

Table 13.2: Cross sections for the test points in pb at NLO (LO) from PROSPINO1.

Point	$M(\tilde{q})$	$M(\tilde{g})$	$\tilde{g}\tilde{g}$	$\tilde{g}\tilde{q}$	$\tilde{q}\tilde{q}$	$\tilde{q}\tilde{q}$	Total
LM1	558.61	611.32	10.55 (6.489)	28.56 (24.18)	8.851 (6.369)	6.901 (6.238)	54.86 (43.28)
LM2	778.86	833.87	1.443 (0.829)	4.950 (3.980)	1.405 (1.013)	1.608 (1.447)	9.41 (7.27)
LM3	625.65	602.15	12.12 (7.098)	23.99 (19.42)	4.811 (3.583)	4.554 (4.098)	45.47 (34.20)
LM4	660.54	695.05	4.756 (2.839)	13.26 (10.91)	3.631 (2.598)	3.459 (3.082)	25.11 (19.43)
LM5	809.66	858.37	1.185 (0.675)	4.089 (3.264)	1.123 (0.809)	1.352 (1.213)	7.75 (5.96)
LM6	859.93	939.79	0.629 (0.352)	2.560 (2.031)	0.768 (0.559)	0.986 (0.896)	4.94 (3.84)
LM7	3004.3	677.65	6.749 (3.796)	0.042 (0.028)	0.000 (0.000)	0.000 (0.000)	6.79 (3.82)
LM8	820.46	745.14	3.241 (1.780)	6.530 (5.021)	1.030 (0.778)	1.385 (1.230)	12.19 (8.81)
LM9	1480.6	506.92	36.97 (21.44)	2.729 (1.762)	0.018 (0.015)	0.074 (0.063)	39.79 (23.28)
LM10	3132.8	1294.8	0.071 (0.037)	0.005 (0.004)	0.000 (0.000)	0.000 (0.000)	0.076 (0.041)
HM1	1721.4	1885.9	0.002 (0.001)	0.018 (0.016)	0.005 (0.005)	0.020 (0.021)	0.045 (0.043)
HM2	1655.8	1785.4	0.003 (0.002)	0.027 (0.024)	0.008 (0.007)	0.027 (0.028)	0.065 (0.061)
HM3	1762.1	1804.4	0.003 (0.002)	0.021 (0.018)	0.005 (0.004)	0.018 (0.019)	0.047 (0.043)
HM4	1815.8	1433.9	0.026 (0.014)	0.056 (0.043)	0.003 (0.003)	0.017 (0.017)	0.102 (0.077)

- associating the objects to one of these axes according to a certain criterion (hemisphere association method)
- recalculating the axes as the sum of the momenta of all the connected objects
- iterating the association until no objects switch from one group to the other

13.4.2 Seeding methods

Two seeding methods have been tested:

1. the first axis is chosen as the direction of the highest momentum object and the second axis as the direction of the object with the largest $p \times \Delta R$ with respect to the first axis, where ΔR is defined as

$$\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2} \quad (13.18)$$

2. the axes are chosen as the directions of the pair of objects which have the largest invariant mass.

13.4.3 Association methods

Three association methods are available. An object is assigned to a given axis \vec{A} when

1. the scalar product $\vec{p} \cdot \vec{A}$ is maximum, which amounts to choosing the smallest angle
2. the hemisphere squared masses are minimum, i.e. object k is associated to the hemisphere with mass m_i rather than m_j if $m_{ik}^2 + m_j^2 \leq m_i^2 + m_{jk}^2$
3. the Lund distance measure is minimum, i.e.

$$(E_i - p_i \cos \theta_{ik}) \frac{E_i}{(E_i + E_k)^2} \leq (E_j - p_j \cos \theta_{jk}) \frac{E_j}{(E_j + E_k)^2}$$

In order to converge to a stable solution, the axes are only updated after a full iteration is performed.

13.4.4 Results

The performance of the hemisphere assignment was tested on events with production of squarks and/or gluinos. Jets were reconstructed using the Iterative Cone method with $\Delta R = 0.5$ and calibrated with the "GammaJet" procedure. They were selected when $E_T > 30$ GeV and $|\eta| < 3.0$. The momentum vectors used were from the Monte carlo parton level objects which matched with the jets and/or leptons. Some of the CMS test points were used, namely LM1 (dilepton final states via \tilde{l}_R), LM5 (with decay of $\tilde{\chi}_2^0$ to $h \rightarrow b\bar{b}$) and LM9 (with dileptons from 3-body decays).

The efficiencies quoted below are the ratio between the correctly assigned MC objects and their total number. The correct hemisphere was chosen as the one for which the axis matched most closely the original squark or gluino, after subtracting from it the unobserved $\tilde{\chi}_1^0$.

The efficiencies of various types of jets for the different algorithms at the test point LM1 are summarized in Table 13.3.

Table 13.3: Efficiencies for test point LM1.

Type of jet	all jets	quark jets	gluon jets	q from \tilde{q}	q from \tilde{g}
Seed 1, Assoc 1	79%	80%	74%	85%	69%
Seed 1, Assoc 2	80%	80%	77%	85%	72%
Seed 2, Assoc 2	81%	81%	78%	85%	72%
Seed 2, Assoc 3	81%	81%	79%	86%	73%

It is seen that all the algorithms behave nearly in the same way, with the combination (seed 1, hemisphere association 1) being slightly worse and (seed 2, hemisphere association 3) slightly better.

The efficiencies obtained for the different test points are listed in Table 13.4 for the different types of jets by using the (seed 2, hemisphere association 3) method. Note that at point LM9 the \tilde{q} undergoes a direct 3-body decay, the \tilde{q} being heavier than the \tilde{g} .

From these tests it can be concluded that quark jets from \tilde{q} have a rather high efficiency, $\geq 85\%$, to be correctly assigned to a hemisphere, whereas the quark jets from a \tilde{g} reach only

Table 13.4: Efficiencies for test points LM1, LM5 and LM9, using the methods Seed 2 and hemisphere association 3.

Point	all jets	quark jets	gluon jets	q from \tilde{q}	q from \tilde{g}
LM1	81%	81%	79%	86%	73%
LM5	77%	77%	74%	87%	70%
LM9	74%	75%	69%	–	76%

$\gtrsim 70\%$. This reflects the fact that the latter jets are much softer, on average, than the jets from the \tilde{q} decay.

The same procedure was also applied to leptons (e or μ). However, due to their tiny mass, the leptons barely “feel” the boost and are sent in any direction. The results were only slightly better than the expectation from random association. Some improvement could be obtained, e.g. for $\tilde{\chi}_2^0 \rightarrow e^+e^-\tilde{\chi}_1^0$, by treating the lepton pair as a single (massive) object. But this introduces some model dependence.

The power of the hemisphere separation can be further illustrated by the search for Higgs at point LM5. The reconstructed jets selected as above are identified as b-jets by a combined b-tagging method when the discriminant variable is > 1.5 . The invariant mass of all combinations of two b-jets is displayed in Figure 13.4 (left). The peak from $h^0 \rightarrow b\bar{b}$ is visible above




Figure 13.4: $b\bar{b}$ invariant mass distributions for (left) all combinations, (centre) combinations in the same hemisphere, (right) combinations in opposite hemispheres.

a large combinatorial SUSY background, mostly due to the production of $b\bar{b}$ and $t\bar{t}$ (directly or from cascade decays). After applying the hemisphere separation method, the 2b invariant mass combinations are separated into the cases where both b-jets are in the same hemisphere (centre), with a clearly visible Higgs peak, and in opposite hemispheres (right), where almost no sign of Higgs remains. Note that these plots were obtained without selection cuts. This method has been used for the Higgs search in Section 13.10 and in other searches.

13.5 Inclusive analysis with missing transverse energy and jets

The missing transverse energy plus multijets final state has been a canonical signature for SUSY searches. This study is a search for the production and decay of gluinos and scalar quarks in ≥ 3 -jet events with large missing transverse energy. The large missing energy originates from the two LSPs in the final states of the squark and gluino decays. The three or more hadronic jets result from the hadronic decays of the squarks and/or gluinos. The full

analysis is presented in section 4.2. The analysis uses the LM1 test-point at which squark and gluino production has a LO cross section of 49 pb. The major Standard Model background components include production of Z +jets with the Z decaying invisibly, W +jets, top-antitop pairs, dibosons, single top and QCD jets. The trigger path used is the missing energy plus jets both at Level-1 and at HLT.

13.5.1 Analysis path and results

Events that are accepted after clean-up preselection requirements, proceed through the analysis path if they have missing transverse energy $E_T^{miss} > 200$ GeV and at least three jets with $E_T \geq 30$ GeV within $|\eta| < 3$. In addition the leading jet is required to be within the central tracker fiducial volume *i.e.* $|\eta| < 1.7$. These requirements directly define the searched for signal signature. The rest of the analysis path is designed based on elimination of the major classes of backgrounds: the QCD production, top-antitop pairs and the W/Z -QCD associated production. In Table 13.5 the path is shown with a remark indicating the reason and aim of each selection step.

Table 13.5: The E_T^{miss} + multijet SUSY search analysis path

Requirement	Remark
Level 1	L1 trigger efficiency parameterization
HLT, $E_T^{miss} > 200$ GeV	trigger/signal signature
primary vertex ≥ 1	primary cleanup
$F_{em} \geq 0.175, F_{ch} \geq 0.1$	primary cleanup
$N_j \geq 3, \eta_d^{1j} < 1.7$	signal signature
$\delta\phi_{min}(E_T^{miss} - jet) \geq 0.3$ rad, $R1, R2 > 0.5$ rad, $\delta\phi(E_T^{miss} - j(2)) > 20^\circ$	QCD rejection
$I_{SO}^{lead\ trk} = 0$	ILV (I) $W/Z/t\bar{t}$ rejection
$f_{em(j(1))}, f_{em(j(2))} < 0.9$	ILV (II), $W/Z/t\bar{t}$ rejection
$E_{T,j(1)} > 180$ GeV, $E_{T,j(2)} > 110$ GeV	signal/background optimization
$H_T \equiv E_{T(2)} + E_{T(3)} + E_{T(4)} + E_T^{miss} > 500$ GeV	signal/background optimization
SUSY LM1 signal efficiency 13%	

A detailed explanation of the analysis path requirements and variables used is given in section 4.2. The global signal efficiency for the analysis is 13% while the signal to background ratio is ~ 26 . The results are shown in Table 13.6 for 1 fb^{-1} .

In summary the major background components and their uncertainties are as follows:

- $t\bar{t}$ uncertainties: 7% E_T^{miss} shape, 22% JES, 13% statistical;
- $Z \rightarrow \nu\bar{\nu}$ +jets, W/Z +jets: 5% Luminosity (direct candle normalization to the data (*cf.* section 4.2));
- QCD: E_T^{miss} 7% shape, 22% JES, 10% statistical.

Table 13.6: Selected SUSY and Standard Model background events for 1 fb^{-1}

Signal	$t\bar{t}$	single t	$Z(\rightarrow \nu\bar{\nu})$ + jets	$(W/Z, WW/ZZ/ZW)$ + jets	QCD
6319	53.9	2.6	48	33	107

Table 13.7: Standard Model background components and uncertainties for 1fb^{-1}

$t\bar{t}$, single top	$Z(\rightarrow \nu\bar{\nu}) + \text{jets}$	$(W/Z, WW/ZZ/ZW) + \text{jets}$	QCD
$56 \pm 11(\text{sys}) \pm 7.5(\text{stat})$	$48 \pm 3.5(\text{all})$	$33 \pm 2.5(\text{all})$	$107 \pm 25(\text{sys}) \pm 10(\text{stat})$

The number of background events per background component and their uncertainties are tabulated in Table 13.7. Based on the Standard Model background estimates and their uncertainties, a 5σ observation of low mass SUSY at LM1 (gluino mass $600\text{ GeV}/c^2$) is achievable with $\sim 6\text{ pb}^{-1}$ in events with large missing energy plus multijets. However it is to be noted that only after $\sim 1.5\text{ fb}^{-1}$ the $W/Z + \text{jets}$ backgrounds, including the invisible decays of the Z boson which constitutes a large irreducible background component, can be reliably normalized using the $Z \rightarrow \mu\mu$ and $Z \rightarrow ee + \text{multijet}$ data candle. The comparison of the signal, total background estimated and its components for the $M_{\text{eff}} \equiv E_{T(1)} + E_{T(2)} + E_{T(3)} + E_{T(4)} + E_T^{\text{miss}}$ is shown in figure 13.5.





Figure 13.5: M_{eff} signal and background distribution in missing energy plus jets SUSY search.


13.6 Inclusive analysis with missing transverse energy, jets, and muons

Searching for the production and decay of new particles in mSUGRA via inclusive final states including muons, high p_T jets, and large missing transverse energy is experimentally interesting due to the ability of the CMS Detector to identify and reconstruct muons with high efficiency [656]. Requiring at least one muon provides a relatively clean experimental signature (complementing searches involving only inclusive jets and missing energy) and has the anticipated advantage of making use of an efficient and well-understood trigger shortly

after the LHC start-up. In this work  **NOTE:** Refer to CMS Note, the fully simulated and reconstructed LM1 mSUGRA point is taken as the benchmark for selection optimisation and study of systematic effects. Even though the study was performed within the context of mSUGRA, this method is not specific to the mSUGRA framework and should apply equally well in other contexts.

13.6.1 Signal Selection and Backgrounds Considered

Because this work is an inclusive study of mSUGRA signatures involving at least one muon accompanied by multiple jets and large E_T^{miss} , several Standard Model processes contribute as sources of background and must be taken into account. Accordingly, the main backgrounds studied in this analysis correspond to QCD dijet (2.8 million events with $0 < \hat{p}_T < 4 \text{ TeV}/c$), top ($t\bar{t}$) production (3.3 million events), and electroweak single-boson production (4.4 million events with $0 < \hat{p}_T < 4.4 \text{ TeV}/c$) and electroweak diboson production (1.2 million events). All backgrounds used in this work are fully simulated and reconstructed. This work uses only leading order cross-sections, consistently for both signal and all backgrounds. Considering NLO k -factors for the signal and background processes do not change the final results significantly.

The CMS trigger system is described in [?], and the current working trigger menu is described in Appendix  **NOTE:** Need OnSel chapter reference. This work uses an event sample which is triggered by either of two HLT triggers: the inclusive isolated single-muon trigger or the isolated dimuon trigger.

To ensure a minimal understanding of the physics objects used in this work, quality criteria are applied to muons and jets. The leading muon is required to have a transverse momentum above $p_T = 30 \text{ GeV}/c$ which ensures that the muon candidate is reconstructed with good efficiency, well above the trigger thresholds. Further, the leading muon is required to be isolated with less than 10 GeV of calorimeter energy within a cone of radius $R = 0.3$, reducing the effects due to fake muons, whilst preserving reasonable efficiency for signal acceptance. Finally, the three leading jets must each have an E_T of at least 50 GeV which guarantees that jets are reconstructed with good efficiency.

The strategy employed in this analysis is to optimise a set of selection cuts based on an objective function which provides a reasonable estimate of the significance to exclude the Standard Model null-hypothesis while explicitly including systematic uncertainties (thus avoiding regions of cut-space which are systematically challenging). This work uses a Genetic Algorithm (GARCON [55]) for the optimisation of cuts and results in: $E_T^{\text{miss}} > 130 \text{ GeV}$, $E_T^{j1} > 440 \text{ GeV}$, $E_T^{j2} > 440 \text{ GeV}$, $|\eta^{j1}| < 1.9$, $|\eta^{j2}| < 1.5$, $|\eta^{j3}| < 3$, $\cos [\Delta\phi(j1, j2)] < 0.2$, $-0.95 < \cos [\Delta\phi(E_T^{\text{miss}}, j1)] < 0.3$, $\cos [\Delta\phi(E_T^{\text{miss}}, j2)] < 0.85$. Assuming 10 fb^{-1} of collected data, this set of cuts would expect to select a total of 2.54 background events from the Standard Model and 311 signal events from the mSUGRA LM1 benchmark signal point.

13.6.2 Results for 10 fb^{-1} using Full Detector Simulation and Reconstruction

After all selection cuts have been applied, the following effects contribute as systematic uncertainties: jet energy scale (10%), jet energy resolution (5%), luminosity measurement (5%), theory (10%), and full simulation versus fast simulation differences (5%, used to determine the analysis reach in mSUGRA parameters in Section 13.6.3). Since this analysis is performed

consistently at leading order, the inclusion of higher order effects involving ISR/FSR is not taken into account. A generator-level comparison of the parton shower method for inclusive $t\bar{t}$ used by PYTHIA [62] with the matrix element calculation for $t\bar{t} + 1\text{jet}$ from CompHEP [47] suggests a $\approx 10\%$ enhancement in the acceptance of $t\bar{t} + 1\text{jet}$ events (generated via the matrix element method) compared with inclusive $t\bar{t}$. The dominant uncertainty (32%) arises from an inability to precisely predict the number of background events, due to finite Monte Carlo simulation statistics. We note that by the time 10 fb^{-1} of data is collected, many of the contributing background processes will be measured from real data, thereby reducing this uncertainty. If one includes the uncertainty due to finite Monte Carlo simulation statistics, the total systematic uncertainty for this work is 36%. Neglecting Monte Carlo simulation statistics, the total systematic uncertainty for this work is 16%.

Table 13.8: Total number of selected events (for 10 fb^{-1}) and significance (“Signif.”) with systematic uncertainties (but excluding uncertainties due to finite Monte Carlo simulation statistics). “SM” represents the total of all Standard Model backgrounds considered.

Sample(s)	Events	Signif.	Sample	Events	Signif.	Sample	Events	Signif.
SM	2.54	–	LM4	246	29.7	LM6	277	32.0
LM1	311	34.1	LM5	165	23.5	HM1	13	4.3

Table 13.8 shows the main results of this study. For the fully simulated low mass mSUGRA point LM1, and assuming 10 fb^{-1} of data, this work selects an expected 311 signal events (with an efficiency of 0.074%) compared with 2.54 expected background events, comprised of $t\bar{t}$ (0.73 events), W +jets (1.56 events), and Z +jets (0.24 events). The separation of signal from background for the different low mass mSUGRA points range in values from 23 to 34 in significance, including systematic uncertainties (but excluding uncertainties related to the limited number of simulated events). Such large values of significance merely indicate that the low mass mSUGRA region will either have been discovered or excluded, long before 10 fb^{-1} of data is collected. We note that shortly after the LHC start-up, the systematic understanding of the CMS detector is expected to be quite different than what is presented in this work, which assumes $L = 10 \text{ fb}^{-1}$. Nevertheless, if one assumes a similar systematic understanding and extrapolates the results of this work to early running, the expected luminosity required to discover the LM1 mSUGRA study point would be $O(0.1) \text{ fb}^{-1}$. Hence, provided a sufficient understanding of detector systematic uncertainties is quickly achieved, low mass SUSY is a prime candidate for possible discovery during the very early running of the LHC.

13.6.3 CMS Reach using Inclusive Muons with Jets and Missing Energy

Since CMS will have either discovered or excluded the lower mass region well in advance of the time required to collect 10 fb^{-1} of data, the selection cuts for 30 fb^{-1} and 60 fb^{-1} are re-optimised using the Genetic Algorithm (GARCON [55]) to select the HM1 mSUGRA point : $E_T^{\text{miss}} > 210 \text{ GeV}$, $E_T^{j1} > 730 \text{ GeV}$, $E_T^{j2} > 730 \text{ GeV}$, $\cos [\Delta\phi(j1, j2)] < 0.95$, $\cos [\Delta\phi(E_T^{\text{miss}}, j1)] < -0.2$, $\cos [\Delta\phi(E_T^{\text{miss}}, j2)] < 0.95$. To estimate the reach for 30 fb^{-1} and 60 fb^{-1} , this same cut-set is applied in both cases and results in an estimated Standard Model background yield of $N_B = 0.25$ for 30 fb^{-1} , and $N_B = 0.49$ for 60 fb^{-1} . In both cases the uncertainty on the background levels is $\approx 71\%$, primarily due to a limited number of simulated events; if one neglects that uncertainty, the systematic uncertainty is $\approx 16\%$.





Figure 13.6: CMS discovery reach contours in the m_0 – $m_{1/2}$ plane using inclusive muons with jets and missing energy for 10 fb^{-1} (lower contour), 30 fb^{-1} (middle contour), and 60 fb^{-1} (upper contour) including systematics.

Fast simulation and reconstruction was also performed in order to scan the plane of universal scalar (m_0) and gaugino ($m_{1/2}$) masses for fixed mSUGRA parameters: $\tan \beta = 10$, $\mu > 0$ and $A_0 = 0$. Points were generated on a coarse grid with $\Delta m_0 = 100 \text{ GeV}/c^2$ and $\Delta m_{1/2} = 100 \text{ GeV}/c^2$, starting from the point $m_0 = 100 \text{ GeV}$, $m_{1/2} = 100 \text{ GeV}$. Figure 13.6 shows the discovery reach of this analysis (contours correspond to a significance value of 5), plotted in the mSUGRA m_0 – $m_{1/2}$ plane. Assuming 10 fb^{-1} of data, CMS can observe SUSY mass scales of over $\approx 1.5 \text{ TeV}/c^2$; assuming 30 fb^{-1} of integrated luminosity, several of the high mass CMS SUSY benchmark points become interesting for possible discovery; and, assuming 60 fb^{-1} of integrated luminosity, CMS is able to reach SUSY mass scales of up to $\approx 2 \text{ TeV}/c^2$.



13.7 Inclusive analyses with same sign dileptons

The topology of two same sign isolated muons, high p_T jets, and large missing transverse energy is interesting as it allows for an efficient suppression of the Standard Model backgrounds, and at the same time allows much of the mSUGRA signal to be retained. Like-sign leptons can result from several signal processes because the gluino, being a Majorana particle, has equal probability of yielding either a positively or a negatively charged lepton in its decay chain. Squark production is another important source of like-sign dileptons, since the squark charge tends to be determined by the valence quarks in the proton-proton collision. The same-sign muon topology capitalizes on the ability of the CMS detector to identify and reconstruct muons with high efficiency [656]. It provides a clean experimental signature and has the extra advantage of an anticipated efficient and well-understood dimuon trigger soon

after LHC start-up. Even though this study  **NOTE:** Refer to CMS Note is performed within the context of mSUGRA, this method is not specific to the mSUGRA framework and applies equally well in other contexts.

13.7.1 Signal selection and backgrounds

Because this work is an inclusive study of mSUGRA signatures involving at least two like-sign muons accompanied by multiple jets and large missing transverse energy, several Standard Model processes contribute as sources of background and must be taken into account. Accordingly, the main backgrounds studied in this analysis correspond to QCD dijet (2.8 million fully simulated events with $0 < \hat{p}_T < 4 \text{ TeV}/c$), top ($t\bar{t}$) production (3.3 million fully simulated events), electro-weak single boson production (4.4 million fully simulated events with $0 < \hat{p}_T < 4.4 \text{ TeV}/c$) and electro-weak diboson production (1.2 million fully simulated events). This work uses only leading order cross-sections, consistently for both signal and all backgrounds.

The dimuon HLT trigger is required for this analysis, which is both natural and ideal ($\approx 98\%$ efficient) for two same-sign muons arising from SUSY processes. Further, to ensure a minimal understanding of the physics objects used in this work, quality requirements are applied to muons and jets. The two leading muons are required to be of the same sign and to each have a transverse momentum above $10 \text{ GeV}/c$, ensuring that the muon candidate is reconstructed with good efficiency, above the symmetric thresholds of $7 \text{ GeV}/c$ in the dimuon trigger. Also this analysis requires at least three jets in the event, all of which are required to have $E_T > 50 \text{ GeV}$, guaranteeing that jets are reconstructed with good efficiency.

In order to select the particular SUSY diagrams responsible for prompt same-sign dimuons, we apply the following criteria. Each reconstructed muon is required to be separated by at least $\Delta R \geq 0.01$ from the other muons. The muon track fit is required to have $\chi_\mu^2 \leq 3$ and the number of hits associated with the muon must be at least 13. Each muon is required to be isolated, both with respect to the tracker and calorimeter. A combined isolation parameter $\text{Iso} = \text{IsoByTk} + 0.75 \times \text{IsoByCalo}$ is used to account for correlations between the tracker and calorimeter isolation variables: $\text{Iso}_{\mu 1} \leq 10 \text{ GeV}$, $\text{Iso}_{\mu 2} \leq 6 \text{ GeV}$.

A genetic algorithm package known as GARCON [55] is used to determine the optimal set of cuts for each mSUGRA benchmark point. An interval for each physics cut-parameter is then defined corresponding to its minimal cut value and the maximum cut value, determined over all different optimal mSUGRA benchmark point cut-sets. The interval for each cut-parameter is then coarsely binned and the significance systematically calculated for each possible cut combination within this reduced sub-space. In addition to *a priori* requiring three jets in the event, the cut-set maximising the significance to discover the lowest significant fully simulated mSUGRA benchmark point is then chosen as the final optimal cut-set: $E_T^{j1} > 175 \text{ GeV}$, $E_T^{j2} > 130 \text{ GeV}$, $E_T^{j3} > 55 \text{ GeV}$, $E_T^{\text{miss}} > 200 \text{ GeV}$.

13.7.2 Results for full detector simulated mSUGRA samples

After all selection cuts have been applied the main systematic uncertainty is due to the absolute jet energy scale, which is estimated to be 15% after 10 fb^{-1} . In addition, jet energy resolution (10%), muon identification efficiency and fake rate (negligible), luminosity (5%), theory (10% ; cross sections, showering, ISR/FSR, etc.) and full simulation versus fast simulation (5% , used to determine the analysis reach in mSUGRA parameters in Section 13.7.3)

have been evaluated. The total systematic uncertainty is 22%. Since this analysis is performed consistently at leading order, the inclusion of higher order effects involving ISR/FSR is not taken into account. A generator-level comparison of the parton shower method for inclusive $t\bar{t}$ used by PYTHIA [62] with the matrix element calculation for $t\bar{t} + 1\text{jet}$ from CompHEP [47] suggests a $\approx 10\%$ enhancement in the acceptance of $t\bar{t} + 1\text{jet}$ events (generated via the matrix element method) compared with inclusive $t\bar{t}$.

Table 13.9 shows the main results of this study. For the fully simulated low mass mSUGRA point LM1, assuming 10 fb^{-1} of data, this work selects an expected 341 signal events (with an efficiency of 0.081%) compared with 1.5 expected background events (comprised of $t\bar{t}$). For other fully simulated low mass mSUGRA points (excluding LM10) and an integrated luminosity 10 fb^{-1} of data, the selection cuts (collectively optimised over all benchmark points) achieve a separation of signal from background with a statistical significance of between 16σ and greater than 37σ , including systematic uncertainties. Such a large significance merely indicates that the low mass mSUGRA region will either have been discovered or excluded, long before 10 fb^{-1} of data is collected. Hence, low mass SUSY is a prime candidate for possible discovery during the very early running of the LHC. The significance for possible discovery of high mass SUSY, represented by the fully simulated HM1 and HM2 points, is more difficult and requires more than 10 fb^{-1} of data.

Table 13.9: Total number of selected events (for $L = 10 \text{ fb}^{-1}$) and significance (“Signif.”) with systematic uncertainties. “SM” represents the total of all Standard Model backgrounds considered.

Sample(s)	Events	Signif.	Sample	Events	Signif.	Sample	Events	Signif.
SM	1.5	–	LM5	61	12.9	LM10	4	2.2
LM1	341	>37.0	LM6	140	21.6	HM1	4	2.2
LM2	94	16.7	LM7	82	15.3	HM2	2	1.2
LM4	90	16.3	LM8	294	35.6			

All fully simulated mSUGRA benchmark points studied in this analysis are very significant even with the inclusion of systematic uncertainties, demonstrating the robustness of the search with respect to the studied systematic effects.

13.7.3 CMS Inclusive Reach

Fast simulation and reconstruction was also performed in order to scan the plane of universal scalar (m_0) and gaugino ($m_{1/2}$) masses for fixed mSUGRA parameters: $\tan \beta = 10$, $\mu > 0$ and $A_0 = 0$. Points were generated on a coarse grid with $\Delta m_0 = 100 \text{ GeV}/c^2$ and $\Delta m_{1/2} = 100 \text{ GeV}/c^2$, starting from the point $m_0 = 100 \text{ GeV}/c^2$, $m_{1/2} = 100 \text{ GeV}/c^2$.

FIGURE: The luminosities and other labels need to be added to the figure itself.

Figure 13.7 shows the 5σ reach of this analysis, including systematic uncertainties, for different integrated luminosities and assuming no re-optimisation of the selection cuts. By the time CMS collects integrated luminosity 30 fb^{-1} , the high mass point HM1 becomes interesting for possible discovery. For comparison, $L = 1 \text{ fb}^{-1}$ and $L = 100 \text{ fb}^{-1}$ are also shown in the figure. Clearly, the systematics for $L = 1 \text{ fb}^{-1}$ will be higher than that assumed in this work, nevertheless these results strongly suggest (provided systematics can be brought un-




Figure 13.7: CMS reach contours (systematic uncertainties included) in the m_0 – $m_{1/2}$ plane for SUSY processes involving two prompt same-sign muons for $L = 1 \text{ fb}^{-1}$ (dot-dashed line), $L = 10 \text{ fb}^{-1}$ (solid line), $L = 100 \text{ fb}^{-1}$ (dashed line). The other mSUGRA parameters are fixed to $\tan \beta = 10$, $\mu > 0$ and $A_0 = 0$. Points corresponding to the full detector simulation and reconstruction are also shown (solid circles).

der control) that most of the low m_0 mSUGRA points are well within reach of CMS during the early running of the LHC.

13.8 Inclusive analyses with opposite sign dileptons

Final states with opposite sign dileptons, originating from the decay $\tilde{\chi}_2^0 \rightarrow \tilde{l}_R l \rightarrow l^+ l^- \tilde{\chi}_1^0$ in the cascade decays of squarks and gluinos provide a clean signature of SUSY with isolated leptons, high p_T jets and missing transverse energy. In addition, the dilepton invariant mass distribution for this decay is expected to have a triangular shape with a sharp upper edge, which renders this signature striking and useful for further characterization of SUSY.

13.8.1 Signal selection and backgrounds

The analysis is performed at the LM1 mSUGRA test-point using Geant-based detailed simulation of the CMS detector (OSCAR 3.6.5). Reconstruction is performed by ORCA 8.7.4. **FIXME:** cite ORCA and OSCAR. The fast CMS simulation and reconstruction (FAMOS 1.4) is used to evaluate the discovery reach in the mSUGRA parameter space.

Signal events were generated by ISAJET 7.69 interfaced to PYTHIA 6.225 at the test point LM1, where the NLO cross section is about 52 pb, dominated by the production of $\tilde{q}\tilde{q}$, $\tilde{g}\tilde{g}$ and $\tilde{q}\tilde{\bar{q}}$. The gluino is the heaviest particle and decays to $\tilde{q}q$. While right squarks decay almost

Table 13.10: Cross section at NLO, selection efficiencies and number of events surviving cuts for signal and background processes.

Process	σ (pb)	Ev. analysed	ε	N_{ev} in 1 fb^{-1}
SUSY (LM1)	52	478k	0.016	853
$t\bar{t}$	830	913k	$1.9 \cdot 10^{-4}$	155
WW + jets	188	197k	$1.4 \cdot 10^{-4}$	26
Z + jets	$5 \cdot 10^3$	606k	$4.8 \cdot 10^{-6}$	24
$DY \rightarrow 2\mu$	$3.97 \cdot 10^3$	916k	$< 1.1 \cdot 10^{-6}$	< 4
$DY \rightarrow 2\tau$	$3.97 \cdot 10^3$	514k	$1.1 \cdot 10^{-6}$	4.5
$Zbb \rightarrow llbb$ ($l = e, \mu, \tau$) $P_{hat}^T > 60 \text{ GeV}/c$	57.4	621k	$8.4 \cdot 10^{-5}$	4.83
$t\bar{t}bb$	3.3	50k	$9.8 \cdot 10^{-4}$	3.2
ZZ + jets	11	37k	$2.4 \cdot 10^{-4}$	2.7
W + jets	$1.5 \cdot 10^5$	1765k	$6.7 \cdot 10^{-9}$	1

directly to the LSP, due to the bino-like nature of the $\tilde{\chi}_1^0$ at Point LM1, left-handed squarks decay to $\tilde{\chi}_2^0$ with a branching ratio $\sim 30\%$.

The SM backgrounds studied consist of $t\bar{t}$, W +jets, Z +jets, WW + jets, ZZ + jets, Zbb (with leptonic decays of the Z boson), Drell-Yan leptonic events and QCD dijet production processes.

The SUSY final state studied contains at least two high- p_T isolated leptons, at least two high- p_T jets and large missing transverse energy. The event selection path includes the following requirements:

- the Level-1 and HLT path that requires a single isolated lepton (muon or electron);
- at least two same-flavour opposite-sign (SFOS) isolated leptons (e or μ) with $p_T \geq 10 \text{ GeV}/c$ and $\Delta R_{ll} \geq 0.2$ and 0.15 for ee and $\mu\mu$, respectively where ΔR_{ll} is the distance of the two leptons in the $\eta - \phi$ space;
- $E_T^{\text{miss}} > 200 \text{ GeV}$;
- at least two jets with $p_T \geq 100$ and $\geq 60 \text{ GeV}/c$ within $|\eta| < 3$.

The isolation of the leptons is obtained requiring the sum of p_T of the tracks in a cone of $\Delta R = 0.25$ around the lepton track to be less than $5 \text{ GeV}/c$. The E_T^{miss} is computed from the vectorial sum of the jets and leptons.

These selection criteria result in 853 signal events (which correspond to 913 dilepton pairs) for a luminosity of 1 fb^{-1} . The Standard Model background consists of 155 $t\bar{t}$ events, 26 events from WW +jets and 24 events from Z +jets (Table 13.10). All other backgrounds have been found to be negligible and amount in total to at most 20 events.

13.8.2 Results for point LM1

The dilepton invariant mass distribution for 1 fb^{-1} is displayed in Figure 13.8 showing a clear dilepton edge structure.

The presence of two SFOS leptons can also be due to other processes. Two leptons can result from independent leptonic decays, for example from two charginos or two W 's. In that case the final state contains as many SFOS leptons as different-flavour opposite-sign (DFOS) ones




Figure 13.8: Invariant mass distribution of $\mu^+\mu^- + e^+e^-$ and $\mu^\pm e^\mp$ pairs at LM1 for 1 fb^{-1} luminosity. The contribution from the $t\bar{t}$ background is also shown.

and with identical distributions. The background to the SFOS contribution is removed by subtracting the DFOS events, which leads to the dilepton mass distribution of Figure 13.9. The $t\bar{t}$ and WW +jets backgrounds are also strongly reduced by the flavour subtraction. The resulting dilepton invariant mass distribution is fitted using a triangular function smeared (for resolution effects) with a Gaussian to extract the end-point related to the kinematics of the decay $\tilde{\chi}_2^0 \rightarrow \tilde{l}_R l \rightarrow l^+ l^- \tilde{\chi}_1^0$. The value obtained from 1 fb^{-1} of integrated luminosity is:

$$M_{ll}^{max} = 80.42 \pm 0.48 \text{ GeV}/c^2 \quad (13.19)$$

to be compared to the expected value of $81.04 \text{ GeV}/c^2$. The signal-to-background ratio at point LM1 is about 4. A statistical significance of 5 sigma, (calculated using S_{cP} [486]), is achieved with 14 pb^{-1} of integrated luminosity. At this luminosity 12.8 signal events are expected with 3.1 Standard Model background events. Therefore this signature is a strong probe for early discovery of low mass supersymmetry.

Systematic uncertainties have been evaluated under the assumption that control data are used for the Standard Model processes. Hence no uncertainties on the theory cross sections, showering, ISR/FSR, are taken into account. The main systematic uncertainty considered is due to the absolute jet energy scale. A $\simeq 7\%$ uncertainty on the jet energy scale for 1 fb^{-1} of data is used while this is expected to be $\simeq 2\%$ after 10 fb^{-1} . After applying the selection cuts this leads to a $\simeq 20\%$ systematic uncertainty on the $t\bar{t}$ background, and to a $\simeq 8\%$ systematic uncertainty on the SUSY signal. The electron energy scale uncertainty, expected to be 1% ,




Figure 13.9: Invariant mass distribution of $\mu^+\mu^- + e^+e^-$ and $\mu^\pm e^\mp$ pairs at point LM1 for 1 fb^{-1} luminosity after subtracting $e^+\mu^-$ and μ^+e^- pairs. The contribution from the $t\bar{t}$ background is also shown.

leads to a systematic uncertainty of less than 1% on the background, and less than 0.3% on the signal. The effect on the signal of the Tracker and Muon System misalignment in the first months of LHC run has also been evaluated. The number of selected dimuon (dielectron) pairs is lowered by about 30% (10%) while the total signal selection efficiency is decreased by about 20%. The measurement of the distribution end-point is affected by about $1 \text{ GeV}/c^2$. The effect of the electron energy scale uncertainty on the dilepton measurement gives a systematic uncertainty of about $0.8 \text{ GeV}/c^2$.

Taking into account the systematic uncertainties on the Standard Model grounds the 5 sigma discovery can be achieved with 17 pb^{-1} of integrated luminosity. **NOTE:** one sentence to indicate that this is not the first 17 pb^{-1} since the systematics are estimated using benchmark numbers for 1 fb^{-1} ; also one sentence on possible residual QCD (i.e. mention the factorization study and that eventually this will be measured in the real data).

13.8.3 CMS inclusive reach

Using the discussed selection path a scan was performed over the mSUGRA parameters in the $(m_0, m_{1/2})$ plane for $\tan\beta = 10$, $A = 0$, $\mu > 0$ to determine the 5σ discovery reach. The observability of the signal over the Standard Model background uses the dilepton estimates before flavor subtraction. The results of the survey are shown for integrated luminosities of and 30 fb^{-1} in Figures 13.10 and 13.10. It is notable that most of the low mass test-points can be discovered with about 1 fb^{-1} .




Figure 13.10: 5 σ discovery reach for $\tan\beta = 10$, $A = 0$, $\mu > 0$ and 1, 10, 30/fb integrated luminosity (statistical uncertainties only)




Figure 13.11: 5 σ discovery reach for $\tan\beta = 10$ taking into account background systematic uncertainties.



9 Inclusive analyses with di-taus

FIXME: background composition, all analysis cuts-of CMSNOTE, figure sizes In this section, $\tilde{\tau}$ production through the $\tilde{\chi}_2^0$ decays in \tilde{q} or \tilde{g} cascades is investigated. The $\tilde{\tau}$ is produced through $\tilde{\chi}_2^0 \rightarrow \tau^\pm \tilde{\tau}^\mp$, where it further decays to $\tau \tilde{\chi}_1^0$ leaving a final state with two taus of opposite sign. The branching fraction of $\tilde{\tau}$ production through $\tilde{\chi}_2^0$ varying with mSUGRA parameters, the analysis is first carried out at large $\tan\beta$, at the LM2 test point which parameters are given in Section 13.3.2, where the $\tilde{\chi}_2^0$ is predicted to decay 95% of the time into $\tau^\pm \tilde{\tau}^\mp$. Results are then generalized to any choice of mSUGRA parameters.

This section studies the opportunity of discovering such a model in the first year of data taking of LHC, at integrated luminosities as low as 0.1 fb^{-1} and up to 30 fb^{-1} . The possibility of measuring the SUSY mass spectra associated to this cascade decay (in particular $\tilde{\chi}_2^0$, $\tilde{\chi}_1^0$ and $\tilde{\tau}$ masses) is investigated in Section 13.13.

13.9.1 Event selection and background studies

For this analysis, 93.5k events (corresponding to an integrated luminosity of 12.6 fb^{-1}) were generated at the LM2 test point with CMKIN using ISASUGRA. Those events were further passed through the full simulation of the CMS detector (using OSCAR) then digitized and reconstructed with ORCA [15]. The same procedure was applied to the Monte Carlo samples used as SM background in this analysis. However, in some cases, where large statistics were required, the fast simulation program FAMOS [?] was used. All Monte Carlo samples used in this analysis are produced with Leading Order Parton Distribution Functions.

Physics processes responsible for W and Z production which final states may contain several taus and jets are considered as potential background sources. In addition, because of its huge cross section ($1.3 \cdot 10^{-4} \text{ mb}$) QCD jet production is also considered. The latter can also represent an important source of fake taus as well as fake missing transverse energy (E_T^{miss}) due to imprecision in jet energy measurement.

13.9.1.1 Event selection using all reconstructed taus

In this analysis, only events passing the JETMET level1 and HLT trigger bits are accepted. The event selection is then carried out using only the E_T^{miss} , the reconstructed taus and jets. In order to increase the sensitivity of the selection both tau's decaying hadronically and leptonically are considered in this section.

The mSUGRA events are selected with the following requirement:

- E_T^{miss} larger than 150 GeV.
This cut removes a large fraction of Standard Model physics background.
- At least two tau candidates are required.
- At least two jets with $E_T > 150 \text{ GeV}$.
This requirement is very aggressive on the LM2 events, however it allows to remove most of the Standard Model background.
- ΔR between any pair of tau's should be smaller than two.
This cut makes use of the fact that in $\tilde{\chi}_2^0$ decays, taus belonging to a same cascade decay will be produced relatively close to each other while in Standard Model physics processes taus as well as Supersymmetric physics processes such as chargino production (producing one tau in each cascade) will tend to be produced in opposite direction. This cut will also reduce the amount of wrong pairing.

Both theoretical and experimental systematic uncertainties are considered in this analysis. The theoretical systematic uncertainty is estimated for the signal according to standard CMS guidelines and involves changing the PDF [353] and varying generator parameters governing both hard process and fragmentation. Each variation leads to the generation of a new LM2 sample which is then simulated and reconstructed using FAMOS and analyzed in the

same way as the main signal samples. Variations in the number of selected events are then taken as systematic uncertainty. The relative theoretical systematic uncertainty on the signal was found to be 11.6%. The experimental systematic uncertainties are coming from the Jet energy scale, the E_T^{miss} and the tau-jet energy scale. These uncertainties are estimated following standard CMS procedure [498] by varying the jet and tau energies by an amount corresponding to their respective energy scales and redoing the analysis. The uncertainty on E_T^{miss} is estimated in a similar way by varying the energy of the jets used to estimate E_T^{miss} within their energy scale. The experimental systematic uncertainty may affect the selection of signal event by 10.7% for low integrated luminosities (smaller than 1 fb^{-1}) but for large integrated luminosities the systematic effect will be less than 3.2%. The experimental systematic uncertainty for the background is 30.2% for integrated luminosities smaller than 1 fb^{-1} and 10.8% for larger integrated luminosities.

At 12.67 fb^{-1} , $N_s = 2735 \pm 273(\text{sys}) \pm 52(\text{stat})$ events from the signal and $N_{bkg} = 938 \pm 103(\text{sys}) \pm 114(\text{stat})$ events from the background survive the selection. 50% of the remaining background is coming from QCD, 39% from $t\bar{t}$ and 11% from W +jets.

To this selection corresponds to a ratio signal over background $S/B = 2.9$. The global efficiency of the selection of the signal is around 3% (of which 88% are SUSY events with at least two taus), while only 0.001% of the background remains after selection. Using S_{cL} significance*, it is possible to estimate that a 5σ discovery can be achieved with only 0.07 fb^{-1} . If the systematic uncertainty on the background is taken into account, a 5σ discovery can be expected with S_{cp} significance [79] of 0.125 fb^{-1} .

13.9.1.2 Event selection using only reconstructed taus decaying hadronically

If only taus decaying hadronically are used in the selection described in 13.9.1.1, both signal and backgrounds are affected differently.

At 12.67 fb^{-1} , $N_s = 1447 \pm 144(\text{sys}) \pm 38(\text{stat})$ events from the signal and $N_{bkg} = 543 \pm 60(\text{sys}) \pm 112(\text{stat})$ events from the background survive the selection. 70% of the remaining background is coming from QCD, 20% from $t\bar{t}$ and 10% from W +jets. To this selection corresponds to a ratio signal over background $S/B = 2.6$. The global efficiency of the selection of the signal is around 1.5% (of which 88% are SUSY events with at least two taus), while only 0.0006% of the background remains after selection. This time, using S_{cL} a 5σ discovery is achieved with only 0.125 fb^{-1} . If the systematic uncertainty on the background is taken into account, a 5σ discovery can be expected with S_{cp} significance of 0.26 fb^{-1} .

13.9.2 Discovery potential of mSUGRA with di-taus final states

A scan of the mSUGRA $(m_0, m_{1/2})$ parameters plane is performed in order to delimit the mSUGRA parameter region where SUSY could be discovered with this analysis. Because the analysis focuses on di-tau final states and since the respective branching ratio to di-taus and to other leptons from SUSY may vary by large amounts in the mSUGRA parameter space, allowing large contamination from leptons into di-taus final states the scan is performed using only hadronic tau decays as described in 13.9.1.2

This scan is achieved by generating many mSUGRA samples varying m_0 and $m_{1/2}$ values so that the entire region of the plane $(m_0, m_{1/2})$ below $m_0 < 1500 \text{ GeV}$ and $m_{1/2} < 800 \text{ GeV}$

* $S_{cl} = \sqrt{2(N_s + N_b) \ln(1 + N_s/N_b) - 2N_s}$

is covered. These samples were generated with ISASUGRA 7.69 then simulated and reconstructed with FAMOS and analyzed in the same way as the LM2 sample. The resulting number of events surviving the selection were used to estimate the significance at each point of the mSUGRA parameter plane. Two types of significance are estimated here, S_{cl} which accounts only for statistical effects and S_{cp} which accounts for both statistical and systematic effects on the background. The resulting 5σ contours over the mSUGRA ($m_0, m_{1/2}$) parameter plane obtained with S_{cl} for several integrated luminosities between 0.1 and 30 fb^{-1} are shown in figures 13.12 and 13.13 for $\tan\beta = 10$ and $\tan\beta = 35$, respectively. Results obtained with S_{cp} are shown in figures 13.14 and 13.15. The region where a 5σ discovery is possible is somewhat shrunken, especially for the very early measurement at 0.1 fb^{-1} as a precise knowledge of the jet scale and of the measurement of the E_T^{miss} will still be limited. However, a large region is still accessible with larger integrated luminosities.




Figure 13.12: Inclusive di-tau analysis discovery potential for mSUGRA between 0.1 and 30 fb^{-1} for $\tan\beta = 10$ including only statistical effects.




Figure 13.13: Inclusive di-tau analysis discovery potential for mSUGRA between 0.1 and 30 fb^{-1} for $\tan\beta = 35$ including only statistical effects.




Figure 13.14: Inclusive di-tau analysis discovery potential for mSUGRA between 0.1 and 30 fb^{-1} for $\tan\beta = 10$ where both statistical and systematic effects are taken into account.




Figure 13.15: Inclusive di-tau analysis discovery potential for mSUGRA between 0.1 and 30 fb^{-1} for $\tan\beta = 35$ where both statistical and systematic effects are taken into account.

13.10 Inclusive analyses with Higgs

This section describes the potential of the CMS experiment to discover a light supersymmetric Higgs boson (h^0) produced at the end of a cascade of supersymmetric particles starting with the strong production of squarks (\tilde{q}) and gluinos (\tilde{g}). Because of the cascade production mechanism, the events can be efficiently triggered using inclusive SUSY triggers such as $\text{jet} + E_T^{\text{miss}}$, and the dominant $h^0 \rightarrow b\bar{b}$ decay mode of the Higgs boson can be exploited.

This analysis focuses on a full CMS detector simulation and event reconstruction at the mSUGRA point LM5, defined in Section 13.3.2. The total SUSY cross section at this parameter point is about 7.75 pb at NLO.

All SUSY channels leading to a light Higgs boson in the final state have been taken into account. The signal events are characterized by at least two b-tagged jets, an important missing transverse energy (E_T^{miss}) and multiple hard jets. This signature allows to suppress the majority of the $b\bar{b}$ background due to SM processes (mainly top pair production $t\bar{t}$, $W^\pm + \text{jets}$, $Z^0 + \text{jets}$).

13.10.1 Signal selection and backgrounds

This analysis has been developed based on the CMS reconstruction software ORCA version 8.7.4 [15]. The two main algorithms used for the signal reconstruction are the jet reconstruction algorithm (the Iterative cone algorithm with a cone size of 0.5 radians and the GammaJet calibration) and the b-tagging algorithm (Combined b-tagging algorithm).

A first rejection of the Standard Model backgrounds happens at the online trigger stage. The L1 and the High Level Trigger (HLT) efficiencies for the signal and background have been evaluated. The trigger path used for this analysis consists of the L1 and HLT Jet + E_T^{miss} stream. This particular trigger is already an important tool in rejecting Standard Model backgrounds, for example it rejects 96 % of the $t\bar{t}$ background while keeping 79 % of the signal events.

In order to further remove the SM background events and reduce the SUSY background, a number of offline selection cuts are applied: a minimal number of four jets with a transverse energy above 30 GeV is required, of which at least two are b-tagged with high quality (i.e. a b-tag discriminator greater than 1.5).

The mean b-tagging efficiency is found to be 50 % with a mistagging rate of about 1.6 %, for u, d, s quarks and gluons, and 12 % for c quarks. The mean b jet energy originating from the Higgs decay is approximately 70 GeV, corresponding to a b-tagging efficiency of about 50 % at this energy. This means that approximately 25 % of the signal events will pass the double b-tag criterion.


Other variables have been identified in order to improve the signal over background ratio, in particular for the most problematic $t\bar{t}$ background: the E_T^{miss} , the first, second and third highest jet P_t . The selection requires a $E_T^{\text{miss}} > 200$ GeV, the highest jet p_t in the event > 200 GeV/ c , the second highest jet p_t in event > 150 GeV/ c , the third highest jet p_t in event > 50 GeV/ c .

Next, in order to select the b-jet pair coming from the Higgs decay, two methods are used. First, the Hemisphere separation technique (see section 13.4) is applied to identify two groups of jets in the detector, each group associated with an initial squark and/or gluino cascade. Af-

ter that, the b-jet pairing is done only in each of these groups separately, reducing the number of possible combinations by a large factor. In addition, as the Higgs is relatively heavy, its decay products have an important boost leading to a small angle $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2}$ between the two b jets. Therefore, in case of multiple possible combinations inside one hemisphere, the pair with the smallest ΔR value within $\Delta R < 1.5$ is chosen. This procedure gives an efficiency of around 40 % and strongly suppresses the combinatorial background.

The full selection chain leads to a signal efficiency of about 8 % for all SUSY channels yielding a Higgs. The global rejection factor for $t\bar{t}$ events, including the rejection made by the $\text{Jet} + E_T^{\text{miss}}$ trigger, is close to $4.6 \cdot 10^4$. No Z +jets, W +jets nor QCD events from the full simulation sample pass the previously described series of cuts.

13.10.2 Results at LM5 and systematics

The resulting invariant mass distribution, after the selection cuts described above, is shown in Figure 13.16. The  not corresponds to the expected statistics equivalent to 1 fb^{-1} of inte-




Figure 13.16: Invariant mass distribution for 1 fb^{-1} .

grated luminosity. A peak around $116 \text{ GeV}/c^2$ is visible. The main background is due to the remaining SUSY background events and some $t\bar{t}$ events.

A fit was performed representing the background by a fifth order polynomial and approximating the Higgs signal by a Gaussian. The r.m.s of the Gaussian has been fixed to 18 GeV, which is the Higgs mass resolution as estimated using the Monte Carlo truth. In real data, this number will be determined from studying b-rich samples such as $t\bar{t}$. The results of the fit, for the equivalent of 1 fb^{-1} of data, are the following : the Higgs mass is found to be $(112.9 \pm 6.6) \text{ GeV}/c^2$ and the fraction of signal in the distribution is evaluated to be 0.28 ± 0.08 . The significance S_{CL} , directly extracted from the fraction of signal in the histogram, is found to be 4.5. A significance of 5 should be achieved with approximately 1.5 fb^{-1} lumi-

osity (equivalent to approximately 2 months of data taking at low luminosity).

For 1 fb^{-1} , the jet energy scale and E_T^{miss} uncertainties have been estimated assuming a linear evolution from $\pm 15\%$ to $\pm 5\%$ for low energy jets (below 50 GeV) and then fixed at $\pm 5\%$ for higher energy jets. As the E_T^{miss} is computed from the jets, a correction on the jet energy is automatically propagated to its estimation. The effects are about 15% on the SUSY event selection and 17% on the $t\bar{t}$ event rejection respectively. The impact on the Higgs mass measurement have been estimated to be $\pm 7.5 \text{ GeV}/c^2$; on α , the effect is ± 0.04 .

Another systematic uncertainty is introduced by the misalignment of the tracker. Both the short and long term misalignment scenarios have been investigated. The short term misalignment corresponds to a displacement of the tracker (strips/pixels) = $(100 \mu\text{m} / 10 \mu\text{m})$, while the long term misalignment takes the following shift of the tracker (strips/pixels) = $(20 \mu\text{m} / 10 \mu\text{m})$ into account. The misalignment of the tracker reduces the track reconstruction resolution, which results in a reduced b-tagging efficiency and which in its turn causes a reduced signal event selection efficiency. The long term misalignment scenario results in a drop of the signal selection efficiency of ($\sim 10\%$) compared to the case of an aligned detector; for the short term misalignment case, the reduction is ($\sim 17\%$). No effect on the position/width of the Higgs mass peak was observed.


Finally, the systematics due to the choice of the background fit function has been estimated to be small (by changing the background function to a third, fourth, sixth or a seventh order polynomial): $\pm 0.3 \text{ GeV}/c^2$ on the Higgs mass and ± 0.01 on α .

The final result including all the previously discussed systematics for 1 fb^{-1} of integrated luminosity is then $112.9 \pm 6.6 \text{ (stat)} \pm 7.5 \text{ (syst)} \text{ GeV}/c^2$ for the Higgs mass and $0.28 \pm 0.08 \text{ (stat)} \pm 0.04 \text{ (syst)}$ for the signal fraction.

13.10.3 CMS reach for inclusive Higgs production

After establishing the visibility of the signal for the LM5 point, a scan was performed in the $(m_0, m_{1/2})$ plane in order to determine the region where a 5σ discovery could be made with 2, 10 and 30 fb^{-1} .

First, an effective cross section ($\sigma \times BR(h0)$) was used (calculated with PROSPINO and ISAS-UGRA) to obtain an estimate of the reach. Using this first estimate, 40 points were chosen for which the full spectrum was calculated and a fast simulation was performed with FAMOS [?]. The same selection criteria as for LM5 point were applied, and the number of Higgs signal and background events was determined; the same significance definition (S_{CL}) was used in order to determine the 5-sigma contours. Comparing the ORCA/FAMOS results at LM5, the significances obtained with both programs were found to agree well.

The result of the scan is displayed in the reach plot in Figure 13.17. Although for 1 fb^{-1} the sensitivity remains below 5σ everywhere, a sizeable region of the $(m_0, m_{1/2})$ plane, up to 1100 (1600) GeV in m_0 and 600 (650) GeV in $m_{1/2}$, can be covered with 10 (30) fb^{-1} . With 2 fb^{-1} of integrated luminosity, a small region of the  can already be probed. The plot assumes $\tan \beta = 10$, $A_0=0$, and a positive sign of μ .

13.11 Inclusive SUSY search with Z^0




Figure 13.17: Higgs discovery reach in SUSY cascades for 2, 10 and 30 fb^{-1} .

13.11.1 Topology of the signal

SUSY processes leading to final states with Z^0 can be easily detected in CMS using the Z^0 decays into opposite sign same flavor (OSSF) lepton pair. The detection of SUSY in the mSUGRA framework through the decay $\tilde{\chi}_2^0 \rightarrow Z^0 + \tilde{\chi}_1^0$ is the scope of this study. The mSUGRA test-point LM4 with the following parameters: $m_0 = 210$ GeV, $m_{1/2} = 285$ GeV, $A_0 = 0$, $\text{sgn}\mu = +1$, $\tan\beta = 10$ is chosen. The $\tilde{\chi}_2^0$ is produced mainly through the cascade decays of gluinos ($M_{\tilde{g}} = 695$ GeV) and squarks (mainly the \tilde{b}_1 with $M_{\tilde{b}_1} = 601$ GeV.) The decays of the second neutralino are following the $\tilde{\chi}_2^0 \rightarrow Z^0 + \tilde{\chi}_1^0$ chain with a large branching ratio ($\sim 100\%$). The signal events are characterized by large missing E_T (due to the undetectable LSP) and the OSSF lepton pair from Z^0 . The analysis details can be found in CMS NOTE 2006/11. **NAME: SOPHIA.**

The main Standard Model backgrounds originate from the production of one or more Z^0 bosons in association with jets as well as $t\bar{t}$. In addition SUSY events contain dileptons that do not originate from the above neutralino decay chain and large missing transverse energy. These events are considered as signal for SUSY detection but as background for the $\tilde{\chi}_2^0$ detection. The following backgrounds were considered in this study: dibosons ($ZZ + j$, $ZW + j$, $WW + j$), inclusive top ($t\bar{t}$) and Z +jets. The signal events were generated interfacing ISAJET 7.69 with PYTHIA. Unless otherwise stated all events are fully simulated and analyzed using the OSCAR CMS detector simulation and ORCA reconstruction packages. The




to leading order (NLO) cross sections of the relevant processes are shown in Table 13.11. **NAME: table needs cleaning**

13.11.2 Event Selection

The following requirements are imposed in order to efficiently select the signal and reject the background events. All criteria were chosen so that the final SUSY search significance estimator S_{c1} [657][96] for 10 fb^{-1} integrated luminosity is maximized. Very similar requirements maximize also significance estimator S_{L2} [96] used in the case of 1 fb^{-1} integrated luminosity. The effect of the selection requirements to the signal and to each background sample separately can be seen in Table 13.11 for 10 fb^{-1} integrated luminosity.

- Events are required to pass the HLT dielectron or dimuon triggers.
- An e^+e^- or $\mu^+\mu^-$ pair with lepton $p_T > 17 \text{ GeV}$ for electrons and $p_T > 7 \text{ GeV}$ for the muons (as per L1 trigger requirements). Each lepton is required to be within $|\eta| < 2.4$.
- The OSSF lepton pair invariant mass is required to be consistent with the Zz mass, i.e. $81 \text{ GeV} < M_{ll} < 96.5 \text{ GeV}$. The reconstructed masses for the e^+e^- and the $\mu^+\mu^-$ pairs and the mass requirements are shown in Figures 13.18a and b respectively.

 This cut reduces backgrounds not involving a Z^0 ($t\bar{t}$, $WW+j$) and the sample of




Figure 13.18: Reconstructed masses for a) e^+e^- and b) $\mu^+\mu^-$ pairs for the background and for the signal (shaded) events. SUSY events not involving $\tilde{\chi}_2^0$ are considered signal. The vertical lines denote the imposed mass requirement.

SUSY events not involving $\tilde{\chi}_2^0$.

- The missing transverse energy E_T^{miss} is required to be greater 230 GeV. This requirement reduces all backgrounds as seen in Fig. 13.19 a. It allows, however, for enough signal and background events in order to maintain good statistics both for 1 fb^{-1} and for 10 fb^{-1} integrated luminosity.
- The angle $\Delta\phi$ between the two leptons of the lepton pair that reconstructs the mass of Z^0 is required to be less than 2.65 rad. The $\Delta\phi$ distribution is shown in Fig. 13.19 b for signal and background. This requirement targets the remainder of the $t\bar{t}$ and the $WW+j$ backgrounds that survived the E_T^{miss} requirement.

1.3 Results and Systematic Uncertainties




Figure 13.19: E_T^{miss} (a) and $\Delta\phi$ between the two leptons (b) for background (black line) and signal (shaded) events. SUSY events not involving $\tilde{\chi}_2^0$ are considered signal. The vertical lines denote the E_T^{miss} and $\Delta\phi$ requirements.

Table 13.11: Number of events for signal ($\tilde{\chi}_2^0 \rightarrow Z^0 + \tilde{\chi}_1^0$, $Z^0 \rightarrow e^+e^-, \mu^+\mu^-$) and background before and after selection criteria for 10 fb^{-1} .

	LM4 with $\tilde{\chi}_2^0$	LM4 no $\tilde{\chi}_2^0$	ZZj	ZWj	WWj	$t\bar{t}$	Zj 85_150	Zj 150_250
σ NLO (pb)	0.664	17.4	15.5	51.5	270	830	102	14.7
10 fb^{-1}								
total events	6640	173.8 K	155 K	515 K	2.7 M	8.3 M	1.02 M	147 K
L1+HLT	6032	81.7 K	12.6 K	24.4 K	174 K	973 K	387 K	74.7 K
OSLP	4489	7147	9124	14.7 K	26.3 K	268 K	281 K	49.5 K
M_{ll}	3773	804	6999	11.5 K	2406	23.1 K	212 K	37.2 K
E_T^{miss}	1420	306	32	24	70	149	O(1)	44
$\Delta\phi_{ll}$	1289	264	31	22	47	61	O(1)	35

FIXME: needs cleaning and to appendix for the significances; when Albert is done with “Significance” appendix **FIXME:** Give signal/background, total signal efficiency

After the application of the above criteria and for 10 fb^{-1} integrated luminosity we have 1553.0 ± 33.2 SUSY events and 196.5 ± 40.3 Standard Model background events. The significance estimator for the discovery of SUSY is then:

$$S_{c1_SUSY} = \frac{N_{\text{signal}}}{\sqrt{N_{\text{bgr}}}} = 110.8 \quad (13.20)$$

S_{c1_SUSY} does not change significantly if we consider the statistical error on the number of events. If the discovery of $\tilde{\chi}_2^0$ is the aim in the mSUGRA scenario, then the SUSY events not involving a $\tilde{\chi}_2^0$ are considered background and $S_{c1} = 60.1$.

In the case of 1 fb^{-1} integrated luminosity, due to the small number of remaining bgr events

we use Eqn. 13.21 [657]

$$S_{L2} = \sqrt{2 \ln(Q)} \quad \text{with} \quad Q = \left(1 + \frac{N_{signal}}{N_{bgr}}\right)^{N_{obs}} \exp(-N_{signal}) \quad (13.21)$$

Optimizing for this significance estimator changes only the E_T^{miss} requirement which becomes $E_T^{\text{miss}} > 215$ GeV but with negligible effect to the significance which is $S_{L2,SUSY} = 21.5$ for detection of SUSY. For the detection of $\tilde{\chi}_2^0$ the above requirements yield $S_{L2} = 14.7$.

When LHC will start running a lot of the uncertainties will be controlled from data. In this analysis relevant uncertainties are the lepton P_t resolution and the E_T^{miss} uncertainty. The lepton P_t resolution ($\sim 3\%$) introduces an uncertainty of 2.7% in the number of background events. The dominant systematic, however, is the E_T^{miss} uncertainty which is estimated to ($\sim 5\%$) and which introduces a 20% uncertainty in the number of $t\bar{t}$ bgr events. Taking into account systematics the significance for SUSY detection becomes $S_{cl,SUSY} = 83.5$ for 10 fb^{-1} and $S_{L2,SUSY} = 17.1$ for 1 fb^{-1} .

13.11.4 CMS reach for inclusive Z^0 search

We have scanned the mSUGRA $m_0, m_{1/2}$ parametric space in order to see if the above analysis can reveal new physics. The test points were taken at high density in the area where the Z^0 has high production cross section (especially due to the decay $\tilde{\chi}_2^0 \rightarrow Z^0 + \tilde{\chi}_1^0$). This is an almost horizontal band in the $m_0 - m_{1/2}$ plane between $m_{1/2} \sim 240$ GeV and $m_{1/2} \sim 340$ GeV. There were also points taken at higher and lower $m_{1/2}$ values, because there is an excess of lepton pairs created due to SUSY processes. These may have invariant mass close to the Z^0 mass and pass analysis cuts assisting in the detection of SUSY. For each point there were produced 2000 events with an OSLP close to the Z^0 mass. The events were generated interfacing ISAJET 7.69 with PYTHIA 6.227 and they were simulated, reconstructed and analysed using the FAMOS fast simulation package **VIXIME: FAMOS reference**. Systematic uncertainties were taken into account. The 5σ significance contour is shown for integrated luminosities of 10 fb^{-1} and 1 fb^{-1} in Fig. 13.20.





Figure 13.20: The 5σ significance contours for 10 fb^{-1} and 1 fb^{-1} (dashed line) integrated luminosities and taking into account systematic uncertainties.

12 Inclusive analyses with top

 **ME:** fix me for style, language, length and figures- also for the conclusion-

 **AME:** Are comments from Oliver included?

The supersymmetric partner of the top quark in most of the supersymmetric scenarios is the lightest squark. Finding evidence of its existence can be a clear signature for supersymmetry. In the main part of the allowed $m_0 - m_{1/2}$ plane, the stop can decay to a top plus a neutralino. This neutralino can be either the LSP ($\tilde{\chi}_1^0$) or a heavier neutralino which decays in turn to a LSP which appears as missing transverse energy (E_T^{miss}). Hence in the final state there is at least a top quark plus large E_T^{miss} .

The search for top was tuned on test point LM1, where the stop decays according to

$$\tilde{t}_1 \rightarrow t\tilde{\chi}_2^0 \rightarrow t\tilde{l}_R \rightarrow tll\tilde{\chi}_1^0 \quad (13.22)$$

giving rise to a final state which also contains two leptons. Although this analysis consists primarily in a search for an excess of top quarks from any SUSY origin with respect to its SM production, it was also optimized for the selection of events where the top results from the production of \tilde{t} .

13.12.1 Top quark and lepton reconstruction and identification

Electrons and muons are requested to have $p_T \geq 5 \text{ GeV}/c$ and $\eta \leq 2.5$.

Electrons are separated from jets by requiring that the ratio of energy deposited in the HCAL to the ECAL $H_oE \leq 0.1$, the absolute difference in η between the electromagnetic cluster in the ECAL and the associated track $\Delta\eta \leq 0.006$ and the energy weighted spread of the electron shower in η be $\sigma_{\eta\eta} \leq 0.015$.

Leptons were required to be isolated, namely that the ratio of p_T of the lepton to the p_T sum of other particles inside a cone of size $\Delta R = 0.1$ around the lepton track be greater than 2. Jets were reconstructed from ECAL and HCAL towers using an Iterative cone algorithm with cone size $\Delta R = 0.5$ and were selected if their uncalibrated transverse energy $E_T \geq 30 \text{ GeV}$ in the acceptance of $\eta \leq 2.5$. Their energy was calibrated using corrections from photon-jet balancing studies.

In this analysis only hadronic decays of the top quark were considered. A kinematic fit with constraints is utilized to find the best combination of jets to make the top quark. Since the purpose of this analysis is not to measure the top quark mass, its known value was used to constrain the invariant mass of the system of three jets. Among these three jets, one and only one must be tagged as a b-jet and the other two were constrained to be consistent with a hadronically decaying W . The fit then consisted in minimizing the χ^2 as a function of the three jet energies and imposing the top and W mass constraints. The solution was obtained by an iterative method based on Lagrange multipliers. As several combinations may lead to a convergent fit for a given event, only the combination with the best χ^2 was kept, with the additional requirement that its χ^2 probability was greater than 0.1.

13.12.2 Signal selection and backgrounds

All events were fully simulated with the CMS program OSCAR [14], digitized with low luminosity pileup and reconstructed with ORCA [15].

The signal events consisted of an inclusive SUSY sample at the test point LM1 (see Section 13.3.2), where the total cross section at NLO is about $52pb$. Top quarks are found in the decay of \tilde{t} , but other important sources exist, e.g. $\tilde{b} \rightarrow t\tilde{\chi}_1^\pm$. At an integrated luminosity of 1 fb^{-1} , the total SUSY production amounts to 52000 events, out of which 8375 contain a top quark.

The main backgrounds, generated with PYTHIA 6.225 [62], consist of $t\bar{t}$, $WW + jets$, $WZ + jets$ and QCD. In addition, single top generated with TopReX 4.11 [35] and $W + jets$ generated with ALPGEN V2.0 [152] were considered.

The selection of SUSY events containing a top quark was based on the following criteria:

- L1T: every event must pass the first level of the Trigger (L1T) cuts corresponding to "Jet/Met" (a jet with $E_T > 88$ and $E_T^{\text{miss}} > 46$ GeV).
- HLT: events were required to pass High level Trigger (HLT) cuts (a jet with $E_T > 180$ and $E_T^{\text{miss}} > 123$ GeV).
- ≥ 4 jets with $E_T^{\text{raw}} \geq 30$ GeV and $\eta \leq 2.5$
- ≥ 1 b-jet with $E_T^{\text{raw}} \geq 30$ GeV and $\eta \leq 2.5$
- $E_T^{\text{miss}} \geq 150$ GeV to suppress $t\bar{t}$ and other SM backgrounds
- a convergent fit with $P(\chi^2) \geq 0.1$
- $\Delta\Phi$ between the fitted top and $E_T^{\text{miss}} \leq 2.6$ rad to suppress semileptonic $t\bar{t}$ events
- ≥ 1 isolated lepton (e or μ) with $p_T \geq 5$ GeV and $\eta \leq 2.5$ to suppress QCD background
- $\Delta R(l, j) \geq 0.2$ to suppress semileptonic decays of b-quarks **VIXME: Is this cut used???**

These criteria were simultaneously optimized to reject SM backgrounds and to maximize the ratio of events with a top quark at generator level, called SUSY(with top), to events without top at generator level, called SUSY(no top).

The effect of the cuts is shown in Table 13.12. As a result of the selection, the signal events remaining for a 1 fb^{-1} luminosity consist of 38 events SUSY(with top) and 17 events SUSY(no top). The remaining backgrounds are 5 events from $t\bar{t}$. The resulting distributions of E_T^{miss} of the fitted top mass are displayed in Figure 13.21.




Figure 13.21: (left) Distributions of E_T^{miss} and (right) fitted top mass after all selection criteria are applied.

Table 13.12: Effect of different cuts on different samples. In every row, the number of the remaining events after that cut is shown. “No.of.used.events” shows the number of events used in this analysis, “NEve(Nor.xsec)1 fb^{-1} ” is the same number after normalizing to the cross section times 1 fb^{-1} and “wT/noT” means $\frac{SUSY(withTop)}{SUSY(noTop)}$.

cut	SUSY (withTop)	SUSY (noTop)	ttInc	WW	ZW	Single t	wT/noT
x-sec(pb) NLO	52		830	269.91	51.5	250	-
No.of.used.events	494261		1674500	305000	70000	100000	-
NEve(Nor.xsec)1 fb^{-1}	8375	43625	830000	269910	51500	250000	0.19
L1T (Jet/Met)	6269	33582	75806	18498	598	10875	0.19
HLT (Jet/Met)	5070	29427	14430	4733	142	1750	0.17
MET \geq 150 GeV	4183	25677	4930	2312	99	653	0.16
$n_{bj} \geq 1$	3457	14388	3718	792	32	355	0.24
$n_j^{b \text{ or } light} \geq 4$	1789	4576	769	25	0	33	0.39
A convergent Fit	1335	3062	557	12	0	28	0.44
χ^2 probability >0.1	105	69	56	0	0	5	1.52
$\Delta\phi < 2.6$	79	52	12	0	0	5	1.51
$n_l > 0$	38	17	5	0	0	0	2.19

13.12.3 Results at point LM1

The significance of a discovery was computed from statistical uncertainties only using the formula


$$Significance = 2 \times (\sqrt{S+B} - \sqrt{B}) \quad (13.23)$$

where the number of signal events, S , is the sum of SUSY(with top) and SUSY(no top) and B represents the sum of all SM backgrounds. Using this formula, the integrated luminosity required to make a discovery at point LM1 with a significance of 5 amounts to $\sim 210 \text{ pb}^{-1}$.

Many systematic uncertainties (cross section, showering, ISR/FSR, ...) will be rendered very small by using real data. The main uncertainties remaining will be the absolute jet energy scale (estimated to 5% for jets and MET in 1 fb^{-1}), which leads to 5.1% from jets and 18.3% from MET in the $t\bar{t}$ sample and the b-tagging efficiency estimated to 8% for 1 fb^{-1} . Adding them in quadrature yields a total systematic uncertainty of 21%, considered common to all backgrounds. It is seen that this remains negligible compared to the statistical uncertainty.

13.12.4 CMS reach for inclusive top search

The CMS fast simulation, FAMOS, was used to find the reach of CMS in this channel in $m_0, m_{1/2}$ plane. In total 36 points have been tried. The ntuples were generated by using the cms-official ISAPYTHIA. The NLO cross sections were derived by PROSPINO [658].

 Figure 13.22 shows the 5σ reach in $m_0, m_{1/2}$ plane with 1, 10 and 30 fb^{-1}




Figure 13.22: The 5σ reach in $m_0, m_{1/2}$ plane with 1, 10 and 30 fb^{-1} .

13.13 Final states with di-taus

In this section the determination of the sparticle masses using invariant mass distributions in the di-tau final state are analysed. The selection of the events is the same as presented in Section 13.9.

13.13.1 Extraction of mSUGRA mass spectra from the measurement of the end points of invariant mass distributions.

Using the kinematics of the successive two body decays in $\tilde{q} \rightarrow q\tilde{\chi}_2^0 \rightarrow q\tau\tilde{\tau} \rightarrow q\tau\tau\tilde{\chi}_1^0$, it is possible to express the mass of the sparticles involved in that cascade as a fully resolved system of equations which depends only on the end-point of the invariant mass distributions obtained by combining the leptons and quark-jets observed in the final state.

However, the tau-lepton always decays, producing at least one undetected neutrino. Therefore, instead of observing a triangle-shaped distribution like for the di-lepton invariant mass distribution of chapter 13.8, where the end-point coincides with the maximum of the distribution, the absence of the neutrino smears the resulting mass distribution to lower values. Even though the end-point of the distribution remains unchanged, it now lies at the tail of a gaussian-like distribution.

The $\tilde{\chi}_2^0$ cascade always produces a pair of opposite charge τ 's, therefore signal samples are obtained by combining opposite charge tau pairs to the two most energetic jets of the event. In these two jets 75% the quark produced by the decay of the \tilde{q} to $\tilde{\chi}_2^0$, due to the fact that the \tilde{q} is much heavier than the $\tilde{\chi}_2^0$. This large number of tau's and jets will be responsible for a high combinatorial background. A good description of this combinatorial background, more particularly the tail, is essential for extracting the true end-points. The combinatorial background are then estimated by taking same sign tau pairs to reproduce the combinatorial background associated to the opposite sign invariant di-tau mass and by combining all tau pairs to jet taken among the 2 most energetic jet of a previous event selected randomly to insure that jet and tau's are uncorrelated..

Five invariant mass and their associated combinatorial background distributions are then

ined ($M(\tau\tau), M(\tau\tau Jet), M(\tau_1 Jet), M(\tau_2 Jet)^\dagger, M(\tau_1 Jet) + M(\tau_2 Jet)$).




Figure 13.23: Di-tau invariant mass distributions.




Figure 13.24: Difference between di-tau invariant mass distribution and combinatorics fit together with log-normal fit.




Figure 13.25: $\tau_1 Jet + \tau_2 Jet$ invariant mass distributions.




Figure 13.26: Difference between $\tau_1 Jet + \tau_2 Jet$ invariant mass distribution and combinatorics fit together with log-normal fit.

The distributions of combinatorial background are first fitted. Then, the resulting fit parameters are used together with a Log-normal distribution, which gives a good description of the tail of the true distributions, to fit the distributions of the signal. Since it is possible to express the log-normal distribution as a function of the end-point, the end-point can be extracted directly from the fit.

Di-tau invariant mass and $M(\tau_1 Jet) + M(\tau_2 Jet)$ are fitted first, the three other invariant mass distributions are built using only candidates found to have values for the two previous distributions below the measured end-points. Then, they are fitted using the same procedure. The sparticle masses are evaluated by solving the system of four equations giving the end-points as a function of the sparticle mass [659].

In the event, several solutions are possible for the SUSY mass spectrum (as it is the case here, where two valid solutions exist), the choice is made by comparing the measured $M(\tau_1 Jet) + M(\tau_2 Jet)$ end-point, E_5 , value to the one computed with the sparticle masses found by solving the systems of equations.

The most probable mass hypothesis is then chosen as the one for which E_5 computed for each

[†] τ_1 is defined in a tau-pair as the one which maximizes the invariant mass formed by its association with a jet; $M(\tau_1 Jet) < M(\tau_2 Jet)$

mass spectrum is the closest to the measured one. The measured end-point was found to be 780 ± 20 GeV while the calculations for case 1 and case 2 mass hierarchy yield to 815 ± 26 GeV and 765 ± 30 GeV respectively (table ??). This corresponds to χ^2 's of 2.3 and 0.5 for case 1 and 2 respectively. The second hypothesis, which corresponds to the LM2 mass hierarchy gives a results compatible with the measured end-point value. This method works relatively well with large statistics, however at 10 fb^{-1} , it is more difficult to distinguish between the two cases as the measured end-point E_5 can be found further away from the measured one and may have errors compatible with both cases.

Three main systematic uncertainties are considered ,the jet scale and tau-jet scales as well as systematics uncertainties arising from the extraction procedure.

Results obtained with that method are shown in table 13.14 for 40 fb^{-1} , together with LM2 generated sparticle masses. They are found to be in good agreement with the theoretical values. They show that it is possible to measure the SUSY mass spectra and in particular $\tilde{\tau}$ mass with a precision less than 30 GeV. Using a 40 fb^{-1} LM2 sample, it is possible to measure the SUSY mass spectra and in particular $\tilde{\tau}$ mass with a precision of 30 GeV.

Table 13.13: End-point obtained with the lognormal fit together with sparticle masses measured with the end-point technique for LM2 for integrated luminosities around 40 fb^{-1} .

End-points (GeV)	case 1 (GeV)	case 2 (GeV)
$m(\tau_1 \tau_2)^{\max} = 95 \pm 3 \pm 3$	$M(\tilde{\chi}_1^0) = 213 \pm 14$	$M(\tilde{\chi}_1^0) = 147 \pm 23$
$m(\tau_1 Q)^{\max} = 559 \pm 11 \pm 9$	$M(\tilde{\chi}_2^0) = 337 \pm 17$	$M(\tilde{\chi}_2^0) = 265 \pm 10$
$m(\tau_2 Q)^{\max} = 298 \pm 7 \pm 5$	$M(\tilde{\tau}) = 310 \pm 17$	$M(\tilde{\tau}) = 165 \pm 10$
$m(\tau_1 \tau_2 Q)^{\max} = 596 \pm 12 \pm 10$	$M(\tilde{q}) = 839 \pm 19$	$M(\tilde{q}) = 763 \pm 33$
$E_5^{\text{meas}} = 780 \pm 20 \pm 10$	$E_5^{\text{calc}} = 815 \pm 26$	$E_5^{\text{calc}} = 765 \pm 30$

Table 13.14: sparticle masses measured with end-point method for LM2 together with theoretical value

	LM2 benchmark point	
	measured	theory
$M(\tilde{\chi}_1^0)$ (GeV)	$147 \pm 23(\text{stat}) \pm 19(\text{sys})$	138.2
$M(\tilde{\chi}_2^0)$ (GeV)	$265 \pm 10(\text{stat}) \pm 25(\text{sys})$	265.5
$M(\tilde{\tau})$ (GeV)	$165 \pm 10(\text{stat}) \pm 20(\text{sys})$	153.9
$M(\tilde{q})$ (GeV)	$763 \pm 33(\text{stat}) \pm 58(\text{sys})$	753-783 (light \tilde{q})



13.14 Direct $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ production in tri-leptons

The exclusive triplepton final state appears in $pp \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ channel with subsequent three body decays of the second neutralino, $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 ll$, and chargino, $\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 W^* \rightarrow \tilde{\chi}_1^0 l \nu$; or via sleptons in two body decay, $\tilde{\chi}_2^0 \rightarrow \tilde{l} \rightarrow l \tilde{\chi}_1^0 l$, and $\tilde{\chi}_1^\pm \rightarrow \tilde{l} \rightarrow l \tilde{\chi}_1^0 \nu$, $\tilde{\chi}_1^\pm \rightarrow \nu \tilde{l} \rightarrow \nu \tilde{\chi}_1^0 l$. The final signatures are two Opposite-Sign Same-Flavor (OSSF) leptons (e, μ) from the neutralino $\tilde{\chi}_2^0$ decay plus any lepton from the chargino $\tilde{\chi}_1^\pm$. Jets are expected to be only due to gluon state radiation or pile up events. In spite of the escaping $\tilde{\chi}_1^0$, the MET is relatively small

at low $m_{1/2}$ and is comparable with the one of SM backgrounds, especially for three body decays at large m_0 . The invariant mass of the OSSF dileptons exhibits a particular shape with a kinematic end point M_{ll}^{max} that depends upon the event topology, see section 13.3.

13.14.1 Signal selection and backgrounds

The trilepton cross section σ_{3l} was calculated with ISASUGRA 7.69 and PYTHIA 6.225 (CTEQ5L) at LO, the K_{NLO} factor calculated with PROSPINO is in the range of 1.30-1.25 (for $m_{\tilde{\chi}_2^0} = 150 - 300 \text{ GeV}/c^2$) [660]. The σ_{3l} drops rapidly with the neutralino mass $m_{\tilde{\chi}_2^0} \sim 0.8m_{1/2}$, $\sigma_{3l} \sim m_{1/2}^{-4}$. This study is restricted to the low $m_{1/2}$ region, where σ_{3l} contributes, for instance, $\sim 0.5\%$ to the total SUSY cross section at $m_0 > 1000 \text{ GeV}$. The three body decays are dominant in this $m_0, m_{1/2}$ region, except for $m_0 < 150 \text{ GeV}$ and $\tan\beta \leq 20$. The kinematic end point in the invariant mass is approximately $M_{ll}^{max} \sim 0.42 * m_{1/2} - 18.4 \text{ GeV}/c^2$ (at $m_0 \sim 1000 \text{ GeV}/c^2$), thus moving into the Z-peak region at $m_{1/2} > 250 \text{ GeV}/c^2$ where the SM background is high. Among the CMS benchmark points in this region, LM9 ($m_{1/2} = 175, m_0 = 1450, \tan\beta = 50, A_0 = 0$) has the largest cross section, ~ 3700 events are produced for 30 fb^{-1} , and it was used as a reference.

The main background results from the DY, Z+jets, $t\bar{t} \rightarrow WbWb, ZW, ZZ, Wt$ +jets, WW+jets, W+jets and inclusive SUSY channels. For all backgrounds, except ZW and ZZ, some leptons are coming from jets, mostly $b \rightarrow l + j$. The background events have been produced with PYTHIA (+TopRex, Alpgen) and their cross section corrected to NLO. The Z and W bosons are forced to decay leptonically to $e, \mu, \tau \rightarrow e, \mu$. The DY and Z+jets cross section is large ($\sigma_{DY, Zj} \sim 10nb$) and events were preselected by requiring three leptons with $P_t > 5 \text{ GeV}/c$ and $|\eta| < 2.4$ at the generator level. The full data samples of 30 fb^{-1} for the LM9 test point and backgrounds are simulated with the fast (*FAMOS.1.3.2*) simulations verified with the smaller samples produced with the full simulation (*ORCA.8.7.3*). The low luminosity pile-up was included.

All events were required to pass L1 and HLT triggers. The main data streams for LM9 are the dimuons (74%) and dielectrons (25%). The trigger efficiency is $E(L1+HLT)=86+91\%$ for LM9 and is increasing for larger $m_{1/2}$ where the leptons become harder. In the off-line selection, at least three isolated leptons in $|\eta| < 2.4$ and $P_T^{\mu, e} > 10 \text{ GeV}/c$ have been required for each event. The leptons have been reconstructed using standard reconstruction algorithms. Electrons and muons were requested to be isolated, i.e. other tracks may contribute to $\sum P_T < 1.5 \text{ GeV}/c$ inside a cone of $\Delta R < 0.3$. Moreover, for muons the energy deposit in calorimeters should be $E_T < 5 \text{ GeV}$ in a cone of $\Delta R < 0.3$. In addition, to be identified as electron particles were requested to satisfy quality criteria based on a likelihood function, $eleId > 0.65$. The muons and electrons reconstruction efficiency in ORCA was found to be 78% ($P_T^\mu > 5 \text{ GeV}/c$) and 66% ($P_T^e > 10 \text{ GeV}/c$) respectively. The jets have been reconstructed by the iterative cone algorithm with the seed energies $E_T^{seed} > 0.5 \text{ GeV}$ in the cone $\Delta R < 0.5$. The MET was reconstructed from the ECAL and HCAL towers. Since the MET for the signal events is relatively small and its reconstruction at low energy scale is limited by the E_T resolution, by the MET cut is drastically reduced in comparison to other SUSY studies.

The reconstructed events have been selected in two steps. First, sequential cuts have been applied: 1) No central jets with corrected energy $E_T > 30 \text{ GeV}$ in $|\eta| < 2.4$, 2) Two OSSF isolated leptons (e, μ) in $|\eta| < 2.4$ with $P_T^\mu > 10 \text{ GeV}/c, P_T^e > 17 \text{ GeV}/c$ and the di-lepton invariant mass below the Z peak $M_{ll} < 75 \text{ GeV}$. 3) The third lepton is with $P_T^{\mu, e} > 10 \text{ GeV}/c$ in

$|\eta| < 2.4$. The evolution of statistics and the efficiencies of the selection cuts are presented in Table 13.15.

Table 13.15: Evolution of signals and background statistics expected for 30 fb^{-1} .

channel	$N_{ev} 30\text{fb}^{-1}$ ($\sigma \times BR$ [pb])	L1+HLT	No Jets	2 OSSF+l OSSF $M_{ll} < 75\text{GeV}$	NN _{LM9}
LM1	2640 (0.088)	1544 (58%)	864 (56%)	70 (8%)	17 (24%)
LM7	1540 (0.051)	1250 (82%)	738 (59%)	91 (12%)	57 (62%)
LM9	3700 (0.125)	2896 (78%)	1740 (60%)	239 (14%)	158 (68%)
SUSY	$4 \cdot 10^5$ (13.1 ^{NLO})	$2.5 \cdot 10^5$ (63%)	$1.8 \cdot 10^4$ (7%)	34 (0.2%)	22 (65%)
ZW	$5 \cdot 10^4$ (1.68 ^{NLO})	$3.6 \cdot 10^4$ (73%)	$1.9 \cdot 10^4$ (53%)	173 (1%)	44 (25%)
ZZ	$4.8 \cdot 10^3$ (0.16 ^{NLO})	$3.5 \cdot 10^3$ (73%)	$1.7 \cdot 10^3$ (48%)	38 (2.3%)	15 (39%)
$t\bar{t}$	$2.6 \cdot 10^6$ (88 ^{NLO})	$1.8 \cdot 10^6$ (70%)	$1.3 \cdot 10^5$ (7%)	239 (0.2%)	89 (37%)
Z+jets(3l)	$4.6 \cdot 10^5$ (15.4 ^{LO})	$3.7 \cdot 10^5$ (80.5%)	$9.8 \cdot 10^4$ (26.5%)	504 (0.5%)	129 (26%)
DY(3l)	$4.5 \cdot 10^5$ (15.1 ^{LO})	$3.2 \cdot 10^5$ (71%)	$1.4 \cdot 10^5$ (44%)	670 (0.5%)	131 (20%)
Z $b\bar{b}$ (3l)	$8.4 \cdot 10^4$ (2.8 ^{LO})	$7.3 \cdot 10^4$ (87%)	$1.5 \cdot 10^4$ (20%)	69 (0.6%)	18 (26%)
Wt+jets	$3 \cdot 10^5$ (10 ^{NLO})	$2.1 \cdot 10^5$ (70%)	$3.9 \cdot 10^4$ (18.5%)	52 (0.1%)	20 (38%)
WW+jets	$6 \cdot 10^5$ (19.8 ^{LO})	$3.8 \cdot 10^5$ (63%)	$1.9 \cdot 10^4$ (50%)	7 (0.04%)	2 (29%)
Tot. bkg	$\sim 4.9 \cdot 10^6$			1786	470 (26%)

In a second step the background suppression was improved with a Neural Network(NN). Five networks for DY, Z+jets, $t\bar{t}$, ZW and ZZ backgrounds have been trained on the LM9 signal sample using the following variables: $P_T^{1,2,3}$, $\sum P_T$, M_{ll} , P_T^{2l} (transverse momentum of two OSSF leptons), $A = \frac{P_T^1 - P_T^2}{P_T^1 + P_T^2}$, Θ_{ll} (angle between two OSSF leptons), Φ_{ll} (angle in transverse plane), MET , N_{jets} (number of jets passing the jets veto), E_t^{hj} (of the highest E_T jet), η_{hj} (rapidity of the highest jet). The selection cuts on the NN outputs were optimized for the maximum significance at LM9 with the genetic algorithm GARCON. The efficiency of the NN selection is also shown in Table 13.15.

13.14.2 Results at LM9 and systematics

After the selection based on cuts the S_{cp} significance calculated for all OSSF pair combination is 6.1 at point LM9 for an integrated luminosity of 30 fb^{-1} . The NN improves the S_{cp} for all OSSF combinations to 7.8.

In addition to the real trilepton final state, leptons can be produced in the detector volume from π^\pm , K^\pm decays, bremsstrahlung, punch-through or faked by jets. The rate per event of such fake leptons was estimated individually for each background by matching the reconstructed lepton with the generated one and is $\sim 10^{-4}$ for electrons and $\sim 10^{-5}$ for muons. The expected fake leptons substantially increase the background, especially for the preselected channels like DY or Z+jets, by $\sim 221 \pm 48$ events and $\sim 31 \pm 16$ events respectively for the trimuon final state where the fake rate is smaller. The S_{cp} significance including fakes but without other systematic uncertainties for all OSSF combinations and for the trimuon state at LM9 is 6.5 and 5.1 respectively.

The reconstruction uncertainties related to the jet energy scale (5%) and the lepton momentum resolution (2%) contribute 1% to the uncertainties on the background. The average theoretical uncertainty from the PDFs, calculated with the LHPDF subsets using the re-weighting

technique for each background channel, amounts to 1.7%. These uncertainties reduce the significances to 5.8 and 4.8 for the all OSSF pairs and for the trimuon final state, respectively. However the largest uncertainties are coming from the statistical errors in the fake rate estimation which contribute $\sim 7\%$ to the background uncertainties rendering the signal hardly visible, $S_{cp} \sim 3.3$. These fake rates uncertainties can be reduced with larger simulation samples.

13.14.3 CMS reach for the tri-lepton final state

Figure 13.27 shows the 5σ discovery reach in m_0 and $m_{1/2}$ plane at $L_{int}=30 \text{ fb}^{-1}$ for all OSSF




Figure 13.27: Discovery reach of trilepton from the $pp \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ production at $L_{int}=30 \text{ fb}^{-1}$ for all OSSF lepton combinations (dashed) and for the trimuon final state (solid) including systematic uncertainties from reconstruction.

combinations and for the trimuon final state including the systematic uncertainties due to the reconstruction. The signal can be observed at large $m_0 > 1000 \text{ GeV}$ in a narrow band below $m_{1/2} < 180 \text{ GeV}$. At low $m_0 < 100 \text{ GeV}$ the two body decays are visible although a better optimization is possible in this region, see Sections 13.8 and 13.15. The trilepton final state from direct neutralino-chargino production is complementary to the inclusive OSSF dilepton search and provides an additional verification for the leptonic decays of the neutralino at low

$m_{1/2}$.

13.15 Production of \tilde{l}

The aim of this section is the study of the possibility of detecting sleptons. Note the previous related papers where the sleptons detection was studied at the level of a toy detector [661, 662, 663, 664, 665].

13.15.1 Simulation details

ISASUSY 7.69 [653] was used for the calculation of coupling constants and cross sections in the leading order approximation for SUSY processes. For the calculation of the next-to-leading order corrections to the SUSY cross sections the PROSPINO code [658] was used. Cross sections of the background events were calculated with PYTHIA 6.227 [62] and COMPHEP 4.2pl [47]. For considered backgrounds the NLO corrections are known and they were used. Official datasets (DST) production was used for the study of CMS test point LM1 and backgrounds ($t\bar{t}$, ZZ, WW, Wt, Zbb, DY2e, DY2 τ). For WZ, DY2 μ and W+jet backgrounds the events were generated with PYTHIA 6.227 and CMKIN 4.3.1 [13] was used as an interface program. The detector simulation and hits production were made with OSCAR 3.6.5 and for digitization ORCA 8.7.3 was used. The DY2 μ and W+jet backgrounds were simulated with FAMOS 1.4.0.

To reconstruct electrons an official ORCA code OfflineElectronReco was used. An official code GlobalMuonReconstructor was used for muons reconstruction.

An official ORCA algorithm PersistentJetFinder was used for jets reconstruction. A standard way using IterativeConeAlgorithm with cone size 0.5, EcalPlusHcalTowerInput as input and GammaJet calibration was used. E_T^{miss} was calculated using an official algorithm METfromEcalPlusHcalTower with LeptonCorrection.

The events are required to pass the Global Level 1 Trigger (L1) and the High Level Trigger (HLT). The events have to pass at least one of the following triggers: single electron, double electron, single muon, double muon.

Fast simulation code FAMOS 1.3.1 was used for the determination of sleptons discovery plot.

13.15.2 Sleptons production and decays

When sleptons are heavy relative to $\tilde{\chi}_1^\pm$, $\tilde{\chi}_2^0$ sleptons are produced significantly at the LHC through the Drell-Yan mechanism (direct sleptons production), via $q\bar{q}$ annihilation with neutral or charged boson exchange in the s-channel, namely, $pp \rightarrow \tilde{l}_L\tilde{l}_L, \tilde{l}_R\tilde{l}_R, \tilde{\nu}\tilde{\nu}, \tilde{\nu}\tilde{l}, \tilde{l}_L\tilde{l}_R$. The left sleptons decay to charginos and neutralinos via the following (kinematically accessible) decays:

$$\tilde{l}_L^\pm \rightarrow l^\pm + \tilde{\chi}_{1,2}^0, \quad (13.24)$$

$$\tilde{l}_L^\pm \rightarrow \nu_l + \tilde{\chi}_1^\pm, \quad (13.25)$$

$$\tilde{\nu} \rightarrow \nu_l + \tilde{\chi}_{1,2}^0, \quad (13.26)$$

$$\tilde{\nu} \rightarrow l^\pm + \tilde{\chi}_1^\pm \quad (13.27)$$

For right sleptons only decays to neutralino are possible and they decay mainly to LSP:

$$\tilde{l}_R^\pm \rightarrow l^\pm + \tilde{\chi}_1^0 \quad (13.28)$$

If sleptons are light relative to $\tilde{\chi}_1^\pm$, $\tilde{\chi}_2^0$, sleptons can be abundantly produced, besides Drell-Yan mechanism, also from chargino and neutralino decays $\tilde{\chi}_1^\pm$, $\tilde{\chi}_2^0$ (indirect production), equations (13.8), (13.9), (13.13) and (13.14).

13.15.3 Signature and backgrounds

The slepton production and decays described previously lead to the signature with the simplest event topology: *two leptons + E_T^{miss} + jet veto*. This signature arises for both direct and indirect slepton pair production. In the case of indirectly produced sleptons not only the event topology with two leptons but with single, three and four leptons is possible. Besides, indirect slepton production from decays of squarks and gluino through charginos, neutralinos can lead to an event topology *two leptons + E_T^{miss} + ($n \geq 1$) jets*.

The cut set close to the optimal one is the following:

a. for leptons:

- p_T - cut on leptons ($p_T^{\text{lept}} > 20$ GeV/c, $|\eta| < 2.4$) and lepton isolation within $\Delta R < 0.3$ cone containing calorimeter cells and tracker;
- effective mass of two opposite-sign and the same-flavour leptons is outside ($M_Z - 15$ GeV, $M_Z + 10$ GeV) interval;
- $\Phi(l^+l^-) < 140^\circ$ cut on angle between two leptons;

b. for E_T^{miss} :

- $E_T^{\text{miss}} > 135$ GeV cut on missing E_T ;
- $\Phi(E_T^{\text{miss}}, ll) > 170^\circ$ cut on relative azimuthal angle between dilepton and E_T^{miss} ;

c. for jets:

- jet veto cut: $N_{\text{jet}} = 0$ for a $E_T^{\text{jet}} > 30$ GeV (corrected jets) threshold in the pseudorapidity interval $|\eta| < 4.5$.

Such type of cuts is the standard cut set [661, 665].

The Standard Model (SM) backgrounds are: $t\bar{t}$, WW, WZ, ZZ, Wt, Zb \bar{b} , DY, W+jet. The main contributions come from WW and $t\bar{t}$ backgrounds. There are also internal SUSY backgrounds which arise from $\tilde{q}\tilde{q}$, $\tilde{g}\tilde{g}$ and $\tilde{q}\tilde{g}$ productions and subsequent cascade decays with jets outside the acceptance or below the threshold. Note that when we are interested in new physics discovery we have to compare the calculated number of SM background events N_{SMbg} with new physics signal events $N_{\text{new physics}} = N_{\text{slept}} + N_{\text{SUSYbg}}$, so SUSY background events increase the discovery potential of new physics.

13.15.4 Results

For the point LM1 with the used set of cuts for the integral luminosity $\mathcal{L} = 10$ fb $^{-1}$ the number of signal events (direct sleptons plus sleptons from chargino/neutralino decays) is $N_S = 60$, whereas the number of SUSY background events is $N_{\text{SUSYbg}} = 4$ and the number of SM background events is $N_{\text{SMbg}} = 41$. The SUSY background is rather small with compare to the signal, so we can assume $N_S = N_{\text{direct sleptons}} + N_{\text{chargino/neutralino}} + N_{\text{SUSYbg}} = 64$. It corresponds to the significances $S_{c12} = 7.7$ and $S_{cL} = 8.3$ (here $S_{c12} = 2(\sqrt{N_S + N_B} - \sqrt{N_B})$ [41] and $S_{cL} = \sqrt{2(N_S + N_B)\ln(1 + \frac{N_S}{N_B})} - N_S$ [94, 96]).

Taking into account the systematic uncertainty of 23% related with inexact knowledge of backgrounds leads to the decrease of significance S_{c12} from 7.7 to 4.3.

The ratio of the numbers of background events from two different channels $N(e^+e^- + \mu^+\mu^-)/N(e^\pm\mu^\mp) = 1.37$ will be used to keep the backgrounds under control.

The CMS discovery plot for *two leptons* + E_T^{miss} + *jet veto* signature is presented in Figure 8.




Figure 13.28: Discovery plot ($\tan\beta = 10$, $\text{sign}(\mu) = +$, $A = 0$) for l^+l^- .

13.16 Lepton flavour violation in neutralino decay

The aim of this section is the study of the possibility to detect SUSY and Lepton Flavour Violation (LFV) using the $e^\pm\mu^\mp + E_T^{\text{miss}}$ signature.

13.16.1 Signal selection and backgrounds

The simulation details of this study could be found in the Section 13.15.

The SUSY production $pp \rightarrow \tilde{q}\tilde{q}', \tilde{g}\tilde{g}, \tilde{q}\tilde{g}$ with subsequent decays leads to the event topology $e^\pm\mu^\mp + E_T^{\text{miss}}$. Note that in the MSSM with flavour lepton conservation neutralino decays into leptons $\tilde{\chi}_{2,3,4}^0 \rightarrow l^+l^-\tilde{\chi}_1^0$ do not contribute to this signature and contribute only to $l^+l^- + E_T^{\text{miss}}$ signature (here $l = e$ or μ). The main backgrounds which contribute to the $e^\pm\mu^\mp$ events are: $t\bar{t}$, ZZ, WW, WZ, Wt, Zbb, DY2 τ . It has been found that $t\bar{t}$ background is the biggest one and it gives more than 50% contribution to the total background.

Our set of cuts is the following:


- p_T - cut on leptons ($p_T^{\text{lept}} > 20$ GeV/c, $|\eta| < 2.4$) and lepton isolation within $\Delta R < 0.3$ cone.
- $E_T^{\text{miss}} > 300$ GeV cut on missing E_T .

13.16.2 Results at CMS test points and reach

For integrated luminosity $\mathcal{L} = 10 \text{ fb}^{-1}$ the number of background events is $N_B = 104$. The results for this luminosity are presented in Table 13.16.

Table 13.16: Number of signal events and significances S_{c12} [41] and S_{cL} [94, 96] for $\mathcal{L} = 10 \text{ fb}^{-1}$.

Point	N events	S_{c12}	S_{cL}
LM1	329	21.2	24.0
LM2	94	7.7	8.2
LM3	402	24.6	28.2
LM4	301	19.8	22.3
LM5	91	7.5	7.9
LM6	222	15.7	17.3
LM7	14	1.3	1.3
LM8	234	16.4	18.1
LM9	137	10.6	11.4

 CMS discovery plot for the $e^\pm\mu^\mp + E_T^{\text{miss}}$ signature is presented in Figure 13.29.




Figure 13.29: Discovery plot ($\tan\beta = 10$, $\text{sign}(\mu) = +$, $A = 0$) for the luminosities $\mathcal{L} = 1, 10, 30 \text{ fb}^{-1}$ for the $e^\pm\mu^\mp + E_T^{\text{miss}}$ signature.

In the MSSM the off-diagonal components of the slepton mass terms violate lepton flavour conservation. As it was shown in Refs. [666, 667, 668] it is possible to look for lepton flavour violation at supercolliders through the production and decays of the sleptons. For the LFV at the LHC one of the most promising processes is the LFV decay of the second neutralino [669, 670] $\tilde{\chi}_2^0 \rightarrow \tilde{l} l \rightarrow \tilde{\chi}_1^0 l l'$, where the non zero off-diagonal component of the slepton mass matrix leads to the different flavours for the leptons in the final state. By using the above mode, LFV in $\tilde{e} - \tilde{\mu}$ mixing has been investigated in Refs.[669, 670] at a parton model level for a toy detector. In this section we study the perspectives of the LFV detection in CMS on the base of full simulation of both signal and background is studied. To be specific, we study the point LM1. We assume that the LFV is due to nonzero mixing of right-handed smuon and selectron. The signal of the LFV $\tilde{\chi}_2^0$ decay is two opposite-sign leptons ($e^+\mu^-$ or $e^-\mu^+$) in the final state with the characteristic edge structure. In the limit of lepton flavour

conservation, the process $\tilde{\chi}_2^0 \rightarrow \tilde{l}l \rightarrow ll\tilde{\chi}_1^0$ has the edge structure for the distribution of the lepton-pair invariant mass m_{ll} and the edge mass m_{ll}^{max} is expressed by the slepton mass $m_{\tilde{l}}$ and the neutralino masses $m_{\tilde{\chi}_{1,2}^0}$ as follows:

$$(m_{ll}^{max})^2 = m_{\tilde{\chi}_2^0}^2 \left(1 - \frac{m_{\tilde{l}}^2}{m_{\tilde{\chi}_2^0}^2}\right) \left(1 - \frac{m_{\tilde{\chi}_1^0}^2}{m_{\tilde{l}}^2}\right) \quad (13.29)$$

The SUSY background for the LFV comes from uncorrelated leptons from different squark or gluino decay chains. The SM background comes mainly from

$$t\bar{t} \rightarrow bWbW \rightarrow bbl'\nu\nu' \quad (13.30)$$

Drell-Yan background from $pp \rightarrow \tau\tau \rightarrow e\mu\dots$ is negligible. It should be stressed that for the signature with $e^\pm\mu^\mp$ in the absence of the LFV we do not have the edge structure for the distribution on the invariant mass $m_{inv}(e^\pm\mu^\mp)$. As the result of the LFV the edge structure for $e^\pm\mu^\mp$ events arises too. Therefore the signature of the LFV is the existence of an edge structure in the $e^\pm\mu^\mp$ distribution. The rate for a flavour violating decay is

$$Br(\tilde{\chi}_2^0 \rightarrow e^\pm\mu^\mp\tilde{\chi}_1^0) = \kappa Br(\tilde{\chi}_2^0 \rightarrow e^+e^-\tilde{\chi}_1^0, \mu^+\mu^-\tilde{\chi}_1^0), \quad (13.31)$$

where:

$$Br(\tilde{\chi}_2^0 \rightarrow e^+e^-\tilde{\chi}_1^0, \mu^+\mu^-\tilde{\chi}_1^0) = Br(\tilde{\chi}_2^0 \rightarrow e^+e^-\tilde{\chi}_1^0) + Br(\tilde{\chi}_2^0 \rightarrow \mu^+\mu^-\tilde{\chi}_1^0), \quad (13.32)$$

$$\kappa = 2x \sin^2 \theta \cos^2 \theta, \quad (13.33)$$

$$x = \frac{\Delta m_{\tilde{e}\tilde{\mu}}^2}{\Delta m_{\tilde{e}\tilde{\mu}}^2 + \Gamma^2}, \quad (13.34)$$

$$Br(\tilde{\chi}_2^0 \rightarrow e^\pm\mu^\mp) = Br(\tilde{\chi}_2^0 \rightarrow e^+\mu^-) + Br(\tilde{\chi}_2^0 \rightarrow e^-\mu^+). \quad (13.35)$$

Here θ is the mixing angle between \tilde{e}_R and $\tilde{\mu}_R$ and Γ is the sleptons decay width. The parameter x is the measure of the quantum interference effect. There are some limits on $\tilde{e} - \tilde{\mu}$ mass splitting from lepton flavour violating processes but they are not very strong.

For $\kappa = 0.25$, $\kappa = 0.1$ the distributions of the number of $e^\pm\mu^\mp$ events on the invariant mass $m_{inv}(e^\pm\mu^\mp)$ (see Figure 13.30) clearly demonstrates the existence of the edge structure[671], i.e. the existence of the lepton flavour violation in neutralino decays. It appears that for the point LM1 the use of an additional cut

$$m_{inv}(e^\pm\mu^\mp) < 85 \text{ GeV} \quad (13.36)$$

reduces both the SM and SUSY backgrounds and increases the discovery potential in the LFV search. For the point LM1 we found that in the assumption of exact knowledge of the background (both the SM and SUSY backgrounds) for the integrated luminosity $\mathcal{L} = 10 \text{ fb}^{-1}$ it would be possible to detect LFV at 5σ level in $\tilde{\chi}_2^0$ decays for $\kappa \geq 0.04$.




Figure 13.30: The distribution of dilepton invariant mass after selection of two isolated $e^\pm \mu^\mp$ leptons with $p_T^{lept} > 20$ GeV/c and $E_T^{miss} > 300$ GeV for flavour violation parameter $k = 0.25$ (left) and $k = 0.1$ (right). The superimposed curves are fits to the invariant mass distribution for the case of 100% LFV.

13.17 Look beyond MSUGRA

13.17.1 Non Universal Higgs Masses

It was emphasized in Section 13.3 that the signatures of SUSY with a stable LSP result from the fundamental Supersymmetry gauge couplings, together with the composition of the lightest charginos and neutralinos. As all previous analyses were based on MSUGRA, it is interesting to verify their robustness when relaxing some of the assumptions which might affect the signal observability. As full generality, including giving up all universality assumptions, would lead to an intractable model, a choice needs to be made. Here, a mild extension is considered whereby the two higgsino mass parameters at the GUT scale are no longer supposed to be degenerate with the other scalar masses, which is sometimes called the Non Universal Higgs Masses (NUHM [672]) scenario. This scenario is conveniently parameterized in terms of two low scale parameters, the mass of the CP-odd Higgs (m_A) and the parameter μ . More specifically, we will analyze the effect of lowering the value of μ compared to its MSUGRA value on the observability of the signatures, as this modifies the composition of the charginos and neutralinos as a function of the gaugino and higgsino fields. For simplicity, m_A is kept at a fixed value. As exemplified in Figure 13.31 for the test point LM1, lowering μ also lowers the gaugino masses and in particular their splittings, which affect the branching ratios through phase space effects (a similar behaviour is observed for the other test points). The \tilde{q} and \tilde{l} spectra are almost unaffected. As for low values of μ the lightest chargino becomes lighter than the exclusion from LEP, $m(\tilde{\chi}_1^\pm) \geq 103$ GeV, this region is excluded and is indicated on the figures below by a grey (green) shaded strip.




Figure 13.31: Variation of the chargino and neutralino masses as a function of μ for the CMS test point LM1.

13.17.1.1 Signatures at point LM1

The test point LM1 was studied above for its detectability in cascade decays via a $\tilde{\chi}_2^0$ into $\tilde{l}_R l$ and $\tilde{\tau}_1 \tau$ (?). Figure 13.32 shows the variation of some branching ratios from the value of μ near the region where radiative electroweak symmetry breaking is not possible up to its value in MSUGRA.

It is seen that by lowering μ , $B(\tilde{q}_L \rightarrow q \tilde{\chi}_2^0 \rightarrow q \tilde{l}_R l)$ first increases (due to closing the competing decay to $\tilde{\nu} \nu$), then decreases when the $\tilde{\chi}_2^0$ becomes higgsino-like, but it remains considerably larger than its MSUGRA value for all values of μ down to the LEP limit. In addition, some new channels open up, like the decay via $\tilde{\chi}_4^0$ into left and right sleptons and the decay via a $\tilde{\chi}_2^\pm \rightarrow \tilde{\nu}_l \tilde{l}$ followed by $\tilde{\nu}_l \rightarrow \tilde{\chi}_1^\pm l$ (the $\tilde{\chi}_4^0$ and $\tilde{\chi}_2^\pm$ become more Wino-like). Other decays via $\tilde{\chi}_3^0$ might also contribute, but only in the region excluded by LEP.

The branching for the decay to $\tilde{\tau} \tau$ shows qualitatively the same behaviour, but is larger than its MSUGRA value in only a small region of μ . Also here a small contribution from the decay $\tilde{\chi}_2^\pm \rightarrow \tilde{\nu} \tau$ is present at small μ .

It is interesting to note that, although for MSUGRA the \tilde{q}_R decays exclusively directly to the LSP, it may have for lower μ a non negligible branching ratio to $\tilde{\chi}_2^0$ and also contributes to the di-lepton signature.

Finally, there is a non-zero branching ratio for the \tilde{q}_L to the light Higgs via the $\tilde{\chi}_2^\pm$ or $\tilde{\chi}_4^0$ (not shown), but it remains below 1% over the whole range of μ above the LEP limit and will be difficult to detect.

13.17.1.2 Signatures at point LM6

The test point LM6 has many features in common with LM1, but the $\tilde{\chi}_2^0$ decays mainly to $\tilde{l}_L l$ with a small admixture of $\tilde{l}_R l$. Moreover the decay $\tilde{\chi}_2^0 \rightarrow h^0 \tilde{\chi}_1^0$ is kinematically allowed, although suppressed due to the strong gaugino dominance in the $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$. The variation of the branching ratios as a function of μ is displayed in Figure 13.33




Figure 13.32: Decay branching ratios as a function of μ for \tilde{q}_L into ll and $\tau\tau$ and for \tilde{q}_R into ll at the test point LM1.

The cascade decays of \tilde{q}_L to $\tilde{l}l$ and $\tilde{\tau}\tau$ via $\tilde{\chi}_2^0$ show grossly the same behaviour as for LM1, with an increase at intermediate values of μ followed by a decrease at low μ . Again, the contributions from other charginos and neutralinos are non negligible near the LEP exclusion limit. Also \tilde{q}_R decays contribute to the di-lepton signal via $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ intermediate states.

A distinctive feature of LM6 is its production of final states with h^0 . The \tilde{q}_L branching ratio via $\tilde{\chi}_2^0 \rightarrow h^0 \tilde{\chi}_1^0$, which is only 2% for MSUGRA increases drastically for lower μ due to the increased higgsino components in $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$, then it drops as the decay becomes kinematically forbidden. After a gap where the branching ratio is below 1%, a strong increase is again visible for lower μ from the cascade dominated by $\tilde{\chi}_2^\pm \rightarrow h_0 \tilde{\chi}_1^\pm$ down to the LEP limit. Such an effect is not observed at LM1 due to the smaller spacing of the masses.




Figure 13.33: Decay branching ratios as a function of μ for \tilde{q}_L into ll , $\tau\tau$ and h^0 and for \tilde{q}_R into ll at the test point LM6.

13.17.1.3 Signatures at point LM4

At LM4 was chosen for its characteristic decay of $\tilde{\chi}_2^0$ into $Z^0 \tilde{\chi}_1^0$. Figure 13.34

MAXME: here figures for ll, h^0

shows the variation of the branching ratios as a function of μ .

As the decay $\tilde{\chi}_2^0 \rightarrow Z^0 \tilde{\chi}_1^0$ requires higgsino components in both the $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$, its branching ratio remains above 90% for all values of μ allowed by the LEP limit. The branching ratio of the \tilde{q}_L into $Z^{(*)}$ via a $\tilde{\chi}_2^0$ decreases mainly due to the decrease of $B(\tilde{q}_L \rightarrow q \tilde{\chi}_2^0)$ (the $\tilde{\chi}_2^0$ becomes less gaugino-like). This loss is, however, compensated by the contributions from cascades via $\tilde{\chi}_2^\pm \rightarrow W \tilde{\chi}_2^\pm$ and $\tilde{\chi}_2^\pm \rightarrow Z^0 \tilde{\chi}_1^\pm$ and the overall effect is a net increase of the branching ratio of the \tilde{q}_L to final states with a Z^0 .

For low values of μ there is also a contribution to h^0 final states via the decay $\tilde{\chi}_2^\pm \rightarrow h^0 \tilde{\chi}_1^\pm$,




Figure 13.34: Decay branching ratios as a function of μ for \tilde{q}_L into ll and h^0 at the test point LM4.

but it remains small above the limit imposed by LEP.

13.17.1.4 Signatures at point LM5

At point LM5, the main signature for MSUGRA is provided by the cascade via $\tilde{\chi}_2^0 \rightarrow h^0 \tilde{\chi}_1^0$. The variation of the branching ratios with μ are shown in Figure 13.35.

The sharp drop in the branching ratio of $\tilde{\chi}_2^0$ to h^0 below the MSUGRA value of μ results from the decrease in the mass splitting between $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$ which suppresses the decay to h^0 . For lower values of μ , final states with h^0 are again produced mainly via the $\tilde{\chi}_2^\pm \rightarrow h^0 \tilde{\chi}_1^\pm$. In between these two decay chains, a narrow gap is left where the Higgs branching ratio is less than 2% and hence very difficult to detect.

It is seen that this loss of sensitivity to Higgs final states is to some extent compensated by an increase of the di-lepton final states in the region of the gap. The cascade decays of both \tilde{q}_L and \tilde{q}_R contribute in this region, the main contributions being through $\tilde{\chi}_2^0 \rightarrow Z^* \tilde{\chi}_1^0$, $\tilde{\chi}_2^\pm \rightarrow Z^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^\pm \rightarrow W \tilde{\chi}_2^0$. It gives a branching ratio of up to 3.5% for the di-lepton decay of \tilde{q}_L and less than 1% for \tilde{q}_R and hence should be detectable. However, the mixture of intermediate states leading to the di-leptons will make the sparticle mass reconstruction very challenging.

13.17.1.5 Conclusion

It can be concluded that the same flavour di-lepton signatures originating from the decay of \tilde{l} or Z^* are quite robust with respect to the chargino and neutralino composition. Lowering μ with respect to its MSUGRA value, a sizeable increase of the branching ratio is even observed for the test points LM1, LM4 and LM6. The h^0 signature at point LM5 is less robust and a region with low branching ratio exists at intermediate values of μ . It is compensated by an increase of di-lepton final states. It may be noted that the loss of $\tilde{\chi}_2^0$ decay to h^0 is due to the reduction of the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$ mass splitting. It is therefore a consequence of the low mass spectrum chosen and should disappear at larger values of $m_{1/2}$. Another feature of the




Figure 13.35: Decay branching ratios as a function of μ for \tilde{q}_L into h^0 and ll and for \tilde{q}_R into ll at the test point LM5.

NUHM scenario is that for small μ the cascades from \tilde{q}_R also contribute to the signatures, unlike the MSUGRA case. Moreover the signatures at low to intermediate μ tend to be produced by several intermediate neutralino and chargino states. This points to the difficulty of identifying which sparticles are at the origin of the observed end points in the effective mass distributions.

FIXME: We could also have looked into the following:

- look beyond MSUGRA: ex. H^0, A^0 in cascades
- wino-like (AMSB), higgsino-like (NUHM), mixed, NMSSM, ...

large m_0 , split supersymmetry, ...

FIXME: But this will be dropped: no study made

13.18 Summary of the reach with inclusive analyses

13.18.1 Summary of the MSUGRA studies

In previous sections, several characteristic topologies (or signatures) for MSSM were studied and it was shown that many are already detectable with rather low integrated luminosity (few years of LHC running) over a sizeable part of the parameter space, extending well beyond the Tevatron reach. **NOTE:** Summarize the findings **NOTE:** Here we will gather all search areas to give a coherent overview of the results of all inclusive searches.

Chapter 14

Extra Dimensions and new vector boson high mass states



14.1 Introduction

Something on the dualities of the models. Don't forget to point co-existence possibility (if not requirement) with SUSY-like scenarios. Don't forget to fix notation or translate notation to the generators we use. Check with people's private productions of RS's. In particular if anyone uses Herwig. Fix the truncation problem in ADD. Do we need mention or references to previous results from the Tevatron? Figure out what to chop. Separate the ADD graviton exchange from the RS strongly coupled case (final states are the same). Mention how the particles get masses (goldstone boson like business in ADD). Mention the 5 helicities of the massive ones.

14.1.1 Models with heavy vector bosons

14.1.2 Arkani-Hamed Dimopoulos Dvali (ADD) models

ADD refers to the class of models which incorporate the large extra dimensions scenario of Arkani-Hamed, Dvali, and Dimopoulos [673]. These were the first extra dimensions models in which the compactified dimensions can be of macroscopic size, consistent with all current experiments and observations and they are referred to as "large extra dimensions" models. In the most basic version, n extra spatial dimensions are compactified on a torus with common circumference R , and a brane is introduced which extends only in the three infinite spatial directions. Strictly speaking, the brane should have a very small tension (energy per unit volume) in order that it does not significantly warp the extra dimensional space. It is assumed that all standard model fields extend only in the brane. This can be considered as a toy version of what happens in string theory, where chiral gauge theories similar to the standard model are confined to reasonably simple brane configurations in reasonably simple string compactifications [674].

A consequence of these assumptions is that the effective 4d Planck scale is related to the underlying fundamental Planck scale of the $4+n$ -dimensional theory and to the volume of the compactified space. This relation follows from Gauss' Law, or by dimensional truncation:

$$M_{\text{Planck}}^2 = M_*^{2+n} R^n \quad , \quad (14.1)$$

where M_{Planck}^2 is defined by Newton's constant: $M_{\text{Planck}} = 1/\sqrt{G_N} = 1.2 \times 10^{19}$ GeV. M_*^{2+n} is defined as the gravitational coupling which appears in the $4+n$ -dimensional version of the Einstein-Hilbert action. It is the quantum gravity scale of the higher dimensional theory.

If M_{Planck} , M_* and $1/R$ are all of the same order, as is usually assumed in string theory, this relation is not very interesting. But it is plausible and experimentally allowed that M_* is equal to some completely different scale. Taking $M_* \sim 1$ TeV [675] the hierarchy problem of the standard model is translated from an ultraviolet problem to an infrared one. Note that if there is any interface with string theory, ADD-like models must arise from string ground states in which the string scale (and thus the ultraviolet cutoff for gravity) is also in the TeV range. Although this is difficult it has been studied [676].

The ADD scenario renders observations of quantum gravity at the LHC plausible. In such models only the graviton, and possibly some non-SM exotics like the right-handed neutrino, probe the full bulk space. There is a Kaluza-Klein tower of graviton modes, where the massless mode is the standard 4d graviton, and the other KK modes are massive spin 2 particles which also couple to SM matter with gravitational strength.

Whereas bremsstrahlung of ordinary gravitons is a completely negligible effect at colliders, the total cross section to produce *some* massive KK graviton is volume enhanced, and effectively suppressed only by powers of M_* and not M_{Planck} . From Eq. (14.1) it follows :

$$\sigma \sim \frac{1}{M_{\text{Planck}}^2} (ER)^n \sim \frac{1}{M_*^2} (EM_*)^n \quad , \quad (14.2)$$

where E is the characteristic energy of the subprocess.

For graviton phenomenology it is useful to replace the ADD parameter M_* by other rescaled parameters. The two most useful choices are taken from the work of Giudice, Rattazzi and Wells (GRZ) [677], and Han, Lykken and Zhang (HLZ) [678]:

$$M_*^{n+2} = \frac{S_{n-1}}{(2\pi)^n} M_s^{n+2} \quad , \quad (14.3)$$

$$M_*^{n+2} = \frac{8\pi}{(2\pi)^n} M_D^{n+2} \quad , \quad (14.4)$$

where M_s is the HLZ scale, M_D is the GRW scale, and S_{n-1} is the surface area of a unit n -sphere:

$$S_{n-1} = \frac{2\pi^{n/2}}{\Gamma(n/2)} \quad . \quad (14.5)$$

Both notations are equivalent. To obtain a complete dictionary between ADD, GRZ and HLZ, one also needs to relate the ADD parameter R to those used by the other authors: $R = R_{\text{HLZ}} = 2\pi R_{\text{GRW}}$, and take note of the different notations for Newton's constant:

$$\kappa^2 = 16\pi G_N \text{ (HLZ)}; \quad \bar{M}_P^2 = \frac{1}{8\pi G_N} \text{ (GRW)} \quad . \quad (14.6)$$

A Kaluza-Klein (KK) graviton mode has a mass specified by an n -vector of integers \vec{k} :

$$m^2(\vec{k}) = \frac{\vec{k}^2}{R_{\text{GRW}}^2} \quad . \quad (14.7)$$

Let $r = |\vec{k}|$. Then for large r (as is often the relevant case for ADD phenomenology) the number of KK graviton states of a given polarization with $r \leq r_{\max}$ is given by the integral

$$\begin{aligned} S_{n-1} \int_0^{r_{\max}} dr r^{n-1} &= \frac{1}{n} S_{n-1} r_{\max}^n \\ &= \int_0^{m_{\max}} \rho(m) dm, \end{aligned} \quad (14.8)$$

where the KK density of states is

$$\rho(m) = \frac{m^{n-1}}{G_N M_s^{n+2}}. \quad (14.9)$$

M_s is the natural scaling parameter for KK graviton production. The density of states formulation can be applied to a much more general class of models than ADD, and can also include graviton wavefunction factors when the extra dimensions are not flat.

Consider an on-shell production of a KK graviton from a pp or collision. To leading order this is a $2 \rightarrow 2$ process with two massless partons in the initial state, plus a massive KK graviton and a massless parton in the final state. Let p_1, p_2 denote the 4-momenta of the initial state partons, p_3 the 4-momentum of the graviton, and p_4 the 4-momentum of the outgoing parton. The total cross section for any particular variety of partonic subprocess has the form

$$\sigma(1 + 2 \rightarrow \text{KK} + 4) = \int dx_1 dx_2 f_1(x_1, \hat{s}) f_2(x_2, \hat{s}) \int d\hat{t} \int_0^{\sqrt{\hat{s}}} dm \rho(m) \frac{d\sigma_m}{d\hat{t}}(\hat{s}, \hat{t}), \quad (14.10)$$

where $f_1(x_1, \hat{s}), f_2(x_2, \hat{s})$ are the parton distribution functions (pdfs) for the initial state partons, $\hat{s} = x_1 x_2 s = (p_1 + p_2)^2$ is the square of the total center of mass (cm) energy of the subprocess, and $\hat{t} = (p_1 - p_3)^2$ is the usual Mandelstam invariant. The formulae for $d\sigma_m/d\hat{t}$, the differential subprocess cross sections for KK gravitons of mass m , are given in [677].

14.1.2.1 Graviton production above the cutoff

At the LHC, proton–proton collisions will probe a distribution of partonic subprocess energies $\sqrt{\hat{s}}$. This creates a problem for the consistent analysis of missing energy signatures in the framework of ADD models. These models are simple low energy effective theories which are only valid for $\sqrt{\hat{s}}$ below some cutoff. This cutoff is at most $2M_*$, and could be a factor of a few smaller if the ultraviolet completion of the model is weakly coupled string theory [679]. The same is true for the Lykken–Randall model [680], which is a low energy description of gravity in a single infinite warped extra dimension, valid up to a cutoff $\sim M_*$. It is inconsistent to use either type of model to describe LHC collisions with subprocess energies greater than the cutoff.

This problem was first noted by the authors of [677], who suggested replacing the ADD graviton density of states $\rho(m)$ by $\rho(m)\theta(\sqrt{\hat{s}} - M_D)$, where θ is a step function. This introduces a systematic theory error into the analysis. The size of this error is very sensitive to the values of M_D and n . For initial LHC data sets, we will be probing the lower range of M_D values, beginning at the current $\simeq 1$ TeV bounds from Tevatron and LEP. This increases the theory systematic from the cutoff for any fixed n . For fixed M_D , the theory systematic increases rapidly for increasing n . For $n = 2$, the theory uncertainty in the total cross section

remains below about 20% even for M_D approaching 1 TeV.* For $n = 6$ and above, the effect of the cutoff is enormous for modest values of M_D , because the rapid rise in the graviton density of states is not compensated by the rapid falloff of the pdfs. The theory error for the total cross section in this case can be as large as an order of magnitude.

The resolution of this problem depends upon whether or not there is a signal in the missing energy channels (we will not discuss the related problems which arise in channels affected by virtual graviton exchanges). If there is a signal, the optimal procedure is to measure the observables $d^2\sigma/dp_T d\eta$ as accurately as possible, perhaps at more than one collider energy as suggested in [681],[682]. No theory systematic should be included in these analyses. Instead, one should use the data to find the best fit form for $\rho(m, \sqrt{\hat{s}})$. Simple trial forms can be obtained, for example, from multiplying the ADD density of states by the form factors obtained in models with strings [679],[683],[684] or branes [685]. For the lower range of M_D values, the sensitivity to n suggested in [681],[682] will tend to be washed out. This is not a bad outcome, since it is a result of convolving the n dependence with the effects of strings, branes or other new physics. Thus the theory systematic is replaced by likelihood fits to theories of Planck scale physics.

More problematic is the case where there is no graviton signal in a given data set. Since in this case we are trying to set a limit, we need an estimate of the theory systematic. The simplest possibility is to implement the GRW cutoff defined above, and estimate the theory error by varying the cutoff. For ADD with $n \geq 6$, one expects to obtain no lower bound at all on M_D , as noted in [677].

14.1.3 Inverse TeV sized extra dimensions

The possibility of TeV^{-1} -sized extra dimensions naturally arises in braneworld theories [675, ?]. By themselves, they do not allow for a reformulation of the hierarchy problem, but they may be incorporated into a larger structure in which this problem is solved. In these scenarios, the Standard Model fields are phenomenologically allowed to propagate in the bulk. This presents a wide variety of choices for model building: (i) all, or only some, of the Standard Model gauge fields exist in the bulk; (ii) the Higgs field may lie on the brane or in the bulk; (iii) the Standard Model fermions may be confined to the brane or to specific locales in the extra dimension. The phenomenological consequences of this scenario strongly depend on the location of the fermion fields. Unless otherwise noted, our discussion assumes that all of the Standard Model gauge fields propagate in the bulk.

The masses of the excitation states in the gauge boson KK towers depend on where the Higgs boson is located. If the Higgs field propagates in the bulk, the zero-mode state of the Higgs KK tower receives a vacuum expectation value (vev) which is responsible for the spontaneous breaking of the electroweak gauge symmetry. In this case, the resulting mass matrix for the states in the gauge boson KK towers is diagonal and the excitation masses are shifted by the mass of the gauge zero-mode, which corresponds to the Standard Model gauge field, giving

$$m_{\vec{n}} = (m_0^2 + \vec{n} \cdot \vec{n}/R_c^2)^{1/2}. \quad (14.11)$$

However, if the Higgs is confined to the brane, its vev induces mixing, amongst the gauge KK states of order $(m_0 R_c)^2$. The KK mass matrix must then be diagonalized in order to

*To avoid strong astrophysical constraints, $n = 2$ ADD models also require an *ad hoc* infrared cutoff, truncating the massive graviton spectrum for masses below about 20 MeV. This has a negligible effect on LHC analysis.

determine the excitation masses. For the case of 1 extra TeV^{-1} -sized dimension, the coupling strength of the gauge KK states to the Standard Model fermions on the brane is $\sqrt{2}g$, where g is the corresponding Standard Model gauge coupling.

14.1.4 Randall Sundrum (RS) models

Randall-Sundrum refers to a class of scenarios, also known as warped extra dimensions models, originated by Lisa Randall and Raman Sundrum [87, 626]. In these scenarios there is one extra spatial dimension, and the five-dimensional geometry is “warped” by the presence of one or more branes. The branes extend infinitely in the usual three spatial dimensions, but are sufficiently thin in the warped direction that their profiles are well-approximated by delta functions in the energy regime of interest. If we ignore fluctuations of the branes, we can always choose a “Gaussian Normal” coordinate system, such that the fifth dimension is labelled y and the usual 4d spacetime by x^μ . The action for such a theory contains, at a minimum, a 5d bulk gravity piece and 4d brane pieces. The bulk piece has the 5d Einstein-Hilbert action with gravitational coupling M^3 , and a 5d cosmological constant Λ . The brane pieces are proportional to the brane tensions V_i , which may be positive or negative. These act as sources for 5d gravity, contributing to the 5d stress-energy terms proportional to

$$\sum_i V_i \delta(y - y_i) \quad (14.12)$$

where the y_i are the positions of the branes. Combined with a negative Λ , this results in a curved geometry, with a 5d metric of the form:

$$\begin{aligned} g_{\mu\nu}(x^\rho, y) &= a^2(y) \tilde{g}_{\mu\nu}(x^\rho) , \\ g_{\mu y} &= 0 , \quad g_{yy} = 1 , \end{aligned} \quad (14.13)$$

where $a(y)$ is called the warp factor, \tilde{g} is a 4d metric, and I have made a useful choice of coordinates. Warping refers to the fact that a 4d distance d_0 measured at $y = y_0$ is related to an analogous 4d distance d_1 measured at $y = y_1$ by $a(y_0)d_0 = a(y_1)d_1$. Thus in Randall-Sundrum scenarios 4d length, time, energy and mass scales vary with y .

Most collider physics phenomenology done with warped extra dimensions so far is based upon one very specific model, the original simple scenario called RSI. In this model the extra dimension is compactified to a circle of circumference $2L$, and then further orbifolded by identifying points related by $y \rightarrow -y$. The fifth dimension then consists of two periodically identified mirror copies of a curved 5d space extending from $y = 0$ to $y = L$. It is assumed that there is a brane at $y = 0$, with positive tension V_0 ; it is known as the Planck brane - strong gravity resides on that brane. There is another brane at $y = L$, with negative tension V_L , known as the TeV brane - the entire 4d universe is confined to the TeV brane.

Randall and Sundrum showed that, for a tuned choice of input parameters $V_0 = -V_L = -M^2\Lambda$, the 5d Einstein equations have a simple warped solution on $0 < y < L$ with metric:

$$\begin{aligned} g_{\mu\nu}(x^\rho, y) &= e^{-2ky} \eta_{\mu\nu} , \\ g_{\mu y} &= 0 , \quad g_{yy} = 1 , \end{aligned} \quad (14.14)$$

where $\eta_{\mu\nu}$ is the 4d flat Minkowski metric, and $k = \sqrt{-\Lambda}$. Away from the branes, the 5d curvature is constant and negative; it is thus equivalent locally to AdS_5 , with the Anti-de

Sitter radius of curvature given by $1/k$. At the locations of the branes the curvature is discontinuous, due to the fact that the branes are delta function sources for curvature.

The RSI model is completely described by three parameters: k , M , and L . Restricting the scenario to a low energy effective description implies considering $k, 1/L \ll M$. In fact in RSI it is assumed that k is merely parametrically small compared to the 5d Planck scale M , i.e. $k \sim M/10$. The effective 4d Planck scale, which is the same as the coupling of the graviton zero mode, is given by dimensional truncation:

$$M_{\text{Planck}} = \frac{M^3}{2k} \left(1 - e^{-2kL}\right). \tag{14.15}$$

Then, within an order of magnitude, $M \sim k \sim M_{\text{Planck}}$. In RSI the distance L is fixed by requiring that $a(L)M_{\text{Planck}} \simeq 1 \text{ TeV}$, thus $kL \sim 30$. This is *not* a large extra dimension: its inverse size is comparable to the grand unification scale.

Since the standard model fields live on the TeV brane as in ADD models, the phenomenology of RSI is concerned with the effects of the massive KK modes of the graviton. These modes as measured on the TeV brane have their mass splittings of the order of a TeV, and have TeV suppressed couplings to the standard model fields. In RSI, the Standard Model is replaced at the TeV scale by a new effective theory in which gravity is still very weak, but there are exotic heavy spin-two particles.

At the LHC the KK gravitons of RSI would be seen as difermion or diboson resonances, since (unlike the KK gravitons of ADD) the coupling of each KK mode is only TeV suppressed [686]. The width of these resonances is controlled by the ratio $c = k/M$; the resonances become more narrow as $c = k/M$ is reduced, as shown in Figure 14.1.




Figure 14.1: The cross section for $e^+e^- \rightarrow \mu^+\mu^-$ including the exchange of KK gravitons in the RSI model. The narrowest resonances correspond to $k/M = 0.05$, the widest to $k/M = 0.1$. (taken from reference [687]).

2 High mass dielectron final states

FIXME: note:picture environment for labeling figures if larger sized figures needed
 This section presents the CMS experiment discovery potential for new heavy resonances,

decaying into an electron pair. The e^+e^- decay channel provides a clean signature in the CMS detector. The presence of a heavy particle would be detected in CMS by the observation of a resonance peak in the dielectron mass spectrum over the Drell-Yan process ($pp \rightarrow \gamma/Z \rightarrow e^+e^-$) which constitutes the main Standard Model background.

Heavy resonances with mass above $1 \text{ TeV}/c^2$ are predicted by several models beyond the Standard Model. Three models are considered here: Kaluza-Klein (KK) excitations of a Z boson (TeV^{-1} model, see Section ??) and KK excitation of a graviton (Randall-Sundrum (RS) model, see Section ??), both predicted in extra dimensions models, and neutral heavy Z' boson predicted by Grand Unified Theories (GUT) (see Section ??). For the Z' bosons, 6 models are studied, as for the $Z' \rightarrow \mu^+\mu^-$ channel [94] that is discussed in Section 14.3.

Details of the analyses presented in this section can be found in [688] and [689].

14.2.1 Event selection and correction

Two electrons are requested for this analysis. They are reconstructed as super-clusters (SC) in the ECAL calorimeter in the barrel and the endcap regions [656]. For endcap SC, the energy loss in the preshower detector is taken into account. The two SC with highest energies are selected as the electron candidates.

An important characteristic of the signal events is that the final state electrons are very energetic and may have a significant energy leakage beyond the ECAL clusters. An improvement in the energy determination is achieved by including the HCAL measurement, event by event.

Reducible backgrounds (like QCD jets and γ -jets) are suppressed by applying the following requirements:

- The ratio of the HCAL to ECAL energy deposits is required to be $H/E < 10 \%$.
- The two SC must be isolated: the total additional transverse energy in a cone of radius $0.1 < \Delta R < 0.5$ is required to be below 2 % of the SC transverse energy (where $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2}$).
- To identify electrons and reject neutral particles, a track is requested to be associated for each electron candidate. If a track is associated with only one of these SC, the event is however kept if it contains a third SC with $E > 300 \text{ GeV}$ with an associated track and satisfying the H/E and isolation cuts described above.

The selected events are then corrected for the following effects:

- Saturation correction: For very energetic electrons and photons, saturation occurs in the ECAL electronics because of the limited dynamical range of the Multi-Gain-Pre-Amplifier. The saturation threshold has been established to be at 1.7 TeV in crystals of the barrel and 3.0 TeV in the endcaps. A correction method (for barrel only) has been developed using the energy deposit in crystals surrounding the saturated crystal. The correction leads to the correct estimate of energy deposit with a resolution of around 7% [690].
- Energy correction: The ECAL measured electron energy after preshower, HCAL and saturation corrections, is smaller than the generated energy. Dedicated energy correction factors for very energetic electrons have been determined using calibration files. These factors depend on both energy, η and whether saturation

occurs or not. The resolution on the corrected SC energy is 0.6 % at $E = 1000$ GeV.

- z-vertex distribution: The measurement in η takes into account the knowledge of the z-vertex position.
- FSR recovery: Hard photon emission from Final State Radiation can induce the detection in the event of a third energetic SC. If a SC with $E > 300$ GeV satisfying the H/E and isolation cuts is observed very close to the SC of the electron candidates ($\Delta R < 0.1$), this additional SC is associated to the corresponding electron.

14.2.2 Mass peak distributions

The resonance mass is reconstructed from the energies and angles of the 2 electron candidates, after the selection cuts and energy corrections mentioned above. Figures 14.2(a) and (b) show the ratio of the reconstructed and the true masses, M_{ee}/M_{true} , before and after energy corrections for KK Z production with $M = 4$ and 6 TeV/c^2 , respectively. The peaks at low values of M_{ee}/M_{true} correspond to events with saturated ECAL electronics. The final resolution on the resonance mass is around 0.6 % for events with no saturation, and 7 % in case of saturation.

Figure 14.3(a) presents the signal and the Drell-Yan background for KK Z boson production with $M = 4$ TeV/c^2 ; Figure 14.3(b) for Z' boson production with $M = 1.5$ TeV/c^2 ; Figure 14.3(c) for graviton production with $M = 1.5$ TeV/c^2 and coupling parameter $c = 0.01$.

14.2.3 Discovery potential of CMS

The discovery potential of a new physics resonance is determined using the likelihood estimator S based on event counting [96], suited for small event samples:

$$S = \sqrt{2[(N_s + N_b) \log(1 + \frac{N_s}{N_b}) - N_s]}, \quad (14.16)$$

where N_s (resp. N_b) is the number of signal (resp. background) events. The discovery limit is defined by $S > 5$.




Figure 14.2: Ratio M_{ee}/M_{true} before and after corrections for KK Z boson production, for $M = 4$ TeV/c^2 (a) and $M = 6$ TeV/c^2 (b).




Figure 14.3: Resonance signal (white histograms) and Drell-Yan background (shaded histograms) for KK Z boson production with $M = 4.0 \text{ TeV}/c^2$ (a), SSM Z' boson production with $M = 3.0 \text{ TeV}/c^2$ (b), and graviton production with $M = 1.5 \text{ TeV}/c^2$, $c = 0.01$ (c), for an integrated luminosity of 30 fb^{-1} .

The number of signal and background events, N_s and N_b , computed for a given mass window around the peak, are presented in Table 14.1 for the three models, together with the corresponding significance, for an integrated luminosity of 30 fb^{-1} .

The 5σ discovery limits as a function of mass are given in Fig. 14.4(a) and Fig. 14.4(b), for KK Z boson production and Z' production (for the 6 considered models), respectively. In the graviton case, the 5σ discovery plane as a function of the coupling parameter c and the resonance mass is given in Fig. 14.4(c).

For KK Z bosons, a 5σ discovery can be achieved for a resonance mass up to $M = 4.97 \text{ TeV}/c^2$ for an integrated luminosity of 10 fb^{-1} , $M = 5.53 \text{ TeV}/c^2$ for 30 fb^{-1} and $M = 5.88 \text{ TeV}/c^2$ for 60 fb^{-1} . For gravitons, with an integrated luminosity of 30 fb^{-1} , a 5σ discovery can be extracted for masses up to $1.64 \text{ TeV}/c^2$ for $c = 0.01$ and up to $3.81 \text{ TeV}/c^2$ for $c = 0.1$. For Z' boson production, with an integrated luminosity of 30 fb^{-1} , a 5σ discovery can be extracted for masses up to $3.31 \text{ TeV}/c^2$ for model ψ and up to $4.27 \text{ TeV}/c^2$ for model ARLM. The 5σ discovery limits on the resonance masses for 10, 30 and 60 fb^{-1} are summarized in Table 14.2.

For KK Z boson production, the luminosities needed for a five σ discovery are 1.5, 4.0, 10.8, 29.4, and 81.4 fb^{-1} for $M = 4.0, 4.5, 5.0, 5.5$ and $6.0 \text{ TeV}/c^2$, respectively; for SSM Z' boson production, they are 0.015, 3.0 and 260 fb^{-1} for $M = 1, 3$ and $5 \text{ TeV}/c^2$; for graviton produc-

Table 14.1: Number of events for resonant signal, N_s , and for Drell-Yan background, N_b , and corresponding significances as defined by Eq. 14.16, for an integrated luminosity of 30 fb^{-1} . The masses M and the mass windows M_w are in TeV/c^2 .

	KK Z		G, $c = 0.01$	G, $c = 0.1$	SSM Z'	
M	4.0	6.0	1.5	3.5	1.0	5.0
M_w	3.5-4.5	5.0-6.7	1.47-1.52	3.30-3.65	0.92-1.07	4.18-5.81
N_s	50.6	1.05	18.8	7.30	72020	0.58
N_b	0.13	0.005	4.16	0.121	85.5	0.025
S	22.5	3.0	6.39	6.83	225	1.63




Figure 14.4: Five σ discovery limit as a function of the resonance mass for KK Z boson production (a), for the 6 Z' models (b); five σ discovery plane for graviton production as a function of the coupling parameter c and the graviton mass (c).

tion, most of the interesting region of the (mass, coupling) plane is already covered with 10 fb⁻¹.

For KK Z and Z' production, a K factor of 1 was conservatively taken for both the signal and the Drell-Yan background, since heavy Z production interferes with Z/γ Drell-Yan production. For the graviton analysis, as little interference is present with the Standard Model processes, a K factor of 1.0 is used for the signal and of 1.3 for the Drell-Yan background, in order to take into account the higher order terms in the cross section. The latter number comes from the CDF analysis [691] and is compatible with the K factor obtained from theoretical computations [350].

Table 14.2: The 5σ discovery limit on the resonance mass (given in TeV/c^2) for the three models, for an integrated luminosity of 10, 30 and 60 fb^{-1} .

Model	Luminosity (fb^{-1})		
	10	30	60
KK Z	4.97	5.53	5.88
G ($c = 0.01$)	1.38	1.64	1.82
G ($c = 0.1$)	3.34	3.81	4.10
Z' (ψ)	2.85	3.31	3.62
Z' (ALRM)	3.76	4.27	4.60

14.2.4 Systematic uncertainties

The uncertainty coming from the choice of the parton distribution function (PDF) was investigated using the set of 20 positive and 20 negative errors, of the CETQ6.1M "best fit" parameterization [692, 29, 693]. For each event, a weight factor is computed according to the x_1 , x_2 , and Q^2 variables, for each of the 40 PDF errors, in the case of graviton production with $M = 1.5\text{ TeV}/c^2$ ($c = 0.01$) and $M = 3.5\text{ TeV}/c^2$ ($c = 0.1$). The uncertainties on the PDF modify the number of signal events by a factor 1.20 (positive deviations) and 0.86 (negative deviations) for $M = 1.5\text{ TeV}/c^2$ ($c = 0.01$). The corresponding numbers for $M = 3.5\text{ TeV}/c^2$ ($c = 0.1$) are 1.47 and 0.78. For the Drell-Yan background, the re-weighting effects on the numbers of events are 1.065 and 0.941 for masses around $1.5\text{ TeV}/c^2$, and 1.19 and 0.88 for masses around $3.5\text{ TeV}/c^2$. For an integrated luminosity of 30 fb^{-1} , the significances with the "best fit" and with the positive/negative deviations are equal respectively to 6.40 and 7.25/5.78 for $M = 1.5\text{ TeV}/c^2$, and to 6.83 and 8.54/5.93 for $M = 3.5\text{ TeV}/c^2$. The main effect of the variation comes from the gluon-fusion contribution to the graviton production cross section. A lower dependence is observed for the KK Z and Z' channels, which are produced by quark-antiquark annihilation. For KK Z boson production at $M = 4\text{ TeV}/c^2$ with an integrated luminosity of 30 fb^{-1} , the significances with the "best fit" and with the positive/negative errors are equal respectively to 22.5 and 23.3/21.9.

Changing to 1 the value of the K factor of the DY background for RS graviton production increases the significance from 6.39 to 6.87 ($M = 1.5\text{ TeV}/c^2$, $c = 0.01$) and from 6.83 to 7.09 ($M = 3.5\text{ TeV}/c^2$, $c = 0.1$). The discovery limits increase respectively from 1.64 to 1.68 TeV/c^2 and from 3.81 to 3.84 TeV/c^2 .

The data themselves will be used to estimate and cross-check the Drell-Yan background at very high energy. For resonance discovery, the number of events in the side-bands of the resonance and their mass dependence will be used to estimate the number of background events under the resonance peak, provided there is enough data in the side-bands. In this approach, the uncertainties on the background cross-sections, the PDF and the luminosity measurement are highly reduced.

14.2.5 Identification of new particles

Once a resonance is found, information will be gained on its characterization from the study of other decay channels, like $\gamma\gamma$ (see Section 14.6), of angular distributions and of asymmetries, in view of the spin determination (see also Section 14.3).




Figure 14.5: Distributions of $\cos \theta^*$ for graviton production (blue curves) and for Drell-Yan production (red curves) normalized to the signal, for $M = 1.25 \text{ TeV}/c^2$ (a) and $2.5 \text{ TeV}/c^2$ (b), and for Z' boson (ALRM model) (red curves), normalized to the signal, for $M = 1.25 \text{ TeV}/c^2$ (c) and $2.5 \text{ TeV}/c^2$ (d), with an integrated luminosity of 100 fb^{-1} . The error bars represent the "1-experiment" distribution for the graviton production. The expected background is included in the $\cos \theta^*$ distributions.

As an example, RS gravitons with spin 2 can be distinguished from the Standard Model background and Z' bosons with spin 1 using the distribution of the $\cos \theta^*$ variable, computed as the cosine of the polar angle between the electron and the boost direction of the heavy particle in the latter rest frame. In addition to the cuts defined above, the electron and positron candidates are requested to have opposite charges, in order to identify the electron, from which the $\cos \theta^*$ variable is computed.

The $\cos \theta^*$ distributions for graviton production with $M = 1.25 \text{ TeV}/c^2$, $c = 0.01$, and $M = 2.5 \text{ TeV}/c^2$, $c = 0.1$, are presented in Fig. 14.5, for an integrated luminosity of 100 fb^{-1} . The error bars represent the corresponding statistical uncertainties, applied to the signal distribution obtained from a large statistics simulation. The spin-2 hypothesis is compared to the spin-1 hypothesis (red curve in the figures), formed by the Drell-Yan production (Figs. 14.5(a) and (b)) or the ALRM Z' production (Figs. 14.5(c) and (d)). For graviton production, the expected background is included in the $\cos \theta^*$ distributions.

The spin 2 of RS gravitons can be determined by contrast with the Drell-Yan production or the Z' boson production for an integrated luminosity of 100 fb^{-1} up to $1.25 \text{ TeV}/c^2$ for $c = 0.01$ and $2.5 \text{ TeV}/c^2$ for $c = 0.1$.

14.3 High mass dimuon final states

Many scenarios beyond the Standard Model are expected to manifest themselves through modifications in the mass spectrum of high-mass dimuon pairs. The potential of the CMS experiment to discover dimuon decays of a new heavy neutral gauge boson, Z' , is discussed in Section 3.3.4; the discovery reach for a representative set of Z' models was found to be in the range between 2.9 and 3.8 TeV for an integrated luminosity of 10 fb^{-1} . In this section, we discuss the observability of $\mu^+\mu^-$ final states predicted in two classes of large extra dimensions models, RS and ADD. While the RS scenario gives rise to relatively narrow resonances, the ADD model is expected to be observed via non-resonant modifications of the dimuon spectrum; therefore, these two searches require somewhat different experimental approaches. The search for compositeness in the dimuon channel is described in Section 15.2.

Once a new physics is discovered, observables other than dimuon invariant mass can be used in the attempt to identify the theoretical framework to which it belongs. The measurement of the forward-backward asymmetries of leptonic decay products has long been known as a powerful tool to identify Z' ; some aspects of such a measurement at the LHC are discussed in Section 3.3.5. Spin discrimination of new heavy resonances based on an unbinned likelihood ratio statistic incorporating the angles of the decay products is described in Section 3.3.6.



14.3.1 The Randall-Sundrum Model in the dimuon channel

FIXME: reference to the significance appendix when done; fix for technical jargon We consider the range of RS1 graviton masses in the range $1 < m < 4 \text{ TeV}$ and the dimensionless coupling constant in the expected theoretical range $0.01 \leq c \leq 0.1$ [694]. A full simulation with Pythia [62] version 6.227S and with the GEANT4-based OSCAR package [14] and reconstruction with the CMS full-reconstruction ORCA package [15], version 8_13_2 with a pile-up of minimum-bias collision is carried out to derive both the CMS discovery potential for Randall-Sundrum graviton and spin determination performance in this channel (see details in Ref.[111]). The non-reducible backgrounds are the Drell-Yan process, vector boson pair production $ZZ, WZ, WW, t\bar{t}$ production, etc. In the SM the expected leading-order cross section of the Drell-Yan process dominates the other contributions (see the Section 9.2 for details). The trigger simulation is realized on the basis of the ORCA package, version 8_13_2, using the on-line reconstruction algorithm. We require for each event the trigger bit is in position “double muon” or “single muon”, no requirement for calorimeter isolation of high- p_T muon is made. The total trigger + reconstruction efficiency varies between 95% to 90% for dimuons in the mass range $1 < m < 4 \text{ TeV}$. Only the events which passed both the L1 and HLT cuts are selected. Note that the trigger efficiency is significantly decreased after applying of the calorimeter isolation cuts (down to 15 %). The reason for a such drop can be related with muon shower expansion with an energy due to electromagnetic radiation and interactions of a muon with yoke matter. In the following, any cuts on calorimeter isolation of muon track are not applied at the HLT level.

14.3.1.1 The Randall-Sundrum Model Discovery potential

The significance estimators recommended by CMS PRS group are used for the studying discover potential of the RS1 model: S_{cP} [486] defined as probability from the Poisson distribution with mean N_B to observe greater or equal than $(N_S + N_B)$ events, converted to the equivalent number of sigmas of a Gaussian distribution, $S_{cL} = \sqrt{2((N_s + N_b) \ln(1 + N_s/N_b) - N_s)}$ which is suitable for small event samples and known to be numerically close to S_{cP} [96], and $S_L = \sqrt{2 \ln(\mathcal{L}_{s+b}/\mathcal{L}_b)}$, where \mathcal{L}_b and \mathcal{L}_{s+b} are the best-fit likelihoods of the null hypothesis H_0 (no signal present) and the alternative hypothesis H_1 (signal plus background) (see discussion of S_L in Section 3.3.4.1).

Figure 14.6a shows the integrated luminosity required for a 5σ discovery as a function of the dimuon mass. The results for the different values of an integrated luminosity are summarized in Table 14.3 and Figure 14.6b. The CMS experiment can observe the RS1 graviton with mass up to 2.3 TeV using the first LHC data at $\int L dt = 1 \text{ fb}^{-1}$ if c is equal to 0.1. For $c = 0.01$ the mass reach is not more than 1.9 TeV even for asymptotic regime of LHC operation with $\int L dt = 300 \text{ fb}^{-1}$. The asymptotic reach limit for $c = 0.1$ is 4.5 TeV.




Figure 14.6: Discovery limit for RS1 graviton with $\mu^+ \mu^-$ decay mode for different values of RS1 coupling constant $c = 0.01, 0.02, 0.05$ and 0.1 (from top to bottom). Used discovery limit $S > 5$ for the S_{cP} estimator (solid lines), S_L (dashed lines), S_{cL} (dotted lines).

(b) Reach of the CMS experiment as a function of the coupling parameter c and the graviton mass for various values of integrated luminosity. The left part of each curve is the region where significance exceeds 5σ .

A combined analysis [694] in RS1 scenario shows that the value of the coupling constant c is strongly restricted (Figure 14.6b) due to the theoretical constraints to assure that the model does not introduce a new hierarchy (the scale parameter $\Lambda_\pi = M_{Pl} e^{kr_c \pi} < 10 \text{ TeV}$). The direct comparison of results on a mass reach region for c with the data of the Figure 14.6 shows that the luminosity of 100 fb^{-1} is needed to test the RS1 model everywhere in $(c - M_{\text{grav}})$ space of model parameters. However, these conclusions are not definitive since initial theoretical constraints are quite arbitrary.

Table 14.3: CMS discovery potential invariant mass reach (in TeV) to observe the RS1 graviton in $\mu^+\mu^-$ channel.

Coupling constant c	Estimator	1 fb^{-1}	10 fb^{-1}	100 fb^{-1}	300 fb^{-1}
0.01	S_{cP}	0.75	1.20	1.69	1.95
	S_{cL}	0.77	1.21	1.71	1.97
	S_L	0.78	1.23	1.73	1.99
0.02	S_{cP}	1.21	1.72	2.30	2.63
	S_{cL}	1.22	1.72	2.31	2.64
	S_L	1.22	1.74	2.34	2.68
0.05	S_{cP}	1.83	2.48	3.24	3.67
	S_{cL}	1.85	2.49	3.26	3.71
	S_L	1.85	2.51	3.31	3.79
0.1	S_{cP}	2.34	3.11	4.12	4.52
	S_{cL}	2.36	3.13	4.14	4.54
	S_L	2.36	3.16	4.23	4.73

14.3.1.2 Systematic Uncertainties

The results taking into account the systematical uncertainties are shown in Figure 14.7. The expected effects of misalignment are considered in two misalignment scenarios: the First Data and the Long Term scenarios [93] which correspond to different stages of the alignment corrections for the positions of the tracker and muon chambers. The current estimate is that the transition to the Long Term scenario can be achieved at an integrated luminosity of about 1 fb^{-1} [78]. In contrast to Figure 14.6 which assumed K-factor equal to unity, K-factor of $K = 1.30 \pm 0.05$ is used both for the RS1 signal and Drell-Yan background. Additional variations due to EW corrections, hard-scale and PDF uncertainties are considered, the details are given in Ref.[111].

14.3.1.3 Spin Discrimination in Angular Analysis

A study of muon angular distributions can allow to discriminate hypotheses of Graviton (spin-2 particle) and Z' (spin-1 particle) – see the discussion and the results in Section 3.3.6

FIXME: Use LaTeX label.

14.3.2 The ADD model in the dimuon channel

FIXME: fix for technical CMS jargon, fix for structure We consider the fundamental Planck scale of the ADD model in the range of $3.0 < M_S < 10.0 \text{ TeV}$ and numbers of extra dimensions in the range of $3 \leq n \leq 6$ [673]. The contribution of KK-modes of ADD gravitons to the Drell-Yan processes is computed using the leading-order matrix element [695] which was implemented in STAGEN generator collection as external matrix element in Pythia [62] version 6.227. A full simulation with the GEANT4-based OSCAR package [14] and reconstruction with the CMS full-reconstruction ORCA package [15], version 8.13.2 without a pile-up of minimum-bias collision is carried out to derive the CMS discovery potential for ADD virtual graviton (see details in Ref.[?]). The non-reducible backgrounds are the Drell-Yan process, vector boson pair production $ZZ, WZ, WW, t\bar{t}$ production, etc. In the SM the expected leading-order cross section of the Drell-Yan process dominates the other contributions (see




Figure 14.7: (a) Discovery limit for coupling constants $c = 0.01, 0.1$ (upper and lower curves, respectively) after taking into account the systematic uncertainties including misalignment in two scenarios: the curves ending at integrated luminosity of 1fb^{-1} correspond the First Data misalignment scenario, the other ones correspond to the Long Term scenario. The ranges show the expected variations due to the systematic uncertainties. (b) The ranges of the expected variations due to the systematic uncertainties for the mass reach of the CMS experiment.

the Section 9.2 for details). The trigger simulation is realized on the basis of the ORCA package, version 8.13.2, using the on-line reconstruction algorithm. We require for each event the trigger bit is in position “double muon” or “single muon”, no requirement for calorimeter isolation of high- p_T muon is made. The total trigger + reconstruction efficiency varies between 70% to 90% for dimuons in dependence on the model parameters. Only the events which passed both the L1 and HLT cuts are selected.

14.3.2.1 The ADD discovery limit

The significance estimators recommended by CMS PRS group are used for the studying discover potential of the RS1 model: the S_{cP} definition [486] is close to the significance function $S_{cL} = \sqrt{2((N_S + N_B) \ln(1 + N_B/N_B) - N_S)}$ suitable for small event samples [96] which has been shown to give practically the same numerical values as S_{cP} estimator, where N_S is the number of signal events passed through all kinematics cuts and N_B is the number of background events.

The computed significance values for the ideal detector as a function of a fundamental theory scale, M_S , are presented in Fig. 14.8 for integrated luminosities of 0.1, 1.0, 10, 100, 300, 1000 fb^{-1} . The main observations are:

- $\int \mathcal{L} dt = 1 \text{fb}^{-1}$, even a low luminosity regime allows to measure the effect from the virtual contributions of ADD gravitons to Drell-Yan process for the effective fundamental Planck scale of 4.0 TeV for the most unfavorable case with $n = 6$. For more promising scenario where the number of extra dimensions is minimal




Figure 14.8: Significance as a function of M_S for (a) $n = 3$ and (b) $n = 6$.

($n = 3$) the reach limit is extended up to 5.8 TeV.

- $\int \mathcal{L} dt = 10 \text{ fb}^{-1}$, M_S values of 4.8 and 7.2 TeV can be reached for $n = 3$ and $n = 6$ respectively.
- $\int \mathcal{L} dt = 100 \text{ fb}^{-1}$, one year of LHC operation in a high luminosity regime provides observation of the ADD signal at $5.8 \div 8.7$ TeV of model scale in dependence on a number of extra dimensions.
- $\int \mathcal{L} dt = 300 \text{ fb}^{-1}$, in asymptotic regime the CMS sensitivity to fundamental Planck scale is increased up to values of $6.5 \div 9.3$ TeV.

The another common method used in experimental physics to estimate the significance is the estimator S_{c12} which is the most close to the likelihood estimator $S_{\mathcal{L}}$ which is the most optimal for the small-statistics background as was discussed in Ref. [94]. The ambiguousness due to usage of different estimators ($2 \div 40$ %) leads to the uncertainties in the M_S reach up to 8.6 % in dependence of the signal-to-background ratio.

14.3.2.2 Systematics

The results taking into account the systematical uncertainties with the S_{cP} estimator are shown in Figure 14.9. To take into account the misalignment effect two scenario of misalignment were considered during reconstruction procedure: *First Data* scenario [93] for 0.1 and 1.0 fb^{-1} and *Long Term* scenario [93] for 10, 100 300, 1000 fb^{-1} . The K-factor of $K = 1.30 \pm 0.05$ is used both for ADD signal and Drell-Yan background. Additional variations due to hard-scale and PDF uncertainties as well as trigger and selection uncertainties are considered, the results are given in Ref.[?].

4 High energy single lepton final states

FIXME: plots are now updated; fix contribution for length; fix for structure

14.4.1 Introduction

Several theoretical models predict, in addition to the well known electroweak vector bosons γ, W, Z , further heavy gauge bosons. These additional particles are postulated for example in Left-Right Symmetric Models [696, 697, 698, 699], based on the gauge group $SU(3)_C \times$




Figure 14.9: 5σ limit on M_S for the number of extra dimensions $n = 3, 4, 5, 6$.

$SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ (B,L: baryon-, lepton-number) in theories predicting a substructure of the known “elementary particles”, and in Little Higgs Models [84].

Here we investigate the detection capabilities for a hypothetical heavy partner of the W , a charged spin-1 boson W' . We do not assume one of the specific models mentioned above, but derive the W' properties from the Reference Model by Altarelli [700], which has been used in several earlier experiments, so that the resulting limits can be compared easily. In this Reference Model the W' is a carbon copy of the W , with the very same left-handed fermionic couplings (including CKM matrix elements), while there is no interaction with the Standard Model gauge bosons or with other heavy gauge bosons as a Z' . Thus the W' decay modes and corresponding branching fractions are similar to those for the W , with the notable exception of the $t\bar{b}$ channel, which opens for W' masses beyond 180 GeV.

In hadron collisions W' bosons can be created through $q\bar{q}$ annihilation, in analogy to W production. Previous searches for the Reference W' at LEP and at the Tevatron give rise to lower bounds approaching 1 TeV [701].

This analysis is based on the decay $W' \rightarrow \mu\nu$, with a branching ratio of roughly 10%. The resulting signature of a high energy muon accompanied by missing energy allows an easy separation of signal and background reactions.

14.4.2 Data Samples

For this study we assume an integrated luminosity of 10 fb^{-1} and an average instantaneous luminosity of $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ corresponding to an average pile-up of 3.5 pp -collisions per bunch crossing.

Reference Model W' events decaying into muon and neutrino have been generated with Pythia v6.227 [62], based on the leading order cross section and the parton density functions CTEQ 5L (leading order) [692]. In total about 300 000 events have been produced for W'

masses between 1 TeV and 8 TeV, using CMKIN (v4.3.1). The product of LO cross section and branching fraction varies between $3.0 \cdot 10^3$ fb (1 TeV) and $3.3 \cdot 10^{-4}$ fb (8 TeV), to be compared with $1.7 \cdot 10^7$ fb for Standard Model W production and muonic decay. The detector response was simulated with OSCAR (v3.6.5) and ORCA (v8.7.1).

Both the signal events and the following background samples were analysed with ORCA (v8.7.3): $W \rightarrow \mu\nu$, $Z \rightarrow \mu\mu$, WW incl, ZZ incl, ZW incl, $t\bar{t}$ incl. These data sets have been produced in the CMS Data Challenge 2004. On average 3.5 minimum bias reactions have been overlaid to each event.

14.4.3 Event selection and analysis

Events have been preselected requiring at least one globally reconstructed muon which pass the trigger criteria.

The final cuts to select $W' \rightarrow \mu\nu$ candidate events are:

- **muon quality:** at least 13 hits along the global track, $\chi^2/N_{dof} < 50$ for the fit.
- **single muon** requirement
- **muon isolation:** no additional track ($p_T > 0.8$ GeV) within a cone of size $\Delta R = 0.17$.

The cuts have been chosen to maximise the signal/background ratio.




Figure 14.10: **Left:** transverse invariant mass spectrum of signal (1 and 5 TeV, non-stacked) and background (stacked) after applying the selection cuts. **Right:** result of the CL_s -method: with an integrated luminosity of 10 fb^{-1} Reference W' bosons can be excluded up to a mass of 4.7 TeV.

For the selected events the transverse mass

$$M_T = \sqrt{2p_{T\mu} E_T^{\text{miss}} (1 - \cos \Delta\phi_{\mu, E_T^{\text{miss}}})}$$

is calculated from the muon transverse momentum $p_{T\mu}$, the missing energy component in the transverse plane E_T^{miss} and the angular $\Delta\phi_{\mu, E_T^{\text{miss}}}$ between both in this plane. Fig. 14.10 shows the resulting distribution for signal (1 and 5 TeV) and background events. The W' boson

distributions show a Jacobian peak which is spread out for large M_T due to the detector resolution. It can be seen immediately, that a 1 TeV boson can be discovered or excluded easily, while for higher masses a statistical analysis is needed to quantify the sensitivity.

14.4.4 Discovery and Exclusion Potential

To interpret the results, the CL_s method [495] is applied, which is based on the likelihood ratios, calculated for all bins of the M_T distribution. CL_s is defined as ratio of the confidence levels for the signal and background hypotheses, $CL_s = CL_{s+b}/CL_b$.

Figure 14.10 shows, that for an integrated luminosity of 10 fb^{-1} , a limit of 4.7 TeV at the 95% CL is reachable, if no signal is present in the CMS data. Both the expected discovery and exclusion limits are displayed in figure 14.11 as function of integrated luminosity and W'




Figure 14.11: The plots show which integrated luminosity is needed to discover (left) or exclude (right) W' bosons of a certain mass.

mass. To investigate the sensitivity to the signal and background cross sections, they have been varied in a wide range; relative changes by factors of 2 and 10, respectively, lead to a lowering of the accessible mass range by about 0.5 TeV in the worst case.

14.4.5 Systematic Uncertainties

The uncertainties arising from an imperfect knowledge of the PDFs at LHC energies and the error from the hard scale parameters have been investigated by using the Les Houches Accord PDFs [88] and varying the hard scale, respectively. The relative errors on the cross-section of the signal are listed in table 14.4. The error on the background is comparable to that of the W' at the corresponding invariant mass.

The steep falling invariant mass distribution especially of the W background holds a potential danger for the detection of W' bosons: if only a small fraction of these events is reconstructed with a by far too large mass, which might result from a mismeasured muon momentum, the detection of a W' becomes extremely difficult. Such a behavior would be visible in so called non-gaussian tails for example in the p_T resolution distribution. Using a large sample of a W events it could be demonstrated, that the alignment precision expected

Systematic Uncertainties					
Type	1 TeV W'	2 TeV W'	3 TeV W'	4 TeV W'	5 TeV W'
PDF $\Delta\sigma/\sigma$	+3.6 -4.3	+6.8 -5.9	+6.2 -8.3	+17.1 -10.6	+33.7 -18.9
Hard Scale $\Delta\sigma/\sigma$	+4.1 -4.1	+7.5 -6.9	+10.4 -9.2	+13.1 -10.3	+14.8 -12.7
Luminosity $\Delta\mathcal{L}/\mathcal{L}$	$\pm 5\%$	$\pm 5\%$	$\pm 5\%$	$\pm 5\%$	$\pm 5\%$

Table 14.4: Systematic uncertainties arising from an imperfect theoretical knowledge (parton density functions, hard scale) and the expected luminosity error for an intergrated luminosity of 10 fb^{-1} .

after an intergrated luminosity of 10 fb^{-1} has only a small influence on the non-gaussian tails of the muon p_T resolution distribution.

The luminosity uncertainty at the considered integrated luminosity of 10 fb^{-1} is expected to be 5%, while other experimental errors (neutron background, dead detector components, etc.) are expected to be negligible.

14.4.6 Summary

For an integrated luminosity of 10 fb^{-1} , W' bosons of the Reference Model can be disc ed or excluded up to a mass of 4.5–5 TeV, from an analysis of the muonic decay mode.

14.5 High mass di-jet final states

14.5.1 Dijet Resonances and Contact Interactions

Dijet resonances and contact interactions are the two major signals of new physics with dijets. Dijet resonances are direct and compelling observations of a new physical object at a mass M , requiring an incoming parton-parton collision energy equal to the mass. Contact interactions (discussed in section ??) are indirect observations of an energy scale of new physics, Λ , which can be significantly larger than the available collision energy. Resonances are clear signals but contact interactions are often observed first.

14.5.2 Dijet Resonance Search

We search for processes producing narrow resonances, X , decaying to dijets: $pp \rightarrow X \rightarrow \text{jet} + \text{jet}$ (inclusive) [702]. Our experimental motivation is that LHC is a parton-parton collider, and resonances made from partons must decay to the same partons giving two jets in the final state. The theoretical motivation is broad, since there are many models that predict narrow dijet resonances.

14.5.2.1 Dijet Resonance Models

Figure 14.12 we show the cross section times branching ratio calculated to lowest order




Figure 14.12: Left) The total cross section for dijet resonances from eight different models (see text). Right) For resonance masses of 0.7, 2.0, and 5.0 TeV, the fractional difference between an excited quark (solid curve) or an E6 diquark (dashed curve) and the QCD dijet background is compared to the QCD statistical errors (vertical lines).

for eight benchmark models. Here we introduce them in order of descending cross section at low mass. Excited states of composite quarks [703] are strongly produced giving large cross sections ($q\bar{q} \rightarrow q^*$). Axigluons [704] or colorons [705] from an additional color interaction are also strongly produced, but require an antiquark in the initial state ($q\bar{q} \rightarrow A$ or C)

slightly reducing the cross section compared to excited quarks. Diquarks [706] from superstring inspired E_6 grand unified models are produced with electromagnetic coupling from the valence quarks of the proton ($ud \rightarrow D$). The cross section for E_6 diquarks at high mass is the largest of all the models considered, because at high parton momentum the probability of finding a quark in the proton is significantly larger than the probability of finding a gluon or antiquark. Color octet technirhos [707] from topcolor-assisted technicolor are produced for either gluons or quark-antiquark pairs in the initial state through a vector-dominance model of mixing between the gluon and the technirho ($q\bar{q}, gg \rightarrow g \rightarrow \rho_{T8}$). Randall-Sundrum gravitons [87] from a model of large extra dimensions are produced with a significant cross section at masses below $1 \text{ TeV}/c^2$ primarily from gluons in the initial state ($q\bar{q}, gg \rightarrow G$). Heavy W bosons [708] inspired by left-right symmetric grand unified models have electroweak couplings and require antiquarks for their production ($q_1\bar{q}_2 \rightarrow W'$) giving small cross sections. Heavy Z bosons [708] inspired by grand-unified models are widely anticipated by theorists, but they are weakly produced, and require an antiquark in the initial state ($q\bar{q} \rightarrow Z'$), so their production cross section is around the lowest of the models considered. Lower limits from CDF [114] and D0 [115] on the mass of these models are between 0.4 and $1.0 \text{ TeV}/c^2$.

14.2.2 Dijet Resonance Sensitivity Estimates




Figure 14.13: Likelihoods for observing a narrow dijet resonance of mass $2 \text{ TeV}/c^2$ in a 1 fb^{-1} data sample that contains only QCD background (left) and a data sample that also contains a resonance with a significance of 5σ (right) are shown with statistical uncertainties only (dashed) and including systematics (solid).

The signal and background dijet mass distributions for narrow resonances were presented in section 4.1.4. In Figure 14.12 we demonstrate the size of the signal for excited quarks and E_6 diquarks compared to the QCD background and its statistical uncertainty. It is clear that we will be sensitive to such large signals for strongly produced dijet resonances. Here we quantify our sensitivity to any model of narrow dijet resonances. In Figure 14.13 we show examples of likelihoods for excluding or observing a narrow resonance signal on a QCD background as a function of the signal cross section. In the case where the observed sample is QCD only, the signal likelihood peaks around zero cross section, and the 95% CL excluded

signal cross section is shown. In the case where the observed sample is QCD plus a resonance signal, we have varied the signal size until the Gaussian distributed likelihood is 5σ above zero. In figure 14.13 we have included estimates of our systematic uncertainties. For a resonance mass of 0.7 (5.0) TeV the systematic uncertainty on the observable signal cross section due to the jet energy uncertainty in the background rate is 15% (25%), the uncertainty due to jet resolution in the resonance shape is 10% (10%), the uncertainty due to radiation's affect on the resonance shape is 10% (25%), and the uncertainty due to luminosity is 10% (10%). For resonance masses just above the dijet mass thresholds where the trigger prescale decreases, there is an additional systematic uncertainty from the jet energy uncertainty. Systematic uncertainties have a greater effect on discovery than exclusion, because exclusions occur at a smaller signal cross section and are dominated by statistical uncertainties.

Figure 14.14 demonstrates that the 95% CL exclusion and 5σ discovery signal cross sections,




Figure 14.14: Left) For resonances of mass 0.7, 2.0 and 5.0 TeV, the rate as a fraction of QCD that CMS expects to exclude (dashed) or discover (solid) including statistical uncertainties only. Right) The resonance cross section that CMS expects to exclude (boxes) or discover (circles), including systematic uncertainties, is compared to the cross section for eight resonance models.

including statistical uncertainties only, have reasonable values when compared to the size of the QCD statistical errors. Also in Figure 14.14 we present the resonance cross section values for jet $|\eta| < 1$ that CMS can expect to exclude at 95% CL or discover at 5σ significance for an integrated luminosity of 1 fb^{-1} . These can be compared with the cross section of any model of narrow dijet resonances, and here we compare with our benchmark models. From Figure 14.14 we can read off the mass limits or discoveries that are possible with 1 fb^{-1} of data, which are listed in Table 14.5 along with the results of repeating the same analysis for 100 pb^{-1} and 10 fb^{-1} . The resonances that are produced via the color interaction (excited quarks, axigluons, colorons and color octet technirhos) or from the valence quarks of each proton (E_6 diquarks) have large cross sections and can be discovered up to a mass of a few TeV. For every model of dijet resonances we have considered here, even the weakly produced models, there is some range of mass that CMS will be able to exclude at 95% CL with sufficient luminosity. A single search for resonances in the dijet mass distribution

Resonance Model	95% CL Excluded Mass (TeV/ c^2)			5 σ Discovered Mass (TeV/ c^2)		
	100 pb $^{-1}$	1 fb $^{-1}$	10 fb $^{-1}$	100 pb $^{-1}$	1 fb $^{-1}$	10 fb $^{-1}$
Excited Quark	0.7 - 3.6	0.7 - 4.6	0.7 - 5.4	0.7 - 2.5	0.7 - 3.4	0.7 - 4.4
Axigluon or Coloron	0.7 - 3.5	0.7 - 4.5	0.7 - 5.3	0.7 - 2.2	0.7 - 3.3	0.7 - 4.3
E_6 diquark	0.7 - 4.0	0.7 - 5.4	0.7 - 6.1	0.8 - 2.0	0.8 - 3.7	0.8 - 5.1
Color Octet Technirho	0.7 - 2.4	0.7 - 3.3	0.7 - 4.3	0.7 - 1.5	0.7 - 2.2	0.7 - 3.1
Randall-Sundrum Graviton	0.7 - 1.1	0.7 - 1.1 1.3 - 1.6	0.7 - 1.1 1.3 - 1.6 2.1 - 2.3	N/A	N/A	N/A
W'	0.8 - 0.9	0.8 - 0.9 1.3 - 2.0	0.8 - 1.0 1.3 - 3.2	N/A	N/A	N/A
Z'	N/A	N/A	2.1 - 2.5	N/A	N/A	N/A

Table 14.5: Sensitivity to dijet resonances with 100 pb $^{-1}$, 1 fb $^{-1}$, and 10 fb $^{-1}$. For each resonance model, we show the range of masses we expect to be able to exclude at a confidence level of 95% or greater, and the range of masses we expect to be able to discover with a significance of 5 σ or greater. All estimates are with both statistical and systematic uncertainties.

provides CMS with a sensitive test of many different models of the widely anticipated new physics at the TeV scale.

14.6 High mass di-photon final states

14.6.1 Introduction

The study of the Randall-Sundrum(RS) graviton decaying into the two photons is particularly interesting as the detection of such few TeV/ c^2 mass resonance in such channel together with its observation in the di-lepton channel will sign a RS graviton, distinguishing it from a Z' production. The model is governed by two parameters: the graviton mass M and its coupling to Standard Model particles c , the latter being related to the natural width of the resonance.

14.6.2 Event generation and kinematics preselection

The search for the $G \rightarrow \gamma\gamma$ signal at LHC is affected by four types of backgrounds:

- The prompt di-photon production from the quark annihilation and gluon fusion diagrams, which provides an intrinsic or 'irreducible' background.
- The γ + jets production consisting of two parts: i) prompt photon from hard interaction + the second photon coming from the outgoing quark due to final state radiation and ii) prompt photon from hard interaction + the decay of a neutral hadron (mostly isolated π^0) in a jet, which could fake a real photon.
- The background from QCD hadronic jets, where electromagnetic energy deposits result from the decay of neutral hadrons (especially isolated π^0 s) in both jets.
- Drell Yan process with $e + e^-$ in a final state which could mimic photons when correspondent electron tracks will not be assigned to the superclusters during the reconstruction.

Generator-level preselection for QCD and bremsstrahlung backgrounds is described in [709] and [710]. Parameters which were used for these preselections in case of QCD and bremsstrahlung backgrounds are defined in [710].

14.6.3 Offline selection and analysis

The analysis was done as follows:

- 1 To find two super-clusters (SCs) with $E_T > 150$ GeV and two HLT trigger bits triggered at the same time: 2p (two photons) and r2p (two photons relaxed).
- 2 Calorimeter isolation criteria: for each SC the energy in a cone of $\Delta R = 0.5$ (excluding SC itself) should be $< 0.02E_T(SC)$
- 3 $E(HCAL)/E(ECAL) < 0.05$
- 4 Tracker isolation: the sum of the energy of all tracks in a cone $\Delta R = 0.5$ around the SC should be $< 0.01E_T(SC)$
- 5 Photon energy corrections are done in a simple way so far:
 - For E1 energy < 1.7 TeV, only a simple energy dependent part of correction is applied (just a shift of the peak)
 - For E1 energy > 1.7 TeV, the MGPA saturation correction (1d) was applied (see and [711])

14.6.4 K-factors

To produce the final results and to calculate the expected statistical significance for RS-1 graviton search recently calculated next-to-leading order corrections (K factors) to the cross sections of different types of background are used: $K = 1.5$ for quark annihilation [16], $K = 1.2$ for gluon fusion [19], $K = 1$ for the $\gamma +$ hadronic jets [19] and $K = 1$ for QCD jets. For signal, a conservative $K = 1$ value is taken.

14.6.5 Results

The number of events passed through the analysis cuts described above, for the signal and for the backgrounds are presented in Table 14.6 ($1.5 \text{ TeV}/c^2, 0.01$) and in Table 14.7 ($3.5 \text{ TeV}/c^2$,

Table 14.6: Number of events passed through the analysis cuts defined above for $M_G = 1.5 \text{ TeV}/c^2$, $c = 0.01$ and $L = 30 \text{ fb}^{-1}$. Leading column is non-saturated events, all saturated events, passed through the analysis, were added in brackets, where applied.

	signal	Born (k=1.5)	Box (k=1.2)	Brem (k=1)	QCD (k=1)	DY (k=1)
trigger + 2SC	28.9	8.6	0.10	29.2	798.7	4.3
+ EM isolation	24.5	5.5	0.08	20.3	361.8	3.5
+ HCAL/ECAL	24.3	5.4	0.08	4.4	12.8	3.5
+ tracker isolation	17.6	4.2(+0.2)	0.05	0.17	0.0	0.0

0.1).

Final figures (see Figure 14.15) for the number of events passed through the analysis for both

Table 14.7: Number of events passed through the analysis cuts defined above for $M_G = 3.5$ TeV/c^2 , $c = 0.1$ and $L = 30 \text{ fb}^{-1}$. Leading column is non-saturated events, all saturated events, passed through the analysis, were added in brackets, where applied.

	signal	Born (k=1.5)	Box (k=1.2)	Brem (k=1)	QCD (k=1)	DY (k=1)
trigger + 2SC	11.6	0.20	$4.4 * 10^{-4}$	0.78	821.9	0.10
+ EM isolation	10.8	0.14	$3.6 * 10^{-4}$	0.32	164.4	0.095
+ HCAL/ECAL	10.6	0.13	$3.4 * 10^{-4}$	0.016	0.0	0.095
+ tracker isolation	8.9(+1.0)	0.10(+0.02)	$2.7(+0.24) * 10^{-4}$	$1.7 * 10^{-3}$	0.0	$7.2 * 10^{-4}$




Figure 14.15: Number of events passed through all cuts for (1.5 TeV/c^2 , 0.01) (left) and (3.0 TeV/c^2 , 0.1) (right) RS-1 gravitons for 30 fb^{-1} integrated luminosity.

signal and backgrounds are presented for cases (1.5 TeV/c^2 , 0.01), (3.0 TeV/c^2 , 0.1) in case of 30 fb^{-1} luminosity. One year low luminosity 10 fb^{-1} results are presented on Figure 14.16.




Figure 14.16: Number of events passed through all cuts for (1.0 TeV/c^2 , 0.01) (left) and (2.5 TeV/c^2 , 0.1) (right) RS-1 gravitons for 10 fb^{-1} integrated luminosity.

Taking into account the K-factors described above, we have got next number of events for signal and background and calculated significance for $c = 0.01$ and $c = 0.1$, for the $L = 30 \text{ fb}^{-1}$. $S = \sqrt{2 \ln Q}$, with $Q = (1 + n_s/n_b)^{n_s+n_b} \exp(-n_s)$ in Tables 14.8 and 14.9.

Table 14.8: Significance for $c = 0.01$ and $L = 30 fb^{-1}$

	$M_G = 1.0$ TeV/c^2	$M_G = 1.25$ TeV/c^2	$M_G = 1.5$ TeV/c^2	$M_G = 1.75$ TeV/c^2	$M_G = 2.0$ TeV/c^2
N_s	135.8	44.0	17.6	7.3	3.9
N_{bkg}	15.0	8.8	4.6	1.8	1.2
Significance	20.6	10.1	5.9	3.9	2.6

Table 14.9: Significance for $c = 0.1$ and $L = 30 fb^{-1}$

	$M_G = 2.5$ TeV/c^2	$M_G = 3.0$ TeV/c^2	$M_G = 3.5$ TeV/c^2	$M_G = 4.0$ TeV/c^2	$M_G = 4.5$ TeV/c^2
N_s	103.8	31.6	9.9	3.44	1.11
N_{bkg}	1.11	0.35	0.13	0.06	0.02
Significance	27.3	15.0	8.2	4.6	2.6

Significance as a function of the graviton mass (M_G) for low luminosity at $10 fb^{-1}$, $30 fb^{-1}$, $60 fb^{-1}$ integrated luminosities are given in Figure 14.17.




Figure 14.17: Significance as a function of the graviton mass for $10 fb^{-1}$ and $30 fb^{-1}$ integrated luminosities, $c=0.01$ (left) and $c=0.1$ (right)

The discovery region in the plane of the coupling parameter c and the graviton mass is given in Fig. 14.18.

Recent results are :




Figure 14.18: Reach of the CMS experiment in the search for the Randall-Sundrum graviton decaying into diphoton channel as a function of the coupling parameter c and the graviton mass for 10 fb^{-1} , 30 fb^{-1} and 60 fb^{-1} . The left part of each curve is the region where the significance exceeds 5σ .

- 5σ discovery limits for 60 fb^{-1} :
 - $c = 0.01 \quad M_G = 1.82 \text{ TeV}/c^2$
 - $c = 0.1 \quad M_G = 4.27 \text{ TeV}/c^2$
- 5σ discovery limits for 30 fb^{-1} :
 - $c = 0.01 \quad M_G = 1.61 \text{ TeV}/c^2$
 - $c = 0.1 \quad M_G = 3.95 \text{ TeV}/c^2$
- 5σ discovery limits for 10 fb^{-1} :
 - $c = 0.01 \quad M_G = 1.31 \text{ TeV}/c^2$
 - $c = 0.1 \quad M_G = 3.47 \text{ TeV}/c^2$

14.6.6 Limits uncertainties for 30 fb^{-1}

1) Hard scale confidence limits uncertainties for 30 fb^{-1} can be found in Table 14.10

Table 14.10: Hard scale confidence limits uncertainties for 30 fb^{-1}

	$4\hat{s}$	$0.25\hat{s}$
$c = 0.01$	$-62 \text{ GeV}/c^2$	$+56 \text{ GeV}/c^2$
$c = 0.1$	$-47 \text{ GeV}/c^2$	$+42 \text{ GeV}/c^2$

2) LHApdf confidence limits uncertainties are

- $c = 0.01, -55 \text{ GeV}/c^2$
- $c = 0.1, -152 \text{ GeV}/c^2$

- 3) There is another source of uncertainties due to the fact, that we have used K-factor = 1.5 for born process, while Tevatron most recent measures pointed out that this K-factor is closer to 2 [712]:



- $c = 0.01, -50 \text{ GeV}/c^2$
- $c = 0.1, -30 \text{ GeV}/c^2$

7 Single γ final state with E_T^{miss} from Extra Dimensions

FIXME: need a word for the jet+MET analysis of Leonardo which did not make it in ; also fix for figure long empty spaces

14.7.1 Topology of single-photon final states

An introduction to the signals involving direct graviton emission in ADD type of extra dimensions frameworks is given in section ???. The topology of single photon events can be identified by:

- a single high p_T photon in the central η region
- high missing p_T back-to-back to the photon in the azimuthal plane with a similar p_T distribution.

These characteristics are not strongly dependent on the ADD model parameters.

14.7.2 Backgrounds from the Standard Model

All signal and background samples used in the following were simulated using the CMS fast detector simulation FAMOS [287]. Fully simulated reference samples were generated for the signal and the largest irreducible background, $Z^0\gamma \rightarrow \nu\bar{\nu} + \gamma$. A detailed comparison of the resolution, efficiency and purity of all reconstructed objects used in this analysis to the Geant-based CMS simulation confirmed that the fast simulation provides a very good approximation of the expected detector response. All samples were consistently generated using a generator level cut in PYTHIA $\hat{p}_T > 400 \text{ GeV}$. The backgrounds considered in the study are, $Z^0\gamma \rightarrow \nu\bar{\nu} + \gamma$, $W^\pm \rightarrow \ell\nu$ where ℓ is electron, muon or tau, $W^\pm\gamma \rightarrow e\nu + \gamma + \text{Jets}$, QCD, di- γ and $Z^0 + \text{jets}$. Non-beam events like beam halo and cosmic muons are not considered in this study. For the main background, a normalization method from measured data is developed employing the reconstructed leptonic decays of the Z^0 into muon and electron pairs.

The detector acceptance for selecting the leptons is parameterised using a two-dimensional function $\alpha(p_T^\gamma, \eta_\gamma)$. Figure 14.19 shows the measured and the p_T^γ spectrum from $\gamma + Z^0 \rightarrow \mu^+\mu^-$ after the (acceptance \times efficiency) parameterization is applied, in comparison with the generator spectrum for $\gamma + Z^0 \rightarrow \nu_i\bar{\nu}_i$ events. For $p_T^\gamma > 100 \text{ GeV}$ there is $1170 Z^0 \rightarrow \mu^+\mu^-/e^+e^-$ events expected after all selection cuts for 30 fb^{-1} . These can be used as the candle sample that provides a direct normalization of the $\gamma + Z^0 \rightarrow \nu_i\bar{\nu}_i$ with a statistical precision of 3%.




Figure 14.19: Number of expected p_T^γ events per 25 GeV bin at 1 fb^{-1} from measured $\gamma + Z^0 \rightarrow \mu^+ \mu^-$ events before and after transformation compared with the generator distribution for $\gamma + Z^0 \rightarrow \nu_i \bar{\nu}_i$ – the transformed muon distribution models the $\nu_i \bar{\nu}_i$ spectrum well.

14.7.3 Event Selection

The main trigger path for the selection of signal and background events will be the single photon trigger, both at the L1 and the HLT. Presently the single photon trigger has a HLT level threshold of 80 GeV, which is far below the selection cut for events with isolated photons above 400 GeV. Hence the expected trigger efficiency is close to 100% and its efficiency can be monitored from data with a E_T^{miss} trigger which will have a threshold in the range of 200 – 300 GeV, well below the acceptance of the bulk of the signal. Both the topological characteristic and the necessity to reduce the Standard Model background lead to the following selection criteria:

- At least a $E_T^{\text{miss}} > 400 \text{ GeV}$ is required and the photon p_T has to be above 400 GeV.
- $|\eta|$ of the photon < 2.4
- $\Delta\phi(E_T^{\text{miss}}, \gamma) > 2.5$
- A track veto for high p_T tracks $> 40 \text{ GeV}$ is applied. This is a powerful criterion to reduce all backgrounds containing high-energetic charged particles (such as e^\pm , μ^\pm , jets)
- An Isolated Photon Likelihood criterion is applied to remove residual background from hard photon emission from jets as well as fake photons from jets.

Figure 14.20 shows the missing transverse energy spectra for events surviving the selection path for both the signal and the backgrounds. As expected the $Z^0 \gamma$ is by far the most dominant component of the background, followed by $W^\pm \gamma$ while the contributions of the other Standard Model backgrounds are small. For all ADD cross section the hard truncation approach is used (see section 14.1), i.e. events with $M_G < M_D$ are rejected.

14.7.4 Systematic uncertainties and discovery potential

We consider an uncertainty of 2% for the measurement of the photon p_T^γ in the electromagnetic calorimeter and an uncertainty of 5% for the E_T^{miss} measurement. The resulting decrease of the significance is 1.0% and 1.6% respectively. For the main background the systematics can be reduced to the luminosity measurement using the Z^0 candle calibration method. It can thus be measured with a precision of 3% after 30 fb^{-1} . The 5σ discovery reach is achievable for $M_D < 2.5 \text{ TeV}$ and all values of extra dimensions while for $M_D < 3 \text{ TeV}$ 5σ reach is achievable for n between 2 and 4. Figure 14.21 shows the expected significances as function of M_D .

8 Black Holes (BHs)

FIXME: edited may 16- some fixes rmain

14.8.1 Introduction to Higher-dimensional BHs

One of the consequences of large extra dimensions is the possibility to produce microscopic black hole (BH) at LHC energies. Such a BH formed in a $(4+n)$ -dimensional space-time has a Schwarzschild radius

$$r_{s(4+n)} = \frac{1}{\sqrt{\pi} M_{(4+n)}} \left(\frac{M_{BH}}{M_{(4+n)}} \left(\frac{8\Gamma((n+3)/2)}{n+2} \right) \right)^{\frac{1}{n+1}} \quad (14.17)$$




Figure 14.20: Spectrum of the missing E_T for all backgrounds (black histogram) and for an example signal sample ($M_D = 2.5 \text{ TeV}$, $n = 2$). The number of events corresponds to an integrated luminosity of 30 fb^{-1} .

where $M_{(4+n)}$ is the reduced Planck scale and n is the number of large extra dimensions [713]. A high energy collision of two partons can result in the formation of a BH when the impact parameter is smaller than $r_{s(4+n)}$. In the semi-classical approach the BH cross section is given by $\sigma(M_{\text{BH}}) = \pi r_{s(4+n)}^2$ at the parton level. If for low masses $M_{(4+n)}$, ie around 2 TeV, the BH production cross sections at the LHC is in the pb range.

Once produced, these BHs are expected to decay thermally via Hawking radiation [714]. The Hawking temperature for a BH in $4 + n$ dimensions is [715]

$$T_{(4+n)} \sim M_{(4+n)} (M_{(4+n)}/M_{\text{BH}})^{1/(n+1)} \quad (14.18)$$

These BHs have a very short lifetime typically of $\sim 10^{-27}$ seconds.

BH events are expected to evaporate democratically by emission of all particle types that exist in nature, independent of their spin, charge, quantum numbers or interaction properties. Therefore they can be a source of new particles. BH physics at the LHC can provide the possibility of probing quantum gravity in the lab.

14.8.2 Analysis selection path and results

Black hole event samples were produced using the CHARYBDIS event generator [716]. As a benchmark the case which is analyzed has the following parameters: a) 2 TeV effective Planck scale, b) 4 TeV minimum and 14 TeV maximum black hole mass c) 3 extra dimensions. Time evolution during Hawking radiation and gray body effects are included. The detector response was simulated by us using the CMS fast simulation (FAMOS, version 1.4.0) after validation against the detailed CMS Geant-based simulation. The Standard Model backgrounds taken into account include QCD jets, top production and boson plus jet production. The invariant mass of all final state objects (electrons, photons, jets and muons) in the event is found to be correlated with the input black hole mass. In addition since the black hole formation can only occur if $M_{\text{BH}} > M_{(4+n)}$, the event invariant mass can indicate the effective Plack scale $M_{(4+n)}$. In the benchmark scenario the invariant mass is required to be greater than ≈ 1.5 TeV. BH events are characterized by a high multiplicity of the final state particles, which




Figure 14.21: Expected significances as function of M_D for different number n of extra dimensions.

increase as a function of the BH mass (and decreases as a function of Hawking temperature). In particular the ratio of jets to leptons is found to be 5 to 1. In this study with a simple jet and lepton multiplicity counting the jet/lepton ratio is formed. Jets and leptons originating from top quarks and W/Z bosons are not counted. The average value of this ratio is found to be 4.5. In the analysis path the hadron/lepton multiplicity ratio is required to be greater than 4. The thermal nature of Hawking radiation requires the distribution of BH remnants to be spherical as shown and a sphericity of 0.28 is required which eliminates drastically the Standard Model backgrounds. The invariant mass distribution and sphericity for the signal and background events is shown in figure 14.22.




Figure 14.22: (a) Reconstructed invariant mass distribution and (b) event sphericity for black hole and standard model background events



Events are counted when the total sum of the P_T of all reconstructed objects plus the missing transverse energy is larger than 2880 GeV. A study of the Level-1 and HLT trigger ϵ shows that the 4 jet trigger has a 93% for the signal events and is used in the analysis. **VIXME:** has the 7% inefficiency been taken into account in the final number surviving?


The event selection criteria applied to the reconstructed events and the efficiencies of the requirements are listed in Table 14.11. **VIXME:** in the original there is a suspicious line which says $E > mbh/2$ which results in no changes to the signal or backgrounds- i removed it

Table 14.11: Event selection and background rejection for signal events and major background processes

Cut	Signal	tt+nJ	W+nj	Z+nJ	QCD Dijet	WW+nJ
Cross Section (pb)	18.85	371	896	781.84	33076.8	269.91
Events (10 fb-1)	188500	3.71×10^6	8.96×10^6	7.82×10^6	3.31×10^8	2.70×10^6
$M_{Inv} > 2$ TeV	18.71	13.29	6.53	3.85	2634.94	20.53
ratio jet/ $\ell > 4$	17.72	13.25	6.43	3.84	2613.18	20.42
Sphericity > 0.28	9.27	1.60	0.23	0.10	53.74	0.07
Final No.Events (10 fb-1)	92740	15990	2328	982	537391	740

The minimum integrated luminosity needed for 5σ significance and for the benchmark point is $\sim 3 \text{ pb}^{-1}$. A survey of the parameter space using 25 points shows that for effective Planck scale of 2-3 TeV, minimum black hole mass up to 4 TeV and 2-6 extra dimensions the 5 sigma significance can be obtained with luminosity between fraction of pb^{-1} and 100's of pb^{-1} . For

effective Planck scale of 4 TeV a few fb^{-1} will be needed.  **ME:** check table once more. 

 **ME:** in the original there is a discussion of pdf uncertainties that makes no sense; please
check

Chapter 15

Alternative BSM signatures



15.1 Technicolour

15.1.1 The $\rho_{TC} \rightarrow W + Z$ channel

Technicolour (TC) stands as an alternative to the elementary Higgs mechanism of the Standard Model. It introduces a new strong interaction [717] providing a dynamical nature to Electroweak Symmetry Breaking. Technicolour is a QCD-like force, acting on technifermions at an energy scale $\Lambda_{TC} \sim \nu_{weak} = 175$ GeV. A number N_D of technifermion doublet condensates yield the pseudo-Goldstone bosons π_{TC} , together with a wide spectroscopy of excited technimesons. The present simulation is performed under the phenomenology of the lowest-lying technihadrons, commonly referenced as the “Technicolour Straw Man” model (TCSM) [707]. The colour-singlet sector includes the spin-zero π_{TC} and the spin-one technimesons ρ_{TC} and ω_{TC} . The decay cross-section of ρ_{TC} is expressed as an admixture of π_{TC} and the Standard Model Z and W bosons:

$$\rho_{TC} \rightarrow \cos^2 \chi \langle \pi_{TC} \pi_{TC} \rangle + \cos \chi \sin \chi \langle \pi_{TC} W_L \rangle + \sin^2 \chi \langle W_L W_L \rangle \quad (15.1)$$

where W_L is the longitudinal mode of the Z or W and $\sin \chi \simeq 1/\sqrt{N_D} \sim 1/3$. The branching fraction $\text{BR}(\rho_{TC} \rightarrow W + Z)$ is competing with the two first terms in Eq. 15.1, hence with $M(\pi_{TC})$.

The decay channel $\rho_{TC} \rightarrow W+Z$ is the subject of this analysis and has the advantage of a very clean final state, namely $3\ell + \nu$. The background contributions arise mainly from Standard Model processes involving boson production and decays. Other Technicolour decay modes that include jets such as $\rho_{TC} \rightarrow \pi_{TC} + W$, have better branching fractions but are much harder to disentangle from the Standard Model processes.

15.1.1.1 Selection path

All signal and backgrounds samples used in this analysis are generated with PYTHIA 6.2 [30] * with the requirement of at least 3 prompt leptons in the CMS fiducial region †. A set of 14 different ρ_{TC} samples are generated within the $[M(\rho_{TC}), M(\pi_{TC})]$ phase space. The CMS fast simulation (FAMOS.1.4.0 [287]) is used for detector simulation and event reconstruction. Nominal CMS Level-1 and High-Level Trigger requirements are applied [68].

*The $Zb\bar{b}$ background is generated using CompHEP [47] interfaced to PYTHIA

†Contributions from processes of type $Z \rightarrow 2\ell$ plus an additional fake lepton from a jet have been taken into account in the systematic uncertainties, see Sect. 15.1.1.2.

The main reconstructed objects and their efficiencies have been validated against the detailed Geant-based CMS detector simulation [14, 15].

The analysis path is designed to reduce the main Standard Model background contributions WZ , ZZ , $Zb\bar{b}$ and $t\bar{t}$, while retaining high signal efficiency. It is summarized as follows:

- (i) Lepton Selection: 3 high- p_T and isolated electrons or muons.
- (ii) Lepton Trigger: Single- or Two-electron or muon mode (Level-1 and HLT)
- (iii) Z : same-flavor and opposite-charge ℓ -pair closest to $M(Z)$, with $p_T(\ell_{1,2}) > (30,10)$ GeV
- (iv) W : solution to 3rd lepton with $p_T > 10$ GeV + Missing E_T + $M(W)$ constraint
- (v) $|M(\ell^+\ell^-) - M(Z)| \leq 3\sigma_{M_Z} \cong 7.8$ GeV
- (vi) $p_T(Z)$ and $p_T(W) > 30$ GeV[‡]
- (vii) $|\Delta[\eta(Z) - \eta(W)]| \leq 1.2$

The Z and W are reconstructed with a purity of $\sim 99\%$, using the 3 highest- p_T leptons in the event, and the Missing Transverse Energy (MET), obtained as the vector sum of the jets in the event. The $M(W)$ constraint yields a 2 fold ambiguity in the p_Z component of the reconstructed neutrino: it is found that the most efficient choice for the ρ_{TC} signal is the minimum p_Z solution. The kinematic cuts are illustrated in Fig. 15.1. The main $t\bar{t}$ reduction is obtained via the Z -mass window requirement (v). The irreducible background $WZ \rightarrow 3\ell + \nu$ is most efficiently separated from the signal via the $\eta(Z) - \eta(W)$ correlation requirement (vii). The p_T cut on Z and W further improves the signal to background ratio, however it is kept modest in order to preserve the exponential background hypothesis of the $3\ell + \nu$ invariant mass spectrum, used to compute the signal sensitivity. The $\rho_{TC}(300, 300)$ signal and background yields are shown in Fig. 15.1(d) and the corresponding reconstruction efficiencies are listed in Table 15.1.

Table 15.1: $\sigma \times \text{BR}(\ell = e \text{ or } \mu)$, 3-lepton preselection efficiency, total efficiency and final yield within 3σ of the signal region (Nev), for $\int \text{Ldt} = 5 \text{ fb}^{-1}$. $\rho_{TC}(300, 300)$ and the main background contributions are shown. The simulation is repeated for all ρ_{TC} benchmark points.

Sample	$\sigma \times \text{BR}(\text{pb})$	$\epsilon(3\text{-lept})$	$\epsilon(\text{Reco}) (\%)$	Nev(5 fb^{-1})
$\rho_{TC} \rightarrow W + Z \rightarrow 3\ell + \nu$	0.13	0.635	25.88 +- 0.40	103
$WZ \rightarrow 3\ell + \nu$	0.39	0.471	9.91 +- 0.11	27
$ZZ \rightarrow 4\ell$	0.07	0.719	15.80 +- 0.14	10
$Zb\bar{b} \rightarrow 2\ell + X$	332	0.046	0.23 +- 0.01	12
$t\bar{t}$	489.72	0.065	0.019 +- 0.001	8

15.1.1.2 Signal Sensitivity and Systematic uncertainties

The sensitivity of each ρ_{TC} benchmark point is computed by taking into account realistic statistical fluctuations at a given integrated luminosity. The sensitivity estimator is defined as the likelihood-ratio:

$$S_L = \sqrt{2 \ln(\mathcal{L}_{S+B}/\mathcal{L}_B)} \quad (15.2)$$

where \mathcal{L}_{S+B} and \mathcal{L}_B are the best-fit likelihoods of the signal-plus-background hypothesis and the null hypothesis (no signal present). The signal probability density function (p.d.f.)

[‡]For benchmark points with $M(\rho_{TC}) = 200$ GeV, the minimum $p_T(Z)$ and $p_T(W)$ threshold is 10 GeV.




Figure 15.1: (a) $M(\mu^+\mu^-)$ for $\rho_{TC}(300, 300)$ and $t\bar{t}$; (b) $\Delta[\eta(Z) - \eta(W)]$ for $\rho_{TC}(300, 300)$ and WZ ; (c) $p_T(Z)$ for $\rho_{TC}(300, 300)$ and all backgrounds ($p_T(W)$ is similar); (d) Reconstructed $M(3\ell + \nu)$ for $\rho_{TC}(300, 300)$ and all backgrounds. The vertical lines indicate the applied requirements.

is assumed Gaussian (dominated by detector resolution) and the background p.d.f. is Exponential in all ρ_{TC} fit regions. The output of the fitting procedure is shown in the contour plot over the $[M(\rho_{TC}), M(\pi_{TC})]$ phase space in Fig. 15.2(a), for various integrated luminosities. A signal sensitivity above 5 is expected for $\int Ldt = 3 \text{ fb}^{-1}$ (before including systematic uncertainties).

The ρ_{TC} sensitivity has been simulated for the early CMS data taking phase. Expected detector related systematic uncertainties at $\int Ldt = 1 \text{ fb}^{-1}$ are taken into account. While no substantial contribution is found from the tracker and muon system misalignment or the calorimeter miscalibration, the accuracy at which the lepton efficiency will be determined from data affects the result: a 2% uncertainty is considered. Moreover, the lepton fake rate has been simulated on $Zb\bar{b}$ and extrapolated to any $Z+jet(s)$ type background[§], in order to take into account additional contaminations from pion/kaon decays or from wrongly identified lepton candidates: a single lepton fake rate of $O(10^{-3})$ is obtained with FAMOS, affecting the ρ_{TC} sensitivity as shown below. Finally, a 7.5% uncertainty on the missing transverse energy measurement is considered. The above uncertainties result in the following relative ρ_{TC} sensitivity drop:

$$\Delta_{\text{SYS}}^{\text{tot}} = \sqrt{(\Delta_{\text{SYS}}^{\text{Eff}})^2 + (\Delta_{\text{SYS}}^{\text{Fake}})^2 + (\Delta_{\text{SYS}}^{\text{MET}})^2} = \sqrt{(2.7\%)^2 + (8.5\%)^2 + (6.6\%)^2} = 11\% \quad (15.3)$$

[§]A production cross-section of 1047pb per lepton flavor is assumed for $Z + n\text{-jets}$, $n \geq 0$.




Figure 15.2: (a) Signal 5σ Sensitivity curves at various integrated luminosities; (b) sensitivity at $\int Ldt = 4fb^{-1}$: the dotted (resp. dashed) curve shows the sensitivity (resp. the 90% C.L. signal upper limit) after including systematic uncertainties.

Introducing K-factors from Next-to-Leading-Order (NLO) expectations for the signal[¶] and background leads to a relative signal sensitivity increase of 6%; however the latter estimate has not been included in the final result.

In summary, the technicolour signature $\rho_{TC} \rightarrow W + Z$ in the context of the Straw Man model is studied. The 5 sigma discovery reach is obtained at $\int \mathcal{L} \simeq \Delta fb^{-1}$.

15.2 Search for Contact Interactions with dimuons

Contact interactions offer a general framework for describing a new interaction with typical energy scale $\Lambda \gg \sqrt{s}$. The presence of operators with canonical dimension $N > 4$ in the Lagrangian gives rise to effects $\sim 1/\Lambda^{N-4}$. Such interactions can occur for instance, if the SM particles are composite, or when new heavy particles are exchanged.

Table 15.2: Contact interaction models.

Model	LL	RR	LR	RL	VV	AA	LL+RR	LR+RL
	Non-parity conserving							
η_{LL}	± 1	0	0	0	± 1	± 1	± 1	0
η_{RR}	0	± 1	0	0	± 1	± 1	± 1	0
η_{LR}	0	0	± 1	0	± 1	∓ 1	0	± 1
η_{RL}	0	0	0	± 1	± 1	∓ 1	0	± 1

In the following we will consider lepton-pair production. The lowest order flavor-diagonal and helicity-conserving operators have dimension six [117].

The differential cross section takes the form

$$\frac{d\sigma}{d\Omega} = SM(s, t) + \varepsilon \cdot C_{Int}(s, t) + \varepsilon^2 \cdot C_{NewPh}(s, t) \quad (15.4)$$

[¶]A NLO K-factor 1.35 is assumed for $\sigma(\rho_{TC})$, in similarity with the Drell-Yan process.

where the first term is the Standard Model contribution, the second comes from interference between the SM and the contact interaction, and the third is the pure contact interaction effect. The Mandelstam variables are denoted as s , t and u .

Usually the coupling is fixed, and the structure of the interaction is parametrized by coefficients for the helicity amplitudes:

$$\begin{array}{ll} g & \text{coupling (by convention } \frac{g^2}{4\pi} = 1) \\ |\eta_{ij}| \leq 1 & \text{helicity amplitudes } (i, j = \text{L, R}) \\ \varepsilon & \frac{g^2 \text{ sign}(\eta)}{4\pi \Lambda^2} \text{ for } f\bar{f} \end{array}$$

Some often investigated models are summarized in Table 15.2. The models in the second half of the table are parity conserving, and hence not constrained by the very precise measurements of atomic parity violation at low energies. The results presented in this contribution cover the LL model, which has the highest sensitivity at LHC energies from the models in the first half of the table.

15.2.1 Analysis Path

The topology under study is high-mass muon pairs with opposite sign. The Global Muon Reconstructor (GMR) output is used. The di-muon events are triggered by the single and di-muon triggers. We have processed events, generated to cover the whole region of interest up to di-muon masses of 6 TeV, through full simulation with OSCAR and reconstruction with ORCA. The di-muon mass resolution is parametrized in two ways:

- as mass dependent one standard deviation (RMS)
- by fitting the mass resolution with a sum of two Gaussians to account for the long tail of less well reconstructed masses.

The results are remarkably stable as a function of the di-muon mass: the second Gaussian contributes around 14% and has a standard deviation 3.3 times bigger than the first Gaussian.

Our strategy is to generate events with `PYTHIA` and apply parametrizations of the di-muon mass efficiency and resolution obtained from full simulation. We have verified our approach by comparing the resulting mass spectra with the ones obtained with `OSCAR/ORCA` or `FAMOS` for Drell-Yan and selected contact interactions samples, observing good agreement in all cases.

Two mass regions: 500–1000 GeV and 1000–6000 GeV are considered. The total cross section and the forward-backward asymmetry as function of the di-muon mass are studied. Our analysis shows that the sensitivity to contact interactions comes almost exclusively from the cross section measurements for the LL model.

In order to reduce the systematic uncertainties both on the experimental and theory sides a “double ratio” method is developed. The number of observed events for a given bin in invariant mass is

$$N_{obs} = L \cdot \sigma \cdot \varepsilon \quad (15.5)$$

where L is the luminosity, σ the differential cross section for the given mass bin, and ε the experimental efficiency. We select a zeroth “normalization” bin for invariant masses between

250–500 GeV, both well above the Z pole and in an area well covered by the Tevatron, and define the experimental ratios

$$R_i^{DATA} = \frac{N_i^D}{N_0^D} = \frac{\sigma_i^D \cdot \varepsilon_i^D}{\sigma_0^D \cdot \varepsilon_0^D}. \quad (15.6)$$

Here the cross sections and efficiencies are the ones for the real LHC data. The index i runs for all measured bins with masses above 500 GeV. The luminosity cancels in the ratio. The choice of this mass bin is not random. If we compare the flavor composition of partons initiating the hard interaction (Table 15.3), at the Z peak 32.1% are heavier flavors (not u or d quarks), with their own parton density functions (PDF) uncertainties. At 250–500 GeV the u and d quarks are “initiators” already in 85.6% of the cases, increasing to 96.3% above 1 TeV etc. Moreover, at the Z peak d quarks are most abundant, while at higher masses u quarks dominate, asymptotically approaching a ratio 4:1. It is clear that our choice of normalization bin gives flavor composition much closer to the most interesting high mass events, compared to a normalization using Z pole events. The PDF uncertainty on cross sections is estimated using LHAPDF [88, 353]. It is interesting to note that this uncertainty reaches a minimum for masses 250-600 GeV, corresponding to medium values of the parton momentum fractions X , reinforcing our choice of normalization bin.

Table 15.3: Flavor composition of partons initiating the hard Drell-Yan interaction. The PDF uncertainty on the cross sections (positive and negative asymmetric errors) is estimated using LHAPDF.

Mass [GeV]	d	u	s	c	b	PDF + [%]	PDF - [%]
Z peak	35.9	32.1	17.2	9.77	5.10	+4.7	-5.7
250-500	24.3	61.3	6.22	6.64	1.54	+3.4	-4.2
500-600	22.8	68.4	4.03	3.95	0.89	+3.5	-4.1
1000+	21.7	74.6	1.86	1.48	0.33	+5.0	-5.8
2000+	19.9	78.4	0.91	0.63	0.14	+9.0	-7.7

We define similar ratios for the Monte Carlo (theory) predictions. The *absolute* values of the cross sections and efficiencies are not important for the ratios, what matters is *the shape* of these quantities as function of invariant mass. For example, the absolute value of K-factors, a way to compensate for missing higher order N(N)LO terms and enable the comparison of leading order Monte Carlo predictions to data (similarly for the electroweak radiative corrections) disappears from the ratios and only the *shape* of the K-function as depending on invariant mass remains - a much smaller effect. And part of the uncertainties introduced due to our limited knowledge of PDFs cancels in the ratio, leaving smaller residual uncertainties due to the change of phase space for changing masses.

Now let us define the double ratios

$$DR_i = \frac{R_i^{DATA}}{R_i^{MC}}. \quad (15.7)$$

This method is inspired by a study of Drell-Yan events and extraction of proton and pion PDFs at lower masses [718], as well as by the SuperKamiokande double ratio method for

measuring atmospheric neutrino oscillations [719]. If our theory understanding and detector modeling are both perfect, we expect $DR_i \equiv 1$. The experimental or Monte Carlo errors introduced in the ratios from the uncertainties in the zeroth bin are negligible, as due to the steeply falling Drell-Yan spectrum this bin has much more data compared to the high mass bins.




Figure 15.3: Double ratios for contact interactions in the di-muon channel, LL model, scale $\Lambda = 20$ TeV, positive and negative interference, and luminosity 100 fb^{-1} . The errors shown are statistical.

An example of double ratios for positive and negative interference is shown in Figure 15.3. As can be seen, for scale $\Lambda = 20$ TeV the expected effects are quite sizable (please note the log scale), with the sensitivity for negative interference starting around di-muon masses of 750 GeV, while for positive interference masses above 2 TeV are crucial.

The experimental systematic effects in the cross section measurement are estimated to be 2% from the total muon efficiency and no more than 1.4% from momentum resolution. The former can be controlled quite well with the huge sample of Z events decaying to di-muons, and the effects for TeV muons are taken into account on top of this. The latter is important at high mass as smearing from lower masses from the steeply falling Drell-Yan spectrum can contaminate the high mass measurements, especially if the tails of the momentum resolution are not under control. It is estimated by varying the two parametrizations of the mass resolution by $\pm 40\%$, giving consistent results. The main source of systematic uncertainties on the momentum resolution comes from the alignment of the muon chambers and the central tracker, both at start-up and at high luminosity.

The systematic uncertainties from our limited knowledge of PDFs is estimated using the CTEQ6m PDF set from LHAPDF. From Table 15.4 our estimate of the PDF uncertainty on the cross section ratio is $+5.2\%$ above 1 TeV or $+10.7\%$ above 2 TeV.

The genuine electro-weak radiative corrections change by $\sim 10\%$ in the relevant mass range [149, 352]. The K-function changes faster below 250-300 GeV. From our normalization bin to the highest masses first estimates show a change below 8% on the cross section ^{||}. Taking conser-

^{||}Calculations by M.Schmitt with the program PHOZPRMS [350].

Table 15.4: The PDF uncertainty on the cross section ratios (positive and negative asymmetric errors) as estimated using LHAPDF. Clearly normalizing to the 250-500 GeV mass bin is superior compared to a normalization relative to the Z peak (70-120 GeV).

Mass [GeV]	$R(\frac{M}{250-500})$		$R(\frac{M}{Z_{peak}})$	
	PDF + [%]	PDF - [%]	PDF + [%]	PDF - [%]
500-600	+1.5	-1.5	+4.6	-4.2
1000+	+5.2	-4.8	+7.8	-7.1
2000+	+10.7	-7.8	+12.9	-9.4

vatively half of these changes with mass as an upper limit on the systematic uncertainty we arrive at 5% and 4% respectively.

Combining all effects in quadrature, we arrive conservatively at systematic uncertainties below 2.5% experimental, 11.5% from theory, 12% total at nominal conditions, 15% shortly after start-up. With the accumulation of data and improved calculations there is hope to improve this number by making progress in our understanding of PDF, electro-weak radiative corrections and K-functions.




Figure 15.4: Five sigma discovery reach (left) and sensitivity at 95 % CL(right) for contact interactions in the di-muon channel for different luminosities and signs of the interference.

The discovery reach for a given model is determined by constructing a negative log-likelihood function combining the deviations between measurements and predictions, including the contact interaction contributions, for all simulated data points. The error on a deviation consists of three parts, which are combined in quadrature: a statistical error, an experimental systematic error and a theoretical uncertainty. The log-likelihood function is integrated in the physically allowed region (all positive Λ for positive interference and all negative Λ for negative interference) to derive the five standard deviations σ discovery reach and one-sided

lower limits at 95% confidence level on the scale.

The discovery reach is summarized in Figure 15.4. The sensitivity is dominated by the cross section measurement, the contribution of the forward-backward asymmetry is minor. The sensitivity for negative interference is substantially better. Even at the highest luminosities the statistical errors at LHC play a major role, as evident from the comparison of the cases with total systematic uncertainties of 3, 15 and 30 %. This is not surprising as the Drell-Yan process is probing directly masses up to $\sim 4\text{--}5$ TeV, where due to the steeply rising cross sections the statistical errors remain important for all considered luminosities.

15.3 Search for Contact Interactions with dijets

New physics at a scale Λ above the mass of the final state is effectively modeled as a contact interaction. Here the propagator for a particle of mass $M \sim \Lambda$ exchanged between quarks, or exchanged between constituent particles inside two interacting composite quarks, shrinks to a single point and gives a contact interaction. Quark contact interactions, for example those that arise from a left-handed interaction among composite quarks [117, 118], will always produce a rise in rate relative to QCD at high dijet mass or high inclusive jet E_T . However, observation in the mass distribution alone requires precise understanding of the QCD rate as a function of dijet mass, which is complicated by the large systematic uncertainties discussed in section 4.1.6. Angular distributions benefit from much smaller systematic uncertainties. The contact interaction is often more isotropic than the QCD background, since QCD is dominated by t-channel scattering and produces jets predominantly in the forward direction. Our analysis uses the dijet ratio, discussed in section 4.1.5, to measure the angular distribution as a function of dijet mass, and see any contact interactions which affect the dijet angular distribution [720].

15.3.0.1 Contact Interaction Sensitivity Estimates

The QCD background distribution for the dijet ratio was discussed in section 14.5. In Figure 15.5 we show a smooth dijet ratio for QCD, estimated at 0.6 from the fit to the full simulation. The error bars shown in Figure 15.5 are the statistical uncertainties expected with 1 fb^{-1} and the jet trigger prescales discussed in section E.4.3.2. The uncertainties are calculated using Poisson statistics at high dijet mass, where few events are expected and Gaussian statistics is less accurate. In Figure 4.7 we presented a lowest order calculation of both QCD and a contact interaction among left-handed quarks. The signal in Figure 15.5 is estimated by scaling the lowest order contact interaction calculation of Figure 4.7 by the ratio of our full simulation prediction for QCD to the lowest order QCD calculation: $\text{signal} = \text{contact} \times 0.6 / \text{QCD}$. Systematic uncertainties on the dijet ratio are small, as discussed in section 4.1.6 and demonstrated in Figure 4.8. The calculated chisquared between QCD and the contact interaction signal, including all uncertainties on the dijet ratio, is listed in table 15.5. In Figure 15.5 we show the significance in σ , estimated as $\sqrt{\chi^2}$, compared to a smooth fit as a function of $1/\Lambda^+$. The anticipated capability of CMS with 1 fb^{-1} to exclude contact interactions at 95% CL or discover them at 5σ can be read off Figure 15.5, and they are listed in table 15.6. This includes the uncertainty on Λ due to the anticipated 5% uncertainty on the observed jet energy. The same analysis is repeated for 100 pb^{-1} and 10 fb^{-1} and the results are also listed in table 15.6. The systematic uncertainties on the dijet ratio reduced the CMS




Figure 15.5: Left) The expected value and statistical error of the dijet ratio of QCD in the CMS detector for 1 fb^{-1} (solid) is compared with QCD plus a quark contact interaction at a scale Λ^+ of 15 TeV (dashed), 10 TeV (dotted) and 5 TeV (dot-dashed). Right) The significance with statistical uncertainties only (open circles) and with all uncertainties (solid circles) of the difference between QCD alone and QCD plus a quark contact interaction is plotted vs $1/\Lambda^+$ and fit with a quadratic function. Horizontal lines show the 5σ and 95% CL levels.

Luminosity	100 pb^{-1}			1 fb^{-1}			10 fb^{-1}		
Λ^+ (TeV)	5	10	15	5	10	15	5	10	15
χ^2 (Stat)	18.3	.090	.0037	316	5.82	.107	3652	133	4.15
χ^2 (All)	16.7	.082	.0011	240	5.55	.061	1340	124	3.56

Table 15.5: Chisquared between signal and background. For each luminosity and contact interaction scale considered we list the chisquared between QCD alone and QCD plus a contact interaction, for the case where only statistical uncertainties are included (Stat), and for the case where both statistical and systematic uncertainties are included (All).

sensitivity to a contact interaction between 0.1 and 0.3 TeV/c^2 depending on luminosity and level of significance. To see how quickly CMS jet data will extend the search for new physics, we note that with 100 pb^{-1} our anticipated 95% CL sensitivity, $\Lambda^+ < 6.3 \text{ TeV}$, is more than twice the sensitivity of the D0 search ($\Lambda^+ < 2.7 \text{ TeV}$ at 95% CL)[116]. We note that our contact interaction sensitivity to composite quarks in table 15.6 is roughly twice our mass resonance sensitivity to excited states of composite quarks in table 14.5, and is equivalent to observing or excluding a quark radius of order 10^{-18} cm .

Luminosity	95% CL Excluded Scale			5σ Discovered Scale		
	100 pb^{-1}	1 fb^{-1}	10 fb^{-1}	100 pb^{-1}	1 fb^{-1}	10 fb^{-1}
Λ^+ (TeV)	<6.2	<10.4	<14.8	<4.7	<7.8	<12.0

Table 15.6: Sensitivity to contact interactions with 100 pb^{-1} , 1 fb^{-1} , and 10 fb^{-1} . We list the largest value of the contact interaction scale we expect to be able to exclude at a confidence level of 95% or greater, and the largest value we expect to be able to discover with a significance of 5σ or greater. Estimates include both statistical and systematic uncertainties.



4 Heavy Majorana neutrinos and right-handed bosons

FIXME: fix for size, content, flow

15.4.1 Introduction

Among several extensions of the Standard Model (SM) that can be tested at LHC, see e.g. [721], the left-right (LR) symmetric model $SU_C(3) \otimes SU_L(2) \otimes SU_R(2) \otimes U(1)$ [722, 696, 697] is certainly one of the most interesting. The model embeds the SM at the scale of the order of 1 TeV and naturally explains the parity violation in weak interactions as a result of the spontaneously broken parity. The model necessarily incorporates three additional gauge bosons W_R and Z' and the heavy right-handed Majorana neutrino states N . The N s can be the partners (N_l) of the light neutrino states ν_l ($l = e, \mu, \tau$) and can provide their non-zero masses through the see-saw mechanism [698]. This makes LR model very attractive, since recent results from atmospheric, solar and reactor neutrino experiments confirm the existence of neutrino oscillations and therefore provide strong evidence that neutrinos are not massless.

There are several recent papers devoted to the study of the production of heavy neutrinos and new gauge bosons at TeV energy scale, see e.g. Ref. [723, 724, 725, 726].

In the framework of the LR symmetric model, we have studied the production and the experimental signature of heavy Majorana neutrinos and the associated heavy gauge bosons in the CMS detector at the LHC. The preliminary results of this study are presented in [727].

Existing experimental data constrain the Z' mass to the values $O(1)$ TeV [728]. The lower bound on the W' mass derived from the $K_L - K_S$ mass difference is quite stringent, $M_{W'} \gtrsim 1.6$ TeV [729], however with some uncertainties from the low energy QCD corrections to the kaon system. The direct searches for W' at the Tevatron yield bounds $M_{W'} \gtrsim 720$ GeV assuming a light (keV-range) N , and $M_{W'} \gtrsim 650$ GeV assuming $M_N < M_{W'}/2$ [730]. These bounds are less stringent in more general LR models.

15.4.2 Heavy Majorana neutrino production and decay

One can study at LHC two kinds of processes with W_R and N_l : i) $pp \rightarrow W_R \rightarrow l + N_l + X$, and ii) $pp \rightarrow Z' \rightarrow N_l + N_l + X$ followed by the decay $N_l \rightarrow l + j_1 + j_2$.

The cross sections of these processes depend on the following parameters:

- the value of the coupling constant g_R ,
- parameters of the CKM mixing matrix for the right-handed sector,
- the $W_R - W_L$ and $Z' - Z$ mixing strengths, and
- the masses of the partners N_l of the light neutrino state.

To simplify our study, we will use the following standard assumptions:

- the mixing angles are small
- right-handed CKM matrix is identical to the left-handed one
- $g_R = g_L$

Under these conditions, Z' is about 1.7 times heavier than W_R and the production cross-section for $pp \rightarrow W_R \rightarrow eN_e$ is found to be at least one order of magnitude higher than for the $pp \rightarrow Z' \rightarrow N_e N_e$ process.

The focus of our study is on the process i) going through the W_R direct production. The additional assumption is that only the lightest M_{N_e} is reachable at LHC. In the case of degenerated masses of N_l , the channels with μ 's and τ 's are open up resulting in the increase of the cross section of the reaction i) by a factor ~ 1.2 . Finally, only two unknown parameters, M_{W_R} and M_{N_e} , are involved into the analysis. **The point $M_{N_e} = 500 \text{ GeV}/c^2$ and $M_{W_R} = 2000 \text{ GeV}/c^2$ in this parameter space we will call below a reference point (LRRP).**

For the signal event generation and calculation of cross sections, the PYTHIA 6.2 [62], which includes the LR symmetric model with the standard assumptions mentioned above, is used. The default, CTEQ5L parton distribution functions of ref. [731] are used for calculations. The fraction of $pp \rightarrow W_R^+$ ($pp \rightarrow W_R^-$) reactions as a function of M_{W_R} changes from $\simeq 70\%$ ($\simeq 30\%$) at $M_{W_R} \simeq 1 \text{ TeV}/c^2$ to $\simeq 95\%$ ($\simeq 5\%$) at $M_{W_R} \simeq 10 \text{ TeV}/c^2$. For W_R boson masses higher than $M_{W_R} \simeq 2 \text{ TeV}/c^2$ the production of W_R^+ boson dominates.

We study the W_R mass region above $1 \text{ TeV}/c^2$, assuming that smaller masses are excluded by indirect analyses [723]. The signal cross section, defined as the product of the total W_R production cross section and the branching ratio of W_R decay into electron and N_e , for different masses of W_R is shown in Figure 15.6 as a function of M_{N_e} .




Figure 15.6: The dependence of the value $\sigma(pp \rightarrow W_R) \cdot Br(W_R \rightarrow e^\pm N_e)$ on the heavy neutrino mass. For $M_{W_R} = 2 \text{ TeV}/c^2$ the same value in case of degenerated N masses is shown by the dashed line.

The detection of events with LR heavy neutrinos is studied using the full CMS detector simulation and reconstruction chain: CMKIN 4.3.1 (PYTHIA 6.227) \rightarrow OSCAR 3.6.5 \rightarrow ORCA 8.7.3. For details, see ref. [732].

15.4.3 Selection of candidate events and the analysis variables

In the analysis we proceeded through the following steps:

- events with 2 isolated electrons are selected (standard isolation in the tracker required).
- events with at least 2 jets are selected. From these jets, two jets with the maximal transverse momentum (signal jet pair) are chosen.
- using the 4-momenta of the signal jet pair and the 4-momentum of a lepton, the invariant mass $M_{ljj} = M_{N_e}^{cand}$ is calculated. Since there are two electrons - two ljj combinations are considered. Finally, the ljj mass peak is to be searched for.
- from the 4-momenta of the signal jet pair and the 4-momenta of electrons the invariant mass $M_{lljj} = M_{W_R}^{cand}$ is calculated. and a peak in the distribution of this mass is to be searched for.




Figure 15.7: The distribution of the invariant mass $M_{N_e}^{cand}$ for the reconstructed events with a heavy neutrino. All combinations are shown. LRRP: the simulated neutrino mass is 500 GeV/c^2 . The normalization is arbitrary.

Two more variables are used in the analysis:

- the minimal invariant mass of **all** same flavor lepton pairs M_{ll} . There is a lower cut on this value (typically $\simeq 200 \text{ GeV}/c^2$)
- the E_t^{miss} of an event.

In Fig. 15.7 the $M_{N_e}^{cand}$ distribution is shown for the events simulated at LRRP. The additional broad peak or long tail above the heavy neutrino mass is due to the wrong choice of lepton.

In Figure 15.8 the efficiency of the primary selection and the efficiency of reconstruction of W_R decay products are shown. For neutrino masses much smaller than the W_R mass the reconstruction efficiency drops due to the significant overlap of the heavy neutrino decay products in the $\eta - \phi$ space.

15.4.4 The background

The background is expected from the SM processes with a lepton pair and jets (two leading jets (from a hard process) or one leading jet and others from QCD effects) in the final state. In the first approximation most of them can be simulated with PYTHIA.




Figure 15.8: The probability to pass the primary selection for the $pp \rightarrow W_R \rightarrow e + N_e \rightarrow 2e + 2jet$ events as a function of the N_e mass for $M_{W_R} = 2 \text{ TeV}/c^2$. The lower curve is the probability to have all four W_R decay products reconstructed as separate CMS reconstruction objects

Two most important sources of background are listed below.

The $t\bar{t}$ production with only leptonic W decay modes allowed. It was checked that other decay modes do not contribute. The cross section is about two orders of magnitude larger than the signal cross section at LRRP.

The $Z + jet$ process. This reaction has a large cross section: about 5 orders of magnitude larger than the signal cross section at LRRP. This background is suppressed by the lower cut on M_{ll} of 200 GeV. In order to reduce the number of simulated events we used the cut $Q_t > 20 \text{ GeV}$ (PYTHIA parameter CKIN(3)) during the simulation and the generator level preselection of events with sufficiently high M_{ll} .

At LRRP the reconstructed N_e mass peak in the presence of the SM background is well visible, though the background is significant (comparable with the peak height). However, if we require that in the same event there is a possibility to construct a heavy W_R boson candidate ($eejj$ system) with the invariant mass above $1 \text{ TeV}/c^2$, the background under the N_e peak drops dramatically, resulting in the mass distribution shown in Fig. 15.9.

In Fig. 15.10 the reconstructed W_R mass peak in presence of the SM background is shown.

The Majorana nature of the heavy neutrino allows us to switch our analysis to same sign leptons. At the present stage of the analysis we see that in the parameter space to be studied (at least $M_{W_R} > 1 \text{ TeV}/c^2$, lower values safely excluded by other experiments) this change does not improve the sensitivity because half of signal events are lost. However, in case of discovery this will be an excellent cross check.

The only found sources of events with same sign leptons that could be a background to the production of heavy Majorana neutrino are the ZH and WH productions. However, the cross section of these processes is small. For this reason the search for heavy neutrino events with the same sign leptons theoretically can be almost background-free. There is however some background to the same sign analysis from the incorrect reconstruction of charge of




Figure 15.9: The heavy neutrino mass peak reconstructed together with the SM background (shaded histogram). The possibility to construct W_R with invariant mass above 1 TeV/ c^2 in the event is required. $L_t = 30 \text{ fb}^{-1}$




Figure 15.10: The W_R mass peak reconstructed together with the SM background (shaded histogram). $L_t = 30 \text{ fb}^{-1}$

electron tracks. But it is still much smaller than the physical background in the different sign analysis.

15.4.5 Results

The discovery potential of the events with W_R and N_e in CMS was calculated using the following relation [41]:

$$S = 2(\sqrt{N_S + N_B} - \sqrt{N_B}) \geq 5, \quad (15.8)$$

where N_S and N_B are the numbers of signal and background events respectively.

The corresponding discovery contour in the $(M_{W_R}; M_{N_e})$ plane is shown in Figure 15.11. After three years of running at low luminosity (30 fb^{-1}) the use of $pp \rightarrow W_R \rightarrow eN_e \rightarrow eejj$ reaction allows to discover W_R and N_e with masses up to 4 TeV/ c^2 and 2.4 TeV/ c^2 , respectively. For the LRRP point the corresponding significance is about 40.

The uncertainty of these results coming from various background uncertainties is small since the background itself is rather small and the discovery region is usually limited by the fast drop of the signal cross section at high ratios $r = M_{N_e}/M_{W_R}$ or by the fast drop of efficiency at small r . As to the signal cross section uncertainties, we studied the one from PDF (parton density functions). By taking different PDF sets we found that the full amplitude of the cross section change is about 6% in all the discovery region. No change of acceptance was observed. Taking 6% pessimistically as the PDF uncertainty, it is easy to estimate from the cross section slopes of Figure 15.6 that the uncertainty of the upper boundary of the discovery region is 1 - 2%. The uncertainty of its lower boundary coming from PDF is 2 - 3%.

In conclusion, the CMS potential to discover a heavy right-handed Majorana neutrino N_e and a heavy W_R gauge boson is studied. For the integral luminosity of 30 fb^{-1} these new




Figure 15.11: CMS discovery potential of the W_R boson and right-handed Majorana neutrinos of the Left-Right Symmetric model for the integrated luminosity $L_t = 30 \text{ fb}^{-1}$ (outer contour) and for $L_t = 1 \text{ fb}^{-1}$ (inner contour)

particles can be observed if M_{W_R} and M_{N_e} masses are less than 4 and 2.4 TeV/c^2 respectively. At the LRRP point (masses 2 TeV/c^2 and 500 GeV/c^2) W_R and N_e can be discovered already after one month of running at low luminosity.

5 Little Higgs models

FIXME: changed for content, results and plots- still fix plot

15.5.1 Introduction

During the last years, new solutions to the hierarchy problem were proposed in Little Higgs models [636]. These models introduce a set of new particles, with same statistics as the corresponding standard model particles and masses at the TeV scale aiming at compensating the contributions to the Higgs mass from the top quark, the gauge bosons and the Higgs boson itself.

The Littlest Higgs[637] model is the smallest extension of the SM to date that stabilizes the electroweak scale with a naturally light Higgs boson and weakly coupled new physics at TeV energies. A new heavy quark of charge $2/3$, marked as T , introduced to cancel the contribution to m_H from the top quark of SM.

15.5.1.1 Analysis path

The heavy T quark acquires its mass via Yukawa interactions with couplings λ_1 and λ_2 and its three dominant decay modes have branching ratios as follows: $BR(T \rightarrow t h) = BR(T \rightarrow t Z) = \frac{1}{2} BR(T \rightarrow b W)$.

The decay channel $T \rightarrow t Z$ with leptonic decays of Z^0 and W bosons provides a clean signature at the LHC environment. This channel has not been previously studied and the work presented here is a feasibility study.

The signal samples were generated with PYTHIA 6.227[30] generator package and for the T quark production the fourth quark generation matrix element calculation was activated (through the W - g fusion) which gives the correct kinematics of the single T quark. The T quark mass is set to $1 \text{ TeV}/c^2$ and its width to $1 \text{ GeV}/c^2$, to ensure the dominance of single T quark production rate. The data is simulated using a detailed Geant-based simulation (OSCAR) and reconstruction is performed with the CMS reconstruction package ORCA.

The major backgrounds considered in this analysis include : $t\bar{t}$, ZW +jets, ZZ +jets, WW +jets, Zbb , and Z +jets production.

The main selection requirements are summarized below:

- Events are required to pass the “double electron” or “double muon” L1 and HLT trigger criteria.
- Electrons are required to have $p_T > 20 \text{ GeV}/c$ and muons $p_T > 10 \text{ GeV}/c$
- The sum of the P_T of an same flavor opposite sign pair is required to be $P_T^{\ell\ell} > 100 \text{ GeV}/c$. The invariant mass of the pair is required to be consistent with the nominal Z^0 mass within $10 \text{ GeV}/c$.
- A third lepton is required; The sum of the third lepton transverse momentum and the missing transverse energy is required to be greater than $60 \text{ GeV}/c$. In addition the transverse mass of the third lepton with the missing transverse energy is required to be less than $120 \text{ GeV}/c^2$, to be consistent with the W boson transverse mass .
- Exactly one jet compatible with a b -jet and with calibrated $P_t > 30 \text{ GeV}/c$ is required.
- The sum of the W boson P_T and b -jet P_T should have exceeded $150 \text{ GeV}/c$ while their invariant mass is required to be in the range $110 - 220 \text{ GeV}/c^2$.
- The $cZ^0 W b$ system invariant mass is required to be in the range of the searched for heavy top, namely $(850 - 1150) \text{ GeV}/c^2$.

The ZZ diboson background is the only process that survives the above selection with a contribution of $N_B = 0.05$ events at 30 fb^{-1} . The total efficiency for the signal selection is $(9.7 \pm 0.4)\%$. The selected signal cross section is 192 fb for $M_T = 1 \text{ TeV}/c^2$ and $\lambda_1 = \lambda_2$ assuming $BR_{T \rightarrow t Z} = 0.25$. A total of $N_S = 2.1 \pm 0.1$ signal events are expected for 30 fb^{-1} .

The statistical significance (S_{C12} , see section ??) is 2.5 with a signal-to-background ratio of 41. Taking into account systematic uncertainties from the electron energy scale, jet and missing energy scale and b -tagging efficiency uncertainty, the significance drops to 2.0. For $\lambda_1/\lambda_2 = 2$ (Fig. ??), a 5σ significance can be achieved with 40 fb^{-1} as shown in figure 15.12.

15.6 Same Sign top

At the LHC di-leptonic $t\bar{t}(+jets)$ events can be selected with a relatively high signal-to-noise ratio and efficiency. Within the clean sample of such events, both leptons (electrons and muons) have an opposite electric charge. In several models beyond the Standard Model however, $tt/\bar{t}\bar{t}(+jets)$ topologies are predicted where both leptons have an equal electric charge. The signal excess is highly enhanced by the application of a combined likelihood




Figure 15.12: Minimum cross section for a 5 sigma discovery as a function of the luminosity for $M_T = 1 \text{ TeV}/c^2$. The horizontal lines correspond to the cross section for the two values of λ_1/λ_2 .

variable described in [272]. The likelihood variable is designed to differentiate the lepton from the W boson decay from leptons arising for example in QCD jets or from fake reconstructions. The signal of new physics can be diluted by the mis-identification of the electric charge of the leptons in Standard Model $t\bar{t}(+jets)$ events and the mis-identification of the leptons from the W decay themselves. The observability of an excess of same-sign signals above the mis-reconstruction of the Standard Model background is determined. The details of the analysis are mentioned in [?].

The jets in the final state are reconstructed with an Iterative Cone jet clustering algorithm using a cone size of $\Delta R = 0.5$. Input objects for the cones are selected from all calorimeter towers above a pseudo-rapidity dependent energy threshold determined from the average underlying event energy deposits [?]. The energy scale of the reconstructed jets is calibrated with corrections from Monte Carlo studies. The primary vertices in the proton bunch crossing are determined, and the vertex with the highest transverse momentum is taken as the one of the hard scattering. Via a track-based algorithm, jets are rejected if they do not match with this hard primary vertex.

The leptons are reconstructed and identified using the methods described in [272]. A likelihood variable is used to suppress leptons from the heavy flavour quark background exploiting several reconstruction aspects of leptons in the CMS detector. This likelihood is determined for each muon or electron in the final state in order to enhance the purity of choosing the correct lepton from the leptonic W decay. The combined likelihood includes observables as tracker isolation, calorimeter isolation, vertex matching significance, transverse momentum of the lepton and angular distance to the closest jet. For the electron likelihood a variable reflecting the reconstruction quality is added. The two muons or electrons having the largest combined likelihood ratio value are taken as the hard leptons of interest.

The inclusive single-muon, single-electron, double muon and double electron triggers are applied as described in [?]. The event should be triggered in at least one of these streams. In total 88.4%, 77.4% and 79.2% of respectively the $\mu\mu$, the ee and the μe signal events remain after applying this trigger criteria. The event is required to have at least 2 jets with a calibrated E_T above 25 GeV/c. These jets need to have a pseudo-rapidity in the range $|\eta| < 2.4$

	$\mu\mu$	μe and ee	$t\bar{t} \rightarrow \tau + X$	Other $t\bar{t}$	$W^\pm W^\mp$	$Z + jets$	S/N
Before selection	6915.0	20745.0	34606.2	485973.2	189951.7	578033.3	0.0078
Trigger	6114.7	16314.8	17415.6	100137.2	41288.4	266366.7	0.017
Two jets $E_T > 25$ GeV	4398.2	11982.7	13560.9	93858.2	20593.8	66146.7	0.032
b-tag criteria	989.8	2485.4	2289.6	8784.7	133.5	240.0	0.13
Two leptons identified	888.2	30.1	375.8	801.6	1.7	73.3	1.30
Two leptons selected	481.5	0.07	48.4	3.01	0.4	53.3	4.7
Efficiency (in %)	6.96	0.0003	0.14	0.0006	0.00022	0.0092	
Opposite-sign	481.3	0	48.3	2.19	0	53.3	
Same-sign	0.2	0.07	0.1	0.82	0.4	0	
	ee	μe and $\mu\mu$	$t\bar{t} \rightarrow \tau + X$	Other $t\bar{t}$	$W^\pm W^\mp$	$Z + jets$	S/N
Before selection	6915.0	20745.0	34606.2	485973.2	189951.7	578033.3	0.0078
Trigger	5354.8	17074.7	17415.6	100137.2	41288.4	266366.7	0.015
Two jets $E_T > 25$ GeV	3960.9	12420.0	13560.9	93858.2	20593.8	66146.7	0.029
b-tag criteria	802.7	2672.4	2289.6	8784.7	133.5	240.0	0.11
Two leptons identified	724.5	34.6	453.8	2283.6	73.1	126.7	0.57
Two leptons selected	285.0	0.3	37.5	5.2	0.8	53.3	3.1
Efficiency (in %)	4.12	0.0013	0.11	0.0011	0.00044	0.0092	
Opposite-sign	279.6	0.3	36.8	4.1	0.4	46.7	
Same-sign	5.4	0	0.7	1.1	0.4	6.7	
	$e\mu$	$\mu\mu$ and ee	$t\bar{t} \rightarrow \tau + X$	Other $t\bar{t}$	$W^\pm W^\mp$	$Z + jets$	S/N
Before selection	13830.0	13830.0	34606.2	485973.2	189951.73	578033.3	0.016
Trigger	10960.0	11469.5	17415.6	100137.2	41288.4	266366.7	0.030
Two jets $E_T > 25$ GeV	8021.8	8359.1	13560.9	93858.2	20593.8	66146.7	0.061
b-tag criteria	1682.7	1792.5	2289.6	8784.7	133.5	240.0	0.25
Two leptons identified	1500.6	66.4	822.1	3001.6	30.2	20.0	0.88
Two leptons selected	722.7	0.9	85.2	6.3	0.4	0	8.3
Efficiency (in %)	5.23	0.0065	0.25	0.0013	0.00022	0	
Opposite-sign	715.5	0.9	83.8	4.9	0	0	
Same-sign	7.2	0	1.3	1.4	0.4	0	

Table 15.7: Overview of the selection criteria applied on the events using simulated events with pile-up collisions included. The expected number of events are rescaled to a dataset of 1fb^{-1} taking into account the respective Leading-Order cross-sections of the processes.

and a b -tag discriminant larger than 0.5 [?]. The reconstructed hard leptons are required to have transverse momentum p_T exceeding 25 GeV/ c in the pseudo-rapidity range of $|\eta| < 2.4$ and a combined likelihood variable larger than 0.05.

In Table 15.7 the efficiencies and signal-to-noise ratios are shown after each selection step. Applying all cuts a signal-to-noise ratio of 4.7, 3.1 and 8.3 is obtained for respectively the $\mu\mu$, the ee and the $e\mu$ final state. Cross-talk between these three considered final states is by construction not possible. As the amount of selected WW and $Z + jets$ events in Table 15.7 is small, their contribution is alternatively estimated by multiplying the efficiencies of the event selection without the b -tagging and the individual b -tagging selection cut efficiency under the assumption that both selection cuts are uncorrelated.

It is illustrated [?] that from the selected topology of di-lepton $t\bar{t}$ events, a ratio $R = \frac{N_{++--}}{N_{+-}}$ can be determined which is sensitive to physics beyond the Standard Model. In the ratio the total amount of events with equally charged leptons is divided by the total amount of events with opposite charged leptons. As the efficiency of reconstructing the leptons electric charge is very high, we can neglect the amount of selected $pp \rightarrow tt$ or $pp \rightarrow t\bar{t}$ events observed with two opposite-charged leptons. Using the uncertainty on the ratio R , the significance of the observation of new physics channels $pp \rightarrow tt$ or $pp \rightarrow t\bar{t}$ is determined as a function of the cross section (see Figure 15.13). The di-muon channel has a larger sensitivity compared to the decay channels with electrons. This is shown to be induced by the electron reconstruction where a large fraction of electron energy clusters are matched with a wrong track resulting in a charge ambiguity.

It is assumed that the new physics processes beyond the Standard Model have a similar kinematic topology compared to the $t\bar{t}$ process, therefore the selection efficiency of the new physics channels is taken equal to that of the Standard Model $t\bar{t}$ process. Several models predict an excess of events with same-sign leptons in this topology, via the process $pp \rightarrow tt/\bar{t}\bar{t}$ or $pp \rightarrow tt/\bar{t}\bar{t} + b/c$. These models are motivated by Flavour Changing Neutral Currents (FCNC) [?, ?], topcolor-assisted Technicolor (TC2) [?] or supersymmetry [?]. With a measurement of R these kinematically similar processes $pp \rightarrow tt/\bar{t}\bar{t}$ can be observed with 30fb^{-1} of integrated luminosity if they have a cross section above 1 pb. Because a ratio of kinematically similar event topologies is measured, most of the experimental and theoretical systematic uncertainties cancel. The uncertainty on the magnitude of the cross section of the background processes on the significances shown in Figure 15.13 is found to be negligible. A feasibility study is performed to estimate the potential uncertainty on the mis-identification efficiency of the electric charge of electrons and muons from Z boson decays [?]. The effect on the significance of the excess of $tt/\bar{t}\bar{t}$ events is found to be negligible.




Figure 15.13: For an integrated luminosity of 30fb^{-1} the significance of the same-sign tt or $\bar{t}\bar{t}$ excess above the Standard Model events is indicated as a function of the cross-section of the inclusive process $pp \rightarrow tt/\bar{t}\bar{t}$.

Appendix A

95% CL limits and 5σ discoveries



A.1 On the True Significance of a Local Excess of Events

In searching for new phenomena in a wide range of possible signal hypotheses (e.g., a narrow resonance of unknown mass over a broad range background), a special care must be exercised in evaluating the true significance of observing a local excess of events. In past, this fact was given substantial scrutiny by statisticians (e.g., [733, 734]) and physicists alike (e.g., [735, 736, 737, 738, 739]). The purpose of this Appendix is to quantify a possible scope of this effect on an example of a search for the Standard Model Higgs boson in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ decay channel. As the case study, we chose a counting experiment approach widely used in this volume.

The dashed line in Fig. A.1 shows the expected 4μ invariant mass distribution for background at $L=30 \text{ fb}^{-1}$ after applying all the $m_{4\mu}$ -dependent analysis cuts described in Section 3.1. Using this distribution, we played out $\sim 10^8$ pseudo-experiments; an example is shown in Fig. A.1. For each pseudo-experiment, we slid a *signal region window* across the spectrum looking for a local event excess over the expectation. The size of the window $\Delta m = w(m_{4\mu})$ was optimized and fixed *a priori* (about $\pm 2\sigma$) to give close to the best significance for a resonance with a width corresponding to the experimental SM Higgs boson width $\sigma(m_{4\mu})$. The step of probing different values of $m_{4\mu}$ was “infinitesimally” small (0.05 GeV) in comparison to the Higgs boson width of more than 1 GeV. The scanning was performed in *a priori* defined range of 115-600 GeV.

We used a significance estimator $S_{cL} = \text{sign}(s) \sqrt{2n_o \ln(1 + s/b) - 2s}$, where b is the expected number of background events, n_o is the number of observed events, and the signal is defined as $s = n_o - b$. This estimator, based on the Log-Likelihood Ratio, is known to follow very closely the true Poisson significance, only slightly over-estimating it in the limit of small statistics [42]. Figure A.2 presents the results of such a scan for the pseudo-experiment shown in Fig. A.1. The maximum value of S_{cL} , S_{max} , and the corresponding mass of a “Higgs boson candidate” obtained in each pseudo-experiment were retained for further statistical studies.

After performing 10^8 pseudo-experiments, the differential probability density function for S_{max} and its corresponding cumulative probability function $P(S_{max} > S)$ (Fig. A.3) were calculated. From Fig. A.3, one can see that the frequency of observing some large values of S_{cL} (solid line) is much higher than its naive interpretation might imply (dashed line). If desired, the actual probability can be converted to the true significance. The result of such “renormalization” is presented in Fig. A.4. One can clearly see that the required de-rating of significance is not negligible; in fact, it is larger than the effect of including all theoretical and instrumental systematic errors for this channel (see Section 3.1). More details on the various

aspects of these studies can be found in [42].

There are ways of reducing the effect. A more detailed analysis of the shape of the $m_{4\mu}$ distribution will help somewhat. Using the predicted number of signal events $s = s_{theory}$ in the significance estimator to begin with and, then, for validating the statistical consistency of an excess $n_o - b$ with the expectation s_{theory} will reduce the effect further. One can also use a non-flat prior on the Higgs mass as it comes out from the precision electroweak measurements. Whether one will be able to bring the effect to a negligible level by using all these additional constraints on the signal hypotheses is yet to be seen. The purpose of this Appendix is not to give the final quantitative answer, but rather to assert that these studies must become an integral part of all future search analyses when multiple signal hypotheses are tried.




Figure A.1: The background *pdf* and an example of one pseudo-experiment with a statistical fluctuation appearing just like a signal.




Figure A.2: Profile of the S_{cL} scan corresponding to the pseudo-experiment example shown on the left. Green (inner) and yellow (outer) bands denote $\pm 1\sigma$ and $\pm 2\sigma$ intervals. Spikes that can be seen are due to events coming in or dropping off the trial-window, a feature of low-statistics searches.




Figure A.3: S_{cL} cumulative probability density function.




Figure A.4: Local significance “renormalization” from an observed value to the true significance with a proper probabilistic interpretation.

Appendix B

Systematic errors



The simulation of events at the LHC is complex and can be conventionally divided into different parts which either involve the description of the interesting physics process or the description of the initial scattering conditions and the physics environment.

The simulation of the hardest part of the physics process is done via matrix element (ME) calculations at a certain order in the coupling constants and continues with the parton showering (PS) of the resulting partons until a cut-off scale, over which the perturbative evolution stops and the fragmentation of the final partons takes on. This cut-off is often referred to as factorization scale, because it is the scale at which the two processes (showering and fragmentation) are supposed to factorize.

The interesting event is accompanied by the so-called underlying event (UE), term which identifies all the remnant activity from the same proton-proton (p-p) interaction and whose definition often includes ISR as well, and the pile-up, composed by other minimum bias (MB) p-p interactions in the same bunch crossing (up to 25 at high luminosity at the LHC). Moreover, since the initial state is not defined in p-p collisions, a proper description of the proton parton density functions (PDFs) should be included in the calculations.

Each of these effects needs to be modeled to the best of our knowledge, and the associated uncertainties need to be determined and propagated to the physics measurements. Moreover, many of the sources are correlated: for instance, fragmentation and showering are obviously dependent on each other, and in turn they assume a certain description of the underlying event. The task of assessing systematics due to theory and modeling can therefore be a difficult one and can sometime contain a certain degree of arbitrariness.

In what follows we propose some guidelines for the estimation of errors coming from the above, trying to divide the systematics sources into wider categories as much uncorrelated as possible: QCD radiation, fragmentation description, PDFs, UE and MB.

In attributing systematic errors we believe that one should use motivated recipes, avoiding unrealistic scenarios which will lead to unnecessarily conservative errors or, much worse, totally arbitrary assumptions.

B.1 Hard process description and parametric uncertainties

The description of the hard process should be done with Monte Carlo tools which are best suited to the specific analysis. For instance, when precise description of hard gluon emission becomes an issue, then next-to-leading order (NLO) generator tools like MCNLO [740],

or higher leading order (LO) α_s generators like COMPHEP [34], MADGRAPH [73], ALPGEN [152], and SHERPA [185] should be considered. This is in general true for both the signal and the background description.

When adopting a ME tool, one should always keep in mind that its output is often (if not always) supposed to be interfaced to PS Monte Carlo such as HERWIG [187], PYTHIA [30] or ISAJET [653], that treat the soft radiation and the subsequent transition of the partons into observable hadrons. One of the most difficult problems is to eliminate double counting where jets can arise from both higher order ME calculations and from hard emission during the shower evolution. Much theoretical progress has been made recently in this field [741, 742, 743, 744]. For what concerns the ME/PS matched description of multi-jet final states, a rich spectrum of processes is currently available in ALPGEN. However, adopting general purpose generators like PYTHIA can still be the best option for topologies that are better described in the Leading Logarithm Approximation (LLA), for instance in the case of two leading jets and much softer secondary jets. The two different descriptions should be regarded as complementary.

In general, a sensible choice for the selection of the best generation tools can be driven by the HEPCODE data base [745]. However, comparison between different generators is recommended whenever applicable.

Each analysis needs then to make sure that other important effects (e.g. spin correlations in the final state, NLO ME corrections to top decays) are included in the generation mechanism. For example TOPREX [35], as long with some of the Monte Carlo generators already introduced in this section, provides a correct treatment of top quark spin correlations in the final state. Neglecting some of these effects corresponds to introducing an error in the analysis that cannot be considered as coming from a theoretical uncertainty.

For both signal and backgrounds, missing higher orders are a delicate source of uncertainty. Formally, the associated error cannot be evaluated unless the higher order calculation is available. This is often not possible, unless extrapolating by using comparisons with analytical calculations of total or differential cross-sections at the next order, if available. One should keep in mind that simple K-factors are not always enough and that the inclusion of higher orders typically also involves distortions in differential distributions.

Moreover, one should not forget that any Standard Model calculation is performed in certain schemes and that the input parameters are subject to their experimental uncertainties; if the error on most of those and the choice of the renormalization scheme are expected to give negligible effects in comparison with other uncertainties, this might not be so for the choice of the hard process scale, which we will discuss in the next section, and some of the input parameters.

Among the input parameters, by far the one known with less accuracy will be the top mass. The current uncertainty of about 2% [746] enters in the LO calculations for processes which involve top or Higgs production. For instance, the total $t\bar{t}$ cross-section is known to have a corresponding 10% uncertainty due to this [36]. As far as Higgs production (in association or not with tops) is concerned, gluon-gluon fusion proceeds via a top loop and therefore the total cross-section can have a strong dependence on the top mass when $m_H \approx 2m_t$. Analyses which include Higgs bosons or top are encouraged to estimate the dependence of the significant observables on the top mass itself. Effects of m_t variation on acceptances of these analyses should instead be negligible.

B.2 Hard process scale

The hard process under study drives the definition of the Q^2 scale, which directly enters in the parameterization of PDFs and α_s , hence in the expression of the cross sections.

The dependence of the observables on the choice for the Q^2 hard process scale is unphysical and should be regarded as one important contribution to the total uncertainty in the theoretical predictions. The sensitivity of the predicted observables to such choice is expected to decrease with the increasing order in which the calculation is performed, and can be tested by changing the hard process scale parameters in the generation (where applicable) using a set of sound values according to the characteristics of the hard process.

A sensible choice for the hard process scale in $2 \rightarrow 1$ processes is often \hat{s} , which is the default in general purpose generators like PYTHIA. Alternative choices to quote theoretical uncertainties can be $0.25\hat{s}$ and $4.0\hat{s}$. In PYTHIA this can be obtained acting on PARP(34).

For $2 \rightarrow n$ processes, many reasonable alternatives for the Q^2 scale definition exist. The PYTHIA default (MSTP(32)=8), corresponds to the average squared transverse mass of the outgoing objects. It is possible to test the sensitivity on the Q^2 scale switching to different options, for example trying $Q^2 = \hat{s}$ (MSTP(32)=4 in PYTHIA).

B.3 PDF description

The parton distribution functions of interacting particles describe the probability density for partons undergoing hard scattering at the hard process scale Q^2 and taking a certain fraction x of the total particle momentum. Since the Q^2 evolution can be calculated perturbatively in the framework of QCD, PDFs measurements can be cross checked using heterogeneous DIS, Drell-Yan and jet data, and achieve predictivity for points where no direct measurements are available yet, for example in a large region of the (x, Q^2) space for p-p interactions at the LHC energy.

Various approaches are currently available to quote the PDFs of the proton, which propose different solutions for what concerns the functional form, the theoretical scheme, the order of the QCD global analysis (including possible QED corrections), and the samples of data retained in the fits: CTEQ [747], MRST [748], Botje [749], Alekhin [750] etc. The CTEQ and MRST PDFs, including Tevatron jet data in the fits, seem to be well suited for use in Monte Carlo simulations for the LHC.

The best way to evaluate theoretical uncertainties due to a certain proton PDFs is to vary the errors on the parameters of the PDF fit itself. With the Les Houches accord [88] PDF (LHAPDF) errors should be easily propagated via re-weighting to the final observables. However, errors are available only for NLO PDF, whereas in most of the cases only LO tools are available for the process calculation. Correctly performing evaluation of theoretical uncertainties in these cases requires some care. The proposed solution is to adopt CTEQxL (LO) for the reference predictions using CTEQxM (NLO) only to determine the errors.

For analyses which are known to be particularly sensitive to PDFs, like cross-section measurements, it would be also desirable to compare two different sets of PDFs (typically CTEQ vs MRST) taking then the maximum variation as an extra error. This is important since, even considering the error boundaries, different set of PDFs may not overlap in some region of the phase space.

The LHAGLUE interface [88] included from the most recent LHAPDF versions simplifies the use of the Les Houches accord PDF in PYTHIA by the switches $\text{MSTP}(52) = 2$, $\text{MSTP}(51) = \text{LHAPDF}_{id}$.

B.4 QCD Radiation: the Parton Shower Monte Carlo

The showering algorithm is basically a numeric Markov-like implementation of the QCD dynamic in the LLA. After the generation of a given configuration at partonic level, the initial state radiation (ISR) and the final state radiation (FSR) are produced following unitary evolutions with probabilities defined by the showering algorithm.

The probability for a parton to radiate, generating a $1 \rightarrow 2$ branching, are given by the Altarelli-Parisi equations [751], however various implementations of the showering algorithm exist in parton shower Monte Carlo, which mostly differ for the definition of the Q^2 evolution variable (virtuality scale) in the $1 \rightarrow 2$ radiation branching and for the possible prescriptions limiting the phase space accessible to the radiation: PYTHIA, HERWIG, ARIADNE [752], ISAJET etc.

The virtuality scales for both ISR and FSR need to be matched to the hard process scale, the latter setting an upper limit on the former ones; such limit has to be considered in a flexible way, given the level of arbitrariness in the scale definitions. While this matching is somewhat guaranteed if one adopts the same simulation tool for both hard scattering and parton shower, a careful cross check is recommended in all other cases. In general, a critical judgement taking into account the hard process type is needed. Allowing a virtuality scale higher than the hard process scale may give rise to double counting. This is the case of $gg \rightarrow gg$ processes with additional hard gluons added in the showering. However other processes are safer from this point of view, for instance the case of the $q\bar{q} \rightarrow Z$ process at LO.

Quantum interference effects in hadronic collisions have been observed by CDF [753] and D0 [754] studying the kinematical correlations between the third jet (regarded as the result of a soft branching in the LLA) and the second one. The implementation of the so called color coherence in PS Monte Carlo is made in the limit of large number of colors and for soft and collinear emissions, restricting the phase space available to the radiation depending on the developed color configuration. Different implementations of the color coherence are available in HERWIG and PYTHIA, while ISAJET doesn't take into account such effects.

The theoretical uncertainty associated to the parton showering descriptions, includes what is normally referred to as ISR or FSR and their interference. In order to achieve practical examples for the recommended parton shower settings, we will consider PYTHIA as the default tool for showering from now on.

Turning OFF ISR and FSR ($\text{MSTP}(61)=0$, $\text{MSTP}(71)=0$ respectively) or even the interference part ($\text{MSTP}(62)=0$, $\text{MSTP}(67)=0$) is certainly a too crude approach and, to a large extent, a totally arbitrary procedure to assess a systematic error. We believe it is much more realistic to vary, according to sound boundaries, the switches regulating the amount and the strength of the radiation of the showering. These can correspond to Λ_{QCD} and the maximum virtuality scales up to which ISR stops and from which FSR starts. It would be important to switch the parameters consistently going from low to high values in both ISR and FSR.

Notice that the radiation parameters were typically fitted at LEP1 together with the frag-

mentation parameters, benefiting from a much simplified scenario where no ambiguity on the maximum virtuality scale applies, the only relevant energy scale of the problem being $\hat{s} = s$. One has to take into account that while for instance FSR accompanying heavy boson decays at the LHC can be directly related to the LEP experience, FSR in processes like $gg \rightarrow b\bar{b}$ entails additional uncertainties arising from the maximum allowed virtuality scale and ISR/FSR interference. On top of that, additional complications arise from the fact that ISR at hadron machines contributes to the description of the underlying event. Matching two different tunings of the same parameter (in particular PARP(67)) can be very subtle at the LHC.

These are the suggested settings in PYTHIA, which have been cross-checked with the ones adopted by the CDF experiment and also follow the prescription by the main author:

- Λ_{QCD} : PARP(61), PARP(72), PARJ(81) from 0.15 to 0.35 GeV consistently, symmetric with respect to 0.25. Notice that these settings have been optimized for the CTEQ6L PDFs. In general different ranges apply when changing PDFs. In order to give the user full control on these parameters the option MSTP(3)=1 has to be set, otherwise Λ_{QCD} is assumed to be derived from the PDFs parameterization.
- Q_{max}^2 : PARP(67) from 0.25 to 4 and PARP(71) from 1 to 16 going from low to high emission in a correlated way. In doing so one should also make sure that the tuning of the underlying event is not changing at the same time. Possible re-tuning of the underlying event in different radiation scenarios may be needed, in particular for what concerns PARP(82).

B.5 Fragmentation

Perturbative QCD cannot provide the full description of the transition from primary quarks to observable hadrons, but only the part which involves large momentum transfer. The formation of final hadrons involves a range of interactions which goes above the Fermi scale and where the strong coupling constant α_s increases above unity, making it necessary to describe this part in a non-perturbative way, normally referred to as fragmentation or hadronization.

The non-perturbative description of fragmentation is realized via models, which need to be tuned to experimental data. The data correspond, typically, to event shapes and multiplicities at leptonic machines or to the inclusive jet shapes at hadronic machines. A comprehensive overview of the models can be found in [755].

Fragmentation is said to depend only on the factorization scale if jet universality is assumed, i.e. assuming that jets fragment in the same way at hadron and lepton machines. Jet universality will be ultimately verified at the LHC; one should clarify whether instrumental effects and the LHC environment will have an impact on the final observables. For instance, the much larger fraction of gluon jets or the different description of the underlying event can change the values of the parameters that regulate the fragmentation. Moreover, for events with high multiplicity of jets it will also be crucial to properly describe fragmentation in conditions where large jet overlapping is to be expected and where inclusive tunings might not be ideal.

The consequence of jet universality is that, once the PS cut-off scale is fixed, the fragmentation description for light quarks should be universal, and the LEP/SLD tunings (or the Tevatron ones) could be used as they are for the LHC.

It is important to underline that the description of the non-perturbative part of the radiation also depends on the way the perturbative one is described. This means that one should not use a tuning of fragmentation done with LO(+LL) tools (typically PYTHIA at LEP) attached to perturbative calculation which are done at higher -or different- order.

B.5.0.2 Light quarks fragmentation

In the absence of LHC data, the best choice is therefore to use a model tuned to the LEP and SLD data [756, 757, 758]. It is important to choose the tuning in a consistent way from the same experiment, given that a combined LEP/SLD tuning has never been attempted. As a possibility, suggested by the major success in describing the data and by its extensive use in the experimental collaborations, is the use of PYTHIA, which uses the string (or Lund) fragmentation model [759]. The parameters that we consider more relevant in PYTHIA for the description of fragmentation are the following, where the central value is taken by the fit performed by the OPAL Collaboration, as an example:

$$\text{PARJ}(81) = 0.250$$

$$\text{PARJ}(82) = 1.90$$

$$\text{PARJ}(41) = 0.11$$

$$\text{PARJ}(42) = 0.52$$

$$\text{PARJ}(21) = 0.40$$

where $\text{PARJ}(81)$ (Λ_{QCD}) and $\text{PARJ}(82)$ (Q_{min}^2) refer to the radiation part. To properly evaluate a systematic error due to pure fragmentation one should vary only $\text{PARJ}(42)$ and $\text{PARJ}(21)$ by their respective errors (0.04 and 0.03 for OPAL). The variation should account for the proper parameter correlation if the effect is critical for the analysis. $\text{PARJ}(41)$ is totally correlated to $\text{PARJ}(42)$.

Alternatively, or additionally, it would also be important to compare PYHTIA with HERWIG with consistent tunings from LEP [756, 757, 758]; in doing so it is important to factorize the UE description (see next section) that can induce important differences in the results.

B.5.0.3 Heavy quarks fragmentation

The description of the heavy quarks fragmentation is important for top physics and for those processes with large b production in the final states. Exclusive channels are particularly influenced by the description of the fragmentation of the b quark.

The description of the fragmentation of the heavy quarks has been tuned to Z data at LEP and SLD [760, 757, 761, 762] (via a measurement of x_B and x_D) and $b\bar{b}$ data at the Tevatron, using different fragmentation functions like Lund, Bowler [763], Peterson [764], Kartvelishvili [765].

In the spirit of fragmentation universality the LEP/SLD tunings can be adopted for the LHC, but with much care. Significant differences among the fitted values in different experiment can point out that the factorization scale used for the PS is not the same everywhere. One should make sure that the scale used is set consistently with the chosen fragmentation function parameters. This can be done by using the tuning from only one experiment, making sure to also use the main switches of the parton showering, ($\text{PARJ}(81)$ and $\text{PARJ}(82)$ in PYTHIA).

The fragmentation function that best describes heavy flavour data at LEP is Bowler. With the same OPAL tuning reported above the best fit of the Bowler parameters, a and bm_{\perp}^2 , to data gives:

$$bm_{\perp}^2 = 65_{-14}^{+17}$$

$$a = 15.0 \pm 2.3$$

The Bowler model would extend the string model to heavy flavours, describing the corrections in terms of the charm and bottom masses. Unfortunately, no tuning exists in the literature which is capable to describe at the same time light and heavy quark fragmentation, i.e. adopting universal parameters $a = \text{PARJ}(41)$ and $b = \text{PARJ}(42)$ for both light and heavy quarks.

Alternatively, the widely used Peterson function can be used, and its parameters are directly switchable in PYTHIA for just b and c fragmentation:

$$\text{PARJ}(54) = -0.031 \pm 0.011$$

$$\text{PARJ}(55) = -0.0041 \pm 0.0004$$

where the two parameters correspond, respectively, to ϵ_c and ϵ_b fitted in the OPAL tuning. The systematic can then be evaluated by varying the errors on the fitted parameters or by comparing with a different fragmentation function like Kartvelishvili, or Lund.

An important feature of the b fragmentation that should be considered by those analyses in the top sector sensitive to the details of the fragmentation, is the way the b fragments in top decays. At the LHC the b from a t is hadronizing with a beam remnant, introducing potentially worrying differences with respect to the fragmentation at LEP. The main effects are presented in [766] and are known as *cluster collapse*, happening when a very low mass strings quark-remnant directly produces hadrons without fragmenting, hence enhancing the original flavour content, and *beam drag*, which is an angular distortion of hadron distribution toward the end of the string in the remnant. If, under reasonable assumptions on the transverse momentum in top events at the LHC, one can exclude to a large extent the importance of the first effect, beam drag could potentially introduce B meson production asymmetries, even though estimations are keeping the effect at the level of 1% at the LHC [766].

B.6 Minimum bias and Underlying event

Multiple parton interaction models, extending the QCD perturbative picture to the soft regime, turn out to be particularly adequate to describe the physics of minimum bias and underlying event. Examples of these models are implemented in the general purpose simulation programs PYTHIA, HERWIG/JIMMY [184], and SHERPA. Other successful descriptions of underlying event and minimum bias at hadron colliders are achieved by alternative approaches like PHOJET [767], which rely on both perturbative QCD and Double Pomeron Models (DPM).

Huge progress in the phenomenological study of the underlying event in jet events have been achieved by the CDF experiment at Tevatron [768], using the multiplicity and transverse mo-

momentum spectra of charged tracks in different regions in the azimuth-pseudorapidity space defined with respect to the direction of the leading jet. Regions that receive contributions only by the underlying event have been identified. The average charged multiplicity per unit of pseudorapidity in these regions turns out to be significantly higher with respect to the one measured in minimum bias events. This effect, referred to as "pedestal effect", is well reproduced only by varying impact parameters models with correlated parton-parton interactions ($MSTP(82) > 1$ in PYTHIA). Simpler models are definitely ruled out.

The main problem of extrapolating the predictions of the multiple interactions models to the LHC is that some of the parameters are explicitly energy dependent, in particular the color screening p_T cut-off (PARP(82) at the tuning energy PARP(89) in PYTHIA). The CDF tuning, often referred to as Tune-A, is not concentrating on this particular aspect. Other works [188, 769] have put more emphasis on this issue. However, one of their results is that currently only PYTHIA can be tuned to provide at the same time description of CDF and lower energy minimum bias data from UA5. One of these tunings can be summarized as follows:

PARP(82) = 2.9
 PARP(83) = 0.5
 PAPER(84) = 0.4
 PARP(85) = 0.33
 PARP(86) = 0.66
 PARP(89) = 14000
 PARP(90) = 0.16
 PARP(91) = 1.0
 MSTP(81) = 1
 MSTP(82) = 4

Sensible estimation of theoretical uncertainties arising from underlying event and minimum bias modeling can be performed assigning $\pm 3\sigma$ variations to the color screening p_T cut-off parameter tuned on minimum bias CDF and UA5 data and extrapolated to the LHC energy [769], i.e. varying PARP(82) in the range [2.4-3.4], while keeping the other parameters listed above to their tuned values.

As a new tool for the description of UE and MB we would like to mention PYTHIA 6.3 [770], that allows for new interesting features, including the new p_T -ordered initial- and final-state showers and a new very sophisticated multiple interactions model that achieves description of colliding partons in the proton in terms of correlated multi-parton distribution functions of flavours, colors and longitudinal momenta. However, as stressed by the PYTHIA authors, the new model (PYEVNW) is still not so well explored. Therefore the old model (PYEVNT) is retained as the default choice, with full backward compatibility. Moreover, in the use of PYTHIA 6.3, one should be careful when switching to the new p_T -ordered showers and multiple interaction models, as their parameters are not tuned yet, in particular for what concerns the energy dependence, necessary to get meaningful extrapolations at the LHC energy.

B.7 Pile up and LHC cross sections

The design parameters of the LHC at both low and high luminosity are such that, on top of possible signal events, additional minimum bias interactions are produced in the same beam crossing, the so-called Pile-up effect.

Pile-up is a purely statistical effect. The number of minimum bias interactions generated in a single beam crossing is a poissonian distribution that depends on the instantaneous luminosity, which varies of about a factor 2 during a LHC fill. Although luminosity variation is not arising from theoretical uncertainties, it is recommended to cross check the stability of the results against variation of the nominal luminosity.

An issue which can affect the Pile-up is the definition of the minimum bias itself. The latter, indeed, may or may not include the diffractive and elastic contributions, with figures for the total cross section which can vary from 100 mb to 50 mb respectively. If the PYTHIA generator is adopted, these two different options correspond to MSEL 2 and MSEL 1, however, in order to get full control on the different contributions to the cross sections, one can use MSEL 2, setting MSTP(31) = 0, and providing explicit input through SIGT(0,0,J), where the meaning of the index J is described below:

J=0 Total cross section (reference value = 101.3 mb).

J=1 Elastic cross section (reference value = 22.2 mb).

J=2 Single diffractive cross section XB (reference value = 7.2 mb).

J=3 Single diffractive cross section AX (reference value = 7.2 mb).

J=4 Double diffractive cross section (reference value = 9.5 mb).

J=5 Inelastic, non-diffractive cross section (reference value = 55.2 mb).

Where J=0 has to correspond to the sum of the contributions for J=1,...,5. With respect to alternative cross section predictions [771], PYTHIA reference values for diffractive cross sections might be slightly shifted on the high side. A possible sound alternative could be to reduce the diffractive cross sections of around 30%, keeping constant the total cross section.

In order to assess the sensitivity of one analysis to the diffractive variations in the pile-up, at least the two options MSEL 1 and MSEL 2 should be tried. Diffractive contribution will in general result in few additional soft charged particles spiraling in the high magnetic fields of the LHC experiments. This effect is most likely to be relevant in the tracker detectors, where multiple hits in the same layer can be generated by the same track.

B.8 Decays

In contrast to the simple decay models available in the common PS Monte Carlo, alternative hadron decay models exist, for example EvtGen [772], which have huge collections of exclusive hadron decays up to branching ratios as low as 10^{-4} .

EvtGen follows the spin density matrix formalism and has an easily tuneable and upgradeable hadron decay data base which currently constitutes the largest and most refined collection of hadron decay models.

Comparison between the simple default decay models implemented in PS Monte Carlo and those available in EvtGen should be recommended at least for analyses dealing with

B hadrons or relying on b-tagging. However, since switching to a new hadron decay model could have a deep spin-offs on the exclusive description of the final states (multiplicity of kaons, pions, photons and muons, multiplicity of tracks reconstructed in secondary vertices) it might be worth to study also effects on trigger performances.

The LHC version of EvtGen was initially provided by the LHCb experiment and is currently maintained by LCG Generator [773]. It comprises an interface to PYTHIA simulation that solves the technical problems of switching between the two different scenarios (i.e. hadron decays performed by PYTHIA, hadron decays performed by EvtGen).

B.9 LHAPDF and PDF uncertainties

The detailed investigations of processes at LHC required a well understanding of the systematic theoretical uncertainties [192]. One of the important source of such errors is the parton distribution functions (PDFs).

The Les Houches Accord Parton Density Functions (LHAPDF) package [88] is designed to work with the different PDF sets *. In this approach a “fit” to the data is no longer described by a single PDF, but by a PDF set consisting of many individual PDF members. Indeed, PDFs are specified in a parametrized form at a fixed energy scale Q_0 , such as

$$f(x, Q_0) = a_0 x^{a_1} (1-x)^{a_2} (1+a_3 x^{a_4} \dots) \quad (\text{B.1})$$

The PDFs at all higher Q are determined by NLO perturbative QCD evolution equations. The total number of PDF parameters (d) could be large (for example, for popular CTEQ parameterization one has $d = 20$ [29]). Fitting procedure is used for evaluation an effective χ^2 function, which can be used to extract the “best fit” (the global minimum of χ^2) and also to explore the neighborhood of the global minimum in order to quantify the uncertainties. As a result one has the “best-fit” PDF and $2d$ subsets of PDF [29, 88]:

$$f_0(x, Q), f_i^\pm(x, Q) = f(x, Q; \{a_i^\pm\}), \quad i = 1, \dots, d \quad (\text{B.2})$$

B.9.0.4 Master equations for calculating uncertainties

Let $X(\{a\})$ be any variable that depends on the PDFs. It can be a physical quantity such as the W production cross section, or a differential distribution.

Let $X_0 = X(\{a_0\})$ be the estimate for X calculated with the best-fit PDF and X_i^\pm be the the observable X calculated with i -th subset $f_i^\pm(x, Q)$.

Following to CTEQ6 collaboration one can estimate the variation of X by using a master formula [29]:

$$\Delta X = \sqrt{\sum_{i=1}^d (X_i^+ - X_i^-)^2} \quad (\text{B.3})$$

*Note, at CMS it was recommended to use the CTEQ 5L set for PTDR simulation. Since there is only **one** CTEQ 5L PDF set (without corresponding subsets), it was recommended to use CTEQ 6M for evaluation of uncertainties due to PDFs for PTDR estimates and only in a special case one can use another sets (e.g. MRST).

However, very often many X_i^+ and X_i^- have different magnitudes and even signs ! This failure of the master formula is a result of the simple observation that the PDF set that minimizes the uncertainty in a given observable X is not necessarily the same as the one that minimizes the fit to the global data set.

The better estimator for the uncertainty of a generic observable X was proposed in [774]. It is defined as the maximum positive and negative errors on an observable X by

$$\left. \begin{aligned} \Delta X_+ &= \sqrt{\sum_{i=1}^d (\max[(X_i^+ - X_0), (X_i^- - X_0), 0])^2}, \\ \Delta X_- &= \sqrt{\sum_{i=1}^d (\max[(X_0 - X_i^+), (X_0 - X_i^-), 0])^2} \end{aligned} \right\} \quad (\text{B.4})$$

In Eqs. (B.4) one sums the maximum deviations on the observable in each of the parameter directions, and hence retain both maximal sensitivity to the parameters that vary most and estimate the range of allowed values of the cross section. Note, that the errors in Table ?? were evaluated with this Eq. (B.4).

Eq. (B.4) could also be used for calculations of differential distribution. Fig. B.1 presents the differential distribution $d\sigma/dP_T$ for $t\bar{t}$ -pair production at LHC.

Figure B.1: $d\sigma/dP_T$ distribution for $t\bar{t}$ -pair production at LHC. The central histogram corresponds to the 'best-fit' of CTEQ6M PDF, while the shaded area represents the deviation due to PDF uncertainties.

B.9.0.5 How to calculate $X(\{a_i\})$?

The most simple and straightforward method is to simulate a sample with the "best-fit" PDFs and then to repeat a such simulation $2d$ times with different $2d$ PDF subsets. As a results one gets $(1 + 2d)$ samples of **unweighted** events with **different** kinematics for each samples. Then use these samples to calculate $(1 + 2d)$ values for observable:

$$X_0 = \sum_{\text{events}} X_n(\{a_0\}), \quad X_i^\pm = \sum_{\text{events}} X_n(\{a_i^\pm\}), \quad i = 1, \dots, d \quad (\text{B.5})$$

In practice, such method requires a large CPU-time and can be recommended only to be used for very few special cases, when a high accuracy is required.

In the second approach ("**re-weighting**" method) one needs to simulate only **one** sample with the 'best-fit' PDF. In doing so the additional weights, corresponding to all other PDF subsets are evaluated. This weight is the ratio of the parton luminosity [PDF($\{a_i\}$) – the product of PDFs] evaluated with PDF subset to the parton luminosity, calculated with the 'best-fit' PDF. As a result, for any n -event one has $2d$ additional weights:

$$w_{(0)} = 1(\text{best fit PDF}), \quad w_{(i)}^\pm = \frac{\text{PDF}(\{a_i^\pm\})_n}{\text{PDF}(\{a_0\})_n}; \quad w_{(i)}^\pm = \mathcal{O}(1) \quad (\text{B.6})$$

The corresponding $(1 + 2d)$ values for observable X are evaluated as follows:

$$X_0 = \sum_{\text{events}} X_n(\{a_0\}), \quad X_i^\pm = \sum_{\text{events}} w_{(i)}^\pm X_n(\{a_0\}) \quad (\text{B.7})$$

Contrary to the first method (see (B.5)) these $(1 + 2d)$ samples have the events with **different** weights, but with **identical** kinematics for each samples. Note, that all additional samples

have different “total number of events”:

$$N_0 = \sum_{\text{events}} w_{(0)} (= 1), \quad N_i^\pm = \sum_{\text{events}} w_{(i)}^\pm \neq N_0, \quad \text{and} \quad N_i^\pm = \mathcal{O}(N_0) \quad (\text{B.8})$$

Starting from CMKIN.6.0.0 version it is possible for each event the evaluation of the additional weights, corresponding to different PDF subsets (i.e. $w_{(i)}^\pm$, see (B.6)). This option is available for CMKIN run with PYTHIA-like generators (PYTHIA, MadGraph, CompHEP, ALPGEN, TopReX, StaGen, etc) and HERWIG. This information is written in `/mc_param/` user block after all variables filled by CMKIN and a user (by using of `kis_xxx` routines).

Appendix C

Monte Carlo models and generators



C.1 Introduction

This section presents a short description of the basic event generators used in CMS during preparation of PTDR (see CMS “Generator Tools group” for details). The nice review of the present Monte Carlo models and generators is given elsewhere [775]. Note, that only MC generators used in CMS are described here, while a description of several popular packages (like ISAJET or AcerMC, see [775]) is omitted.

There are several available Monte Carlo event generators for pp , pA , and AA collisions, namely HERWIG [187], HIJING [776], ISAJET [653], PYTHIA [62], and SHERPA [777]. Each of these simulates a hadronic final state corresponding to some particular model of the underlying physics. The details of the implementation of the physics are different in each of these generators, however the underlying philosophy of the generators is the same.

The cross section values and the differential distribution for almost all processes are evaluated with parton model with the basic equation as follows:

$$\sigma(pp \rightarrow CX) = \sum_{ij} \int f_i^p(x_1, Q^2) f_j^p(x_2, Q^2) \hat{\sigma}(ij \rightarrow C) dx_1 dx_2, \quad (\text{C.1})$$

where $f_i^p(x, Q^2)$ are the Parton Distribution Functions (PDF) of i th parton, that carried out the fraction x of the initial proton momentum at specific scale (Q^2); $\sigma(ij \rightarrow C)$ is the cross section for the hard process (i.e. describing two partons, i and j , interaction).

A general scheme of event generation assumes evaluation of the hard process (the cross section value, the incoming and outgoing particle’s momenta and colours), then to evolve the event through a parton showering and hadronization the coloured products of shower, and the decay of the unstable particles. As a result an event information (stored in /HEPEVT/common block [62]) contains the momenta of the final hadrons, leptons and photons and positions of their decay vertexes. Typically such information contains also the characteristics (momenta, colours, KF-codes, mother’s and daughter’s relations) of all intermediate partons (quarks, gluons, gauge bosons, unstable physical particles, etc) that provide to trace-back the history of particles production insight of event. Any such generator by using an acceptance-rejection method (Von Neumann) returns weighted event.

The parton showering is based on the expansions around the soft and collinear limits and can be ascribed to either the initial or final state. As compared to PYTHIA the algorithm used by HERWIG and SHERPA includes some effects due to quantum interference. The events

that have more energy in the parton process have more showering, and consequently more jet activity.

The collection of quarks and gluons must then be hadronised into mesons and baryons. This is done differently in each of the event generators, but is described by a set of (fragmentation) parameters that must be adjusted to agree with experimental results. HERWIG looks for colour singlet collections of quarks and gluons with low invariant mass and groups them together; this set then turns into hadrons. PYTHIA splits gluons into quark-antiquark pairs and turns the resulting set of colour singlet quark-antiquark pairs into hadrons via a string model. ISAJET simply fragments each quark independently paying no attention to the colour flow.

The dominant cross-section at the LHC consists of events with no hard scattering. There is little detailed theoretical understanding of these minimum-bias events and the event generators must rely on data at current energies. These minimum-bias events are important at LHC, particularly at design luminosity, as they overlap interesting hard-scattering events such as the production of new particles. The generators use a different approach in this case. HERWIG uses a parametrization of data mainly from the CERN $p\bar{p}$ Collider. PYTHIA uses a mini-jet model where the jet cross-section is used at very low transverse momenta, i.e. the hard scattering process is extrapolated until it saturates the total cross-section. CMS has used the PYTHIA approach with dedicated modifications that agree with present data from Tevatron [62]. The model of the hadronic interactions implemented in the physics generator has a direct impact on physical observables such as jet multiplicity, their average transverse momentum, internal structure of the jets and their heavy flavour content. That was one of the reasons why, whenever possible, PYTHIA was used enabling a consistent set of signal and background simulations to be generated.

Table ?? presents the predicted cross-section values for the basic SM processes, as used in the simulations for PTDR. The cross-section values (at leading order) were calculated by using PYTHIA 6.327 with CTEQ5L (default PDF for PTDR) and with CTEQ6M PDFs. α_s at 1st (2nd) order is used with CTEQ5L (CTEQ6M) PDFs. For CTEQ6M the quoted errors are related to the uncertainties due to PDFs (see Subsection B.9).

C.2 General scheme of generator's usage in CMS

All event generators, included in CMS simulation software, can be separated into two groups.

The first group (HERWIG, HIJING, ISAJET, PYTHIA) provides the *full simulation* of events. The basic package explored in CMS is PYTHIA and only few specific processes were simulated with HERWIG or HIJING.

Pure schematically a data flow in PYTHIA and HERWIG is presented on Fig. C.1.

After initialization the package (HERWIG or PYTHIA) calls "hard process" routines (see "1" arrow lines on Fig. C.1). Then an information (the momenta of initial and final partons, the colours and KF-codes) is passed to package for further showering, hadronization, fragmentation and decays of the unstable particles.

However, all these "full event simulation" generators have very limited number of the hard process matrix elements (typically for $2 \rightarrow 2$ reaction at LO). Therefore, several special generators are used for simulation of many other LO processes. In fact, such packages generate the




Figure C.1: Pure schematically data flow in PYTHIA and HERWIG.

hard processes kinematic quantities, such as masses and momenta, the spin, the colour connection, and the flavour of initial- and final-state partons. Then such information is stored in the “Les Houches” format [778] (`/HEPEUP/` common block) and is passed to full event simulation package like PYTHIA or HERWIG (see thick “output” line on Fig. C.1).

Three generators, namely ALPGEN [152], CompHEP [47], and MadGraph [73, 482], are widely used for simulation of many processes, especially for generation of the hard processes with multi-jet final states. For example, the ALPGEN allows to generate $Q\bar{Q}$ pair production with up to 6 jets. Due to a complexity of the matrix elements, describing the multi-jet processes, and re-weighting procedure the generation of events is very CPU-time consuming. As a result, the information with kinematics is stored in the output files. (see “2” lines on Fig. C.1). Then, like in generic PYTHIA process, such information is passed to PYTHIA (see thick “output” line on Fig. C.1).

There are several “dedicated generators”, TopReX [35], StaGen [779], SignleTop, COSMIC [779], SIMUB [779], PHASE [779], PYQUEN [780, 781], HYDJET [782], EDDE. These generators are used for simulation of several specific process (see below for a short description of these codes). The information with hard processes kinematic quantities is stored in `/HEPEUP/` common block [778] and is passed to “full event simulation” package (see “3” lines on Fig. C.1).

After full simulation of event with PYTHIA or HERWIG the output information is stored in `/HEPEVT/` common block. In addition two *special functionality* codes provide a better description of photon radiation from a charge final particles (PHOTOS [28]) and τ^\pm -lepton decays (TAUOLA [146]). Typically, these codes read an information from `/HEPEVT/` common, perform simulation and then add generated information (new particles) into `/HEPEVT/` common block (see Fig. C.1).

C.3 CMKIN

Almost all generators available in CMS could be used with the CMKIN package [779]. Now the CMKIN is used for OSCAR and FAMOS detector simulation input. This software package provides a common interface between physics event generators and CMS detector sim-

ulation (see Fig. C.2). It also provides an environment to make physics plots of generated events. The CMKIN provides an interface to a number of physics generators like PYTHIA, ISAJET and HERWIG. It also offers the possibility to use different 'external generators' like ALPGEN [152], CompHEP [47], MadGraph [73, 482] and TopReX [35]. Cosmics simulation is available as well. Simple particle generation is also included, i.e. single and double particles as well as simple multi particle events. The interface is based on a common block HEPEVT - a HEP standard to store particle kinematics information for one event [62]. The /HEPEVT/ common block is converted to HBOOK n-tuples. The event output format follows the HEPEVT standard and additional information can be included by the user in the `MC_PARAM/`.




Figure C.2: Illustration of the CMKIN interface.

There is a unified compilation script which is used as follows:

```
kine_make_ntpl.com <generator> [lhpdf]
```

where the first parameter can have one of the following values: *pythia*, *herwig*, *isajet*, *simple*, *single*, *double*, *simplemulti*, *cosmic*, *comphep*, *alpgen*, *madgraph*, *phase*, *toprex* or *stagen*. The optional second parameter *lhpdf* is given when the user wants to use LHAPDF library [88].

The resulting executable is run with the corresponding command:

```
kine_make_ntpl.run <generator>
```

C.4 Full event simulation generators

C.4.1 PYTHIA

The PYTHIA package [62] is a general-purpose generator for hadronic events in pp , e^+e^- and ep colliders. It contains a subprocess library and generation machinery, initial- and final-state parton showers, underlying event, hadronization and decays, and analysis tools. PYTHIA contains around 240 different $2 \rightarrow 2$ (and some $2 \rightarrow 1$ or $2 \rightarrow 3$) subprocesses, all

at leading order. The subsequent decays of unstable resonances (W , Z , top, Higgs, SUSY, ...) brings up the partonic multiplicity, for many processes with full spin correlations in the decays. The external processes can be evolved through the showering and hadronization (like internal ones).

The final-state shower is based on forward evolution in terms of a decreasing timelike virtuality m^2 , with angular ordering imposed by veto. The framework is leading-log, but includes many NLL aspects such as energy-momentum conservation, $\alpha_s(p_{\perp}^2)$ and coherence. Further features include gluon polarization effects and photon emission.

The initial-state shower is based on backwards evolution, i.e. starting at the hard scattering and moving backwards in time to the shower initiators, in terms of a decreasing spacelike virtuality Q^2 . Initial and final showers are matched to each other by maximum emission cones.

The composite nature of hadrons (and resolved photons) allows for several partons from each of the incoming hadrons to undergo scatterings. Such multiple parton-parton interactions are instrumental in building up the activity in the underlying event, in everything from charged multiplicity distributions and long-range correlations to minijets and jet pedestals. The interactions are described by perturbation theory, approximated by a set of more or less separate $2 \rightarrow 2$ scatterings; energy conservation and other effects introduce (anti)correlations. The scatterings are colour-connected with each other and with the beam remnants.

The Lund string model, used for hadronization, is based on a picture with linear confinement, where (anti)quarks or other colour (anti)triplets are located at the ends of the string, and gluons are energy and momentum carrying kinks on the string. The string breaks by the production of new $q\bar{q}$ pairs, and a quark from one break can combine with an antiquark from an adjacent one to form a colour singlet meson.

Unstable particles are allowed to decay. In cases where better decay models are available elsewhere, e.g. for τ^{\pm} with spin information or for B hadrons, such decays can be delegated to specialized packages.

At present the parameters from almost all PYTHIA common blocks (see BLOCK DATA PYDATA) could be set via data cards. With the CMKIN these parameters could be set in data card file with the following format (note, that only capital letters should be used):

PYTHIA	CMKIN	comment
parameter		
MSEL = 6	MSEL 6	$t\bar{t}$ production
one- and two-dimensional arrays		
CKIN(1) = 100.	CKIN 1 = 100.	min. $\sqrt{\hat{s}}$
i.e. PMAS(6,1) = 178.	PMAS 6,1 = 178.	top-quark mass

- *Common cards for CMKIN*

Below we present the list of PYTHIA parameters used for full event simulation for PTDR. Some of these parameters correspond to the old multiple interactions scenario, namely *Rick Field's Tune A* [783].

MSTP (2) = 1 : 1(first)/2(second) order running α_s

MSTP (33) = 0 : do not include of K -factors in hard cross sections
 MSTP (51) = 7 : PDF set (here is CTEQ5L)
 MSTP (81) = 1 : multiple parton interactions is switched ON
 MSTP (82) = 4 : defines the multiple parton interactions model
 PARP (67) = 1. : amount of initial-state radiation
 PARP (82) = 1.9 : P_t cut-off for multiparton interactions
 PARP (83) = 0.5 : fraction of total hadronic matter in core
 PARP (84) = 0.4 : radius of core
 PARP (85) = 0.33 : gluon production mechanism in multiple interactions
 PARP (86) = 0.66 : gluon prod. mechanism in multiple interactions
 PARP (88) = 0.5
 PARP (89) = 1000. : reference energy scale for which PARP (82) is set
 PARP (90) = 0.16 : effective P_t cut-off = [PARP(82)/PARP(89)]**PARP(90)
 PARP (91) = 1.0 : width of Gaussian primordial k_{\perp} distribution inside hadron
 PARJ (71) = 10. : maximum average $c\tau$ for particles allowed to decay
 MSTJ (11) = 3 : choice of the fragmentation function
 MSTJ (22) = 2 : allow to decay those unstable particles
 PMAS (5, 1) = 4.8 : the mass of b -quark
 PMAS (6, 1) = 175.0 : the mass of t -quark

C.4.2 HERWIG

HERWIG contains a wide range of Standard Model, Higgs and supersymmetric processes [187]. HERWIG uses the parton-shower approach for initial- and final-state QCD radiation, including colour coherence effects and azimuthal correlations both within and between the jets.

In the treatment of supersymmetric processes, HERWIG itself doesn't calculate the SUSY mass spectrum or decay rates, but reads in an input file containing the low-energy parameters (masses, couplings, decays, ...). This file can be written by hand or more conveniently be generated with the ISAWIG program. This program provides an interface to ISAJET (and therefore to all models in ISASUSY and ISASUGRA), to HDECAY (for NLO Higgs decays), and can also add R-parity violating decays.

Colour coherence effects of (initial and final) partons are taken into account in all hard sub-processes, including the production and decay of heavy quarks and supersymmetric particles. HERWIG uses the angular ordered parton shower algorithm which resums both soft and collinear singularities. HERWIG includes spin correlation effects in the production and decay of top quarks, tau leptons and supersymmetric particles. For the SUSY decays, there is an option for using either the matrix elements (fast) or the full spin correlations. HERWIG uses a cluster hadronization model based on non-perturbative gluon splitting, and a similar

cluster model for soft and underlying hadronic events. This model gives a good agreement with the LEP data on event shapes, but doesn't fit the identified particle spectrum well.

C.4.3 ISAJET

ISAJET is a Monte Carlo program which simulates pp , $\bar{p}p$, e^+e^- interactions at high energies [653]. ISAJET is based on perturbative QCD plus phenomenological models for parton and beam jet fragmentation. At CMS ISAJET is used for calculations of SUSY parameters.

C.4.4 HIJING

It is expected that hard or semi-hard parton scatterings with transverse momentum of a few GeV are expected to dominate high energy heavy ion collisions. The HIJING (Heavy Ion Jet INteraction Generator) Monte Carlo model [776] was developed by M. Gyulassy and X.-N. Wang with special emphasis on the role of minijets in pp , pA and AA reactions at collider energies.

Detailed systematic comparison of HIJING results with a very wide range of data demonstrates that a quantitative understanding of the interplay between soft string dynamics and hard QCD interaction has been achieved. In particular, HIJING reproduces many inclusive spectra two particle correlations, and can explain the observed flavor and multiplicity dependence of the average transverse momentum.

C.5 Tree level matrix element generators

C.5.1 ALPGEN

ALPGEN is designed for the generation of Standard Model processes in hadronic collisions, with emphasis on final states with large jet multiplicities [152]. It is based on the exact leading order evaluation of partonic matrix elements and t and gauge boson decays with helicity correlations. The code generates events in both a weighted and unweighted mode. Weighted generation allows for high-statistics parton-level studies. Unweighted events can be processed in an independent run through shower evolution and hadronization programs.

The current available processes are:

- $W/Z/H Q\bar{Q} + N$ jets ($Q = c, b, t$) with $N \leq 4$
- $Q\bar{Q} + N$ jets, with $N \leq 6$
- $QQ'Q'\bar{Q}' + N$ jets, with $N \leq 4$
- $W + \text{charm} + N$ jets, with $N \leq 5$
- N jets, $W/Z + N$ jets, with $N \leq 6$
- $nW + mZ + lH + N$ jets, with $n + m + l + N \leq 8$, $N \leq 3$
- $N\gamma + M$ jets, with $N \geq 1$, $N + M \leq 8$ and $M \leq 6$
- $H + N$ jets ($N \leq 4$), with the Higgs produced via ggH vertex
- single top production.

C.5.2 CompHEP

CompHEP [784] is a package for evaluating Feynman diagrams, integrating over multi-particle phase space and generating events with a high level of automation.

CompHEP computes squared Feynman diagrams symbolically and then numerically calculates cross sections and distributions. After numerical computation one can generate with CompHEP the unweighted events with implemented colour flow information. The events are in the form of the Les Houches Accord event record [778] to be used in the PYTHIA program for showering and hadronization.

CompHEP allows for the computation of scattering processes with up to 6 particles and decay processes with up to 7 particles in the final state. CompHEP is a the tree level program. In addition it is possible an inclusion of partial (approximate) NLO corrections: NLO tree level $2 \rightarrow N + 1$ real emission corrections to the $2 \rightarrow N$ process, NLO structure functions, loop relations between parameters, known K -factors, and the known loop contributions as effective vertices.

C.5.3 MadGraph and MadEvent

MadEvent [73] is a multi-purpose, tree-level event generator which is powered by the matrix element generator MadGraph [482]. Given a user process, MadGraph automatically generates the amplitudes for all the relevant subprocesses and produces the mappings for the integration over the phase space. This process-dependent information is packaged into MadEvent, and a stand-alone code is produced. It allows the user to calculate cross sections and to obtain unweighted events automatically. Once the events have been generated – event information, (e.g. particle id's, momenta, spin, color connections) is stored in the “Les Houches” format [778]. Events may be passed directly to a shower Monte Carlo program (interfaces are available for HERWIG and PYTHIA).

Limitations of the code are related to the maximum number of final state QCD particles. Currently, the package is limited to ten thousand diagrams per subprocess. So, for example, $W+5$ jets which has been calculated, is close to its practical limit. At present, only the Standard Model Feynman rules are implemented and the user has to provide his/her own rules for beyond Standard Model physics, such as MSSM.

C.5.4 TopReX

The event generator TopReX [35] provides the simulation of several important processes in pp and $p\bar{p}$ collisions, not implemented in PYTHIA. In the matrix elements used in TopReX the decays of the final t -quarks, W^\pm , Z and charged Higgs bosons are also included. The final top quark could decay into SM channel ($t \rightarrow qW^+$, $q = d, s, b$), b -quark and charged Higgs ($t \rightarrow bH^+$) and the channels with flavor changing neutral current (FCNC): $t \rightarrow u(c) V$, $V = g, \gamma, Z$. The implemented matrix elements take into account spin polarizations of the top quark, that provides a correct description of the differential distributions and correlations of the top quarks decay products.

C.6 Supplementary packages

C.6.1 PHOTOS

PHOTOS is a universal package to simulate QED photon radiative corrections in decays [28]. The precision of the generation may in some cases be limited, in general it is not worse than the complete double bremsstrahlung in LL approximation. The infrared limit of the distributions is also correctly reproduced. The action of the algorithm consists of generating, with internally calculated probability, bremsstrahlung photon(s), which are later added to the `/HEPEVT/` record. Kinematic configurations are appropriately modified. Energy-momentum conservation is assured. When using PHOTOS, the QED bremsstrahlung of the principal generator must be switched off. For example in case of PYTHIA one has to use `MSTJ 41=1`.

C.6.2 TAUOLA


TAUOLA is a package for simulation of the τ^\pm -lepton decays [146]. It uses the PHOTOS package to simulate radiative corrections in the decay. The TAUOLA interface is made with the the PYTHIA generator [779] This interface evaluates also the position of τ -lepton decay (i.e. the information on the production vertex of the decay products of τ -lepton).

C.6.3 PYQUEN

The event generator PYQUEN (PYthia QUENched) [780, 781] provides the simulation of rescattering and energy loss of hard partons in dense QCD-matter (quark-gluon plasma) created in ultrarelativistic heavy ion collisions. The approach relies on an accumulative energy losses, when gluon radiation is associated with each scattering in expanding medium together including the interference effect by the modified radiation spectrum $dE/d\ell$ as a function of decreasing temperature T . The model is realized as fast Monte-Carlo tool implemented to modify standard PYTHIA jet event.

C.6.4 HYDJET

The event generator HYDJET [782] (HYDroynamics + JETs) provides the fast simulation of heavy ion events at LHC energy including longitudinal, transverse and elliptic flow effects together with jet production and jet quenching (rescattering and energy loss of hard partons in dense QCD-matter, quark-gluon plasma). The model merges a fast generator of flow effects HYDRO [785] with PYTHIA (for jet production) and PYQUEN [780, 781] (for jet quenching) by simulating full heavy ion event as a superposition of soft, hydro-type state and hard multi-jets.

First of all, HYDJET calculates the number N^{hard} of hard nucleon-nucleon sub-collisions and number N^{Part} nucleons-participants (at given impact parameter b of AA collision and minimum P_T of hard parton scattering) and generates the initial parton spectra by calling PYTHIA N^{hard} times (fragmentation off). After each jet parton affected by medium-induced rescattering and energy loss according with PYQUEN model. In the end of each PYTHIA sub-event adding new (in-medium emitted) gluons into PYTHIA parton list and rearrangements of partons to update string formation are performed. Then PYQUEN forms final hadrons with PYEXEC subroutine (fragmentation on). Finally, HYDJET calculates the multiplicity of soft, hydro-induced part of the event and add new particles in the end of the event record. 

Appendix D

GARCON: Genetic Algorithm for Rectangular Cuts Optimization

Typically HEP analysis has quite a few selection criteria (cuts) to optimize for example a significance of the “signal” over “background” events: transverse energy/momenta cuts, missing transverse energy, angular correlations, isolation and impact parameters, etc. In such cases simple scan over multi-dimensional cuts space (especially when done on top of a scan over theoretical predictions parameters space like for SUSY e.g.) leads to CPU time demand varying from days to many years... One of the alternative methods, which solves the issue is to employ a Genetic Algorithm (GA), see e.g. [786, 787, 788].

We wrote a code, GARCON [55], which automatically performs an optimization and results stability verification effectively trying $\sim 10^{50}$ cut set parameters/values permutations for millions of input events in hours time. Examples of analyses are presented in the Physics TDR, see for example sub-sections 3.1, 8.4, 13.6, 13.7, 13.14 and recent papers [42, 789, 790, 307].

The GARCON program among many other features allows user:

- to select an optimization function among known significance estimators, as well as to define user’s own formula, which may be as simple as signal to background ratio, or a complicated one including different systematic uncertainties separately on different signal and background processes, different weights per event and so on;
- to define a precision of the optimization;
- to restrict the optimization using different kind of requirements, such us minimum number of signal/background events to survive after final cuts, variables/processes to be used for a particular optimization run, number of optimizations inside one run to ensure that optimization converges/finds not just a local maximum(s), but a global one as well (in case of a complicated phase space);
- to automatically verify results stability.

GARCON, GA-based programs in general exploit evolution-kind algorithms and uses evolution-like terms:


- Individual is a set of qualities, which are to be optimized in a particular environment or set of requirements. In HEP analysis case Individual is a set of lower and upper rectangular cut values for each of variables under study/optimization.
- Environment or set of requirements of evolutionary process in HEP analysis case

is a Quality Function (QF) used for optimization of individuals. The better QF value the better is an Individual. Quality Function may be as simple as S/\sqrt{B} , where S is a number of signal events and B is a total number of background events after cuts, or almost of any degree of complexity, including systematic uncertainties on different backgrounds, etc.

- A given number of individuals constitute a Community, which is involved in evolution process.
- Each individual involved in the evolution: breeding with possibility of mutation of new individuals, death, etc. The higher is the QF of a particular individual, the more chances this individual has to participate in breeding of new individuals and the longer it lives (participates in more breeding cycles, etc.), thus improving community as a whole.
- Breeding in HEP analysis example is a producing of a new individual with qualities (set of min/max cut values) taken in a defined way from two "parent" individuals.
- Death of an individual happens, when it passes over an age limit for it's quality: the bigger it's quality, the more it lives.
- Cataclysmic Updates may happen in evolution after a long period of stagnation in evolution, at this time the whole community gets renewed and gets another chance to evolve to even better quality level. In HEP analysis case it corresponds to a chance to find another local and ultimately a global maximum in terms of quality function. Obviously, the more complicated phase space of cut variables is used the more chances exist that there are several local maximums in quality function optimization.
- There are some other algorithms involved into GAs. For example mutation of a new individual. In this case newly "born" individual has not just qualities of its "parents", but also some variations, which in terms of HEP analysis example helps evolution to find a global maximum, with less chances to fall into a local one. There are also random creation mechanisms serving the same purpose.

There is nothing special involved in GARCON input preparation. One would need to prepare a set of arrays for each background and a signal process of cut variable values for optimization. Similar to what is needed to have to perform a classical eye-balling cut optimization.

In comparison to other automatized optimization methods GARCON output is transparent to user: it just says what rectangular cut values are optimal and recommended in an analysis. Interpretation of these cut values is absolutely the same as with eye-balling cuts when one selects a set of rectangular cut values for each variable in a "classical" way by eye.

All-in-all it is a simple yet powerful ready-to-use tool with flexible and transparent optimization and verification parameters setup. It is publicly available  along with a paper on it [55] consisting of an example case study and user's manual.

Appendix E

Online Selection



E.1 Introduction

The CMS trigger menu depends upon the luminosity delivered by the LHC and the available bandwidth between and out of the systems. The LHC luminosity is expected to start at $\mathcal{L} = 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$ in 2007 and gradually rise to $\mathcal{L} = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ by 2010. The CMS data acquisition can be operated with one to eight slices of Event Filter Farms that execute High-Level Trigger (HLT) algorithms. It is expected that we start with one slice in 2007, allowing a bandwidth of 12.5 kHz between L1 and HLT, and build up to the full eight slices by 2010, when the L1 to HLT bandwidth can be raised to 100 kHz. It is assumed that the data logging capability after the HLT selection will remain constant at a rate between 100 Hz to 150 Hz*. The L1 and HLT algorithms will be configured to operate with the lowest possible thresholds making the best use of the available bandwidth.

Here we focus solely on trigger studies for $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. The scenario of operation assumes that CMS uses four DAQ slices capable of 50 kHz. While the actual choice of trigger thresholds, especially at HLT, depends strongly upon the physics of interest at the time of operation, we propose here an example set of trigger menus within the constraints of the data acquisition system. An effort has been made to optimize the L1 and HLT thresholds coherently, taking into account possible bandwidth limitations.

The structure of this note is as follows: first we overview the object-identification algorithms used for these studies. The emphasis is given to the changes that have been introduced since a similar study was performed in the DAQ TDR [68]. We then introduce a series of new trigger paths, aiming at increasing the event yield for various physics analyses. The central idea is to exploit various multi-object (or *cross-channel*) triggers in an attempt to improve the rejection and, at the same time, lower the kinematic thresholds of the corresponding objects. We finally present the performance of the triggers, and we calculate the overlap among them the total HLT output rate.

E.2 Description of trigger tools

*At the time of the writing of this document, several scenarios for the HLT output rate, the disk requirements for the storage manager and the associated cost are under discussion.

E.2.1 L1 reconstruction

There have been no significant changes in the L1 algorithms since the DAQ TDR. We have introduced an H_T algorithm which sums the corrected jet E_T of all the jets found above a programmable threshold, within $|\eta| < 5$. It does not account for E_T carried by muons and neutrinos.

The L1 strategy is the following: We have made an effort to keep the thresholds at the same levels, or even reduce them in order to be able to study cross-channel triggers (typically appearing with lower kinematic cuts). The notable exception is the tau triggers, where an increase in the HCAL noise and the usage of a new pile-up model in the simulation do affect the L1 τ identification tools, and therefore the related trigger rates. We have introduced additional L1 conditions for all HLT paths. The determination of thresholds and prescales is a compromise between the desire to distribute reasonably the available L1 bandwidth to the various triggers, and the need to optimize the L1 and HLT thresholds coherently in well-defined trigger paths.

E.2.2 HLT reconstruction

Well defined L1 terms are used in order to obtain triggers whose behavior and efficiency can be studied with real data. We have replaced some of the L1 conditions with respect to the DAQ TDR with new L1 terms when this leads to more reasonable trigger paths or triggers that are more stable and carry less of a bias. The optimization of the thresholds for the various triggers has been a compromise between the physics needs of the CMS experiment and the total HLT rate available. This study serves only as an intermediate step in a long-term trigger study project. Further improvements in the reconstruction tools, better optimization of the thresholds, implementation of additional triggers and a CMS-wide discussion of the allocation of the HLT bandwidth to the physics groups according to the priorities of the experiment, are foreseen.

A general and detailed description of the HLT system can be found in Ref. [68]. Here we summarize the recent modifications of the HLT tools, and the expected changes in the rates of the various triggers with respect to the earlier studies.

- Muons: The muon algorithm has not changed, with the exception of the drift-tube local reconstruction and segment building. Therefore, no significant changes in the rates of single- and di-muon trigger paths are expected. The option of constructing muon triggers without isolation has been added.
- Electrons-Photons: Here the most important change is that all saturated trigger towers at L1 are now considered isolated. This increases both the signal efficiency and the background. At HLT, the photon rate can be reduced by increasing the thresholds or by applying some isolation cuts. For the electrons the options include a matching with pixel lines and tracks, as well as isolation requirements in the hadron calorimeter and the tracker. A study of the algorithm optimization can be found in Ref. [7]. An improvement of the rejection power of the electron-photon algorithms is achieved with a simultaneous decrease of the HLT thresholds. Similar enhancements are expected for cross-channel triggers where one of the objects under consideration is an electron or a photon.
- Jets and E_T^{miss} : The main jet-finder algorithm (Iterative Cone with $R = 0.5$) has not been modified. Some optimizations of the tower thresholds have been added,

and the jet corrections have been updated (“Scheme C”). Similarly, there are no major algorithm changes for E_T^{miss} , however it has been ensured that all triggers including a E_T^{miss} object do not have any off-line corrections applied. Another improvement that has been recently introduced is the ability to construct *acoplanar* triggers by combining two jets, or a jet and a E_T^{miss} object that do not lie “back-to-back”. Details of the physics algorithms can be found in Refs. [156] and [142].

- ***b*-jets:** The algorithm now uses muon information for fast rejection. Further improvements have been made for faster decisions and for an increased efficiency in fully hadronic final states. The documentation for the *b*-jet HLT algorithm can be found in Ref. [279].
- ***Taus:*** The HLT τ algorithm has not changed. However, the increase in the L1 rate does propagate into the HLT. The isolation parameters for the electromagnetic calorimeter and the tracker have been tuned after recent studies performed by the Higgs group, described in Ref. [268]. The overall rate for τ -related triggers is expected to be slightly increased.

A new addition to the HLT reconstruction tools is the H_T algorithm. It sums the corrected jet E_T of all the $E_T > 5$ GeV jets found within $|\eta| < 5$, along with the energy of the $p_T > 5$ GeV/c HLT muons found in the event, and the E_T^{miss} computed using the calorimeter deposits. It is meant to be driven off the corresponding L1 H_T term.

E.3 Triggering with Forward Detectors

E.3.1 Objective

We discuss [†] the feasibility of a special forward detectors trigger stream, with target output rate of $\mathcal{O}(1)$ kHz at L1 and $\mathcal{O}(1)$ Hz on the HLT, as well as the potential of the already foreseen CMS L1 trigger streams for retaining events with diffractive processes.

The proposed forward detectors trigger stream combines the information of the central CMS detector with that from detectors further downstream of the CMS IP. The forward detectors considered are the TOTEM T1 and T2 tracker telescopes as well as the TOTEM Roman Pot (RP) detectors up to 220 m downstream of CMS [791, 792]. Information from TOTEM will be available to the CMS L1 trigger. We also consider detectors at a distance of 420 m, in the cryogenic region of the LHC ring, currently being studied by the FP420 project [242].

Topologically, diffractive events are characterized by a gap in the rapidity distribution of final-state hadrons. In addition, the fractional momentum loss, ξ , of diffractively scattered protons peaks at $\xi = 0$ (“diffractive peak”). The TOTEM RP detectors will permit to measure protons in the region $0.2 > \xi > 0.02$. Detectors at a distance of 420 m from the IP would provide a coverage of $0.02 > \xi > 0.002$, complementary to that of the TOTEM detectors, but cannot be included in the L1 trigger without an increase in the L1 latency of $3.2 \mu\text{s}$ (though a special, long latency running mode might be feasible at lower luminosities).

The studies discussed in the following assume that the RP detectors are 100% efficient in detecting all particles that emerge at a distance of at least $10 \sigma_{\text{beam}} + 0.5$ mm from the beam axis (1.3 mm at 220 m, 4 mm at 420 m). Their acceptance was calculated for the nominal

[†]These studies were carried out in collaboration with TOTEM.

LHC optics ($\beta^* = 0.55$ m), version V6.5 [793, 794], and by way of a simulation program that tracks particles through the accelerator lattice [795]. LHC bunches with 25 ns spacing were assumed.

The results presented below do not depend on the specific hardware implementation of the TOTEM T1, T2 and RP detectors; they hold for any tracker system with the T1, T2 η coverage in conjunction with RPs at 220 m from the IP.

E.3.2 Level-1 Trigger Rates for Forward Detectors Trigger Stream

E.3.2.1 2-Jet Conditions

A particularly interesting and challenging diffractive channel is the central exclusive production of a Higgs Boson, $pp \rightarrow pHp$, with Higgs mass close to the current exclusion limit. The dominant decay of a SM Higgs Boson of mass ~ 120 GeV/ c^2 is into two b -quarks and generates 2 jets with at most 60 GeV/ c transverse momentum each. In order to retain as large a signal fraction as possible, as low an E_T threshold as possible of the L1 2-jet trigger is desirable. In practice, the threshold value cannot be chosen much lower than 40 GeV per jet. The L1 trigger applies cuts on the calibrated E_T value of the jet. Thus, a threshold of 40 GeV corresponds to 20-25 GeV in reconstructed E_T , i.e. to values where noise starts becoming sizable.

For luminosities of 10^{32} cm $^{-2}$ s $^{-1}$ and above, the L1 rate from standard QCD processes for events with at least 2 central jets ($|\eta| < 2.5$) with $E_T > 40$ GeV exceeds by far the target output rate of $\mathcal{O}(1)$ kHz. Thus additional conditions need to be employed to reduce the rate from QCD processes. The efficacy of several conditions was investigated [796, 797, 239, 798, 799]. In the following, the corresponding rate reduction factors are always quoted with respect to the rate of QCD events that contain at least 2 central jets with $E_T > 40$ GeV per jet.

The QCD background events were generated with the Pythia Monte Carlo generator. In order to assess the effect when the signal is overlaid with pile-up, a sample of 500,000 pile-up events was generated with Pythia. This sample includes inelastic as well as elastic and single diffractive events. Pythia underestimates the number of final state protons in this sample. The correction to the Pythia leading proton spectrum described in [800] was used to obtain the results discussed in the following.

Given a L1 target rate for events with 2 central L1 jets of $\mathcal{O}(1)$ kHz, a total rate reduction between a factor 20 at 1×10^{33} cm $^{-2}$ s $^{-1}$ and 200 at 1×10^{34} cm $^{-2}$ s $^{-1}$ is necessary. Table E.1 summarizes the situation for luminosities between 10^{32} cm $^{-2}$ s $^{-1}$ and 10^{34} cm $^{-2}$ s $^{-1}$, and for different RP detector conditions: a track at 220 m on one side of the IP (single-arm 220 m), without and with a cut on ξ ; a track at 420 m on one side of the IP (single-arm 420 m); a track at 220 m and 420 m (asymmetric); a track at 420 m on both sides of the IP (double-arm 420 m). Because the detectors at 220 m and 420 m have complementary coverage in ξ , the asymmetric condition in effect selects events with two tracks of very different ξ value, in which one track is seen at 220 m on one side of the IP and a second track is seen on the other side at 420 m. If not by the L1 trigger, these asymmetric events can be selected by the HLT and are thus of highest interest. At luminosities where pile-up is present, the rate reduction achievable with the RP detector conditions decreases because of the diffractive component in the pile-up.

Table E.1: Reduction of the rate from standard QCD processes for events with at least 2 central L1 jets with $E_T > 40$ GeV, achievable with requirements on the tracks seen in the RP detectors. Additional rate reductions can be achieved with the H_T condition and with a topological condition. Each of them yields, for all luminosities listed, an additional reduction by about a factor 2.

Lumi nosity [$\text{cm}^{-2} \text{s}^{-1}$]	Pile-up events per BX	L1 2-jet rate [kHz] for $E_T >$ 40GeV	Total reduc tion needed	Reduction when requiring track in RPs at						
				220 m $\xi < 0.1$	420 m	220 & 420 m (asymmetric) $\xi < 0.1$		420 & 420 m		
1×10^{32}	0	2.6	2	370						
1×10^{33}	3.5	26	20	7	15	27	160	380	500	
2×10^{33}	7	52	40	4	10	14	80	190	150	
5×10^{33}	17.5	130	100	3	5	6	32	75	30	
1×10^{34}	35	260	200	2	3	4	17	39	10	

A collimator located in front of the LHC magnet Q5, planned to be operative at higher luminosities, will have an effect on the acceptance of the RP detectors resembling that of a ξ cut. This effect has not been taken into account in Table E.1.

Using T1 and T2 as vetoes in events with 2 central L1 jets was found to be effective only in the absence of pile-up [801].

In addition to the E_T values of individual L1 jets, the CMS Calorimeter Trigger has at its disposal the scalar sum, H_T , of the E_T values of all jets. Requiring that essentially all the E_T be concentrated in the two central L1 jets with highest E_T , i.e. $[E_T^1 + E_T^2]/H_T > 0.9$ (H_T condition), corresponds to imposing a rapidity gap of at least 2.5 units with respect to the beam direction. This condition reduces the rate of QCD events by approximately a factor 2, independent of the presence of pile-up and with only a small effect on the signal efficiency.

A further reduction of the QCD rate could be achieved with the help of a topological condition. The 2-jet system has to balance the total momentum component of the two protons along the beam axis. In signal events with asymmetric ξ values, the proton seen on one side in the RP detectors at 220 m distance is the one with the larger ξ and thus has lost more of its initial momentum component along the beam axis. Hence the jets tend to be located in the same η -hemisphere as the RP detectors that detect this proton. A trigger condition requiring that $[\eta^{jet1} + \eta^{jet2}] \times \text{sign}(\eta^{220m RP}) > 0$ reduces the QCD background by a factor 2, independent of pile-up, and with no loss in signal efficiency.

A reduction of the QCD rate to levels compatible with a L1 output target rate of $\mathcal{O}(1)$ kHz by including RP detectors at a distance of 220 m from the CMS IP thus appears feasible for luminosities up to $2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$, as long as a ξ cut can be administered in the L1 trigger.

E.3.2.2 Other Conditions

The effect of combining already foreseen L1 trigger conditions with conditions on the RP detectors is illustrated in Table E.2 [798]. Single- and double-arm RP detector conditions are indicated with 's' and 'd' endings, respectively. Entries marked with a '(c)' indicate thresholds applicable if a cut on $\xi < 0.1$ is implemented for the RP detectors at 220 m. The jet

conditions consider all L1 jets with $|\eta| < 5$.

Table E.2: Estimated threshold values that result in a L1 output rate of ~ 1 kHz, for various conditions on central CMS detector quantities and on tracks seen in the RP detectors at 220 m and 420 m.

L1 condition	L1 E_T or p_T threshold [GeV] at $\mathcal{O}(1)$ kHz L1 output rate for luminosity [$\text{cm}^{-2} \text{s}^{-1}$]			
	1×10^{33}	2×10^{33}	5×10^{33}	1×10^{34}
1 Jet	115	135	160	190
2 Jet	90	105	130	150
1 Jet+220s	90	115	155	190
2 Jet+220s	65	90	125	150
1 Jet+220d	55	85	130	175
2 Jet+220d	30	60	100	140
1 Jet+220s(c)	70	90	150	185
2 Jet+220s(c)	60	70	115	145
1 Jet+220d(c)	30	65	110	155
2 Jet+220d(c)	20	45	85	125
1 Jet+420s	65	90	125	165
2 Jet+420s	45	70	100	130
1 Jet+420d	20	40	80	115
2 Jet+420d	< 10	30	60	90
1 μ +220s	12	16	23	> 100
1 μ +220d	4	9	17	80
1 μ +220s(c)	–	11	22	100
1 μ +220d(c)	–	6	13	30
1 μ +420s	7	11	14	37
1 μ +420d	< 2	4	7	14

A further rate reduction by approximately a factor two can be obtained at luminosities with negligible pile-up by imposing a rough large rapidity gap cut at L1. This was implemented by requiring that there be no forward jets, i.e. jets in the HF, in either hemisphere in the event.

E.3.3 Level-1 Signal Efficiencies

Of the L1 conditions discussed so far, only those based on the RP detectors have a significant impact on the signal efficiency. Of further interest is the question how many signal events are being retained by the already foreseen trigger streams, notably the muon trigger.

E.3.3.1 Central Exclusive Higgs Production ($H(120 \text{ GeV}/c^2) \rightarrow b\bar{b}$)

In order to study the effect of the L1 trigger selection on the Higgs signal, signal samples of 100,000 events with central exclusive production of a Higgs Boson were generated with the Monte Carlo programs EDDE [249] (version 1.1) and Exhume [247] (version 1.0).

Figure E.1 shows the L1 selection efficiency as a function of the E_T threshold values when at least 2 central L1 jets with E_T above threshold are required [798]. For a threshold of 40 GeV




Figure E.1: L1 selection efficiency for $pp \rightarrow pHp$ and $H(120 \text{ GeV}/c^2) \rightarrow b\bar{b}$ as function of the E_T threshold value when at least 2 central L1 jets with E_T above threshold are required. All plots are for the non-pile-up case and the H_T condition has been applied. Left: Comparison between the EDDE and Exhume Monte Carlo generators, without applying any additional RP conditions. Right: Comparison of the effect of different RP conditions on the efficiency in the Exhume Monte Carlo sample.

per jet, Exhume and EDDE both yield an efficiency of about 20%. The plot on the right-hand side overlays the efficiency curves obtained with Exhume when the 2-jet condition is combined with RP detector conditions. With an E_T threshold of 40 GeV per jet, the single-arm 220 m (420 m) condition results in an efficiency of the order 12% (15%), the double-arm 420 m condition in one of 8% and the asymmetric condition in one of 6%. This also means that, even without the possibility of including the RP detectors at 420 m from the CMS IP in the L1 trigger, 6% of the signal events can be triggered on with the single-arm 220 m condition, but will have a track also in the 420 m detectors that can be used in the HLT.

An alternative trigger strategy is to exploit the relatively muon-rich final state from B -decays: about 20% of the events have at least a muon in the final state. Requiring at least one (two) L1 muon(s) with p_T above 14 GeV (3 GeV) yields an efficiency of 6% (2%). Demanding at least 1 muon and 1 jet, the latter with $E_T > 40$ GeV, is a condition not yet foreseen in the CMS trigger tables. For a muon p_T threshold of 3 GeV, the rate at a luminosity of 10^{33} cm^{-2} is slightly less than 3 kHz, and about half of the decays with muons in the final state (i.e. 9 %) are retained [799].

E.3.3.2 Central Exclusive Higgs Production ($H(140 \text{ GeV}/c^2) \rightarrow WW$)

For SM Higgs Boson masses above $120 \text{ GeV}/c^2$, the $H \rightarrow WW$ branching ratio becomes sizable; in this case the final state contains high- p_T leptons that can be used for triggering. Efficiencies are in general high [799]. About 23% of the events have at least one muon in the final state. Approximately 70% of these (i.e. 16%) are retained by requiring at least one muon with a p_T threshold of 14 GeV. An extra $\approx 10\%$ (i.e. 2%) would be retained by implementing the muon/jet slot discussed above with thresholds of 3 GeV on the muon p_T and 40 GeV on the jet E_T .

E.3.3.3 Single diffractive hard processes

Double-Pomeron exchange processes constitute only a small part of the diffractive cross section. Hard single-diffraction, $pp \rightarrow pX$, where only one proton remains intact and the other is diffractively excited, have much higher cross sections than hard double-Pomeron exchange events. Efficiencies have been studied for $pp \rightarrow pX$, with X containing a W or a Z boson that decay to jets and to muons, as well as with X containing a dijet system. Samples of 100,000 signal events each were generated with the Pomwig Monte Carlo generator [802] (version 1.3).

For two example processes, Figure E.2 shows the efficiency as a function of the L1 threshold value, normalized to the number of events where for the diffractively scattered proton $0.001 < \xi < 0.2$ holds [798]. Three different trigger conditions are considered: trigger on central detector quantities alone (i), trigger on central detector quantities in conjunction (ii) with the single-arm 220 m condition, and (iii) with the single-arm 420 m condition. Also shown is the number of events expected to pass the L1 selection per pb^{-1} of LHC running. A significant part of events is retained when a proton is required in the 220 m RPs.




Figure E.2: L1 selection efficiency as function of the E_T threshold value for $pp \rightarrow pWX$ (left) and $pp \rightarrow pjX$ (right), when at least one (left) or two (right) L1 jets ($|\eta| < 5$) above threshold are required. All plots are for the non-pile-up case.

E.3.4 Effect of pile-up, beam-halo and beam-gas backgrounds

Pile-up effects are included in all rate and efficiency studies presented. In the 220 m stations, 0.055 protons/pile-up event are expected on average, in the 420 m stations, 0.012 protons/pile-up event. At a luminosity of $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$, there are 35 pile-up events on average; this entails, on average, 2 extra tracks in the 220 m stations and less than one in the 420 m stations.

The effect from beam-halo and beam-gas events on the L1 rate is not yet included in the studies discussed here. Preliminary estimates suggest that they are chiefly a concern for any trigger condition based solely on the forward detectors. For any trigger condition that includes a requirement on central CMS detector quantities the size of their contribution is such that they do not lead to a significant increase of the L1 output rate.

E.3.5 HLT strategies

Jets are reconstructed at the HLT with an iterative cone ($R < 0.5$) algorithm. The L1 selection cuts are repeated with HLT quantities. The following conditions are imposed [798]:

- A The event pass the single-arm 220 m L1 condition with $\xi < 0.1$ cut. As demonstrated in Table E.1, this condition reduces the L1 output rate to below $\mathcal{O}(1)$ kHz. Additional rate reduction factors of ~ 300 (~ 1000) at $1(2) \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ are needed to reach the HLT target output rate of $\mathcal{O}(1)$ Hz.
- B The two jets are back-to-back in the azimuthal angle ϕ ($2.8 < \Delta\phi < 3.48$ rad), and have $(E_T^1 - E_T^2)/(E_T^1 + E_T^2) < 0.4$, and $E_T > 40$ GeV for each jet.
- C The proton fractional momentum loss ξ is evaluated with the help of calorimeter quantities [803, 804, 805]:

$$\xi_{+-} = (1/\sqrt{s})\sum_i E_{Ti} \exp(\mp\eta_i), \quad (\text{E.1})$$

where the sum runs over the two jets and the $+$, $-$ signs denote the two hemispheres. The result is compared with the ξ value measured by the RP detectors. At present, no simulation of the RP reconstruction is available. As estimate of the ξ resolution, 15% (10%) is assumed at 220 m (420 m) [?]. Events are rejected if the difference between the two values of ξ is larger than 2σ .

- D At least one of the two jets is b -tagged.
- E A proton is seen at 420 m.

The case without pile-up presents no difficulty: essentially no QCD background events survive the selection. If conditions A+B+C are applied, the signal efficiency for $pp \rightarrow pHp$ with $H(120 \text{ GeV}/c^2) \rightarrow b\bar{b}$ is at 11% essentially unchanged with respect to the L1 selection, but the HLT output rate exceeds the target output rate, see Table E.3. If b -tagging is required but no ξ matching (conditions A+B+D), the efficiency drops to 7%, without any improvement in the rate reduction. The combination of conditions A+B+C+E finally leads to the targeted HLT output rate of $\mathcal{O}(1)$ Hz, without any loss in signal efficiency compared to L1.

Table E.3: Results of HLT selection.

HLT selection condition	A+B+C	A+B+D	A+B+C+E
HLT rate at $1 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$	15 Hz	20 Hz	< 1 Hz
line HLT rate at $2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$	60 Hz	80 Hz	1 Hz
e Signal eff. $H(120) \text{ GeV}/c^2 \rightarrow b\bar{b}$	11%	7%	6%



E.4 High-Level Trigger paths

We are starting with the DAQ-TDR trigger table as the baseline. This includes single- and double-triggers for the basic objects (e , γ , μ , τ) along with jets and b -jets. Some cross-channel triggers are also present. We are expanding the cross-channel “menu” by introducing additional triggers. We introduce an H_T algorithm, which we combine with other objects. We are also adding a series of central single-jets, non-isolated muons, and a diffractive trigger discussed earlier.

E.4.1 L1 conditions

Table E.4 summarizes the L1 conditions used to drive all the trigger paths. A pseudo “L1 bit number” has been assigned for easy reference in the following sections.

L1 bit #	Trigger	Thresholds (GeV)	Prescale
0	Single μ	14	1
1	Double μ	3	1
2	Single isolated $e\gamma$	23	1
3	Double isolated $e\gamma$	11	1
4	Double $e\gamma$ (isolated & non-isolated)	11	1
8	Single central jet	177	1
9	Single forward jet	177	1
10	Single τ -jet	100	1
11	2 central jets	130	1
12	2 forward jets	130	1
13	2 τ -jets	66	1
14	3 central jets	86	1
15	3 forward jets	86	1
16	3 τ -jets	40	1
17	4 central jets	70	1
18	4 forward jets	70	1
19	4 τ -jets	30	1
26	(isolated) $e\gamma + \tau$	14, 52	1
31	H_T	300	1
32	E_T^{miss}	60	1
33	Single jet (central, forward or τ)	140	10
34	Single jet (central, forward or τ)	60	1 000
35	Single jet (central, forward or τ)	20	100 000
36	Single jet (central, forward or τ)	150	1
37	2 jets (central, forward or τ)	100	1
38	3 jets (central, forward or τ)	70	1
39	4 jets (central, forward or τ)	50	1

Table E.4: L1 conditions used in High Level Trigger paths.

E.4.2 Evolution of DAQ-TDR triggers

The trigger paths that have been studied in Ref. [68] have been inherited and constitute the “bulk” of this next iteration of the CMS Trigger Menu for $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. Modifications (optimization of isolation cuts and thresholds) have been made for certain of the triggers, to reflect changes in the physics algorithms, or the improved understanding of the background from Monte Carlo (MC) simulations. The proposed Trigger Tables includes:

- **Muons:** The standard muon triggers include calorimeter-based isolation at L2, and both calorimeter and tracker isolation at L3. The p_T thresholds remain at 19 GeV/ c for the single-muon and (7, 7) GeV/ c for the di-muon trigger. A sec-

ond set of relaxed single- and double-muons has been added with $p_T > 37 \text{ GeV}$ and $p_T > 10 \text{ GeV}$, respectively. The main motivation here is Drell-Yan studies. In general, physics analyses that do not need a low p_T muon but do suffer from the isolation requirement on the muon. The reduced rejection caused by the removal of the isolation cuts is compensated by the higher- p_T thresholds on the muons, without affecting the event yield for the physics signal. The relaxed triggers have the advantage that the muons here are immune to radiative losses for the higher energy spectrum ($p_T > 500 \text{ GeV}/c$). Both isolated and relaxed triggers run off the corresponding non-isolated single- and double-muon bits at L1.

- Electrons: The p_T threshold remains at $26 \text{ GeV}/c$ for the single electron trigger and has a new value of $(12, 12) \text{ GeV}/c$ for the di-electron trigger. An additional relaxed di-electron trigger appears with $p_T > 19 \text{ GeV}/c$. The single-electron and double-electron triggers run off the corresponding L1 bits.
- Photons: The new p_T thresholds are $80 \text{ GeV}/c$ for the single-photon trigger and $(30, 20) \text{ GeV}/c$ for the di-photon trigger (both relaxed and non-relaxed flavors). A few prescaled single- and double-photon triggers have also been introduced, for the purpose of studying trigger efficiencies. The photon HLT algorithms run off the corresponding L1 $e\gamma$ bits (single- and double-triggers).
- Taus: The single- τ trigger runs off the corresponding L1 bit. The double- τ trigger is driven by the .OR.-ing of the single- and double- τ trigger bits at L1. There is no explicit kinematic cut on the tau at HLT. There is, however, a match-to-track requirement in addition to the $p_T > 100 (66) \text{ GeV}/c$ L1 precondition for the inclusive (double) tau trigger. The single- τ has also a $E_T^{\text{miss}} > 65 \text{ GeV}$ requirement at HLT.
- Tau and electron: The L1 condition is the corresponding $\tau+e\gamma$ trigger. The p_T threshold remains at $16 \text{ GeV}/c$ for the electron. There is no explicit p_T cut for the τ at HLT, but there is the match-to-track requirement for the τ candidate.
- Jets: The L1 conditions for the single-, double-, triple- and quadruple-jet triggers have been simplified considerably. Single jet triggers run off an .OR. of a central-, forward- or tau-jet trigger at L1. Double-, triple- and quadruple-jet triggers use an .OR. of the all the L1 terms requiring the same number of jets or less. For example, the triple-jet trigger is driven by an .OR. of the single-, double- and triple-jet L1 bits. In all cases, jets can be found in either the central or the forward region of the detector, and they include the τ candidates. The additional p_T cuts at HLT are: 400 (single), 350 (double), 195 (triple) and 80 (quadruple) GeV . The new double-jet trigger is expected to have a large overlap with the single-jet trigger path. However, it is useful for testing the additional bias introduced by the requirement for a second jet in the event. A series of prescaled triggers have also been introduced, which are discussed later (Sec. E.4.3.2).
- b-jet: This trigger is also based on the logical .OR. of the single-, double-, triple- and quadruple-jet L1 terms. At HLT, we have the additional requirement that the event is consistent with b -content. The E_T cut for the HLT jets is one of the following: 350 GeV if the event has one jet, 150 GeV if the event has three jets, or 55 GeV if the event has four jets.
- Jet and E_T^{miss} : The E_T thresholds are 180 and 80 GeV , respectively. The L1 condition is a single E_T^{miss} object above 60 GeV .

E.4.3 New triggers

E.4.3.1 Cross-channel triggers

The trigger studies presented in the DAQ TDR [68] have been the most comprehensive CMS effort to date to calculate rates for various trigger paths across many physics channels. For those studies the focus has been the optimization of the rejection of the individual object-id algorithms (muon, electron, tau, etc.) rather than the combination of them into more powerful trigger tools. However, single (or even double) trigger objects are limited by the rate and, therefore, have their thresholds often higher than desired for many physics analyses. If the signal contains more than one trigger objects, using trigger paths combining different objects may yield a considerable gain by allowing lower trigger thresholds and higher efficiency. Cross-channel triggers can be much more stable and less prone to rate fluctuations from operating conditions. The correlations among trigger objects can help reduce difficult backgrounds and instrumental fakes. The additional advantage is that such cross-channel triggers have noticeably lower rates than the single trigger channels and therefore contribute fairly little to the overall bandwidth.

Some cross-channel triggers have already been considered and their rates estimated [68], such as $\tau + e$ and $\tau + E_T^{\text{miss}}$, motivated by the Higgs searches with hadronic decays of τ and leptons, and $\text{jet} + E_T^{\text{miss}}$, important for searches of super-symmetric particles. The new addition to the Trigger Menu, expanding the scope of Higgs searches, is a combined $\tau + \mu$ trigger with p_T thresholds at 40 and 15 GeV/ c , respectively. It is driven by the single- μ L1 bit.

We are presenting here a few additional cross-channel triggers, along with the physics motivation.

- A new category of triggers introduced here is the acoplanar dijet and $\text{jet} + E_T^{\text{miss}}$ for SUSY signals. The gain is the lower thresholds that become possible because of the topology constraint. Possible biases should be studied, so these triggers are meant to run in parallel with the standard jet and $\text{jet} + E_T^{\text{miss}}$ triggers without the acoplanarity requirements. We introduce a double-jet trigger with E_T thresholds at (200, 200) GeV and $|\Delta\phi| < 2.1$, and a new Jet + E_T^{miss} trigger with E_T thresholds at (100, 80) GeV and $|\Delta\phi| < 2.1$. The former is driven by an .OR. of the single- and double-jet requirements at L1 (bits 36, 37). The latter is driven by a simple $E_T^{\text{miss}} > 60$ GeV L1 requirement.
- “ $E_T^{\text{miss}} + X$ ” triggers: A combination of an E_T^{miss} object with an H_T cut, one (or more) jet or lepton may be the only way to access E_T^{miss} -enhanced triggers if there are problems (e.g. instrumental fakes) that prevent CMS from running an inclusive E_T^{miss} trigger. At this point we have implemented:
 - Multi-jets and E_T^{miss} : These will be useful for SUSY studies, just like the series of jet triggers. However, the additional E_T^{miss} requirement allows us to lower the thresholds on the jets, and therefore increase the sensitivity of the analyses. We introduce here a dijet + E_T^{miss} trigger with $E_T^{\text{jet}} > 155$ GeV, $E_T^{\text{miss}} > 80$ GeV, a triple-jet + E_T^{miss} trigger with $E_T^{\text{jet}} > 85$ GeV, $E_T^{\text{miss}} > 80$ GeV and a quadruple-jet + E_T^{miss} trigger with $E_T^{\text{jet}} > 35$ GeV, $E_T^{\text{miss}} > 80$ GeV. These all run off the single L1 requirement for $E_T^{\text{miss}} > 60$ GeV.

- $H_T + E_T^{\text{miss}}$ and $H_T + e$: It is difficult to contain the rate for an inclusive H_T trigger without any additional cuts. The requirement for a E_T^{miss} cut or an additional electron in the event allows us to access events with lower E_T^{miss} or softer electrons. This can give an increased efficiency for W +jets, top physics, SUSY cascades, and other similar physics channels. Here we propose an $H_T + E_T^{\text{miss}}$ trigger with $H_T > 350$ GeV, $E_T^{\text{miss}} > 80$ GeV and an $H_T + e$ trigger with $H_T > 350$ GeV and $p_T > 20$ GeV/ c for the electron. They are both driven by the $E_T^{\text{miss}} > 60$ GeV condition at L1.

Some additional cross-channel triggers that have not been included in this Trigger Table iteration but should be considered in future trigger studies are

- An $e + \mu$ trigger is of interest in many studies, for example:
 - $qqH, H \rightarrow \tau\tau \rightarrow 2\ell$, with an expected gain thanks to the lower lepton thresholds compared to the single-electron and single-muon trigger paths,
 - many SUSY decays including leptons in the final state,
 - top measurements in the double leptonic channel ($t\bar{t} \rightarrow b\bar{b}\ell\nu\ell\nu$), gaining sensitivity at the lower p_T spectrum, and
 - $B_s \rightarrow \ell\ell$, to allow for the lepton-number-violating channel to be studied.
- $E_T^{\text{miss}} + \ell$: The idea here is to exploit the presence of a W boson or a top decay in many channels. This could be used in many SM channels where lowering the lepton threshold extends the range of the measurement. For example:
 - top measurement in the double leptonic and semi-leptonic channels,
 - single top production, and
 - W measurements.

Furthermore, this is a typical signature of an event containing super-symmetric particles.

- Triggers combining a lepton and a jet, or a lepton and a b -jet could be of interest for top measurements. The $\ell + \text{jet}$ signature is also very common in super-symmetric events.
- Finally, a combination of a lepton and a photon ($e + \gamma$ and $\mu + \gamma$) is ideal for Flavor Changing Neutral Current analyses, exploiting the extraordinary capabilities of CMS in detecting photons. These triggers allow to lower the thresholds on the lepton and the photon, increasing the event yield compared to the single- e , μ or γ trigger paths.



E.4.3.2 Single-jet triggers

In this section we describe the single jet trigger paths. These have been driven by the needs of the inclusive jet and dijet analysis. The full study can be found in Ref. [112]. Here we summarize the conclusions, along with a short description of the strategy for adjusting thresholds and prescales as the luminosity changes. This study looks at the evolution of the single-jet triggers for various luminosities. It serves as an example of how to preserve the long-term continuity of the triggers used for physics analyses. It is, therefore, interesting and instruc-




Figure E.3: The integrated trigger rates at L1 (left) and HLT (right) above the E_T thresholds for the highest E_T jet is plotted versus the E_T threshold for three luminosity scenarios: $\mathcal{L} = 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$ (solid), and $\mathcal{L} = 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ (dashed), and $\mathcal{L} = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ (dot-dashed). HLT thresholds that give 2.5 Hz are shown by vertical dotted lines.

tive beyond the strict scope of the single-jet trigger suite.

To measure jet spectra down to low jet E_T and dijet mass requires multiple triggers, of roughly equal total rate, and with appropriately chosen E_T thresholds and prescales. In Fig. E.3 we show estimates of the L1 and HLT single jet trigger rates vs. corrected jet E_T . In Table E.5 we show the single jet trigger paths from L1 to HLT including thresholds, prescales and estimates of the rates. We find that the maximum allowed HLT rate is the constraining factor for triggering on jets. For luminosities $\mathcal{L} = 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$, $\mathcal{L} = 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ and $\mathcal{L} = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ the highest E_T threshold at HLT was chosen to give a rate of roughly 2.5 Hz, as illustrated in Fig. E.3, so that four triggers would saturate an allowed jet rate of roughly 10 Hz at HLT.

The highest E_T threshold at any luminosity is never prescaled. Lower thresholds are prescaled and are chosen at roughly half the E_T of the next highest threshold. This allows reasonable statistics in the overlap between the two samples, necessary for measuring trigger efficiencies and producing a continuous jet spectrum. The total L1 jet rate required is only around 0.3 kHz, a small fraction of the L1 total bandwidth. Since these triggers are limited by HLT, not L1, for each trigger path the L1 thresholds are chosen low enough to have a L1 trigger efficiency of more than 95% at the corresponding HLT threshold in the path, as shown in Fig. E.4. This strategy utilizes ten times more bandwidth at L1 than at HLT to insure that all of the resulting HLT sample has high enough trigger efficiency to be useful for analysis.

Table E.5 illustrates an example trigger strategy to maintain the continuity of jet analysis as the luminosity increases over a time span of years. The most important reason for introducing a series of multiple triggers with varying thresholds and prescales is the continuity of the trigger path for the physics analyses. For every new “luminosity era” CMS enters, it is fundamental that we maintain the thresholds introduced in the previous months or years, allowing combination of trigger samples over time. For the prescaled thresholds, we may increase the prescales, either in discrete steps or dynamically, to maintain the allowed HLT rate with increasing luminosity. However, to maintain maximum sensitivity to new physics, the highest E_T threshold must never be prescaled. For example, in Table E.5 when the luminosity increases by only a factor of 2 from $\mathcal{L} = 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ to $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$, we double the prescales on the lower- E_T triggers but don’t change either the threshold or the

Path	L1				HLT	
	E_T Cut (GeV)	Unpres. Rate (KHz)	Prescale (N)	Presc. Rate (kHz)	E_T Cut GeV	Rate (Hz)
Single Jet Triggers in Scenario 1: $\mathcal{L} = 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$						
High	140	0.044	1	0.044	250	2.8
Med	60	3.9	40	0.097	120	2.4
Low	25	2.9×10^2	2,000	0.146	60	2.8
Single Jet Triggers in Scenario 2: $\mathcal{L} = 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$						
Ultra	270	0.019	1	0.019	400	2.6
High	140	0.44	10	0.044	250	2.8
Med	60	39	400	0.097	120	2.4
Low	25	2.9×10^3	20,000	0.146	60	2.8
Single Jet Triggers in Scenario 3: $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$						
Ultra	270	0.038	1	0.038	400	5.2
High	140	0.88	20	0.044	250	2.8
Med	60	78	800	0.097	120	2.4
Low	25	5.8×10^3	40,000	0.146	60	2.8
Single Jet Triggers in Scenario 4: $\mathcal{L} = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$						
Super	450	0.014	1	0.014	600	2.8
Ultra	270	0.19	10	0.019	400	2.6
High	140	4.4	100	0.044	250	2.8
Med	60	3.9×10^2	4,000	0.097	120	2.4
Low	25	2.9×10^4	200,000	0.146	60	2.8

Table E.5: Single jet trigger table showing path names, trigger thresholds in corrected E_T , prescales, and estimated rates at L1 and HLT for four different luminosity scenarios.

prescale of the highest E_T trigger labeled Ultra. This allows us to maintain stability of the trigger thresholds, and analyses that depend on them, with only modest increases in the total rate for single object triggers. When the HLT rate in the unprescaled trigger becomes intolerably high, a higher E_T threshold unprescaled trigger is introduced, and the old unprescaled trigger can then be prescaled as necessary.

For the particular case of single-jet triggers: To commission the calorimeters, or perform a one-time jet study, it may be desirable to have more jets. If we want to write more than roughly 10 Hz of single jets at HLT, we can still use the same suite of single-jets, but lower the prescales to obtain more jets at low E_T . This is preferable to moving the threshold for the unprescaled trigger, or any of the triggers, and ending up with a special trigger that is only applicable for a given running period and difficult to combine with other samples.

For $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$, the suggested jet thresholds have been studied again in the scope of the global High-Level trigger analysis (Sec. E.5) and new L1 prescales and rates have been determined. For the trigger table proposed in this study, we have chosen four triggers, with E_T thresholds of 400, 250, 120 and 60 GeV, and prescales of 1, 10, 1000 and 100 000, respectively.




Figure E.4: The efficiency for passing the L1 jet trigger is shown as a function of HLT corrected jet E_T for each of the trigger paths shown in Table E.5. The L1 thresholds were chosen to give an efficiency of greater than 95% at the corresponding HLT threshold.

E.4.3.3 Other triggers

The remaining triggers that have been introduced since the DAQ TDR are:

- Inclusive E_T^{miss} trigger: As discussed earlier, this is a difficult trigger that is subject to the good understanding and control of the detector noise. We suggest here a single E_T^{miss} trigger with $E_T > 91$ GeV, driven by the $E_T^{\text{miss}} > 60$ GeV L1 condition. This is just an indicative value, rather on the low side, as E_T^{miss} rates appear lower compared to Ref. [68]. It is foreseen that additional E_T^{miss} triggers with different thresholds and prescales will be introduced in the future.
- Diffractive trigger: This trigger is different than all others described earlier in that it uses the TOTEM detector [791, 792]. At L1 we ask for two central jets with $E_T > 40$ GeV, along with a proton tagged with the 220 m Roman Pot. At HLT, a similar di-jet cut and a “back-to-back” azimuthal condition are applied. We also require that we have a consistent measurement of the proton energy loss ξ in the two hemispheres (within 2σ , measured at the Roman Pots). A final condition for a tagged proton seen by the 420 m Roman Pot brings the HLT rate down to $\mathcal{O}(1)$ Hz.



This trigger is discussed in detail in Sec. E.3.

E.5 Performance

The performance of the trigger system is studied by using simulated data that has been digitized with appropriate pileup[‡], taking into account both the inelastic (55.2 mb) and the

[‡]We have estimated the average number of in-time interactions per bunch crossing to be 5 for $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. Additional, out-of-time interactions have been ignored.

diffractive (24.1 mb) cross sections. To reduce the amount of simulation time, about 50 million minimum bias events were simulated and reused in random combinations. It was ensured that these events do not cause triggers by themselves to avoid over estimating the rates due to this reuse of events.

In the following sections we list trigger rates along with their statistical uncertainties. These take into account the luminosity-dependent weight of the events from the different samples, the corresponding cross sections and the \hat{p}_T of the main interaction and the pile-up contribution. They do *not* take into account the uncertainties of these individual factors, *i.e.* no systematic effects are studied here.

The L1 calorimeter trigger object rate studies are performed using QCD data that has been generated in several bins of \hat{p}_T . A special event-weighting procedure [806] has been applied to properly take into account the cross sections of the sub-samples. The L1 muon and E_T^{miss} rate studies are performed using a purely minimum bias sample.

The HLT rates are estimated using specially enriched samples. For the triggers invoking muons, electrons and photons we have used a minimum bias sample enriched in muons, as well as $W \rightarrow e/\mu\nu$, $Z \rightarrow ee/\mu\mu$ and jet(s) + γ MC datasets. For the triggers including jets we have used QCD samples. These samples also contribute to the electron and photon triggers. Events triggered exclusively with muons have been excluded from the QCD samples, to avoid double-counting with the muon-enriched sample. Table E.6 summarizes the MC samples used for the trigger studies, and their corresponding contribution to the HLT rate. A more detailed breakdown of the contributions to the electron, photon and muon trigger rates from the different samples is discussed later (Sec. E.5.3). For our calculations, we have used the standard HLT physics algorithms (ORCA_8_13_3 [15]) for the implementation of all trigger paths. At the time of this writing, this includes the latest algorithms and jet calibrations. For the global evaluation of the trigger rates we have used the ‘‘HLT steering code’’ [807, 154].

E.5.1 L1 Rates

The background at L1 is entirely dominated by strong interactions. The muon rates at L1 are dominated by low p_T muons which are reconstructed as high p_T muons due to limited resolution at the trigger level. For the electron/photon trigger the rate is dominated by jets that fragment to high E_T π^0 s. The jet rates are dominated by true jets in the QCD events. The E_T^{miss} background is due to the limited energy resolution, and pile-up of minimum bias interactions.

We first produce a trigger table with L1 rates for DAQ TDR chosen thresholds for comparison. For the calculations we use a sample of 2 million minimum bias crossings with an average of 5 events per crossing, constructed from the minbias events, without reuse of events. The out-of-time pile-up is neglected. Even though there are small differences for the individual triggers, the integral rate is consistent with the rates reported in Ref. [68]. This comparison serves as a cross-check and is a necessary intermediate step before the introduction of new trigger terms. Table E.7 summarizes the L1 rate calculations for the DAQ TDR triggers with the new MC samples. Besides the ‘‘95% efficiency points’’ (used throughout the DAQ TDR), the applied L1 thresholds are also given.

For the new trigger table: We select several thresholds for each trigger object type and quote corresponding rates and prescales for $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. For the single objects we have

added a series of prescaled triggers to determine the efficiency turn-on. For the multi-object triggers we have picked the lowest common threshold that is allowed for the allocated bandwidth. For the cross-channel triggers we have attempted to keep the lepton thresholds as low as possible, within the allocated bandwidth based on the physics needs of the experiment. The prescales are chosen such that the simulated rate at all times falls below the DAQ bandwidth taking into account a safety factor of 3. Detailed studies with the selected thresholds, pre-scales and the individual trigger rates for the different sets of L1 trigger objects can be found in Ref. [808]. The total L1 rate for all triggers (including prescaled ones) is 22.3 ± 0.6 kHz.

E.5.2 L1 Trigger Object Corrections

The trigger decisions are based on E_T of the objects reconstructed by various algorithms. Unfortunately, the energy deposition in the calorimeter and the size of the trigger towers, are not entirely uniform. We have used fits to the reconstructed-to-generated E_T ratios to correct for non-uniformity of the response for jets and electron/photon candidates found at all levels of trigger [799]. This correction procedure adjusts the mean response to the generated level.

The energy response of the calorimeters and the limited number of bits used in trigger calculations result in a finite resolution for the reconstructed trigger objects. Similarly, misalignments of the tracking systems and the limited number of patterns in the muon trigger look-up-tables also result in a finite resolution. To avoid systematic problems in understanding the trigger efficiency turn-on with the E_T of the trigger objects, it is envisioned that only data where high trigger efficiency is assured is used for analysis.

E.5.3 HLT Rates

A rough comparison of the HLT bandwidth given to various triggers, calculated with the latest algorithms and the ones reported in Ref. [68] is shown in Table E.8. It must be noted that not only thresholds but also other cuts are different in the two trigger studies. Furthermore, additional changes in the HLT algorithms (summarized in Sec. E.2.2) must be taken into account. This comparison serves only as a consistency check. It reaffirms that despite the evolution of the CMS reconstruction algorithms over the years, trigger rates remain under control and that no major bandwidth changes are expected.

The contributions to the single and double electron and photon trigger rates at HLT from the various MC samples is given at Table E.9. The main contributions to the single electron trigger come from the QCD and $W \rightarrow e\nu$ samples, whereas for the single photon trigger the primary source is the jet(s) + γ events.

Table E.10 shows in a similar way the contributions to the single and double standard and relaxed muon rates from the various MC samples.

E.5.4 Trigger Tables

Table E.11 summarizes the L1 triggers used in this study, their kinematic thresholds, the individual and cumulative rates. We have assumed a DAQ capability of 50 kHz, taking into account a safety factor of 3.




Figure E.5: Heuristic comparison of HLT bandwidth assigned to various trigger paths calculated in this study with the DAQ TDR. For the triggers introduced in this study the DAQ TDR entries appear empty. See text for details on different kinematic cuts and changes in the HLT algorithms.

Table E.12 gives the full list of trigger paths proposed for $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ that have been described earlier for an HLT output rate of approximately 120 Hz.

Fig. E.5 shows a graphic representation of the HLT bandwidth assigned to all trigger paths presented in this study. For the triggers that appeared in the DAQ TDR, the corresponding rates are overlaid, in a heuristic comparison.

Table E.6: Description and sizes of MC Samples used for the trigger studies. The contribution to the HLT rate does not include prescaled triggers.

Sample description	Cuts (Momenta in GeV/c)	Cross section (mb)	# of events	HLT rate (Hz)
Minimum bias with in-time pile-up; <# of interactions>= 5	—	79.3	50 000 000	—
QCD	$\hat{p}_T \in [15, 20]$	1.46×10^0	49 491	
QCD	$\hat{p}_T \in [20, 30]$	6.32×10^{-1}	49 244	
QCD	$\hat{p}_T \in [30, 50]$	1.63×10^{-1}	49 742	
QCD	$\hat{p}_T \in [50, 80]$	2.16×10^{-2}	99 486	
QCD	$\hat{p}_T \in [80, 120]$	3.08×10^{-3}	96 238	
QCD	$\hat{p}_T \in [120, 170]$	4.94×10^{-4}	99 736	
QCD	$\hat{p}_T \in [170, 230]$	1.01×10^{-4}	99 226	
QCD	$\hat{p}_T \in [230, 300]$	2.45×10^{-5}	99 481	
QCD	$\hat{p}_T \in [300, 380]$	6.24×10^{-6}	98 739	
QCD	$\hat{p}_T \in [380, 470]$	1.78×10^{-6}	46 491	
QCD	$\hat{p}_T \in [470, 600]$	6.83×10^{-7}	47 496	
QCD	$\hat{p}_T \in [600, 800]$	2.04×10^{-7}	48 986	
QCD	$\hat{p}_T \in [800, 1000]$	3.51×10^{-8}	45 741	
<i>Partial total</i>			930 099	55.3 ± 6.9
$W \rightarrow e\nu$	1 electron with $ \eta < 2.7, p_T > 25$	7.9×10^{-6}	3 944	9.7 ± 0.2
$Z \rightarrow ee$	2 electrons with $ \eta < 2.7, p_T > 5$	8.2×10^{-7}	4 000	1.4 ± 0.0
$pp \rightarrow \text{jet}(s) + \gamma,$ $\hat{p}_T > 30 \text{ GeV}/c$	jet: $p_T > 20,$ $\gamma: p_T > 30$	2.5×10^{-6}	4 000	1.0 ± 0.0
$W \rightarrow \mu\nu$	1 muon with $ \eta < 2.5, p_T > 14$	9.8×10^{-6}	4 000	14.0 ± 0.3
$Z \rightarrow \mu\mu$	2 muons with $ \eta < 2.5, p_T > 20, 10$	7.9×10^{-7}	2 941	1.5 ± 0.0
$pp \rightarrow \mu + X$	1 muon with $p_T > 3$	2.4×10^{-2}	839 999	25.5 ± 1.2

Table E.7: Trigger table showing L1 rates for DAQ TDR chosen thresholds for $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. Whenever the “95% efficiency point” is reported in DAQ TDR, we also give the actual kinematic threshold that has been applied.

Trigger	95% Eff. point	Threshold (GeV)	Rate (kHz)	Cumulative Rate (kHz)
Single $e\gamma$	29	23.4	3.38 ± 0.23	3.4 ± 0.2
Double $e\gamma$	17	11.5	0.85 ± 0.12	4.0 ± 0.3
Single μ	—	14	2.53 ± 0.20	6.5 ± 0.3
Double μ	—	3	4.05 ± 0.26	10.3 ± 0.4
Single τ	86	93	3.56 ± 0.24	9.7 ± 0.4
Double τ	59	66	1.97 ± 0.18	10.6 ± 0.4
1-, 3-, 4-jets	177, 86, 70	135, 58, 45	2.43 ± 0.20	11.9 ± 0.4
Jet + E_T^{miss}	—	88, 46	1.07 ± 0.13	12.2 ± 0.4
$e\gamma + \tau$	—	21, 45	3.64 ± 0.24	12.9 ± 0.5
L1 Trigger Total				12.9 ± 0.5

Table E.8: Comparison of HLT bandwidth given to various trigger paths calculated in this study with the DAQ TDR. See text for details on different kinematic cuts and changes in the HLT algorithms.

Trigger	DAQ TDR Rate (Hz)	New Rate (Hz)
Inclusive e	33.0	23.5 ± 6.7
$e-e$	1.0	1.0 ± 0.1
Relaxed $e-e$	1.0	1.3 ± 0.1
Inclusive γ	4.0	3.1 ± 0.2
$\gamma-\gamma$	5.0	1.6 ± 0.7
Relaxed $\gamma-\gamma$	5.0	1.2 ± 0.6
Inclusive μ	25.0	25.8 ± 0.8
$\mu-\mu$	4.0	4.8 ± 0.4
$\tau + E_T^{\text{miss}}$	1.0	0.5 ± 0.1
$\tau + e$	2.0	< 1.0
Double Pixel τ	1.0	4.1 ± 1.1
Double Tracker τ	1.0	6.0 ± 1.1
Single jet	1.0	4.8 ± 0.0
Triple jet	1.0	1.1 ± 0.0
Quadruple jet	7.0	8.9 ± 0.2
jet + E_T^{miss}	5.0	3.2 ± 0.1
b -jet (leading jet)	5.0	10.3 ± 0.3
b -jet (2 nd leading jet)	5.0	8.7 ± 0.3

Table E.9: Contributions to the HLT rates for the electron and photon triggers from the various MC datasets.

Trigger	Threshold (GeV)	Rates (Hz)			
		QCD	$W \rightarrow e\nu$	$Z \rightarrow ee$	jet(s) + γ
Inclusive e	26	12.6 ± 6.7	9.7 ± 0.2	1.2 ± 0.0	—
e - e	12, 12	0.1 ± 0.1	—	1.0 ± 0.0	—
Relaxed e - e	19, 19	0.3 ± 0.1	—	1.0 ± 0.0	—
Inclusive γ	80	1.1 ± 0.2	—	—	2.0 ± 0.1
γ - γ	30, 20	1.3 ± 0.8	—	—	0.3 ± 0.0
Relaxed γ - γ	30, 20	0.9 ± 0.6	—	—	0.3 ± 0.0

Table E.10: Contributions to the HLT rates for the muon triggers from the various MC datasets.

Trigger	Threshold (GeV)	Rates (Hz)		
		Enriched- μ sample	$W \rightarrow \mu\nu$	$Z \rightarrow \mu\mu$
Inclusive μ	19	10.9 ± 0.8	13.4 ± 0.3	1.5 ± 0.0
Relaxed μ	37	5.1 ± 0.5	5.7 ± 0.1	1.1 ± 0.0
μ - μ	7, 7	3.4 ± 0.4	—	1.3 ± 0.0
Relaxed μ - μ	10, 10	7.1 ± 0.5	—	1.4 ± 0.0

Table E.11: The L1 Trigger Menu at $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. Individual and cumulative rates are given for the different trigger paths and selected kinematic thresholds.

Trigger	L1 Threshold (GeV)	L1 Rate (kHz)	Cumulative L1 Rate (kHz)
Inclusive $e\gamma$	22	3.9 ± 0.3	3.9 ± 0.3
Double $e\gamma$	11	1.0 ± 0.1	4.6 ± 0.3
Inclusive μ	14	2.5 ± 0.2	7.1 ± 0.3
Double μ	3	4.0 ± 0.3	11.0 ± 0.4
Inclusive τ	100	2.2 ± 0.2	12.9 ± 0.5
Double τ	60	3.0 ± 0.2	14.9 ± 0.5
1-,2-,3-,4-jets	150,100,70,50	2.2 ± 0.2	15.8 ± 0.5
H_T	275	2.0 ± 0.2	16.2 ± 0.5
E_T^{miss}	60	0.4 ± 0.1	16.3 ± 0.5
$H_T + E_T^{\text{miss}}$	200, 40	1.1 ± 0.1	16.6 ± 0.5
jet + E_T^{miss}	100, 40	1.1 ± 0.1	16.7 ± 0.5
$\tau + E_T^{\text{miss}}$	60, 40	2.7 ± 0.2	18.8 ± 0.5
$\mu + E_T^{\text{miss}}$	5, 30	0.3 ± 0.1	19.0 ± 0.6
$e\gamma + E_T^{\text{miss}}$	15, 30	0.5 ± 0.1	19.1 ± 0.6
$\mu + \text{jet}$	7, 100	0.2 ± 0.1	19.1 ± 0.6
$e\gamma + \text{jet}$	15, 100	0.6 ± 0.1	19.2 ± 0.6
$\mu + \tau$	7, 40	1.2 ± 0.1	19.8 ± 0.6
$e\gamma + \tau$	15, 60	2.6 ± 0.2	20.5 ± 0.6
$e\gamma + \mu$	15, 7	0.2 ± 0.1	20.5 ± 0.6
Prescaled	See Ref. [808]		22.3 ± 0.6
Total L1 Rate			22.3 ± 0.6

Table E.12: The High-Level Trigger Menu at $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ for an output of approximately 120 Hz. The E_T values are the kinematic thresholds for the different trigger paths.




Trigger	L1 bits used	L1 Prescale	HLT Threshold (GeV)	HLT Rate (Hz)
Inclusive e	2	1	26	23.5 ± 6.7
e - e	3	1	12, 12	1.0 ± 0.1
Relaxed e - e	4	1	19, 19	1.3 ± 0.1
Inclusive γ	2	1	80	3.1 ± 0.2
γ - γ	3	1	30, 20	1.6 ± 0.7
Relaxed γ - γ	4	1	30, 20	1.2 ± 0.6
Inclusive μ	0	1	19	25.8 ± 0.8
Relaxed μ	0	1	37	11.9 ± 0.5
μ - μ	1	1	7, 7	4.8 ± 0.4
Relaxed μ - μ	1	1	10, 10	8.6 ± 0.6
$\tau + E_T^{\text{miss}}$	10	1	65 (E_T^{miss})	0.5 ± 0.1
Pixel τ - τ	10, 13	1	—	4.1 ± 1.1
Tracker τ - τ	10, 13	1	—	6.0 ± 1.1
$\tau + e$	26	1	52, 16	< 1.0
$\tau + \mu$	0	1	40, 15	< 1.0
b -jet (leading jet)	36, 37, 38, 39	1	350, 150, 55 (see text)	10.3 ± 0.3
b -jet (2 nd leading jet)	36, 37, 38, 39	1	350, 150, 55 (see text)	8.7 ± 0.3
Single-jet	36	1	400	4.8 ± 0.0
Double-jet	36, 37	1	350	3.9 ± 0.0
Triple-jet	36, 37, 38	1	195	1.1 ± 0.0
Quadruple-jet	36, 37, 38, 39	1	80	8.9 ± 0.2
E_T^{miss}	32	1	91	2.5 ± 0.2
jet + E_T^{miss}	32	1	180, 80	3.2 ± 0.1
acoplanar 2 jets	36, 37	1	200, 200	0.2 ± 0.0
acoplanar jet + E_T^{miss}	32	1	100, 80	0.1 ± 0.0
2 jets + E_T^{miss}	32	1	155, 80	1.6 ± 0.0
3 jets + E_T^{miss}	32	1	85, 80	0.9 ± 0.1
4 jets + E_T^{miss}	32	1	35, 80	1.7 ± 0.2
Diffractive	Sec. E.3	1	40, 40	< 1.0
$H_T + E_T^{\text{miss}}$	31	1	350, 80	5.6 ± 0.2
$H_T + e$	31	1	350, 20	0.4 ± 0.1
Inclusive γ	2	400	23	0.3 ± 0.0
γ - γ	3	20	12, 12	2.5 ± 1.4
Relaxed γ - γ	4	20	19, 19	0.1 ± 0.0
Single-jet	33	10	250	5.2 ± 0.0
Single-jet	34	1 000	120	1.6 ± 0.0
Single-jet	35	100 000	60	0.4 ± 0.0
Total HLT rate				119.3 ± 7.2

Appendix F

References

Notes:

- a) CMS Notes are available at <http://cms.cern.ch/iCMS/> unless otherwise noted.
- b) References marked **doi** should be prefixed with <http://dx.doi.org/>.

-
- [1] C. Seez et al., "Photon decay modes of the intermediate mass Higgs," in *Proceedings of the Large Hadron Collider Workshop*, Jarlskog, G. and Rein, D., ed., volume II. CERN 90-10, Aachen, 4-9 October, 1990.
 - [2] CMS Collaboration, "The Compact Muon Solenoid Technical Proposal," *CERN/LHCC 94-38* (1994). LHCC/P1.
 - [3] CMS Collaboration, "The Electromagnetic Calorimeter Technical Design Report," *CERN/LHCC 97-033* (1997). CMS TDR 4, Addendum CERN/LHCC 2002-027.
 - [4] A. Favara and M. Pieri, "Confidence Level Estimation and Analysis Optimization," *Preprint DFF 278/4/1997* (1997).
 - [5] **L3 Collaboration** Collaboration, M. Acciarri et al., "Search for the Standard Model Higgs Boson in e+e- Interactions at $161 \text{ GeV} < \sqrt{s} < 172 \text{ GeV}$," *Phys. Lett. B* **411** (1997) 373.
 - [6] **CMS Collaboration**, M. Pieri ¹, "Inclusive search for the Higgs boson in the $H \rightarrow \gamma\gamma$ channel," *CMS AN ~~XXXX~~ME: No AN's AN 2006/069* (2006).
 - [7] **CMS Collaboration** ¹, "The CMS Physics Technical Design Report, Volume 1," *CERN/LHCC ~~XXXX~~ME: need final url2006-001* (2006). CMS TDR 8.1.
 - [8] M. Spira, "QCD effects in Higgs physics," *Fortsch. Phys.* **46** (1998) 203–284, [arXiv:hep-ph/9705337](https://arxiv.org/abs/hep-ph/9705337).
 - [9] A. Djouadi, J. Kalinowski, and M. Spira, "HDECAY: a program for Higgs boson decays in the Standard Model and its supersymmetric extension," *Comput. Phys. Commun.* **108** (1998) 56–74. doi:10.1016/S0010-4655(97)00123-9.
 - [10] **CMS Collaboration** ¹ Dubinin et al., "The vector boson fusion production with $H \rightarrow \gamma\gamma$," *CMS AN ~~XXXX~~ME: No AN's AN-2006/050* (2006).

- [11] CMS Collaboration, M. Leterrier, "Search for SM Higgs boson with W/Z+H, $H \rightarrow 2\gamma$ channel," *CMS AN* **2006/058** (2006). **VXAME: No AN's 2006/058**
- [12] CMS Collaboration, J. Agram et al., "Associated $t\bar{t}H$ Production with $H \rightarrow \gamma\gamma$ in CMS," *CMS AN* **2006/064** (2006). **VXAME: Analysis Note 2006/064**
- [13] "CMSIM home page." Located at <http://cmsdoc.cern.ch/cmsim/cmsim.html>.
- [14] "OSCAR: CMS Simulation Package Home Page." <http://cmsdoc.cern.ch/oscar>.
- [15] "ORCA: CMS Reconstruction Package." Site located at <http://cmsdoc.cern.ch/orca>.
- [16] W. Giele et al., "The QCD/SM working group: Summary report," arXiv:hep-ph/0204316. **VXAME: To V.Litvin: is this ref. OK?**
- [17] M. Dobbs et al., "The QCD/SM working group: Summary report," arXiv:hep-ph/0403100.
- [18] T. Binoth, "Two photon background for Higgs boson searches at the LHC," arXiv:hep-ph/0005194.
- [19] Z. Bern, L. J. Dixon, and C. Schmidt, "Isolating a light Higgs boson from the di-photon background at the LHC," *Phys. Rev.* **D66** (2002) 074018, arXiv:hep-ph/0206194.
- [20] CMS Collaboration, L. Agostino et al., "HLT Selection of Electrons and Photons," *CMS AN* **2005/064** (2005).
- [21] CMS Collaboration, E. Meschi et al., "Electron Reconstruction in the CMS Electromagnetic Calorimeter," *CMS Note* **2001/034** (2001).
- [22] CMS Collaboration, N. Marinelli, "Track Finding and Identification of Converted Photons," *CMS Note* **2006/005** (2006).
- [23] CMS Collaboration, V. Litvin et al., "The Rejection of Background to the $H \rightarrow \gamma\gamma$ Process Using Isolation Criteria Based on Information from the Calorimeter and Tracker," *CMS Note* **2002-030** (2002).
- [24] CMS Collaboration, M. Pieri et al., "Distinguishing Isolated Photos from Jets," *CMS Note* **2006/007** (2006).
- [25] The LEP Collaborations: ALEPH, DELPHI, L3 and OPAL, "Lower Bound for the Standard Model Higgs Boson Mass from Combining the Results of the Four LEP Experiments," *CERN EP* **98-046** (1998).
- [26] CMS Collaboration, S. Baffioni et al., "Discovery potential for the SM Higgs boson in the $H \rightarrow ZZ^{(*)} \rightarrow e^+e^-e^+e^-$ decay channel," *CMS AN* **2006/070** (2006). **VXAME: No AN's 2006/070**
- [27] E. Barberio, B. van Eijk, and Z. Was, "PHOTOS: A Universal Monte Carlo for QED radiative corrections in decays," *Comput. Phys. Commun.* **66** (1991) 115–128.


- [28] E. Barberio and Z. Was, "PHOTOS: A Universal Monte Carlo for QED radiative corrections. Version 2.0," *Comput. Phys. Commun.* **79** (1994) 291–308.
doi:10.1016/0010-4655(94)90074-4.
- [29] J. Pumplin et al., "New generation of parton distributions with uncertainties from global QCD analysis," *JHEP* **07** (2002) 012, arXiv:hep-ph/0201195.
- [30] T. Sjostrand, L. Lonnblad, and S. Mrenna, "PYTHIA 6.2: Physics and manual," arXiv:hep-ph/0108264.
- [31] M. Spira, "HIGLU: A Program for the Calculation of the Total Higgs Production Cross Section at Hadron Colliders via Gluon Fusion including QCD Corrections," arXiv:hep-ph/9510347.
- [32] A. Djouadi, J. Kalinowski, and M. Spira, "HDECAY: A Program for Higgs Boson Decays in the Standard Model and its Supersymmetric Extension," *Comput. Phys. Commun.* **108** (1998) 56–74, arXiv:hep-ph/9704448.
- [33] C. Zecher et al., "Leptonic Signals from off-shell Z Boson Pairs at Hadron Colliders," arXiv:hep-ph/9404295.
- [34] **CompHEP** Collaboration, E. Boos et al., "CompHEP 4.4: Automatic computations from Lagrangians to events," *Nucl. Instrum. Meth.* **A534** (2004) 250–259, arXiv:hep-ph/0403113.
- [35] S. Slabospitsky and L. Sonnenschein, "TopReX generator (version 3.25): Short manual," *Comput. Phys. Commun.* **148** (2002) 87–102.
doi:10.1016/S0010-4655(02)00471-X.
- [36] M. Beneke et al., "Top quark physics," arXiv:hep-ph/0003033.
- [37] **CMS** Collaboration, S. Baffioni, C. Charlot, F. Ferri, D. Futyan, P. Meridiani, I. Puljak, C. Rovelli, R. Salerno, and Y. Sirois, "Electron Reconstruction in CMS," *CMS Note* **2006/040** (2006).
- [38] S. Abdullin et al., "Study of PDF and QCD scale uncertainties in $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ events at the LHC," *CMS Note* **2006/068** (2006). See also hep-ph/0604120, Les Houches Physics at TeV Colliders 2005, Standard Model and Higgs working group: Summary report. April 2006.
- [39] P. Meridiani and R. Paramatti, "Use of $Z \rightarrow e^+e^-$ events for ECAL calibration," *CMS Note* **2006/039** (2006).
- [40] V. Bartsch and G. Quast, "Expected signal observability at future experiments," *CMS Note* **2005-004** (2006).
- [41] S. Bitjukov and N. Krasnikov, "New physics discovery potential in future experiments," *Mod. Phys. Lett.* **A13** (1998) 3235–3249.
- [42] S. Abdullin et al., "Search for $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ Using $M(4\mu)$ -Dependent Cuts," *CMS Note* **2006-004**: in preparation (2006).
- [43] V. Karimaki et al., "CMKIN v3 User's Guide," *CMS IN* **2004-016** (2004).


- [44] M. Spira, "SM Higgs boson production cross sections at NLO' and 'SM Higgs boson BRs'." Available at <http://cmsdoc.cern.ch/~anikiten/cms-higgs/>, 2005.
- [45] **Particle Data Group** Collaboration, S. Eidelman et al., "Review of particle physics," *Phys. Lett.* **B592** (2004) 1. doi:10.1016/j.physletb.2004.06.001.
- [46] F. Maltoni, "Theoretical Issues and Aims at the Tevatron and LHC," in *Proceedings of the 1st Hadron Collider Physics Symposium (HCP 2005)*. Les Diablerets, Switzerland, July, 2005. Available at <http://hcp-2005.web.cern.ch/HCP-2005>.
- [47] A. Pukhov et al., "CompHEP: A package for evaluation of Feynman diagrams and integration over multi-particle phase space. User's manual for version 33," arXiv:hep-ph/9908288.
- [48] J. Campbell, "W/Z + B anti-B / jets at NLO using the Monte Carlo MCFM," arXiv:hep-ph/0105226.
- [49] S. Abdullin et al., "Relative Contributions of t- and s-Channels to the $ZZ \rightarrow 4\mu$ Process," *CMS Note* **2006/057** (2006). See also hep-ph/0604120, Les Houches Physics at TeV Colliders 2005, Standard Model and Higgs working group: Summary report. April 2006.
- [50] P. Bartalini et al., "NLO vs. LO: kinematic differences for signal and background in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ analysis," *CMS Note* **2006/058** (2006). **NAME: in preparation**
- [51] D. Acosta et al., "Measuring Muon Reconstruction Efficiency from Data," *CMS Note* **2006/060** (2006).
- [52] M. Aldaya et al., "Search for the Standard Model Higgs boson in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ decay channel using a mass-independent analysis," *CMS Note* **2006/061** (2006). **NAME: in preparation**
- [53] A. Read, "Modified Frequentist Analysis of Search Results (The CLs Method)," *1st Workshop on Confidence Limits, CERN, Geneva* (2000). CERN-EP/2000-005.
- [54] **LEP Working Group for Higgs boson searches** Collaboration, R. Barate et al., "Search for the standard model Higgs boson at LEP," *Phys. Lett.* **B565** (2003) 61–75, arXiv:hep-ex/0306033.
- [55] S. Abdullin, A. Drozdetskiy, et al., "GARCON: Genetic Algorithm for Rectangular Cuts Optimization. User's manual for version 2.0," arXiv:hep-ph/0605143. See also <http://drozdets.home.cern.ch/drozdets/home/genetic/>.
- [56] S. Abdullin et al., "Sensitivity of the Muon Isolation Cut Efficiency to the Underlying Event Uncertainties," *CMS Note* **2006/033** (2006). See also hep-ph/0604120, Les Houches Physics at TeV Colliders 2005, Standard Model and Higgs working group: Summary report. April 2006.
- [57] M. Aldaya et al., "Determination of the mass, cross-section and width of the Standard Model Higgs boson using the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ decay channel," *CMS Note* **2006/062** (2006). **NAME: in preparation**
- [58] V. Bartsch, "Simulation of Silicon Sensors and Study of the Higgs Decay," *CMS Note* **2006/063** (2006). **NAME: need ref,**

- [59] M. Dittmar and H. Dreiner, "How to find a Higgs boson with a mass between 155 GeV to to 180 GeV at the CERN LHC," *Phys. Rev. D* **55** (1997) 167–172, arXiv:hep-ph/9608317. doi:10.1103/PhysRevD.55.167.
- [60] M. Dittmar and H. Dreiner, "LHC Higgs Search with $l^+\nu l^-\bar{\nu}$ final states," *CMS Note* **1997/083** (1997).
- [61] CMS Collaboration, "The CMS Physics Technical Design Report, Volume 1," CERN/LHCC **2006-001** (2006).
- [62] T. Sjostrand et al., "High-energy-physics event generation with PYTHIA 6.1," *Comput. Phys. Commun.* **135** (2001) 238–259. doi:10.1016/S0010-4655(00)00236-8.
- [63] T. Binoth, M. Ciccolini, N. Kauer, and M. Kramer, "Gluon-induced $W W$ background to Higgs boson searches at the LHC," *JHEP* **03** (2005) 065, arXiv:hep-ph/0503094.
- [64] G. Davatz et al., "Effective k-factors for $H \rightarrow WW^{(*)} \rightarrow 2\mu 2\nu$ at the LHC," *JHEP* **0405** (2004) 009, arXiv:hep-ph/0402218. doi:10.1088/1126-6708/2004/05/009.
- [65] S. Frixione and B. Webber, "Matching NLO QCD computations and parton shower simulations," *JHEP* **0206** (2002) 029, arXiv:hep-ph/0204244.
- [66] G. Davatz, M. Dittmar, and A. Giolo-Nicollerat, "Standard Model Higgs Discovery Potential of CMS in the $H \rightarrow WW^{(*)} \rightarrow l\nu l\nu$ Channel," *CMS Note* **2006/047** (2006).
- [67] J. Campbell and F. Tramontano, "Next-to-leading order corrections to Wt production and decay," *Nucl. Phys.* **B726** (2005) 109–130. doi:10.1016/j.nuclphysb.2005.08.015.
- [68] CMS Collaboration, "The TriDAS Project Technical Design Report, Volume 2: Data Acquisition and High-Level Trigger," CERN/LHCC **2002-26** (2002). CMS TDR 6.2.
- [69] V. Drollinger, U. Gasparini, E. Torassa, and M. Zanetti, "Physics study of the Higgs decay channel $H \rightarrow WW^{(*)} \rightarrow 2\mu 2\nu$," *CMS Note* **2006/055** (2006).
- [70] S. V. Chekanov, "Jet algorithms: a mini review," in *Proceedings of the 14th Topical Conference on Hadron Collider Physics (HCP 2002)*. Karlsruhe, Germany, Sep-Oct, 2002. arXiv:hep-ph/0211298.
- [71] S. Bityukov and N. Krasnikov, "Uncertainties and discovery potential in planned experiments," *CMS CR* **2002/05** (2002).
- [72] G. Davatz, A. Giolo-Nicollerat, and M. Zanetti, "Systematics uncertainties of the top background in the $H \rightarrow WW$ channel," *CMS Note* **2006/048** (2006).
- [73] F. Maltoni and T. Stelzer, "MadEvent: Automatic event generation with MadGraph," *JHEP* **02** (2003) 027. doi:10.1088/1126-6708/2003/02/027.
- [74] N. Kauer, "Top background extrapolation for $H \rightarrow WW$ searches at the LHC," arXiv:hep-ph/0404045.
- [75] J. D'Hont, J. Heynink, S. Lowette, and P. Vanlear, "Offline Calibration of b-Jet Identification Efficiency," *CMS Note* **2006/013** (2006).

- [76] V. Drollinger et al., "Modeling the production of W pairs at the LHC," *CMS Note* **2005/024** (2005).
- [77] C. Anastasiou et al., "High precision QCD at hadron colliders: Electroweak gauge boson rapidity distribution at NNLO," *Phys. Rev.* **D 69** (2004).
- [78] I. Belotelov et al., "Simulation of misalignment scenarios for CMS tracking devices," *CMS Note* **2006-008** (2006).
- [79] S. Bityukov, "Uncertainty, Systematics, Limits." Available at <http://cmsdoc.cern.ch/bityukov>, 2005.
- [80] M. Cvetič and P. Langacker, "Implications of Abelian Extended Gauge Structures From String Models," *Phys. Rev.* **D54** (1996) 3570–3579, arXiv:hep-ph/9511378.
- [81] M. Cvetič and P. Langacker, "New Gauge Bosons from String Models," *Mod. Phys. Lett.* **A11** (1996) 1247–1262, arXiv:hep-ph/9602424.
- [82] A. Leike, "The phenomenology of extra neutral gauge bosons," *Phys. Rept.* **317** (1999) 143–250, arXiv:hep-ph/9805494.
- [83] C. T. Hill and E. H. Simmons, "Strong dynamics and electroweak symmetry breaking," *Phys. Rept.* **381** (2003) 235–402, arXiv:hep-ph/0203079.
- [84] T. Han, H. Logan, B. McElrath, and L.-T. Wang, "Phenomenology of the little Higgs model," *Phys. Rev.* **D67** (2003) 095004. doi:10.1103/PhysRevD.67.095004.
- [85] M. Cvetič and S. Godfrey, "Discovery and identification of extra gauge bosons," in *Electroweak symmetry breaking and new physics at the TeV scale*, T. Barlow and S. Dawson, eds., pp. 383–415. World Scientific, 1996. arXiv:hep-ph/9504216.
- [86] S. Godfrey, "Update of discovery limits for extra neutral gauge bosons at hadron colliders," in *Proceedings of Snowmass 2001*, p. P344. Snowmass, Colorado, June–July, 2001. arXiv:hep-ph/0201093.
- [87] L. Randall and R. Sundrum, "A large mass hierarchy from a small extra dimension," *Phys. Rev. Lett.* **83** (1999) 3370–3373. doi:10.1103/PhysRevLett.83.3370.
- [88] M. R. Whalley, D. Bourilkov, and R. C. Group, "The Les Houches accord PDFs (LHAPDF) and LHAGLUE," arXiv:hep-ph/0508110.
- [89] J. L. Rosner, "Off peak lepton asymmetries from new Zs," *Phys. Rev.* **D35** (1987) 2244.
- [90] J. L. Rosner, "Forward-Backward Asymmetries in Hadronically Produced Lepton Pairs," *Phys. Rev.* **D54** (1996) 1078–1082, arXiv:hep-ph/9512299.
- [91] R. N. Mohapatra, S. A. Goldstein, and D. Money, eds., "Unification and supersymmetry: The Frontiers of quark - lepton physics". Springer-Verlag, 2002. 3rd Ed.
- [92] M. Schmitt, "Private communication."
- [93] I. Belotelov et al., "Influence of misalignment scenarios on muon reconstruction," *CMS Note* **2006-017** (2006).

- [94] R. Cousins, J. Mumford, and S. Valuev, "Detection of Z' gauge bosons in the dimuon decay mode in CMS," *CMS Note* **2005-002** (2005).
- [95] R. Cousins, J. Mumford, and V. Valuev, "Detection of Z' gauge bosons in the dimuon decay mode in CMS," *CMS CR* **2004/050** (2006).
- [96] V. Bartsch and G. Quast, "Expected signal observability at future experiments," *CMS Note* **2005-004** (2005).
- [97] S. Wilks, "The large-sample distribution of the likelihood ratio for testing composite hypotheses," *Annals of Math. Stat.* **9** (1938) 60.
- [98] U. Baur, O. Brein, W. Hollik, C. Schappacher, and D. Wackerroth, "Electroweak radiative corrections to neutral-current Drell-Yan processes at hadron colliders," *Phys. Rev.* **D65** (2002) 033007, arXiv:hep-ph/0108274.
- [99] V. Zykunov, "Weak radiative corrections to Drell-Yan process for large invariant mass of di-lepton pair," arXiv:hep-ph/0509315.
- [100] A. De Roeck and S. Slabospitsky, "Evaluation of the PDF Uncertainties (in CMS)." Available at <http://cmsdoc.cern.ch/cms/PRS/gentools/www/pdfuncert/uncert.html>.
- [101] G. Cerminara. Private communication.
- [102] P. Langacker, R. W. Robinett, and J. L. Rosner, "New heavy gauge bosons in pp and $p\bar{p}$ collisions," *Phys. Rev.* **D30** (1984) 1470.
- [103] J. L. Rosner, "Observability of charge asymmetries for lepton pairs produced in present collider experiments," *Phys. Lett.* **B221** (1989) 85.
- [104] T. P. Cheng and L. F. Li, "Gauge theory of elementary particle physics". Oxford University Press, 1988.
- [105] R. Cousins, J. Mumford, and V. Valuev, "Measurement of Forward-Backward Asymmetry of Simulated and Reconstructed $Z' \rightarrow \mu^+ \mu^-$ Events in CMS," *CMS Note* **2005-022** (2005).
- [106] M. Dittmar, "Neutral current interference in the TeV region: The experimental sensitivity at the LHC," *Phys. Rev.* **D55** (1997) 161–166, arXiv:hep-ex/9606002.
- [107] C.-E. Wulz, " Z' at LHC," in *Proceedings of the 1996 DPF/DPB Summer Study on New Directions in High-Energy Physics*. 1996. Available at <http://www.slac.stanford.edu/pubs/snowmass96>.
- [108] J. C. Collins and D. E. Soper, "Angular distribution of dileptons in high-energy hadron collisions," *Phys. Rev.* **D16** (1977) 2219.
- [109] B. C. Allanach, K. Odagiri, M. A. Parker, and B. R. Webber, "Searching for narrow graviton resonances with the ATLAS detector at the Large Hadron Collider," *JHEP* **09** (2000) 019, arXiv:hep-ph/0006114.
- [110] R. Cousins, J. Mumford, J. Tucker, and V. Valuev, "Spin discrimination of new heavy resonances at the LHC," *JHEP* **11** (2005) 046. doi:10.1088/1126-6708/2005/11/046.


- [111] I. Belotel  al., “Search for Randall-Sandrum Graviton decay into muon pairs,” *CMS AN* **2006/041** (2006).
- [112] S. Esen and R. M. Harris, “Jet Triggers and Dijet Mass,” *CMS Note* **2006/069** (2006).
- [113] **CMS Collaboration**, D. Acosta et al., “CMS Physics TDR Volume 1, Section 11.3: Monte Carlo Corrections,” *CERN/LHCC* **2006-001** (2006) 409.
- [114] **CDF Collaboration**, F. Abe et al., “Search for new particles decaying to dijets at CDF,” *Phys. Rev.* **D55** (1997) 5263–5268. doi:10.1103/PhysRevD.55.R5263.
- [115] **DØ Collaboration**, V. M. Abazov et al., “Search for new particles in the two-jet decay channel with the DØ detector,” *Phys. Rev.* **D69** (2004) 111101, arXiv:hep-ex/0308033.
- [116] **DØ Collaboration**, B. Abbott et al., “The dijet mass spectrum and a search for quark compositeness in $\bar{p}p$ collisions at $\sqrt{s} = 1.8\text{--}TeV$,” *Phys. Rev. Lett.* **82** (1999) 2457–2462. doi:10.1103/PhysRevLett.82.2457.
- [117] E. Eichten, K. D. Lane, and M. E. Peskin, “New Tests for Quark and Lepton Substructure,” *Phys. Rev. Lett.* **50** (1983) 811–814. doi:10.1103/PhysRevLett.50.811.
- [118] K. D. Lane, “Electroweak and flavor dynamics at hadron colliders,” arXiv:hep-ph/9605257.
- [119] **CDF Collaboration**, F. Abe et al., “Measurement of dijet angular distributions at CDF,” *Phys. Rev. Lett.* **77** (1996) 5336–5341. doi:10.1103/PhysRevLett.77.5336.
- [120] **CMS Collaboration**, D. Acosta et al., “CMS Physics TDR Volume 1, Section 11.6.1: Data-driven calibration strategy,” *CERN/LHCC* **2006-001** (2006) 421.
- [121] **CMS Collaboration**, D. Acosta et al., “CMS Physics TDR Volume 1, Section 11.6.2: Dijet Balancing,” *CERN/LHCC* **2006-001** (2006) 422.
- [122] **CDF Collaboration**, D. Acosta et al., “Measurement of the Lifetime Difference between B_s Mass Eigenstates,” *Phys. Rev. Lett.* **94** (2005) 101803, arXiv:hep-ex/0412057. doi:10.1103/PhysRevLett.94.101803.
- [123] **DØ Collaboration**, “Measurement of the Lifetime Difference in the B_s System,” *DØ Conference note* 5052 (2006).
- [124] A. Belkov and S. Shulga, “Studies of angular correlations in the decays $B_s^0 \rightarrow J/\psi\phi$ by using the SIMUB generator,” *Comput. Phys. Commun.* **156** (2004) 221–240, arXiv:hep-ph/0310096. doi:10.1016/S0010-4655(03)00465-X.
- [125] A. S. Dighe, I. Dunietz, and R. Fleischer, “Extracting CKM phases and $B_s - \bar{B}_s$ mixing parameters from angular distributions of non-leptonic B decays,” *Eur. Phys. J.* **C6** (1999) 647–662. doi:10.1007/s100529800954.
- [126] A. S. Dighe, I. Dunietz, H. J. Lipkin, and J. L. Rosner, “Angular distributions and lifetime differences in $B_s \rightarrow J/\psi\phi$ decays,” *Phys. Lett.* **B369** (1996) 144–150. doi:10.1016/0370-2693(95)01523-X.


- [127] CDF Collaboration, F. Abe et al., "Production of J/ψ mesons from χ_c meson decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **79** (1997) 578–583.
doi:10.1103/PhysRevLett.79.578.
- [128] B. Cano-Coloma and M. A. Sanchis-Lozano, "Charmonia production in hadron colliders and the extraction of colour-octet matrix elements," *Nucl. Phys.* **B508** (1997) 753–767, arXiv:hep-ph/9706270. doi:10.1016/S0550-3213(97)00660-3.
- [129] S. Cucciarelli, M. Konecki, D. Kotlinski, and T. Todorov, "Track reconstruction, primary vertex finding and seed generation with the Pixel Detector," *CMS Note* **2006/026** (2006).
- [130] K. Prokofiev and T. Speer, "A kinematic fit and a decay chain reconstruction library," *CERN Yellow Report* **2005-002** (2005). Proc. of the 2004 Conference for Computing in High-Energy and Nuclear Physics (CHEP 04), Interlaken, Switzerland, 2004, available at <http://indico.cern.ch>.
- [131] I. Dunietz, R. Fleischer, and U. Nierste, "In pursuit of new physics with B_s decays," *Phys. Rev.* **D63** (2001) 114015. doi:10.1103/PhysRevD.63.114015.
- [132] BABAR Collaboration, B. Aubert et al., "Ambiguity-free measurement of $\cos(2\beta)$: Time-integrated and time-dependent angular analyses of $B \rightarrow J/\psi K\pi$," *Phys. Rev.* **D71** (2005) 032005. doi:10.1103/PhysRevD.71.032005.
- [133] I. Dunietz, H. Quinn, A. Snyder, W. Toki, and H. J. Lipkin, "How to extract CP-violating asymmetries from angular correlations," *Phys. Rev.* **D43** (1991) 2193–2208. doi:10.1103/PhysRevD.43.2193.
- [134] P. Vanlaer et al., "Impact of CMS Silicon Tracker Misalignment on Track and Vertex Reconstruction," *CMS Note* **2006/029** (2006).
- [135] T. Speer et al., "Vertex Fitting in the CMS Tracker," *CMS Note* **2006/032** (2006).
- [136] S. Heinemeyer, W. Hollik, and G. Weiglein, "FeynHiggs: A program for the calculation of the masses of the neutral CP-even Higgs bosons in the MSSM," *Comput. Phys. Commun.* **124** (2000) 76–89, arXiv:hep-ph/9812320.
- [137] S. Heinemeyer, W. Hollik, and G. Weiglein, "The masses of the neutral CP-even Higgs bosons in the MSSM: Accurate analysis at the two-loop level," *Eur. Phys. J.* **C9** (1999) 343–366, arXiv:hep-ph/9812472.
- [138] G. Degrossi, S. Heinemeyer, W. Hollik, P. Slavich, and G. Weiglein, "Towards high-precision predictions for the MSSM Higgs sector," *Eur. Phys. J.* **C28** (2003) 133–143, arXiv:hep-ph/0212020.
- [139] S. Lehti, "Study of bbZ as a benchmark for MSSM bbH ," *CMS AN 2005/068*  **NAME: NEEDS TO BE CMS-NOTE** (2005).
- [140] G. Bagliesi, S. Dutta, S. Gennai, A. Kalinowski, M. Konecki, D. Kotlinski, F. Moortgat, A. Nikitenko, W. L., and S. Wakefield, "Tau jet reconstruction and tagging at High Level Trigger and off-line," *CMS Note* **2006/028** (2006).
- [141] S. Kunori, R. Kinnunen, and A. Nikitenko, "Missing Transverse Energy Measurement with Jet Energy Corrections," *CMS Note* **2001/040** (2001).

- [142] H. Pi et al., "Measurement of Missing Transverse Energy with the CMS detector at the LHC," *CMS Note* **2006/035** (2006).
- [143] A. Kalinowski and A. Nikitenko, "Measurement of the b -tagging efficiency using the $Z \rightarrow \tau\tau \rightarrow \mu + \text{hadrons} + X$ events," *CMS AN 2006/002* **NAME: NEEDS TO BE CMS-NOTE** (2006).
- [144] A. Rizzi, F. Palla, and G. Segneri, "Track impact parameter based b -tagging with CMS," *CMS Note* **2006/019** (2006).
- [145] A. Kalinowski, M. Konecki, and D. Kotliński, "Search for M_{Higgs} heavy neutral Higgs boson in $\tau + \tau \rightarrow \mu + \text{jet}$ decay mode," *CMS AN 2006/022* **NAME: NEEDS TO BE CMS-NOTE** (2005).
- [146] S. Jadach, Z. Was, R. Decker, and J. H. Kuhn, "The tau decay library TAUOLA: Version 2.4," *Comput. Phys. Commun.* **76** (1993) 361–380.
doi:10.1016/0010-4655(93)90061-G.
- [147] A.-S. Giolo-Nicollerat *CMS Note* **NAME: under preparation**.
- [148] C. Weiser, "A Combined Secondary Vertex Based B-Tagging Algorithm in CMS," *CMS Note* **2006/014** (2006).
- [149] S. Haywood et al., "Electroweak physics," arXiv:hep-ph/0003275. In 'Standard Model Physics (and more) at the LHC', Geneva, 1999.
- [150] D. Chakraborty, J. Konigsberg, and D. L. Rainwater, "Review of top quark physics," *Ann. Rev. Nucl. Part. Sci.* **53** (2003) 301–351, arXiv:hep-ph/0303092.
- [151] D. Benedetti, S. Cucciarelli, C. Hill, J. Incandela, S. Koay, C. Riccardi, A. Santocchia, A. Schmidt, P. Torre, and C. Weiser, "Search for $H \rightarrow bb$ in association with a $t\bar{t}$ pair at CMS," *CMS Analysis Note* **NAME: AN 2006/093** (2006).
- [152] M. L. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A. D. Polosa, "ALPGEN, a generator for hard multiparton processes in hadronic collisions," *JHEP* **07** (2003) 001, arXiv:hep-ph/0206293. doi:10.1088/1126-6708/2003/07/001.
- [153] W. Beenakker, S. Dittmaier, M. Kraemer, B. Pluemper, M. Spira, and P. Zerwas, "NLO QCD corrections to $t\bar{t}H$ production in hadron collisions," *Nucl. Phys.* **B653** (2003) 151–203, arXiv:hep-ph/0211352.
- [154] V. Roberfroid et al., "Validation and status of the HLT Steering Code and default menu," *CMS IN* **2006/001** (2006).
- [155] E. James, Y. Maravin, M. Mulders, and N. Neumeister, "Muon Identification in CMS," *CMS Note* **2006/010** (2006).
- [156] A. Heister, O. Kodolova, V. Konopliankov, S. Petrushanko, J. Rohlf, C. Tully, and A. Ulyanov, "Measurement of Jets with the CMS Detector at the LHC," *CMS Note* **2006/036** (2006).
- [157] A. Santocchia, "Optimization of Jet Reconstruction Settings and Parton-Level Correction for the $t\bar{t}H$ Channel," *CMS Analysis Note* **NAME: Needs to be note 2006/030** (2006).



- [158] J. D'Hondt, S. Lowette, O. Buchmuller, S. Cucciarelli, F. Schilling, M. Spiropulu, S. Mehdiabadi, D. Benedetti, and L. Pape, "Fitting of Event Topologies with External Kinematic Constraints in CMS," *CMS Note* **2006/023** (2006).
- [159] M. Mangano. Private communication.
- [160] X.-N. Wang and M. Gyulassy, "HIJING: A Monte Carlo model for multiple jet production in p p, p A and A A collisions," *Phys. Rev.* **D44** (1991) 3501–3516.
- [161] M. Bedjidian et al., "Hard probes in heavy ion collisions at the LHC: Heavy flavour physics," arXiv:hep-ph/0311048.
- [162] K. Eskola, V. Kolhinen, and C. Salgado, "The scale dependent nuclear effects in parton distributions for practical applications," *Eur. Phys. J.* **C9** (1999) 61–68, arXiv:hep-ph/9807297.
- [163] O. Kodolova, M. Bedjidian, and S. Petrouchanko, "Dimuon reconstruction in heavy ion collisions using a detailed description of CMS geometry," *CMS Note* **1999-004** (1999).
- [164] M. Bedjidian and O. Kodolova, "Quarkonia measurements in heavy-ion collisions in CMS," *CMS Analysis Note* **2006-011** (2006).
- [165] G. Sterman and S. Weinberg, "Jets from Quantum Chromodynamics," *Phys. Rev. Lett.* **39** (1977) 1436.
- [166] JADE Collaboration, W. Bartel et al., "Experimental Studies on Multi-Jet Production in e+ e- Annihilation at Petra Energies," *Z. Phys.* **C33** (1986) 23.
- [167] JADE Collaboration, S. Bethke et al., "Experimental Investigation of the Energy Dependence of the Strong Coupling Strength," *Phys. Lett.* **B213** (1988) 235.
- [168] S. Catani, Y. L. Dokshitzer, M. Olsson, G. Turnock, and B. R. Webber, "New clustering algorithm for multi-jet cross-sections in e+e- annihilation," *Phys. Lett.* **B269** (1991) 432–438.
- [169] CDF Run II Collaboration, A. Abulencia et al., "Measurement of the inclusive jet cross section in $p\bar{p}$ interactions at $\sqrt{s} = 1.96$ TeV using a cone-based jet algorithm," arXiv:hep-ex/0512020.
- [170] DØ Collaboration, R. Strohmer, "Inclusive jet cross-sections and dijet azimuthal decorrelations with DØ," *PoS HEP2005* (2006) 051, arXiv:hep-ex/0601016.
- [171] J. Butterworth, J. Couchman, B. Cox, and B. Waugh, "KtJet: A C++ implementation of the K(T) clustering algorithm," *Comput. Phys. Commun.* **153** (2003) 85–96, arXiv:hep-ph/0210022.
- [172] U. Baur, R. K. Ellis, and D. Zeppenfeld, eds., "QCD and Weak Boson Physics in Run II". FERMILAB, Batavia, IL, USA, 2000. Prepared for Physics at Run II: QCD and Weak Boson Physics Workshop: Final General Meeting, Batavia, Illinois, 4-6 Nov 1999.
- [173] S. D. Ellis, J. Huston, and M. Tonnesmann, "On building better cone jet algorithms," *eConf* **C010630** (2001) P513, arXiv:hep-ph/0111434.

- [174] M. Wobisch, "Private communication."
- [175] T. Sjostrand, "High-energy physics event generation with PYTHIA 5.7 and JETSET 7.4," *Comput. Phys. Commun.* **82** (1994) 74–90.
- [176] Z. Nagy, "Three-jet cross sections in hadron hadron collisions at next-to-leading order," *Phys. Rev. Lett.* **88** (2002) 122003, arXiv:hep-ph/0110315.
- [177] T. Kluge, K. Rabbertz, and M. Wobisch, "fast NLO - fast pQCD calculations for hadron-induced processes." Available at <http://hepforge.cedar.ac.uk/fastnlo>. To be published.
- [178] S. Moretti, M. R. Nolten, and D. A. Ross, "Weak corrections and high E(T) jets at Tevatron," arXiv:hep-ph/0503152.
- [179] DØ Collaboration, V. M. Abazov et al., "The inclusive jet cross-section in p anti-p collisions at $s^{1/2} = 1.8$ -TeV using the k(T) algorithm," *Phys. Lett.* **B525** (2002) 211–218, arXiv:hep-ex/0109041.
- [180] DØ Collaboration, B. Abbott et al., "High- p_T jets in $\bar{p}p$ collisions at $\sqrt{s} = 630$ GeV and 1800 GeV," *Phys. Rev.* **D64** (2001) 032003. doi:10.1103/PhysRevD.64.032003.
- [181] CDF Collaboration, A. A. Affolder et al., "Charged jet evolution and the underlying event in proton anti-proton collisions at 1.8-TeV," *Phys. Rev.* **D65** (2002) 092002.
- [182] CDF Collaboration, D. Acosta et al., "The underlying event in hard interactions at the Tevatron anti-p p collider," *Phys. Rev.* **D70** (2004) 072002, arXiv:hep-ex/0404004.
- [183] T. Sjostrand and M. van Zijl, "A Multiple Interaction Model for the Event Structure in Hadron Collisions," *Phys. Rev.* **D36** (1987) 2019.
- [184] J. Butterworth, J. R. Forshaw, and M. Seymour, "Multiparton interactions in photoproduction at HERA," *Z. Phys.* **C72** (1996) 637–646, arXiv:hep-ph/9601371.
- [185] T. Gleisberg et al., "SHERPA 1.alpha, a proof-of-concept version," *JHEP* **02** (2004) 056, arXiv:hep-ph/0311263.
- [186] F. W. Bopp, R. Engel, and J. Ranft, "Rapidity gaps and the PHOJET Monte Carlo," arXiv:hep-ph/9803437.
- [187] G. Corcella et al., "HERWIG 6: An event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)," *JHEP* **01** (2001) 010, arXiv:hep-ph/0011363. doi:10.1088/1126-6708/2001/01/010.
- [188] C. M. Buttar, D. Clements, I. Dawson, and A. Moraes, "Simulations of minimum bias events and the underlying event, MC tuning and predictions for the LHC," *Acta Phys. Polon.* **B35** (2004) 433–441.
- [189] CDF Collaboration, R. Field, "Min-bias and the underlying event in Run 2 at CDF," *Acta Phys. Polon.* **B36** (2005) 167–178.
- [190] D. Acosta et al., "The Underlying Event at the LHC," *CMS Note* (2006), no. 2006/067.

- [191] CDF Collaboration, F. Abe et al., "Measurement of the Z -boson p_T distribution in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **67** (1991) 2937–2941. doi:10.1103/PhysRevLett.67.2937.
- [192] P. Bartalini, R. Chierici, and A. De Roeck, "Guidelines for the estimation of theoretical uncertainties at the LHC," *CMS Note* **2005/013** (2005).
- [193] W. Adam, B. Mangano, T. Speer, and T. Todorov, "Track reconstruction in the CMS tracker," *CMS Note* **2006/041** (2006).
- [194] V. Andreev, D. Cline, and S. Otwinowski *CMS Note* **2006/000**  **NAME: need real number** (2006).
- [195] CDF Collaboration, F. Abe et al., "Measurement of the bottom quark production cross section using semileptonic decay electrons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **71** (1993) 500. doi:10.1103/PhysRevLett.71.500.
- [196] CDF Collaboration, F. Abe et al., "Measurement of the B meson and b quark cross sections at $\sqrt{s} = 1.8$ TeV using the exclusive decay $B^0 \rightarrow J/\psi K^*(892)^0$," *Phys. Rev. D* **50** (1994) 4252. doi:10.1103/PhysRevD.50.4252.
- [197] CDF Collaboration, F. Abe et al., "Measurement of the B Meson Differential Cross Section $d\sigma/dp_T$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **75** (1995) 1451. doi:10.1103/PhysRevLett.75.1451.
- [198] CDF Collaboration, F. Abe et al., "Measurement of the B^+ total cross section and B^+ differential cross section $d\sigma/dp_T$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. D* **65** (2002) 052005. doi:10.1103/PhysRevD.65.052005.
- [199] CDF Collaboration, F. Abe et al., "Measurement of the ratio of b quark production cross sections in $\bar{p}p$ collisions at $\sqrt{s} = 630$ GeV and $\sqrt{s} = 1800$ GeV," *Phys. Rev. D* **66** (2002) 032002. doi:10.1103/PhysRevD.66.032002.
- [200] DØ Collaboration, S. Abachi et al., "Inclusive μ and b -quark production cross-sections in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **74** (1995) 3548. doi:10.1103/PhysRevLett.74.3548.
- [201] DØ Collaboration, B. Abbott et al., "The $b\bar{b}$ production cross section and angular correlations in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Lett.* **B487** (2000) 264. doi:10.1016/S0370-2693(00)00844-3.
- [202] DØ Collaboration, B. Abbott et al., "Cross section for b -Jet production in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **85** (2000) 5068. doi:10.1103/PhysRevLett.85.5068.
- [203] H1 Collaboration, C. Adloff et al., "Measurement of open beauty production at HERA," *Phys. Lett.* **B 467** (1999) 156. doi:10.1016/S0370-2693(99)01099-0.
- [204] H1 Collaboration, C. Adloff et al., "Erratum," *Phys. Lett.* **B 518** (2001) 331.
- [205] ZEUS Collaboration, J. Breitweg et al., "Measurement of open beauty production in photoproduction at HERA," *Eur. Phys. J.* **C18** (2001) 625–637. doi:10.1007/s100520100571.


- [206] H1 Collaboration, A. Aktas et al., "Measurement of F2(c anti-c) and F2(b anti-b) at high Q^2 using the H1 vertex detector at HERA," *Eur. Phys. J.* **C40** (2005) 349–359, arXiv:hep-ex/0411046.
- [207] ZEUS Collaboration, S. Chekanov et al., "Bottom photoproduction measured using decays into muons in dijet events in e p collisions at $s^{1/2} = 318$ GeV," *Phys. Rev.* **D70** (2004) 012008.
- [208] ZEUS Collaboration, S. Chekanov et al., "Measurement of beauty production in deep inelastic scattering at HERA," *Phys. Lett.* **B599** (2004) 173–189, arXiv:hep-ex/0405069.
- [209] H1 Collaboration, A. Aktas et al., "Measurement of beauty production at HERA using events with muons and jets," *Eur. Phys. J.* **C41** (2005) 453–467, arXiv:hep-ex/0502010.
- [210] L3 Collaboration, M. Acciarri et al., "Measurements of the cross sections for open charm and beauty production in $\gamma\gamma$ collisions at $\sqrt{s} = 189$ GeV - 202 GeV," *Phys. Lett.* **B503** (2001) 10. doi:10.1016/S0370-2693(01)00134-4.
- [211] L3 Collaboration, P. Achard et al., "Measurement of the cross section for open-beauty production in photon photon collisions at LEP," *Phys. Lett.* **B619** (2005) 71.
- [212] P. Nason, S. Dawson, and R. K. Ellis, "The Total Cross-Section for the Production of Heavy Quarks in Hadronic Collisions," *Nucl. Phys.* **B303** (1988) 607.
- [213] P. Nason, S. Dawson, and R. K. Ellis, "The One Particle Inclusive Differential Cross-Section for Heavy Quark Production in Hadronic Collisions," *Nucl. Phys.* **B327** (1989) 49–92.
- [214] W. Beenakker, W. Van Neerven, R. Meng, G. Schuler, and J. Smith, "QCD corrections to heavy quark production in hadron-hadron collisions," *Nucl. Phys.* **B351** (1991) 507–560. doi:10.1016/S0550-3213(05)80032-X.
- [215] M. Cacciari, S. Frixione, M. L. Mangano, P. Nason, and G. Ridolfi, "QCD analysis of first b cross section data at 1.96-TeV," *JHEP* **07** (2004) 033. See also included references. doi:10.1088/1126-6708/2004/07/033.
- [216] M. L. Mangano, "The saga of bottom production in proton antiproton collisions," *AIP Conf. Proc.* **753** (2005) 247–260, arXiv:hep-ph/0411020. See also included references.
- [217] S. Frixione, "Bottom production," 2004.
- [218] J. Rohlf and C. Tully *CMS Internal Note*  **NOTE: Needs to be a Note/No. 2006/000** (2006).
- [219] S. Weinberg, "Unitarity Constraints on CP Nonconservation in Higgs Exchange," *Phys. Rev.* **D42** (1990) 860–866.
- [220] L. Lavoura, "Maximal CP violation via Higgs boson exchange," *Int. J. Mod. Phys.* **A8** (1993) 375–390.



- [221] E. Jenkins, M. E. Luke, A. V. Manohar, and M. J. Savage, "Semileptonic B_c decay and heavy quark spin symmetry," *Nucl. Phys.* **B390** (1993) 463–473, arXiv:hep-ph/9204238.
- [222] V. V. Kiselev, A. E. Kovalsky, and A. K. Likhoded, " B_c decays and lifetime in QCD sum rules," *Nucl. Phys.* **B585** (2000) 353–382. doi:10.1016/S0550-3213(00)00386-2.
- [223] X.-G. Wu, C.-H. Chang, Y.-Q. Chen, and Z.-Y. Fang, "The meson B_c annihilation to leptons and inclusive light hadrons," *Phys. Rev.* **D67** (2003) 094001. doi:10.1103/PhysRevD.67.094001.
- [224] C.-H. Chang and Y.-Q. Chen, "The Production of B_c or \bar{B}_c meson associated with two heavy quark jets in Z^0 boson decay," *Phys. Rev.* **D46** (1992) 3845–3855. doi:10.1103/PhysRevD.46.3845.
- [225] C.-H. Chang and Y.-Q. Chen, "Hadronic production of the B_c meson at TeV energies," *Phys. Rev. D* **48** (1993) 4086. doi:10.1103/PhysRevD.48.4086.
- [226] C.-H. Chang, Y.-Q. Chen, G.-P. Han, and H.-T. Jiang, "On hadronic production of the B_c meson," *Phys. Lett. B* **364** (1995) 78. doi:10.1016/0370-2693(95)01235-4.
- [227] C.-H. Chang, Y.-Q. Chen, and R. Oakes, "Comparative study of the hadronic production of B_c mesons," *Phys. Rev. D* **54** (1996) 4344. doi:10.1103/PhysRevD.54.4344.
- [228] C.-H. Chang and X.-G. Wu, "Uncertainties in estimating hadronic production of the meson B/c and comparisons between TEVATRON and LHC," *Eur. Phys. J.* **C38** (2004) 267–276. doi:10.1140/epjc/s2004-02015-0.
- [229] CDF Collaboration, F. Abe et al., "Observation of the B_c meson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **81** (1998) 2432–2437. doi:10.1103/PhysRevLett.81.2432.
- [230] CDF Collaboration, D. Acosta et al., "Evidence for the exclusive decay $B_c^\pm \rightarrow J/\psi\pi^\pm$ and measurement of the mass of the B_c meson," *Phys. Rev. Lett.* **96** (2006) 082002. doi:10.1103/PhysRevLett.96.082002.
- [231] A. V. Berezhnoi, V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, "B/c meson at LHC," *Phys. Atom. Nucl.* **60** (1997) 1729–1740, arXiv:hep-ph/9703341.
- [232] I. P. Gouz, V. V. Kiselev, A. K. Likhoded, V. I. Romanovsky, and O. P. Yushchenko, "Prospects for the B_c studies at LHCb," *Phys. Atom. Nucl.* **67** (2004) 1559–1570. doi:10.1134/1.1788046.
- [233] C.-H. Chang and Y.-Q. Chen, "Decays of the B_c meson," *Phys. Rev.* **D49** (1994) 3399–3411. doi:10.1103/PhysRevD.49.3399.
- [234] W.-k. Kwong and J. L. Rosner, "Masses of new particles containing b quarks," *Phys. Rev.* **D44** (1991) 212–219. doi:10.1103/PhysRevD.44.212.
- [235] E. J. Eichten and C. Quigg, "Mesons with beauty and charm: Spectroscopy," *Phys. Rev.* **D49** (1994) 5845–5856, arXiv:hep-ph/9402210.

- [236] Y.-Q. Chen and Y.-P. Kuang, "Improved QCD motivated heavy quark potentials with explicit $\Lambda_{\overline{MS}}$ dependence," *Phys. Rev.* **D46** (1992) 1165–1171.
doi:10.1103/PhysRevD.46.1165.
- [237] G. Chen, X. Meng, and J. Tao, "Feasibility to study the B_c meson at CMS," *CMS AN* **FIXME: No AN's 2006/079** (2006). 
- [238] T. Sjostrand, L. Lonnblad, S. Mrenna, and P. Skands, "PYTHIA 6.3: Physics and manual," arXiv:hep-ph/0308153.
- [239] M. Grothe et al., "Triggering on Forward Physics," *CMS Note* **2006/054** (2006). Also available as TOTEM NOTE 2006-01.
- [240] CMS diffractive group and TOTEM collaboration, "Diffractive and Forward Physics," *CMS AN* **FIXME: No AN's 2006/062** (2006). 
- [241] M. Arneodo and M. Diehl, "Diffraction for non-believers,"
arXiv:hep-ph/0511047.
- [242] **FP420** Collaboration, M. Albrow et al., "FP420: A proposal to investigate the feasibility of installing proton tagging detectors in the 420 m region of the LHC," *CERN/LHCC* **2005-025** (2005).
- [243] **CDF** Collaboration, A. A. Affolder et al., "Diffractive dijets with a leading antiproton in anti-p p collisions at $\sqrt{s} = 1800$ GeV," *Phys. Rev. Lett.* **84** (2000) 5043–5048.
- [244] A. De Roeck, V. A. Khoze, A. D. Martin, R. Orava, and M. G. Ryskin, "Ways to detect a light Higgs boson at the LHC," *Eur. Phys. J.* **C25** (2002) 391–403,
arXiv:hep-ph/0207042.
- [245] V. Khoze, A. Martin, and M. . Ryskin, "Prospects for new physics observations in diffractive processes at the LHC and Tevatron," *Eur. Phys. J.* **C23** (2002) 311–327,
arXiv:hep-ph/0111078.
- [246] A. Kisselev, V. Petrov, and R. Ryutin, "5-dimensional quantum gravity effects in exclusive double diffractive events," *Phys. Lett.* **B630** (2005) 100–107.
doi:10.1016/j.physletb.2005.09.059.
- [247] J. Monk and A. Pilkington, "ExHuME: A Monte Carlo event generator for exclusive diffraction," arXiv:hep-ph/0502077.
- [248] M. Boonekamp et al., "Monte-Carlo generators for central exclusive diffraction," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at <http://www.desy.de/~heralhc/proceedings/wg4montecarlo.pdf>.
- [249] R. Ryutin, "EDDE Monte Carlo event generator," arXiv:hep-ph/0409180.
- [250] A. Kaidalov, V. Khoze, A. Martin, and M. Ryskin, "Central exclusive diffractive production as a spin parity analyser: From hadrons to Higgs," *Eur. Phys. J.* **C31** (2003) 387–396, arXiv:hep-ph/0307064.
- [251] M. Carena, S. Heinemeyer, C. Wagner, and G. Weiglein, "MSSM Higgs boson searches at the Tevatron and the LHC: Impact of different benchmark scenarios," *Eur. Phys. J.* **C45** (2006) 797–814, arXiv:hep-ph/0511023.

- [252] J. R. Ellis, J. S. Lee, and A. Pilaftsis, "Diffraction as a CP and lineshape analyzer for MSSM Higgs bosons at the LHC," *Phys. Rev.* **D71** (2005) 075007, arXiv:hep-ph/0502251.
- [253] V. Petrov, R. Ryutin, A. Sobol, and J.-P. Guillaud, "Azimuthal angular distributions in EDDE as spin-parity analyser and glueball filter for LHC," *JHEP* **06** (2005) 007, arXiv:hep-ph/0409118.
- [254] K. Piotrkowski, "Tagging two-photon production at the LHC," *Phys. Rev.* **D63** (2001) 071502, arXiv:hep-ex/0009065.
- [255] V. Budnev, I. Ginzburg, G. Meledin, and V. Serbo, "The Two photon particle production mechanism. Physical problems. Applications. Equivalent photon approximation," *Phys. Rept.* **15** (1974) 181–281. doi:10.1016/0370-1573(75)90009-5.
- [256] V. Khoze, A. Martin, and M. Ryskin, "Photon-exchange processes at hadron colliders as a probe of the dynamics of diffraction," *Eur. Phys. J.* **C24** (2002) 459–468, arXiv:hep-ph/0201301. doi:10.1007/s10052-002-0964-4.
- [257] A. R. White, "The physics of a sextet quark sector," *Phys. Rev.* **D72** (2005) 036007, arXiv:hep-ph/0412062.
- [258] K. Piotrkowski, "High energy two-photon interactions at the LHC," arXiv:hep-ex/0201027.
- [259] N. N. Kalmykov, S. S. Ostapchenko, and A. I. Pavlov, "Quark-gluon string model and EAS simulation problems at ultra-high energies," *Nucl. Phys. Proc. Suppl.* **52B** (1997) 17–28.
- [260] R. Engel, T. K. Gaisser, T. Stanev, and P. Lipari, "Air shower calculations with the new version of SIBYLL." Prepared for 26th International Cosmic Ray Conference (ICRC 99), Salt Lake City, Utah, 17-25 Aug 1999.
- [261] S. Roesler, R. Engel, and J. Ranft, "The Event generator DPMJET-III at cosmic ray energies." Prepared for 27th International Cosmic Ray Conference (ICRC 2001), Hamburg, Germany, 7-15 Aug 2001.
- [262] F. Karsch, "Lattice QCD at high temperature and density," *Lect. Notes Phys.* **583** (2002) 209–249, arXiv:hep-lat/0106019.
- [263] E. Iancu and R. Venugopalan, "The color glass condensate and high energy scattering in QCD," arXiv:hep-ph/0303204.
- [264] D. J. Schwarz, "The first second of the universe," *Annalen Phys.* **12** (2003) 220–270, arXiv:astro-ph/0303574.
- [265] A. O. Starinets, "Transport coefficients of strongly coupled gauge theories: Insights from string theory," arXiv:nucl-th/0511073.
- [266] R. Bonciani, S. Catani, M. L. Mangano, and P. Nason, "NLL resummation of the heavy-quark hadroproduction cross-section," *Nucl. Phys.* **B529** (1998) 424–450, arXiv:hep-ph/9801375.


- [267] M. Davids et al., “Measurement of top-pair cross section and top-quark mass in the di-lepton and full-hadronic channels with CMS,” *CMS Note* **2006/077** (2006).
- [268] S. Gennai et al., “Tau jet reconstruction and tagging at High Level Trigger and off-line,” *CMS Note* **2006/028** (2006).
- [269] J. D’Hondt, J. Heyninck, and S. Lowette, “Measurement of the cross section of single leptonic $t\bar{t}$ events,” *CMS Note* **2006/064** (2006).
- [270] C. Buttar et al., “Les Houches Physics at TeV Colliders 2005, Standard Model and Higgs working group: Summary report,” arXiv:hep-ph/0604120.
- [271] V. Konopliankov, O. Kodolova, and A. Ulyanov, “Jet Calibration using γ +jet Events in the CMS Detector,” *CMS Note* **2006/042** (2006).
- [272] J. D’Hondt et al., “Electron and muon reconstruction in single leptonic $t\bar{t}$ events,” *CMS Note* **2006/024** (2006).
- [273] CMS Collaboration, D. Acosta et al., “CMS Coll. Physics TDR Vol.I, Section 12.2.8,” *CERN/LHCC* **2006-001** (2006).
- [274] S. Lowette, J. D’Hondt, J. Heyninck, and P. Vanlaer, “Offline Calibration of b-jet Identification Efficiency,” CERN-CMS-NOTE-2006-013.
- [275] J. D’Hondt et al., “Tagging of Event Topologies with External Kinematic Constraints in CMS,” *CMS AN* **2006/023** (2006). **NOTE: No AN’s**
- [276] J. D’Hondt, S. Lowette, J. Heyninck, and S. Kasselmann, “Light quark jet energy scale calibration using the W mass constraint in single-leptonic $t\bar{t}$ events,” CERN-CMS-NOTE-2006-025.
- [277] The CDF Collaboration, D. Acosta et al., “Measurement of the $t\bar{t}$ cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV using kinematic characteristics of lepton plus jets events,” *Phys. Rev.* **D71** (2005) 072005. doi:10.1103/PhysRevD.71.072005.
- [278] DØ Collaboration, V. Abazov et al., “Measurement of the $t\bar{t}$ cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV using kinematic characteristics of lepton plus jets events,” *Phys. Lett.* **B626** (2005) 55. doi:10.1016/j.physletb.2005.08.105.
- [279] M. Vos and F. Palla, “B-tagging in the High Level Trigger,” *CMS Note* **2006-030** (2006).
- [280] CDF and DØ Collaboration, D. Wicke, “Top pair production cross-section measurement in the all-hadronic channel at CDF and DØ,” *Int. J. Mod. Phys. A* **20** (2005) 3183–3186, arXiv:hep-ex/0411009. doi:10.1142/S0217751X05026091.
- [281] J. D’Hondt, J. Heyninck, and S. Lowette, “Top Quark mass measurement in single-leptonic $t\bar{t}$ events,” *CMS Note* **2006/066** (2006).
- [282] CDF Collaboration, F. Abe et al., “First Observation of the All-Hadronic Decay of $t\bar{t}$ pairs,” *Phys. Rev. Lett.* **79** (1997) 1992–1997. doi:10.1103/PhysRevLett.79.1992.
- [283] T. E. W. Group, “Combination of CDF and DØ results on the mass of the top quark,” arXiv:hep-ex/0603039.

- [284] A. Kharchilava, "Top mass determination in leptonic final states with J/ψ ," *Phys. Lett.* **B476** (2000) 73–78, arXiv:hep-ph/9912320.
- [285] P. Grenier *ATLAS Physics Note* **2001-023** (2001).
- [286] R. Chierici and A. Dierlamm, "Determination of the top mass in exclusive J/ψ decays," *CMS Note* **2006-058** (2006).
- [287] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 2.6: Fast simulation," *CERN/LHCC* **2006-001** (2006) 55.
- [288] T. E. W. Group, "Combination of CDF and DØ results on the mass of the top quark," arXiv:hep-ex/0603039.
- [289] C. S. Hill, J. R. Incandela, and J. M. Lamb, "A method for measurement of the top quark mass using the mean decay length of b hadrons in t anti-t events," *Phys. Rev.* **D71** (2005) 054029, arXiv:hep-ex/0501043.
- [290] I. Borjanovic et al., "Investigation of top mass measurements with the ATLAS detector at LHC," *Eur. Phys. J.* **C39S2** (2005) 63–90, arXiv:hep-ex/0403021.
- [291] L. Lyons, D. Gibaut, and P. Clifford, "How to combine correlated estimates of a single physical quantity," *Nucl. Instrum. Meth.* **A270** (1988) 110.
- [292] A. Valassi, "Combining correlated measurements of several different physical quantities," *Nucl. Instrum. Meth.* **A500** (2003) 391–405.
- [293] G. Mahlon and S. Parke, "Angular Correlation in Top Quark Pair Production and Decay at Hadron Collider," *Phys. Rev.* **D 53** (1996) 4886–4896, arXiv:hep-ph/9512264.
- [294] T. Stelzer and S. Willenbrock, "Spin correlation in top-quark production at hadron collider," *Phys. Lett.* **B 374** (1996) 169–172, arXiv:hep-ph/9512292.
- [295] A. Brandenburg, "Spin-Spin Correlations of Top Quark Pairs at Hadron Colliders," *Phys. Lett.* **B 387** (1996) 626–632, arXiv:hep-ph/9606379.
- [296] M. Baarmand, H. Mermerkaya, and I. Vodopiyanov, "Measurement of Spin Correlation in Top Quark Pair Production in Semi-Leptonic Final State," *CMS Note* **FIXME: CMS Note number and URL** (2006) 
- [297] T. Tait, "The tW^- mode of single top production," *Phys. Rev.* **D61** (2000) 034001, arXiv:hep-ph/9909352. doi:10.1103/PhysRevD.61.034001.
- [298] A. Belyaev and E. Boos, "Single top quark $tW + X$ production at the LHC: A closer look," *Phys. Rev.* **D 63** (2001) 034012, arXiv:hep-ph/0003260.
- [299] S. Zhu, "Next-to-leading order QCD corrections to $b g \rightarrow t W^-$ at the CERN Large Hadron Collider," *pl* **B 524** (2002) 283–288.
- [300] E. Boos, V. Bunichev, L. Dudko, V. Savrin, and A. Sherstnev, "A simulation method of the Electroweak Top Quark Production Events in the NLO Approximation. A Monte-Carlo Generator 'SingleTop'," 2005. Abstract available at http://www.npi.msu.su/eng/science.php3?sec=preprint&ref_pp=1262.

- [301] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 9.1.2: Global muon reconstruction," *CERN/LHCC 2006-001* (2006) 333.
- [302] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 9.3: Muon identification," *CERN/LHCC 2006-001* (2006) 351.
- [303] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 10.4: Electron reconstruction and selection," *CERN/LHCC 2006-001* (2006) 390.
- [304] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 11.2.1: Iterative cone," *CERN/LHCC 2006-001* (2006) 408.
- [305] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 11.6.3: γ +jet events," *CERN/LHCC 2006-001* (2006) 423.
- [306] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 12.2: b-tagging tools," *CERN/LHCC 2006-001* (2006) 461.
- [307] V. Abramov  al., "Selection of single top events with the CMS detector at LHC," *CMS Note* **NAME: in preparation?** (2006).
- [308] P. Ye  al., "Search for W-associated Production of Single Top Quarks in CMS," *CMS AN* **NAME: No AN's 2006/059** (2006).
- [309] J. Campbell, R. K. Ellis, and D. L. Rainwater, "Next-to-leading order QCD predictions for W + 2jet and Z + 2jet production at the CERN LHC," *Phys. Rev.* **D68** (2003) 094021. doi:10.1103/PhysRevD.68.094021.
- [310] CMS Collaboration, D. Acosta et al., "CMS Coll. Physics TDR Vol.I, Section 11.6.5," *CERN/LHCC 2006-001* (2006).
- [311] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 8.5: Sources of systematic effects," *CERN/LHCC 2006-001* (2006) 229.
- [312] R. A. Fisher, "The use of multiple measurements in taxonomic problems," *Annals of Eugenics* **7** (1936) 179–188.
- [313] G. Eilam, J. L. Hewett, and A. Soni, "Rare decays of the top quark in the standard and two-Higgs doublet models," *Phys. Rev.* **D44** (1991) 1473–1484. doi:10.1103/PhysRevD.44.1473.
- [314] B. Mele, S. Petrarca, and A. Soddu, "A new evaluation of the $t \rightarrow cH$ decay width in the standard model," *Phys. Lett.* **B435** (1998) 401–406, arXiv:hep-ph/9805498.
- [315] C.-H. Chang, X.-Q. Li, J.-X. Wang, and M.-Z. Yang, "The production of $t\bar{c}$ or $\bar{t}c$ quark pair by e^+e^- collision based on the standard model and its extensions," *Phys. Lett.* **B313** (1993) 389–394. doi:10.1016/0370-2693(93)90008-6.
- [316] C.-S. Huang, X.-H. Wu, and S.-H. Zhu, "Top-charm associated production at high energy e^+e^- colliders in standard model," *Phys. Lett.* **B452** (1999) 143–149, arXiv:hep-ph/9901369.
- [317] H. E. Haber and G. L. Kane, "The Search for Supersymmetry: Probing Physics Beyond the Standard Model," *Phys. Rept.* **117** (1985) 75.

- [318] C. S. Li, R. J. Oakes, and J. M. Yang, "Rare decay of the top quark in the minimal supersymmetric model," *Phys. Rev.* **D49** (1994) 293–298.
- [319] G. Couture, C. Hamzaoui, and H. Konig, "Flavor changing top quark decay within the minimal supersymmetric standard model," *Phys. Rev.* **D52** (1995) 1713–1716, arXiv:hep-ph/9410230.
- [320] J. L. Lopez, D. Nanopoulos, and R. Rangarajan, "New supersymmetric contributions to $t \rightarrow cV$," *Phys. Rev.* **D56** (1997) 3100–3106, arXiv:hep-ph/9702350.
- [321] G. M. de Divitiis, R. Petronzio, and L. Silvestrini, "Flavour changing top decays in supersymmetric extensions of the standard model," *Nucl. Phys.* **B504** (1997) 45–60, arXiv:hep-ph/9704244.
- [322] J. Guasch and J. Sola, "FCNC top quark decays: A door to SUSY physics in high luminosity colliders?," *Nucl. Phys.* **B562** (1999) 3–28, arXiv:hep-ph/9906268.
- [323] G. Eilam, A. Gemintern, T. Han, J. M. Yang, and X. Zhang, "Top quark rare decay $t \rightarrow ch$ in R-parity-violating SUSY," *Phys. Lett.* **B510** (2001) 227–235, arXiv:hep-ph/0102037. doi:10.1016/S0370-2693(01)00598-6.
- [324] C. S. Li, X.-m. Zhang, and S. H. Zhu, "SUSY-QCD effect on top charm associated production at linear collider," *Phys. Rev.* **D60** (1999) 077702, arXiv:hep-ph/9904273.
- [325] J. Diaz-Cruz, R. Martinez, M. Perez, and A. Rosado, "Flavor changing radiative decay of the t quark," *Phys. Rev.* **D41** (1990) 891–894. doi:10.1103/PhysRevD.41.891.
- [326] B. Ananthanarayan, Q. Shafi, and X. M. Wang, "Improved predictions for top quark, lightest supersymmetric particle, and Higgs scalar masses," *Phys. Rev.* **D50** (1994) 5980–5984, arXiv:hep-ph/9311225.
- [327] D. Atwood, L. Reina, and A. Soni, "Phenomenology of two Higgs doublet models with flavor changing neutral currents," *Phys. Rev.* **D55** (1997) 3156–3176, arXiv:hep-ph/9609279.
- [328] S. Bar-Shalom, G. Eilam, A. Soni, and J. Wudka, "Probing the flavor-changing tc vertex via tree-level processes: $e^+e^- \rightarrow t\bar{c}v/e\bar{v}/e, t\bar{c}e^+e^-$ and $t \rightarrow cW^+W^-$," *Phys. Rev. Lett.* **79** (1997) 1217–1220, arXiv:hep-ph/9703221.
- [329] W.-S. Hou, G.-L. Lin, and C.-Y. Ma, "Flavor changing neutral Higgs couplings and top charm production at next linear collider," *Phys. Rev.* **D56** (1997) 7434–7443, arXiv:hep-ph/9708228.
- [330] S. Bejar, J. Guasch, and J. Sola, "FCNC top quark decays beyond the standard model," arXiv:hep-ph/0101294.
- [331] R. A. Diaz, R. Martinez, and J. Alexis Rodriguez, "The rare decay $t \rightarrow c\gamma$ in the general 2HDM type III," arXiv:hep-ph/0103307.
- [332] T. Han and J. L. Hewett, "Top charm associated production in high energy e^+e^- collisions," *Phys. Rev.* **D60** (1999) 074015, arXiv:hep-ph/9811237.

- [333] F. del Aguila, J. A. Aguilar-Saavedra, and R. Miquel, "Constraints on top couplings in models with exotic quarks," *Phys. Rev. Lett.* **82** (1999) 1628–1631, arXiv:hep-ph/9808400.
- [334] J. A. Aguilar-Saavedra and B. M. Nobre, "Rare top decays $t \rightarrow c\gamma$, $t \rightarrow cg$ and CKM unitarity," *Phys. Lett.* **B553** (2003) 251–260, arXiv:hep-ph/0210360.
- [335] **CDF and DØ Collaboration**, M. Paulini, "Heavy flavor physics from top to bottom," arXiv:hep-ex/9701019.
- [336] **CDF Collaboration**, F. Abe et al., "Search for Flavor-Changing Neutral Current Decays of the Top Quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **80** (1998) 2525–2530. doi:10.1103/PhysRevLett.80.2525.
- [337] A. Heinson, "Future top physics at the Tevatron and LHC," arXiv:hep-ex/9605010.
- [338] J. A. Aguilar-Saavedra and G. C. Branco, "Probing top flavour-changing neutral scalar couplings at the CERN LHC," *Phys. Lett.* **B495** (2000) 347–356, arXiv:hep-ph/0004190.
- [339] J. A. Aguilar-Saavedra, "Top flavour-changing neutral coupling signals at a linear collider," *Phys. Lett.* **B502** (2001) 115–124, arXiv:hep-ph/0012305.
- [340] J. A. Aguilar-Saavedra and T. Riemann, "Probing top flavor-changing neutral couplings at TESLA," arXiv:hep-ph/0102197.
- [341] K. Karafasoulis et al., "Study of Flavour Changing Neutral Currents in Top Quark Decays with the CMS Detector," *CMS AN* **NAME: No AN's 2006/074** (2006).
- [342] **CMS Collaboration**, D. Acosta et al., "CMS Coll. Physics TDR Vol.I, Chapter 8," *CERN/LHCC* **2006-001** (2006).
- [343] G. Dissertori et al., "How accurately can we count the number of $pp \rightarrow ZX$ and $pp \rightarrow WX$ events using decays to electrons," *CMS AN* **NAME: No AN's 2006/054** (2006).
- [344] J. Alcaraz, "Measurement of $Z \rightarrow \mu^+\mu^-$ and $W \rightarrow \mu\nu$ rates in CMS," *CMS AN* **NAME: No AN's 2006/068** (2006).
- [345] S. Frixione and B. R. Webber, "The MC@NLO 3.2 event generator," arXiv:hep-ph/0601192.
- [346] For further details see Proceedings of the 2004/2005 HERA-LHC workshop.
- [347] M. Dittmar, F. Pauss, and D. Zurcher, "Towards a precise parton luminosity determination at the CERN LHC," *Phys. Rev.* **D56** (1997) 7284–7290, arXiv:hep-ex/9705004.
- [348] S. D. Drell and T.-M. Yan, "Massive Lepton Pair Production in Hadron-Hadron Collisions at High-Energies," *Phys. Rev. Lett.* **25** (1970) 316–320. doi:10.1103/PhysRevLett.25.316.
- [349] I. Belotelov et al., "Study of Drell-Yan Di-muon Production with the CMS Detector," *CMS Note* **NAME: CMS Note number and URL** (2006).

- [350] R. Hamberg, W. L. van Neerven, and T. Matsuura, "A Complete calculation of the order α_s^2 correction to the Drell-Yan K -factor," *Nucl. Phys.* **B359** (1991) 343–405. doi:10.1016/0550-3213(91)90064-5.
- [351] D. Bourilkov, "Compositeness Search with Di-muons in CMS," 2006. CMS AN 2006/016, CERN.
- [352] U. Baur and D. Wackerth, "Electroweak radiative corrections to weak boson production at hadron colliders," *Nucl. Phys. Proc. Suppl.* **116** (2003) 159–163, arXiv:hep-ph/0211089.
- [353] D. Bourilkov, "Study of Parton Density Function Uncertainties with LHAPDF and PYTHIA at LHC," 2003.
- [354] V. Buege et al., "Prospects for the precision measurement of the W mass with the CMS detector," *CMS Note* **2006-061** (2006).
- [355] W. T. Giele and S. Keller, "Determination of W boson properties at hadron colliders," *Phys. Rev.* **D57** (1998) 4433–4440, arXiv:hep-ph/9704419.
- [356] V. e. Brigt , "Study of Di-Boson Production with the CMS Detector at the LHC," *CMS AN* **2006/063** (2006). **NOTE: No AN's 2006/063**
- [357] P. W. Higgs, "Broken symmetries, massless particles and gauge fields," *Phys. Lett.* **12** (1964) 132–133.
- [358] P. W. Higgs, "Spontaneous Symmetry Breakdown Without Massless Bosons," *Phys. Rev.* **145** (1966) 1156–1163.
- [359] F. Englert and R. Brout, "Broken Symmetry and the Mass of Gauge Vector Mesons," *Phys. Rev. Lett.* **13** (1964) 321–322.
- [360] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, "Global Conservation Laws and Massless Particles," *Phys. Rev. Lett.* **13** (1964) 585–587.
- [361] J. M. Cornwall, D. N. Levin, and G. Tiktopoulos, "Uniqueness of spontaneously broken gauge theories," *Phys. Rev. Lett.* **30** (1973) 1268–1270. doi:10.1103/PhysRevLett.30.1268.
- [362] J. M. Cornwall, D. N. Levin, and G. Tiktopoulos, "Derivation of Gauge Invariance from High-Energy Unitarity Bounds on the S - Matrix," *Phys. Rev.* **D10** (1974) 1145.
- [363] C. H. Llewellyn Smith, "High-Energy Behavior and Gauge Symmetry," *Phys. Lett.* **B46** (1973) 233–236.
- [364] S. D. Joglekar, "S-matrix derivation of the Weinberg model," *Ann. Phys.* **83** (1974) 427.
- [365] M. Veltman, "Perturbation theory of massive Yang-Mills fields," *Nucl. Phys.* **B7** (1968) 637–650.
- [366] G. 't Hooft, "Renormalization of Massless Yang-Mills Fields," *Nucl. Phys.* **B33** (1971) 173–199.
- [367] G. 't Hooft, "Renormalizable Lagrangians for Massive Yang-Mills Fields," *Nucl. Phys.* **B35** (1971) 167–188.

- [368] G. 't Hooft and M. J. G. Veltman, "Regularization and Renormalization of Gauge Fields," *Nucl. Phys.* **B44** (1972) 189–213.
- [369] G. 't Hooft and M. J. G. Veltman, "Combinatorics of gauge fields," *Nucl. Phys.* **B50** (1972) 318–353.
- [370] A. Djouadi, "The anatomy of electro-weak symmetry breaking. I: The Higgs boson in the standard model," *arXiv:hep-ph/0503172*.
- [371] A. Djouadi, "The anatomy of electro-weak symmetry breaking. II: The Higgs bosons in the minimal supersymmetric model," *arXiv:hep-ph/0503173*.
- [372] N. Cabibbo, L. Maiani, G. Parisi, and R. Petronzio, "Bounds on the Fermions and Higgs Boson Masses in Grand Unified Theories," *Nucl. Phys.* **B158** (1979) 295.
- [373] M. S. Chanowitz, M. A. Furman, and I. Hinchliffe, "Weak Interactions of Ultraheavy Fermions," *Phys. Lett.* **B78** (1978) 285.
- [374] R. A. Flores and M. Sher, "Upper Limits to Fermion Masses in The Glashow-Weinberg-Salam Model," *Phys. Rev.* **D27** (1983) 1679.
- [375] M. Lindner, "Implications of Triviality for the Standard Model," *Zeit. Phys.* **C31** (1986) 295.
- [376] M. Sher, "Electroweak Higgs Potentials and Vacuum Stability," *Phys. Rept.* **179** (1989) 273–418.
- [377] M. Sher, "Precise vacuum stability bound in the standard model," *Phys. Lett.* **B317** (1993) 159–163, *arXiv:hep-ph/9307342*.
- [378] G. Altarelli and G. Isidori, "Lower limit on the Higgs mass in the standard model: An Update," *Phys. Lett.* **B337** (1994) 141–144.
- [379] J. Espinosa and M. Quiros, "Improved metastability bounds on the standard model Higgs mass," *Phys. Lett.* **B353** (1995) 257–266, *arXiv:hep-ph/9504241*.
- [380] A. Hasenfratz, K. Jansen, C. B. Lang, T. Neuhaus, and H. Yoneyama, "The Triviality Bound of the Four Component Φ^4 Model," *Phys. Lett.* **B199** (1987) 531.
- [381] J. Kuti, L. Lin, and Y. Shen, "Upper Bound on the Higgs Mass in the Standard Model," *Phys. Rev. Lett.* **61** (1988) 678.
- [382] M. Luscher and P. Weisz, "Scaling Laws and Triviality Bounds in the Lattice Φ^4 Theory. 3. n Component Model," *Nucl. Phys.* **B318** (1989) 705.
- [383] **Higgs Working Group** Collaboration, M. Carena et al., "Report of the Tevatron Higgs working group," *arXiv:hep-ph/0010338*.
- [384] ATLAS Collaboration, "Technical Design Report," *CERN/LHCC 94-14* (1994). ATLAS TDR 14.
- [385] S. Abdullin et al., "Summary of the CMS potential for the Higgs boson discovery," *Eur. Phys. J.* **C39S2** (2005) 41–61. doi:10.1140/epjcd/s2004-02-003-9.

- [386] E. Braaten and J. P. Leveille, "Higgs Boson Decay and the Running Mass," *Phys. Rev.* **D22** (1980) 715.
- [387] N. Sakai, "Perturbative QCD Corrections to the Hadronic Decay Width of the Higgs Boson," *Phys. Rev.* **D22** (1980) 2220.
- [388] T. Inami and T. Kubota, "Renormalization Group Estimate of the Hadronic Decay Width of the Higgs Boson," *Nucl. Phys.* **B179** (1981) 171.
- [389] S. G. Gorishnii, A. L. Kataev, and S. A. Larin, "The Width of Higgs Boson Decay into Hadrons: Three Loop Corrections of Strong Interactions," *Sov. J. Nucl. Phys.* **40** (1984) 329–334.
- [390] M. Drees and K.-i. Hikasa, "Heavy Quark Thresholds in Higgs Physics," *Phys. Rev.* **D41** (1990) 1547.
- [391] M. Drees and K.-i. Hikasa, "Note on QCD Corrections to Hadronic Higgs Decay," *Phys. Lett.* **B240** (1990) 455.
- [392] K. G. Chetyrkin, "Correlator of the quark scalar currents and $\Gamma(\text{tot})(H \rightarrow \text{hadrons})$ at $O(\alpha(s)^3)$ in pQCD," *Phys. Lett.* **B390** (1997) 309–317, arXiv:hep-ph/9608318.
- [393] J. Fleischer and F. Jegerlehner, "Radiative Corrections to Higgs Decays in the Extended Weinberg-Salam Model," *Phys. Rev.* **D23** (1981) 2001–2026.
- [394] D. Y. Bardin, B. M. Vilensky, and P. K. Khristova, "Calculation of the Higgs boson decay width into fermion pairs," *Sov. J. Nucl. Phys.* **53** (1991) 152–158.
- [395] A. Dabelstein and W. Hollik, "Electroweak corrections to the fermionic decay width of the standard Higgs boson," *Z. Phys.* **C53** (1992) 507–516.
- [396] B. A. Kniehl, "Radiative corrections for $H \rightarrow f\bar{f}(\gamma)$ in the standard model," *Nucl. Phys.* **B376** (1992) 3–28. doi:10.1016/0550-3213(92)90065-J.
- [397] H.-Q. Zheng and D.-D. Wu, "First order QCD corrections to the decay of the Higgs boson into two photons," *Phys. Rev.* **D42** (1990) 3760–3763.
- [398] A. Djouadi, M. Spira, J. van der Bij, and P. Zerwas, "QCD corrections to gamma gamma decays of Higgs particles in the intermediate mass range," *Phys. Lett.* **B257** (1991) 187–190.
- [399] S. Dawson and R. P. Kauffman, "QCD corrections to $H \rightarrow \gamma\gamma$," *Phys. Rev.* **D47** (1993) 1264–1267.
- [400] A. Djouadi, M. Spira, and P. Zerwas, "Two photon decay widths of Higgs particles," *Phys. Lett.* **B311** (1993) 255–260, arXiv:hep-ph/9305335.
- [401] K. Melnikov and O. I. Yakovlev, "Higgs \rightarrow two photon decay: QCD radiative correction," *Phys. Lett.* **B312** (1993) 179–183, arXiv:hep-ph/9302281.
- [402] M. Inoue, R. Najima, T. Oka, and J. Saito, "QCD corrections to two photon decay of the Higgs boson and its reverse process," *Mod. Phys. Lett.* **A9** (1994) 1189–1194.

- [403] M. Steinhauser, "Corrections of $O(\alpha(s)^2)$ to the decay of an intermediate-mass Higgs boson into two photons," [arXiv:hep-ph/9612395](https://arxiv.org/abs/hep-ph/9612395).
- [404] A. Djouadi, P. Gambino, and B. A. Kniehl, "Two-loop electroweak heavy-fermion corrections to Higgs-boson production and decay," *Nucl. Phys.* **B523** (1998) 17–39, [arXiv:hep-ph/9712330](https://arxiv.org/abs/hep-ph/9712330).
- [405] U. Aglietti, R. Bonciani, G. Degrossi, and A. Vicini, "Two-loop light fermion contribution to Higgs production and decays," *Phys. Lett.* **B595** (2004) 432–441, [arXiv:hep-ph/0404071](https://arxiv.org/abs/hep-ph/0404071).
- [406] G. Degrossi and F. Maltoni, "Two-loop electroweak corrections to the Higgs-boson decay $H \rightarrow \gamma \gamma$," [arXiv:hep-ph/0504137](https://arxiv.org/abs/hep-ph/0504137).
- [407] B. A. Kniehl, "Radiative corrections for $H \rightarrow ZZ$ in the standard model," *Nucl. Phys.* **B352** (1991) 1–26. doi:10.1016/0550-3213(91)90126-I.
- [408] B. A. Kniehl, "Radiative corrections for $H \rightarrow W^+W^-(\gamma)$ in the standard model," *Nucl. Phys.* **B357** (1991) 439–466. doi:10.1016/0550-3213(91)90476-E.
- [409] H. Georgi, S. Glashow, M. Machacek, and D. Nanopoulos, "Higgs Bosons from Two Gluon Annihilation in Proton Proton Collisions," *Phys. Rev. Lett.* **40** (1978) 692. doi:10.1103/PhysRevLett.40.692.
- [410] D. Graudenz, M. Spira, and P. M. Zerwas, "QCD corrections to Higgs boson production at proton proton colliders," *Phys. Rev. Lett.* **70** (1993) 1372–1375. doi:10.1103/PhysRevLett.70.1372.
- [411] M. Spira, A. Djouadi, D. Graudenz, and P. Zerwas, "SUSY Higgs production at proton colliders," *Phys. Lett.* **B318** (1993) 347–353. doi:10.1016/0370-2693(93)90138-8.
- [412] M. Spira, A. Djouadi, D. Graudenz, and P. Zerwas, "Higgs boson production at the LHC," *Nucl. Phys.* **B453** (1995) 17–82, [arXiv:hep-ph/9504378](https://arxiv.org/abs/hep-ph/9504378). doi:10.1016/0550-3213(95)00379-7.
- [413] M. Kramer, E. Laenen, and M. Spira, "Soft gluon radiation in Higgs boson production at the LHC," *Nucl. Phys.* **B511** (1998) 523–549, [arXiv:hep-ph/9611272](https://arxiv.org/abs/hep-ph/9611272). doi:10.1016/S0550-3213(97)00679-2.
- [414] A. Djouadi, M. Spira, and P. Zerwas, "Production of Higgs bosons in proton colliders: QCD corrections," *Phys. Lett.* **B264** (1991) 440–446.
- [415] S. Dawson, "Radiative corrections to Higgs boson production," *Nucl. Phys.* **B359** (1991) 283–300.
- [416] R. P. Kauffman and W. Schaffer, "QCD corrections to production of Higgs pseudoscalars," *Phys. Rev.* **D49** (1994) 551–554, [arXiv:hep-ph/9305279](https://arxiv.org/abs/hep-ph/9305279).
- [417] S. Dawson and R. Kauffman, "QCD corrections to Higgs boson production: nonleading terms in the heavy quark limit," *Phys. Rev.* **D49** (1994) 2298–2309, [arXiv:hep-ph/9310281](https://arxiv.org/abs/hep-ph/9310281).

- [418] R. V. Harlander and W. B. Kilgore, "Next-to-next-to-leading order Higgs production at hadron colliders," *Phys. Rev. Lett.* **88** (2002) 201801, arXiv:hep-ph/0201206.
- [419] R. V. Harlander and W. B. Kilgore, "Production of a pseudo-scalar Higgs boson at hadron colliders at next-to-next-to leading order," *JHEP* **10** (2002) 017, arXiv:hep-ph/0208096.
- [420] C. Anastasiou and K. Melnikov, "Higgs boson production at hadron colliders in NNLO QCD," *Nucl. Phys.* **B646** (2002) 220–256, arXiv:hep-ph/0207004.
- [421] V. Ravindran, J. Smith, and W. L. van Neerven, "NNLO corrections to the total cross section for Higgs boson production in hadron hadron collisions," *Nucl. Phys.* **B665** (2003) 325–366, arXiv:hep-ph/0302135.
- [422] S. Catani, D. de Florian, M. Grazzini, and P. Nason, "Soft-gluon resummation for Higgs boson production at hadron colliders," *JHEP* **07** (2003) 028, arXiv:hep-ph/0306211.
- [423] A. Djouadi and P. Gambino, "Leading electroweak correction to Higgs boson production at proton colliders," *Phys. Rev. Lett.* **73** (1994) 2528–2531, arXiv:hep-ph/9406432.
- [424] K. G. Chetyrkin, B. A. Kniehl, and M. Steinhauser, "Virtual top-quark effects on the $H \rightarrow b$ anti- b decay at next-to-leading order in QCD," *Phys. Rev. Lett.* **78** (1997) 594–597, arXiv:hep-ph/9610456.
- [425] K. G. Chetyrkin, B. A. Kniehl, and M. Steinhauser, "Three-loop $O(\alpha_s)^2 G(F) M(t)^2$ corrections to hadronic Higgs decays," *Nucl. Phys.* **B490** (1997) 19–39, arXiv:hep-ph/9701277.
- [426] A. Ghinculov and J. J. van der Bij, "The Higgs resonance shape in gluon fusion: Heavy Higgs effects," *Nucl. Phys.* **B482** (1996) 59–72, arXiv:hep-ph/9511414.
- [427] R. K. Ellis, I. Hinchliffe, M. Soldate, and J. J. van der Bij, "Higgs Decay to $\tau^+ \tau^-$: A Possible Signature of Intermediate Mass Higgs Bosons at the SSC," *Nucl. Phys.* **B297** (1988) 221.
- [428] U. Baur and E. W. N. Glover, "Higgs Boson Production at Large Transverse Momentum in Hadronic Collisions," *Nucl. Phys.* **B339** (1990) 38–66.
- [429] C. R. Schmidt, " $H \rightarrow ggg(gg\bar{q})$ at two loops in the large- M_t limit," *Phys. Lett.* **B413** (1997) 391–395. doi:10.1016/S0370-2693(97)01102-7.
- [430] D. de Florian, M. Grazzini, and Z. Kunszt, "Higgs production with large transverse momentum in hadronic collisions at next-to-leading order," *Phys. Rev. Lett.* **82** (1999) 5209–5212, arXiv:hep-ph/9902483.
- [431] V. Ravindran, J. Smith, and W. L. Van Neerven, "Next-to-leading order QCD corrections to differential distributions of Higgs boson production in hadron hadron collisions," *Nucl. Phys.* **B634** (2002) 247–290, arXiv:hep-ph/0201114.
- [432] C. J. Glosser and C. R. Schmidt, "Next-to-leading corrections to the Higgs boson transverse momentum spectrum in gluon fusion," *JHEP* **12** (2002) 016, arXiv:hep-ph/0209248.

- [433] C. Anastasiou, K. Melnikov, and F. Petriello, "Fully differential Higgs boson production and the di-photon signal through next-to-next-to-leading order," *Nucl. Phys.* **B724** (2005) 197–246, arXiv:hep-ph/0501130.
- [434] S. Catani, E. D'Emilio, and L. Trentadue, "The Gluon Form-Factor to Higher Orders: Gluon Gluon Annihilation at Small Q-Transverse," *Phys. Lett.* **B211** (1988) 335–342.
- [435] I. Hinchliffe and S. F. Novaes, "On the Mean Transverse Momentum of Higgs Bosons at the SSC," *Phys. Rev.* **D38** (1988) 3475–3480.
- [436] R. P. Kauffman, "Higgs boson p(T) in gluon fusion," *Phys. Rev.* **D44** (1991) 1415–1425.
- [437] R. Kauffman, "Higher order corrections to Higgs boson p(T)," *Phys. Rev.* **D45** (1992) 1512–1517.
- [438] C. Balazs and C. P. Yuan, "Higgs boson production at the LHC with soft gluon effects," *Phys. Lett.* **B478** (2000) 192–198, arXiv:hep-ph/0001103.
- [439] E. L. Berger and J.-w. Qiu, "Differential cross section for Higgs boson production including all-orders soft gluon resummation," *Phys. Rev.* **D67** (2003) 034026, arXiv:hep-ph/0210135.
- [440] A. Kulesza and W. J. Stirling, "Non-perturbative effects and the resummed Higgs transverse momentum distribution at the LHC," *JHEP* **12** (2003) 056, arXiv:hep-ph/0307208.
- [441] A. Kulesza, G. Sterman, and W. Vogelsang, "Joint resummation for Higgs production," *Phys. Rev.* **D69** (2004) 014012, arXiv:hep-ph/0309264.
- [442] A. Gawron and J. Kwiecinski, "Resummation effects in Higgs boson transverse momentum distribution within the framework of unintegrated parton distributions," *Phys. Rev.* **D70** (2004) 014003, arXiv:hep-ph/0309303.
- [443] G. Watt, A. Martin, and M. Ryskin, "Unintegrated parton distributions and electroweak boson production at hadron colliders," *Phys. Rev.* **D70** (2004) 014012, arXiv:hep-ph/0309096.
- [444] A. Lipatov and N. Zotov, "Higgs boson production at hadron colliders in the k(T)-factorization approach," *Eur. Phys. J.* **C44** (2005) 559–566, arXiv:hep-ph/0501172.
- [445] D. de Florian and M. Grazzini, "Next-to-next-to-leading logarithmic corrections at small transverse momentum in hadronic collisions," *Phys. Rev. Lett.* **85** (2000) 4678–4681, arXiv:hep-ph/0008152.
- [446] S. Catani, D. De Florian, and M. Grazzini, "Higgs production at hadron colliders in (almost) NNLO QCD," in *Bologna 2001, Deep inelastic scattering*, pp. 518–521. Bologna, Italy, April, 2001. arXiv:hep-ph/0106049.
- [447] S. Catani, D. de Florian, and M. Grazzini, "Universality of non-leading logarithmic contributions in transverse momentum distributions," *Nucl. Phys.* **B596** (2001) 299–312, arXiv:hep-ph/0008184.
- [448] G. Bozzi, S. Catani, D. de Florian, and M. Grazzini, "The q(T) spectrum of the Higgs boson at the LHC in QCD perturbation theory," *Phys. Lett.* **B564** (2003) 65–72, arXiv:hep-ph/0302104.

- [449] G. Bozzi, S. Catani, D. de Florian, and M. Grazzini, "Transverse-momentum resummation and the spectrum of the Higgs boson at the LHC," *Nucl. Phys.* **B737** (2006) 73–120, arXiv:hep-ph/0508068.
- [450] R. N. Cahn and S. Dawson, "Production of Very Massive Higgs Bosons," *Phys. Lett.* **B136** (1984) 196.
- [451] K.-i. Hikasa, "Heavy Higgs Production in $e^+ e^-$ and $e^- e^-$ Collisions," *Phys. Lett.* **B164** (1985) 385.
- [452] G. Altarelli, B. Mele, and F. Pitolli, "Heavy Higgs Production at Future Colliders," *Nucl. Phys.* **B287** (1987) 205–224.
- [453] T. Han, G. Valencia, and S. Willenbrock, "Structure function approach to vector boson scattering in $p p$ collisions," *Phys. Rev. Lett.* **69** (1992) 3274–3277, arXiv:hep-ph/9206246.
- [454] T. Figy, C. Oleari, and D. Zeppenfeld, "Next-to-leading order jet distributions for Higgs boson production via weak-boson fusion," *Phys. Rev.* **D68** (2003) 073005, arXiv:hep-ph/0306109.
- [455] S. Glashow, D. Nanopoulos, and A. Yildiz, "Associated Production of Higgs Bosons and Z Particles," *Phys. Rev.* **D18** (1978) 1724–1727.
- [456] Z. Kunszt, Z. Trocsanyi, and W. J. Stirling, "Clear signal of intermediate mass Higgs boson production at LHC and SSC," *Phys. Lett.* **B271** (1991) 247–255.
- [457] T. Han and S. Willenbrock, "QCD correction to the $p p \rightarrow W H$ and $Z H$ total cross-sections," *Phys. Lett.* **B273** (1991) 167–172.
- [458] O. Brein, A. Djouadi, and R. Harlander, "NNLO QCD corrections to the Higgs-strahlung processes at hadron colliders," *Phys. Lett.* **B579** (2004) 149–156, arXiv:hep-ph/0307206.
- [459] M. L. Ciccolini, S. Dittmaier, and M. Kramer, "Electroweak radiative corrections to associated $W H$ and $Z H$ production at hadron colliders," *Phys. Rev.* **D68** (2003) 073003, arXiv:hep-ph/0306234.
- [460] R. Raitio and W. W. Wada, "Higgs Boson Production at Large Transverse Momentum in QCD," *Phys. Rev.* **D19** (1979) 941.
- [461] J. N. Ng and P. Zakarauskas, "A QCD Parton Calculation of Conjoined Production of Higgs Bosons and Heavy Flavors in p anti- p Collision," *Phys. Rev.* **D29** (1984) 876.
- [462] Z. Kunszt, "Associated Production of Heavy Higgs Boson with Top Quarks," *Nucl. Phys.* **B247** (1984) 339.
- [463] J. F. Gunion, "Associated top anti-top Higgs production as a large source of $W H$ events: Implications for Higgs detection in the lepton neutrino gamma gamma final state," *Phys. Lett.* **B261** (1991) 510–517.
- [464] W. J. Marciano and F. E. Paige, "Associated production of Higgs bosons with t anti- t pairs," *Phys. Rev. Lett.* **66** (1991) 2433–2435.

- [465] W. Beenakker, S. Dittmaier, M. Kraemer, B. Pluemper, M. Spira, and P. Zerwas, "Higgs radiation off top quarks at the Tevatron and the LHC," *Phys. Rev. Lett.* **87** (2001) 201805, arXiv:hep-ph/0107081.
- [466] S. Dawson, L. H. Orr, L. Reina, and D. Wackerroth, "Associated top quark Higgs boson production at the LHC," *Phys. Rev.* **D67** (2003) 071503, arXiv:hep-ph/0211438.
- [467] S. Dawson and L. Reina, "QCD corrections to associated Higgs boson production," *Phys. Rev.* **D57** (1998) 5851–5859, arXiv:hep-ph/9712400.
- [468] S. Abdullin et al., "Search for $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ using $M(4\mu)$ dependent cuts," *CMS AN 2006/056* **ATLAS: NEEDS TO BE CMS-NOTE** (2006).
- [469] D. Acosta et al., "NLO vs. LO: kinematical differences for signal and background in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ analysis," *CMS AN 2006/053* **ATLAS: NEEDS TO BE CMS-NOTE** (2006).
- [470] S. Baffioni et al., "Discovery potential for the Higgs boson in the $H \rightarrow ZZ^{(*)} \rightarrow e^+e^-e^+e^-$ decay channel," *CMS AN 2006/070* **ATLAS: NEEDS TO BE CMS-NOTE: use CMS Note 2006-HZZ4e 2006/070** (2006).
- [471] M. Dittmar and H. K. Dreiner, "How to find a Higgs boson with a mass between 155-GeV to 180-GeV at the LHC," *Phys. Rev.* **D55** (1997) 167–172, arXiv:hep-ph/9608317.
- [472] G. Davatz, G. Dissertori, M. Dittmar, M. Grazzini, and F. Pauss, "Effective K-factors for $gg \rightarrow H \rightarrow WW \rightarrow \ell\nu\ell\nu$ at the LHC," *JHEP* **05** (2004) 009, arXiv:hep-ph/0402218.
- [473] J. Campbell and F. Tramontano, "Next-to-leading order corrections to Wt production and decay," *Nucl. Phys.* **B726** (2005) 109–130, arXiv:hep-ph/0506289.
- [474] CDF Collaboration, D. Acosta et al., "Measurement of the $t\bar{t}$ Production Cross Section in pp bar Collisions at $\sqrt{s} = 1.96$ TeV using Lepton + Jets Events with Secondary Vertex b-tagging," *Phys. Rev.* **D71** (2005) 052003.
- [475] A. Djouadi and S. Ferrag, "PDF Uncertainties In Higgs Production At Hadron Colliders," arXiv:hep-ph/0310209.
- [476] Higgs Working Group Collaboration, K. Assamagan et al., "The Higgs working group: Summary report 2003," arXiv:hep-ph/0406152.
- [477] G. Belanger, F. Boudjema, and K. Sridhar, "SUSY Higgs at the LHC: Large stop mixing effects and associated production," *Nucl. Phys.* **B568** (2000) 3–39, arXiv:hep-ph/9904348.
- [478] M. Dubinin, V. Ilyin, and V. Savrin, "Light Higgs Boson Signal at LHC in the reaction $pp \rightarrow \gamma\gamma + \text{jet}$ and $pp \rightarrow \gamma\gamma + \text{lepton}$," *CMS Note* **1997/101** (1997).
- [479] R. Kinnunen and D. Denegri, "Expected SM/SUSY Higgs Observability in CMS," *CMS Note* **1997/057** (1997).
- [480] ATLAS Collaboration, "ATLAS detector and physics performance. Technical design report. Vol. 2, ". CERN-LHCC-99-15.

- [481] P. Beauchemin, G. Azuelos, and C. Burgess, "Dimensionless coupling of bulk scalars at the LHC," *J. Phys.* **G30** (2004) N17. doi:10.1088/0954-3899/30/10/N01.
- [482] T. Stelzer and W. Long, "Automatic generation of tree level helicity amplitudes," *Comput. Phys. Commun.* **81** (1994) 357–371. doi:10.1016/0010-4655(94)90084-1.
- [483] H. Murayama, I. Watanabe, and K. Hagiwara, "HELAS: HELicity amplitude subroutines for Feynman diagram evaluations," KEK-91-11.
- [484] M. Mangano, M. Moretti, and R. Pittau, "Multijet matrix elements and shower evolution in hadronic collisions: $Wb\bar{b} + n$ -jets as a case study," *Nucl. Phys.* **B632** (2002) 343–362, arXiv:hep-ph/0108069. doi:10.1016/S0550-3213(02)00249-3.
- [485] F. Caravaglios, M. Mangano, M. Moretti, and R. Pittau, "A new approach to multi-jet calculations in hadron collisions," *Nucl. Phys.* **B539** (1999) 215–232, arXiv:hep-ph/9807570.
- [486] S. Bityukov, S. Erofeeva, N. Krasnikov, and A. Nikitenko, "Program for evaluation of significance, confidence intervals and limits by direct calculation of probabilities," *Proceedings of PhyStat 2005* (2005).
- [487] M. Lethuillier et al., "Search for neutral Higgs boson with WH / ZH, $H \rightarrow \gamma\gamma$ channel," CMS AN-2006/058 ~~NAME: NEEDS TO BE CMS-NOTE~~ (2006).
- [488] O. Ravat, "Etude du Calorimètre électromagnétique de l'expérience CMS et recherche de bosons de Higgs neutres dans le canal de production associée". PhD thesis, IPN, Lyon, 2004. Thesis LYCEN-T2004-29.
- [489] D. Graham, "An algorithm using tracks to locate the two photon vertex at high luminosity," *CMS TN* **1995/115** (1995).
- [490] A. Djouadi, "Squark effects on Higgs boson production and decay at the LHC," *Phys. Lett.* **B435** (1998) 101–108, arXiv:hep-ph/9806315.
- [491] G. Eynard, "Study of associated Higgs boson production $HW, Ht\bar{t}, HZ \rightarrow \gamma\gamma + e^\pm/\mu^\pm + X$ with the ATLAS detector at LHC. (In French)". PhD thesis, Annecy, 1998. CERN-THESIS-2000-036.
- [492] P.-H. Beauchemin and G. Azuelos, "Search for the Standard Model Higgs Boson in the $\gamma\gamma + E_T^{miss}$ channel," *ATL-PHYS-2004-028* (2004).
- [493] CMS Collaboration, S. Dasu et al., "CMS. The TriDAS project. Technical design report, vol. 1: The trigger systems," CERN-LHCC-2000-038.
- [494] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 10.3.2: Photon isolation," *CERN/LHCC* **2006-001** (2006) 376.
- [495] A. L. Read, "Presentation of search results: The CL(s) technique," *J. Phys.* **G28** (2002) 2693–2704.
- [496] T. Junk, "Confidence level computation for combining searches with small statistics," *Nucl. Instrum. Meth.* **A434** (1999) 435–443, arXiv:hep-ex/9902006.

- [497] A. Djouadi and S. Ferrag, "PDF uncertainties in Higgs production at hadron colliders," *Phys. Lett.* **B586** (2004) 345–352, arXiv:hep-ph/0310209.
- [498] J. Rohlf and C. Tully, "Recommendations for Jet and Missing Transverse Energy Reconstruction Settings and Systematics Treatment," *CMS IN PROGRESS: NEEDS TO BE CMS-NOTE 2006/XXX* (2006).
- [499] J. R. Dell'Aquila and C. A. Nelson, "Simple tests for CP or P violation by sequential decays: V_1V_2 modes with decays into $\bar{\ell}_A\ell_B$ and/or \bar{q}_Aq_B ," *Phys. Rev.* **D33** (1986) 101. doi:10.1103/PhysRevD.33.101.
- [500] A. Skjold and P. Osland, "Signals of CP violation in Higgs decay," *Phys. Lett.* **B329** (1994) 305–311, arXiv:hep-ph/9402358.
- [501] S. Y. Choi, D. J. Miller, M. M. Muhlleitner, and P. M. Zerwas, "Identifying the Higgs spin and parity in decays to Z pairs," *Phys. Lett.* **B553** (2003) 61–71, arXiv:hep-ph/0210077.
- [502] C. P. Buszello, I. Fleck, P. Marquard, and J. J. van der Bij, "Prospective analysis of spin- and CP-sensitive variables in $H \rightarrow ZZ \rightarrow l(1)+l(1)-l(2)+l(2)-$ at the LHC," *Eur. Phys. J.* **C32** (2004) 209–219, arXiv:hep-ph/0212396.
- [503] J. Wess and B. Zumino, "Supergauge Transformations in Four-Dimensions," *Nucl. Phys.* **B70** (1974) 39–50.
- [504] P. Fayet and S. Ferrara, "Supersymmetry," *Phys. Rept.* **32** (1977) 249–334.
- [505] H. P. Nilles, "Supersymmetry, Supergravity and Particle Physics," *Phys. Rept.* **110** (1984) 1.
- [506] R. Barbieri, "Looking Beyond the Standard Model: the Supersymmetric Option," *Riv. Nuovo Cim.* **11N4** (1988) 1–45.
- [507] E. Witten, "Mass Hierarchies in Supersymmetric Theories," *Phys. Lett.* **B105** (1981) 267.
- [508] S. Dimopoulos, S. Raby, and F. Wilczek, "Supersymmetry and the Scale of Unification," *Phys. Rev.* **D24** (1981) 1681–1683.
- [509] L. E. Ibanez and G. G. Ross, "Low-Energy Predictions in Supersymmetric Grand Unified Theories," *Phys. Lett.* **B105** (1981) 439.
- [510] L. E. Ibanez and G. G. Ross, "SU(2)-L \times U(1) Symmetry Breaking as a Radiative Effect of Supersymmetry Breaking in GUTs," *Phys. Lett.* **B110** (1982) 215–220.
- [511] K. Inoue, A. Kakuto, H. Komatsu, and S. Takeshita, "Aspects of Grand Unified Models with Softly Broken Supersymmetry," *Prog. Theor. Phys.* **68** (1982) 927.
- [512] L. Alvarez-Gaume, M. Claudson, and M. B. Wise, "Low-Energy Supersymmetry," *Nucl. Phys.* **B207** (1982) 96.
- [513] J. R. Ellis, D. V. Nanopoulos, and K. Tamvakis, "Grand Unification in Simple Supergravity," *Phys. Lett.* **B121** (1983) 123.

- [514] S. Dimopoulos and H. Georgi, "Softly Broken Supersymmetry and SU(5)," *Nucl. Phys.* **B193** (1981) 150.
- [515] N. Sakai, "Naturalness in Supersymmetric 'GUTS'," *Zeit. Phys.* **C11** (1981) 153.
- [516] Y. Okada, M. Yamaguchi, and T. Yanagida, "Upper bound of the lightest Higgs boson mass in the minimal supersymmetric standard model," *Prog. Theor. Phys.* **85** (1991) 1–6.
- [517] H. E. Haber and R. Hempfling, "Can the mass of the lightest Higgs boson of the minimal supersymmetric model be larger than $m(Z)$?" *Phys. Rev. Lett.* **66** (1991) 1815–1818.
- [518] J. R. Ellis, G. Ridolfi, and F. Zwirner, "Radiative corrections to the masses of supersymmetric Higgs bosons," *Phys. Lett.* **B257** (1991) 83–91.
- [519] R. Barbieri, M. Frigeni, and F. Caravaglios, "The Supersymmetric Higgs for heavy superpartners," *Phys. Lett.* **B258** (1991) 167–170.
- [520] A. Yamada, "Radiative corrections to the Higgs masses in the minimal supersymmetric standard model," *Phys. Lett.* **B263** (1991) 233–238.
- [521] A. Brignole, J. R. Ellis, G. Ridolfi, and F. Zwirner, "The Supersymmetric charged Higgs boson mass and LEP phenomenology," *Phys. Lett.* **B271** (1991) 123–132.
- [522] P. H. Chankowski, S. Pokorski, and J. Rosiek, "Charged and neutral supersymmetric Higgs boson masses: Complete one loop analysis," *Phys. Lett.* **B274** (1992) 191–198.
- [523] J. Espinosa and M. Quiros, "Two loop radiative corrections to the mass of the lightest Higgs boson in supersymmetric standard models," *Phys. Lett.* **B266** (1991) 389–396.
- [524] R. Hempfling and A. H. Hoang, "Two loop radiative corrections to the upper limit of the lightest Higgs boson mass in the minimal supersymmetric model," *Phys. Lett.* **B331** (1994) 99–106, [arXiv:hep-ph/9401219](https://arxiv.org/abs/hep-ph/9401219).
- [525] J. Casas, J. Espinosa, M. Quiros, and A. Riotto, "The Lightest Higgs boson mass in the minimal supersymmetric standard model," *Nucl. Phys.* **B436** (1995) 3–29, [arXiv:hep-ph/9407389](https://arxiv.org/abs/hep-ph/9407389).
- [526] M. Carena, J. Espinosa, M. Quiros, and C. Wagner, "Analytical expressions for radiatively corrected Higgs masses and couplings in the MSSM," *Phys. Lett.* **B355** (1995) 209–221, [arXiv:hep-ph/9504316](https://arxiv.org/abs/hep-ph/9504316).
- [527] M. Carena, M. Quiros, and C. Wagner, "Effective potential methods and the Higgs mass spectrum in the MSSM," *Nucl. Phys.* **B461** (1996) 407–436, [arXiv:hep-ph/9508343](https://arxiv.org/abs/hep-ph/9508343).
- [528] S. Heinemeyer, W. Hollik, and G. Weiglein, "QCD corrections to the masses of the neutral CP-even Higgs bosons in the MSSM," *Phys. Rev.* **D58** (1998) 091701, [arXiv:hep-ph/9803277](https://arxiv.org/abs/hep-ph/9803277).
- [529] S. Heinemeyer, W. Hollik, and G. Weiglein, "Precise prediction for the mass of the lightest Higgs boson in the MSSM," *Phys. Lett.* **B440** (1998) 296–304, [arXiv:hep-ph/9807423](https://arxiv.org/abs/hep-ph/9807423).

- [530] S. Heinemeyer, W. Hollik, and G. Weiglein, "Constraints on $\tan(\beta)$ in the MSSM from the upper bound on the mass of the lightest Higgs boson," *JHEP* **06** (2000) 009, arXiv:hep-ph/9909540.
- [531] J. R. Espinosa and R.-J. Zhang, "Complete two-loop dominant corrections to the mass of the lightest CP-even Higgs boson in the minimal supersymmetric standard model," *Nucl. Phys.* **B586** (2000) 3–38, arXiv:hep-ph/0003246.
- [532] A. Brignole, G. Degrassi, P. Slavich, and F. Zwirner, "On the $\mathcal{O}(\alpha_s^2)$ two-loop corrections to the neutral Higgs boson masses in the MSSM," *Nucl. Phys.* **B631** (2002) 195–218. doi:10.1016/S0550-3213(02)00184-0.
- [533] A. Brignole, G. Degrassi, P. Slavich, and F. Zwirner, "On the two-loop sbottom corrections to the neutral Higgs boson masses in the MSSM," *Nucl. Phys.* **B643** (2002) 79–92, arXiv:hep-ph/0206101.
- [534] S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, "High-precision predictions for the MSSM Higgs sector at $\mathcal{O}(\alpha_s \alpha_f)$," *Eur. Phys. J.* **C39** (2005) 465–481. doi:10.1140/epjc/s2005-02112-6.
- [535] J. A. Coarasa, R. A. Jimenez, and J. Sola, "Strong effects on the hadronic widths of the neutral Higgs Bosons in the MSSM," *Phys. Lett.* **B389** (1996) 312–320, arXiv:hep-ph/9511402.
- [536] P. H. Chankowski, S. Pokorski, and J. Rosiek, "Complete on-shell renormalization scheme for the minimal supersymmetric Higgs sector," *Nucl. Phys.* **B423** (1994) 437–496, arXiv:hep-ph/9303309.
- [537] A. Dabelstein, "The One loop renormalization of the MSSM Higgs sector and its application to the neutral scalar Higgs masses," *Z. Phys.* **C67** (1995) 495–512, arXiv:hep-ph/9409375.
- [538] A. Brignole, "Radiative corrections to the supersymmetric neutral Higgs boson masses," *Phys. Lett.* **B281** (1992) 284–294.
- [539] H. E. Haber, R. Hempfling, and A. H. Hoang, "Approximating the radiatively corrected Higgs mass in the minimal supersymmetric model," *Z. Phys.* **C75** (1997) 539–554, arXiv:hep-ph/9609331.
- [540] R.-J. Zhang, "Two-loop effective potential calculation of the lightest CP-even Higgs-boson mass in the MSSM," *Phys. Lett.* **B447** (1999) 89–97, arXiv:hep-ph/9808299.
- [541] J. R. Espinosa and R.-J. Zhang, "MSSM lightest CP-even Higgs boson mass to $\mathcal{O}(\alpha_s \alpha_t)$: The effective potential approach," *JHEP* **03** (2000) 026, arXiv:hep-ph/9912236.
- [542] G. Degrassi, P. Slavich, and F. Zwirner, "On the neutral Higgs boson masses in the MSSM for arbitrary stop mixing," *Nucl. Phys.* **B611** (2001) 403–422, arXiv:hep-ph/0105096.
- [543] R. Hempfling, "Yukawa coupling unification with supersymmetric threshold corrections," *Phys. Rev.* **D49** (1994) 6168–6172.

- [544] L. J. Hall, R. Rattazzi, and U. Sarid, "The Top quark mass in supersymmetric SO(10) unification," *Phys. Rev.* **D50** (1994) 7048–7065, arXiv:hep-ph/9306309.
- [545] M. Carena, M. Olechowski, S. Pokorski, and C. Wagner, "Electroweak symmetry breaking and bottom-top Yukawa unification," *Nucl. Phys.* **B426** (1994) 269–300, arXiv:hep-ph/9402253.
- [546] J. Guasch, P. Haflinger, and M. Spira, "MSSM Higgs decays to bottom quark pairs revisited," *Phys. Rev.* **D68** (2003) 115001, arXiv:hep-ph/0305101.
- [547] A. Dedes, G. Degrossi, and P. Slavich, "On the two-loop Yukawa corrections to the MSSM Higgs boson masses at large $\tan(\beta)$," *Nucl. Phys.* **B672** (2003) 144–162, arXiv:hep-ph/0305127.
- [548] H. Eberl, K. Hidaka, S. Kraml, W. Majerotto, and Y. Yamada, "Improved SUSY QCD corrections to Higgs boson decays into quarks and squarks," *Phys. Rev.* **D62** (2000) 055006, arXiv:hep-ph/9912463.
- [549] M. Carena, D. Garcia, U. Nierste, and C. E. Wagner, "Effective Lagrangian for the $\bar{t}bH^+$ interaction in the MSSM and charged Higgs phenomenology," *Nucl. Phys.* **B577** (2000) 88–120. doi:0.1016/S0550-3213(00)00146-2.
- [550] S. Heinemeyer, W. Hollik, and G. Weiglein, "Electroweak precision observables in the minimal supersymmetric standard model," *Phys. Rept.* **425** (2006) 265–368, arXiv:hep-ph/0412214.
- [551] M. Carena and H. E. Haber, "Higgs boson theory and phenomenology. ((V))," *Prog. Part. Nucl. Phys.* **50** (2003) 63–152, arXiv:hep-ph/0208209.
- [552] **ALEPH, DELPHI, L3 and OPAL Collaborations. The LEP Working Group for Higgs Boson Searches** Collaboration, "Search for neutral MSSM Higgs bosons at LEP," arXiv:hep-ex/0602042.
- [553] **LEP Higgs Working Group for Higgs boson searches** Collaboration, "Search for charged Higgs bosons: Preliminary combined results using LEP data collected at energies up to 209- GeV," arXiv:hep-ex/0107031.
- [554] A. Dabelstein, "Fermionic decays of neutral MSSM Higgs bosons at the one loop level," *Nucl. Phys.* **B456** (1995) 25–56, arXiv:hep-ph/9503443.
- [555] S. Heinemeyer, W. Hollik, and G. Weiglein, "Decay widths of the neutral CP-even MSSM Higgs bosons in the Feynman-diagrammatic approach," *Eur. Phys. J.* **C16** (2000) 139–153, arXiv:hep-ph/0003022.
- [556] A. Djouadi, P. Janot, J. Kalinowski, and P. M. Zerwas, "SUSY Decays of Higgs Particles," *Phys. Lett.* **B376** (1996) 220–226, arXiv:hep-ph/9603368.
- [557] A. Djouadi, J. Kalinowski, P. Ohmann, and P. M. Zerwas, "Heavy SUSY Higgs bosons at e^+e^- linear colliders," *Z. Phys.* **C74** (1997) 93–111, arXiv:hep-ph/9605339.
- [558] S. Dawson, A. Djouadi, and M. Spira, "QCD Corrections to SUSY Higgs Production: The Role of Squark Loops," *Phys. Rev. Lett.* **77** (1996) 16–19, arXiv:hep-ph/9603423.

- [559] R. V. Harlander and M. Steinhauser, "Hadronic Higgs production and decay in supersymmetry at next-to-leading order," *Phys. Lett.* **B574** (2003) 258–268, arXiv:hep-ph/0307346.
- [560] R. Harlander and M. Steinhauser, "Effects of SUSY-QCD in hadronic Higgs production at next- to-next-to-leading order," *Phys. Rev.* **D68** (2003) 111701, arXiv:hep-ph/0308210.
- [561] R. V. Harlander and M. Steinhauser, "Supersymmetric Higgs production in gluon fusion at next-to- leading order," *JHEP* **09** (2004) 066, arXiv:hep-ph/0409010.
- [562] R. V. Harlander and F. Hofmann, "Pseudo-scalar Higgs production at next-to-leading order SUSY-QCD," arXiv:hep-ph/0507041.
- [563] U. Langenegger, M. Spira, A. Starodumov, and P. Trueb, "SM and MSSM Higgs boson production: Spectra at large transverse momentum," arXiv:hep-ph/0604156.
- [564] A. Djouadi and M. Spira, "SUSY-QCD corrections to Higgs boson production at hadron colliders," *Phys. Rev.* **D62** (2000) 014004, arXiv:hep-ph/9912476.
- [565] P. Wu et al., "NLO supersymmetric QCD corrections to t anti- t h_0 associated production at hadron colliders," arXiv:hep-ph/0505086.
- [566] S. Dittmaier, M. Kramer, and M. Spira, "Higgs radiation off bottom quarks at the Tevatron and the LHC," *Phys. Rev.* **D70** (2004) 074010, arXiv:hep-ph/0309204.
- [567] S. Dawson, C. B. Jackson, L. Reina, and D. Wackerroth, "Exclusive Higgs boson production with bottom quarks at hadron colliders," *Phys. Rev.* **D69** (2004) 074027, arXiv:hep-ph/0311067.
- [568] D. A. Dicus and S. Willenbrock, "Higgs Boson Production from Heavy Quark Fusion," *Phys. Rev.* **D39** (1989) 751.
- [569] D. Dicus, T. Stelzer, Z. Sullivan, and S. Willenbrock, "Higgs boson production in association with bottom quarks at next-to-leading order," *Phys. Rev.* **D59** (1999) 094016, arXiv:hep-ph/9811492.
- [570] C. Balazs, H.-J. He, and C. P. Yuan, "QCD corrections to scalar production via heavy quark fusion at hadron colliders," *Phys. Rev.* **D60** (1999) 114001, arXiv:hep-ph/9812263.
- [571] R. V. Harlander and W. B. Kilgore, "Higgs boson production in bottom quark fusion at next-to- next-to-leading order," *Phys. Rev.* **D68** (2003) 013001, arXiv:hep-ph/0304035.
- [572] M. A. G. Aivazis, F. I. Olness, and W.-K. Tung, "Leptoproduction of heavy quarks. 1. General formalism and kinematics of charged current and neutral current production processes," *Phys. Rev.* **D50** (1994) 3085–3101, arXiv:hep-ph/9312318.
- [573] M. A. G. Aivazis, J. C. Collins, F. I. Olness, and W.-K. Tung, "Leptoproduction of heavy quarks. 2. A Unified QCD formulation of charged and neutral current processes from fixed target to collider energies," *Phys. Rev.* **D50** (1994) 3102–3118, arXiv:hep-ph/9312319.


- [574] J. Campbell et al., “Higgs boson production in association with bottom quarks,” [arXiv:hep-ph/0405302](https://arxiv.org/abs/hep-ph/0405302).
- [575] J. Campbell, R. K. Ellis, F. Maltoni, and S. Willenbrock, “Higgs boson production in association with a single bottom quark,” *Phys. Rev.* **D67** (2003) 095002, [arXiv:hep-ph/0204093](https://arxiv.org/abs/hep-ph/0204093).
- [576] A. C. Bawa, C. S. Kim, and A. D. Martin, “Charged Higgs Production at Hadron Colliders,” *Z. Phys.* **C47** (1990) 75–82.
- [577] F. Borzumati, J.-L. Kneur, and N. Polonsky, “Higgs-strahlung and R-parity violating slepton-strahlung at hadron colliders,” *Phys. Rev.* **D60** (1999) 115011, [arXiv:hep-ph/9905443](https://arxiv.org/abs/hep-ph/9905443).
- [578] A. Belyaev, D. Garcia, J. Guasch, and J. Sola, “Prospects for heavy supersymmetric charged Higgs boson searches at hadron colliders,” *JHEP* **06** (2002) 059, [arXiv:hep-ph/0203031](https://arxiv.org/abs/hep-ph/0203031).
- [579] P. Wu et al., “NLO supersymmetric QCD corrections to the t anti- b H^- associated production at hadron colliders,” *Phys. Rev.* **D73** (2006) 015012, [arXiv:hep-ph/0601069](https://arxiv.org/abs/hep-ph/0601069). doi:10.1103/PhysRevD.73.015012.
- [580] S.-H. Zhu, “Complete next-to-leading order QCD corrections to charged Higgs boson associated production with top quark at the CERN Large Hadron Collider,” *Phys. Rev.* **D67** (2003) 075006. doi:10.1103/PhysRevD.67.075006.
- [581] T. Plehn, “Charged Higgs boson production in bottom-gluon fusion,”. Prepared for 10th International Conference on Supersymmetry and Unification of Fundamental Interactions (SUSY02), Hamburg, Germany, 17-23 Jun 2002.
- [582] E. L. Berger, T. Han, J. Jiang, and T. Plehn, “Associated production of a top quark and a charged Higgs boson,” *Phys. Rev.* **D71** (2005) 115012, [arXiv:hep-ph/0312286](https://arxiv.org/abs/hep-ph/0312286).
- [583] G. Gao, G. Lu, Z. Xiong, and J. M. Yang, “Loop effects and non-decoupling property of SUSY QCD in $gb \rightarrow tH^-$,” *Phys. Rev.* **D66** (2002) 015007. doi:10.1103/PhysRevD.66.015007.
- [584] S. S. D. Willenbrock, “Pair Production of Supersymmetric Charged Higgs Bosons,” *Phys. Rev.* **D35** (1987) 173.
- [585] A. Krause, T. Plehn, M. Spira, and P. M. Zerwas, “Production of charged Higgs boson pairs in gluon gluon collisions,” *Nucl. Phys.* **B519** (1998) 85–100, [arXiv:hep-ph/9707430](https://arxiv.org/abs/hep-ph/9707430).
- [586] Y. Jiang, L. Han, W.-G. Ma, Z.-H. Yu, and M. Han, “Pair production of charged Higgs bosons in gluon gluon collisions,” *J. Phys.* **G23** (1997) 385–400, [arXiv:hep-ph/9703275](https://arxiv.org/abs/hep-ph/9703275).
- [587] O. Brein and W. Hollik, “Pair production of charged MSSM Higgs bosons by gluon fusion,” *Eur. Phys. J.* **C13** (2000) 175–184, [arXiv:hep-ph/9908529](https://arxiv.org/abs/hep-ph/9908529).
- [588] A. A. Barrientos Bendezu and B. A. Kniehl, “ H^+H^- pair production at the Large Hadron Collider,” *Nucl. Phys.* **B568** (2000) 305–318. doi:10.1016/S0550-3213(99)00732-4.

- [589] H.-S. Hou et al., "Pair production of charged Higgs bosons from bottom-quark fusion," *Phys. Rev.* **D71** (2005) 075014. doi:10.1103/PhysRevD.71.075014.
- [590] A. A. Barrientos Bendezu and B. A. Kniehl, " $W^\pm H^\mp$ associated production at the Large Hadron Collider," *Phys. Rev.* **D59** (1999) 015009, arXiv:hep-ph/9807480. doi:10.1103/PhysRevD.59.015009.
- [591] O. Brein, W. Hollik, and S. Kanemura, "The MSSM prediction for $W^{+-} H^+$ production by gluon fusion," *Phys. Rev.* **D63** (2001) 095001, arXiv:hep-ph/0008308.
- [592] A. A. Barrientos Bendezu and B. A. Kniehl, "Squark loop correction to $W^{+-} H^+$ associated hadroproduction," *Phys. Rev.* **D63** (2001) 015009, arXiv:hep-ph/0007336.
- [593] D. Dicus, J. Hewett, C. Kao, and T. Rizzo, " $W^\pm H^\mp$ production at hadron colliders," *Phys. Rev.* **D40** (1989) 787.
- [594] W. Hollik and S.-h. Zhu, " $\mathcal{O}(\alpha_f)$ corrections to $b\bar{b} \rightarrow W^\pm H^\mp$ at the CERN Large Hadron Collider," *Phys. Rev.* **D65** (2002) 075015, arXiv:hep-ph/0109103. doi:10.1103/PhysRevD.65.075015.
- [595] J. Zhao, C. S. Li, and Q. Li, "SUSY-QCD corrections to $W^\pm H^\mp$ associated production at the CERN Large Hadron Collider," *Phys. Rev.* **D72** (2005) 114008, arXiv:hep-ph/0509369.
- [596] S. Lehti, "Study of $gg \rightarrow b\bar{b} H_{SUSY}, H_{SUSY} \rightarrow \tau\tau \rightarrow \ell\ell + X$," *CMS Note* **2002/035** (2002).
- [597] D. Bloch. Private communication.
- [598] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 12.2.3: Combined secondary vertex tag," *CERN/LHCC* **2006-001** (2006) 466.
- [599] GEANT4 Collaboration, S. Agostinelli et al., "GEANT4: A simulation toolkit," *Nucl. Instrum. Meth.* **A506** (2003) 250–303.
- [600] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 12.2.6: HLT b tag," *CERN/LHCC* **2006-001** (2006) 474.
- [601] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 11.6.4: Parton-level corrections," *CERN/LHCC* **2006-001** (2006) 428.
- [602] DØ Collaboration, V. M. Abazov et al., "Search for neutral supersymmetric Higgs bosons in multijet events at $\sqrt{s} = 1.96$ TeV," *Phys. Rev. Lett.* **95** (2005) 151801. doi:10.1103/PhysRevLett.95.151801.
- [603] D. Roy, "The hadronic tau decay signature of a heavy charged Higgs boson at LHC," *Phys. Lett.* **B459** (1999) 607–614, arXiv:hep-ph/9905542.
- [604] R. Kinnunen, "Study of $A/H \rightarrow \tau\tau$ and $H^\pm \rightarrow \tau\nu$ in CMS," *Czech. J. Phys.* **54** (2004) A93–A101.
- [605] K. A. Assamagan and Y. Coadou, "The hadronic tau decay of a heavy H^{+-} in ATLAS," *Acta Phys. Polon.* **B33** (2002) 707–720.


- [606] D. Roy, "Looking for the charged Higgs boson," *Mod. Phys. Lett.* **A19** (2004) 1813–1828, arXiv:hep-ph/0406102.
- [607] J. Alwall and J. Rathsman, "Improved description of charged Higgs boson production at hadron colliders," *JHEP* **12** (2004) 050, arXiv:hep-ph/0409094.
- [608] T. Plehn, "Charged Higgs boson production in bottom gluon fusion," *Phys. Rev.* **D67** (2003) 014018, arXiv:hep-ph/0206121.
- [609] S. Lowette, J. D'Hondt, and P. Vanlaer, "Charged MSSM Higgs Boson Observability in the $H^\pm \rightarrow tb$ Decay," 2006. CMS AN 2006/012, CERN.
- [610] CMS Collaboration, G. Anagnostou and G. Daskalakis, "Search for the MSSM $A \rightarrow Zh$ decay with $Z \rightarrow \ell^+ \ell^-$, $h \rightarrow b\bar{b}$," *CMS Note* **2006/063** (2006).
- [611] B. C. Allanach et al., "The Snowmass points and slopes: Benchmarks for SUSY searches," *Eur. Phys. J.* **C25** (2002) 113–123, arXiv:hep-ph/0202233.
- [612] M. Battaglia et al., "Updated post-WMAP benchmarks for supersymmetry," *Eur. Phys. J.* **C33** (2004) 273–296, arXiv:hep-ph/0306219.
- [613] M. Carena, S. Heinemeyer, C. Wagner, and G. Weiglein, "Suggestions for improved benchmark scenarios for Higgs-boson searches at LEP2," arXiv:hep-ph/9912223.
- [614] M. Carena, S. Heinemeyer, C. Wagner, and G. Weiglein, "Suggestions for benchmark scenarios for MSSM Higgs boson searches at hadron colliders," *Eur. Phys. J.* **C26** (2003) 601–607, arXiv:hep-ph/0202167.
- [615] LHC/LC Study Group Collaboration, G. Weiglein et al., "Physics interplay of the LHC and the ILC," arXiv:hep-ph/0410364.
- [616] ECFA/DESY LC Physics Working Group Collaboration, J. A. Aguilar-Saavedra et al., "TESLA Technical Design Report Part III: Physics at an e+e- Linear Collider," arXiv:hep-ph/0106315.
- [617] C. L. Bennett et al., "First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Preliminary Maps and Basic Results," *Astrophys. J. Suppl.* **148** (2003) 1, arXiv:astro-ph/0302207.
- [618] WMAP Collaboration, D. N. Spergel et al., "First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters," *Astrophys. J. Suppl.* **148** (2003) 175, arXiv:astro-ph/0302209.
- [619] H. Goldberg, "Constraint on the photino mass from cosmology," *Phys. Rev. Lett.* **50** (1983) 1419.
- [620] J. R. Ellis, J. Hagelin, D. Nanopoulos, K. A. Olive, and M. Srednicki, "Supersymmetric relics from the big bang," *Nucl. Phys.* **B238** (1984) 453–476.
- [621] Heavy Flavor Averaging Group Collaboration, E. Barberio et al., "Averages of b-hadron properties at the end of 2005," arXiv:hep-ex/0603003.
- [622] H. M. Asatrian, A. Hovhannisyan, V. Poghosyan, C. Greub, and T. Hurth, "Towards the NNLL precision in anti-B \rightarrow X/s gamma," arXiv:hep-ph/0512097.

- [623] Muon Collaboration, G. W. Bennett, "Final report of the muon E821 anomalous magnetic moment measurement at BNL," [arXiv:hep-ex/0602035](https://arxiv.org/abs/hep-ex/0602035).
- [624] A. Czarnecki and W. J. Marciano, "The muon anomalous magnetic moment: A harbinger for 'new physics'," *Phys. Rev.* **D64** (2001) 013014, [arXiv:hep-ph/0102122](https://arxiv.org/abs/hep-ph/0102122).
- [625] M. Carena et al., "Reconciling the two-loop diagrammatic and effective field theory computations of the mass of the lightest CP-even Higgs boson in the MSSM," *Nucl. Phys.* **B580** (2000) 29–57, [arXiv:hep-ph/0001002](https://arxiv.org/abs/hep-ph/0001002).
- [626] L. Randall and R. Sundrum, "An alternative to compactification," *Phys. Rev. Lett.* **83** (1999) 4690–4693, [arXiv:hep-th/9906064](https://arxiv.org/abs/hep-th/9906064), [doi:10.1103/PhysRevLett.83.4690](https://doi.org/10.1103/PhysRevLett.83.4690).
- [627] G. F. Giudice, R. Rattazzi, and J. D. Wells, "Graviscalars from higher-dimensional metrics and curvature- Higgs mixing," *Nucl. Phys.* **B595** (2001) 250–276, [arXiv:hep-ph/0002178](https://arxiv.org/abs/hep-ph/0002178).
- [628] M. Chaichian, A. Datta, K. Huitu, and Z.-h. Yu, "Radion and Higgs mixing at the LHC," *Phys. Lett.* **B524** (2002) 161–169, [arXiv:hep-ph/0110035](https://arxiv.org/abs/hep-ph/0110035).
- [629] J. L. Hewett and T. G. Rizzo, "Shifts in the properties of the Higgs boson from radion mixing," *JHEP* **08** (2003) 028, [arXiv:hep-ph/0202155](https://arxiv.org/abs/hep-ph/0202155).
- [630] D. Dominici, B. Grzadkowski, J. F. Gunion, and M. Toharia, "The scalar sector of the Randall-Sundrum model," *Nucl. Phys.* **B671** (2003) 243–292, [arXiv:hep-ph/0206192](https://arxiv.org/abs/hep-ph/0206192).
- [631] M. Battaglia, S. De Curtis, A. De Roeck, D. Dominici, and J. F. Gunion, "On the complementarity of Higgs and radion searches at LHC," *Phys. Lett.* **B568** (2003) 92–102, [arXiv:hep-ph/0304245](https://arxiv.org/abs/hep-ph/0304245).
- [632] G. Azuelos, D. Cavalli, H. Przysieznik, and L. Vacavant, "Search for the radion using the ATLAS detector," *Eur. Phys. J. direct* **C4** (2002) 16.
- [633] D. Dominici, G. Dewhurst, A. Nikitenko, S. Gennai, and L. Fanò, "Search for radion decays into Higgs boson pairs in the $\gamma\gamma b\bar{b}$, $\tau\tau b\bar{b}$ and $b\bar{b}b\bar{b}$ final states," *CMS Note* **2005/007** (2005).
- [634] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, "(De)constructing dimensions," *Phys. Rev. Lett.* **86** (2001) 4757–4761, [arXiv:hep-th/0104005](https://arxiv.org/abs/hep-th/0104005).
- [635] H.-C. Cheng, C. T. Hill, S. Pokorski, and J. Wang, "The standard model in the latticized bulk," *Phys. Rev.* **D64** (2001) 065007, [arXiv:hep-th/0104179](https://arxiv.org/abs/hep-th/0104179).
- [636] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, "Electroweak symmetry breaking from dimensional deconstruction," *Phys. Lett.* **B513** (2001) 232–240, [arXiv:hep-ph/0105239](https://arxiv.org/abs/hep-ph/0105239).
- [637] N. Arkani-Hamed, A. G. Cohen, E. Katz, and A. E. Nelson, "The littlest Higgs," *JHEP* **07** (2002) 034, [arXiv:hep-ph/0206021](https://arxiv.org/abs/hep-ph/0206021).


- [638] J. Schechter and J. Valle, "Neutrino masses in $SU(2) \times U(1)$ Theories," *Phys. Rev.* **D22** (1980) 2227. doi:10.1103/PhysRevD.22.2227.
- [639] E. Ma and U. Sarkar, "Neutrino masses and leptogenesis with heavy Higgs triplets," *Phys. Rev. Lett.* **80** (1998) 5716–5719, arXiv:hep-ph/9802445.
- [640] E. Ma, M. Raidal, and U. Sarkar, "Verifiable model of neutrino masses from large extra dimensions," *Phys. Rev. Lett.* **85** (2000) 3769–3772, arXiv:hep-ph/0006046.
- [641] E. Ma, M. Raidal, and U. Sarkar, "Phenomenology of the neutrino-mass-giving Higgs triplet and the low-energy seesaw violation of lepton number," *Nucl. Phys.* **B615** (2001) 313–330, arXiv:hep-ph/0012101.
- [642] E. Ma and U. Sarkar, "Connecting dark energy to neutrinos with an observable Higgs triplet," arXiv:hep-ph/0602116.
- [643] G. Marandella, C. Schappacher, and A. Strumia, "Little-Higgs corrections to precision data after LEP2," *Phys. Rev.* **D72** (2005) 035014, arXiv:hep-ph/0502096.
- [644] K. Huitu, J. Maalampi, A. Pietila, and M. Raidal, "Doubly charged Higgs at LHC," *Nucl. Phys.* **B487** (1997) 27–42, arXiv:hep-ph/9606311.
- [645] T. Han, H. E. Logan, and L.-T. Wang, "Smoking-gun signatures of little Higgs models," arXiv:hep-ph/0506313.
- [646] G. Azuelos et al., "Exploring little Higgs models with ATLAS at the LHC," *Eur. Phys. J.* **C39S2** (2005) 13–24, arXiv:hep-ph/0402037.
- [647] J. F. Gunion, J. Grifols, A. Mendez, B. Kayser, and F. I. Olness, "Higgs bosons in left-right symmetric models," *Phys. Rev.* **D40** (1989) 1546. doi:10.1103/PhysRevD.40.1546.
- [648] M. Muhlleitner and M. Spira, "A note on doubly-charged Higgs pair production at hadron colliders," *Phys. Rev.* **D68** (2003) 117701, arXiv:hep-ph/0305288.
- [649] G. Bagliesi, S. Dutta, S. Gennai, A. Kalinowski, M. Konecki, D. Kotlinski, F. Moortgat, A. Nikitenko, L. Wendland, and S. Wakefield, "Tau jet reconstruction and tagging at High Level Trigger and off-line," *CMS Note* **2006/028** (2006).
- [650] S. Bityukov, N. Krasnikov, and V. Taperechkina, "Confidence intervals for Poisson distribution parameter," arXiv:hep-ex/0108020.
- [651] S. P. Martin, "A supersymmetry primer," in *Perspectives in Supersymmetry*, ed. G. Kane, *World Scientific* (1997) arXiv:hep-ph/9709356.
- [652] CMS Collaboration, S. Abdullin et al., "Discovery potential for supersymmetry in CMS," *J. Phys.* **G28** (2002) 469. doi:10.1088/0954-3899/28/3/401.
- [653] F. E. Paige, S. D. Protopopescu, H. Baer, and X. Tata, "ISAJET 7.69: A Monte Carlo Event Generator for pp , $\bar{p}p$, and e^+e^- Reactions," arXiv:hep-ph/0312045.
- [654] A. De Roeck et al., "Supersymmetric benchmarks with non-universal scalar masses or gravitino dark matter," arXiv:hep-ph/0508198.

- [655] W. de Boer et al., "Excess of EGRET galactic gamma ray data interpreted as dark matter annihilation," arXiv:astro-ph/0408272.
- [656] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume1, Section 10," CERN/LHCC 2006-001 (2006).
- [657] A. Nikitenko, G. Quast, and V. Ciulli, "Minimal requirements for significance estimates of the physics result." Info  **NAME: Need more than this.** located at <http://cmsdoc.cern.ch/~anikiten/cms-higgs/stat.txt>.
- [658] W. Beenakker, R. Hopker, and M. Spira, "PROSPINO: A program for the PROduction of Supersymmetric Particles In Next-to-leading Order QCD," arXiv:hep-ph/9611232.
- [659] B. C. Allanach, C. G. Lester, M. A. Parker, and B. R. Webber, "Measuring sparticle masses in non-universal string inspired models at the LHC," *JHEP* **09** (2000) 004, arXiv:hep-ph/0007009.
- [660] W. Beenakker et al., "The production of charginos/neutralinos and sleptons at hadron colliders," *Phys. Rev. Lett.* **83** (1999) 3780–3783. doi:10.1103/PhysRevLett.83.3780.
- [661] F. del Aguila and L. Ametller, "On the detectability of sleptons at large hadron colliders," *Phys. Lett.* **B261** (1991) 326–333. doi:10.1016/0370-2693(91)90336-0.
- [662] H. Baer, C.-h. Chen, F. Paige, and X. Tata, "Detecting sleptons at hadron colliders and supercolliders," *Phys. Rev.* **D49** (1994) 3283–3290, arXiv:hep-ph/9311248.
- [663] D. Denegri, L. Rurua, and N. Stepanov, "Detection of Sleptons in CMS, Mass Reach," *CMS TN* **96-059** (1996).
- [664] Y. M. Andreev, S. I. Bityukov, and N. V. Krasnikov, "Sleptons at post-WMAP benchmark points at LHC(CMS)," *Phys. Atom. Nucl.* **68** (2005) 340–347, arXiv:hep-ph/0402229.
- [665] S. Bityukov and N. Krasnikov, "The search for sleptons and lepton-flavor-number violation at LHC (CMS)," *Phys. Atom. Nucl.* **62** (1999) 1213–1225, arXiv:hep-ph/9712358.
- [666] N. Krasnikov, "Flavor lepton number violation at LEP-2," *Mod. Phys. Lett.* **A9** (1994) 791–794.
- [667] N. Arkani-Hamed, H.-C. Cheng, J. L. Feng, and L. J. Hall, "Probing Lepton Flavor Violation at Future Colliders," *Phys. Rev. Lett.* **77** (1996) 1937–1940, arXiv:hep-ph/9603431. doi:10.1103/PhysRevLett.77.1937.
- [668] N. Krasnikov, "Search for flavor lepton number violation in slepton decays at LHC," *JETP Lett.* **65** (1997) 148. doi:10.1134/1.567315.
- [669] K. Agashe and M. Graesser, "Signals of supersymmetric lepton flavor violation at the LHC," *Phys. Rev.* **D61** (2000) 075008, arXiv:hep-ph/9904422.

- [670] J. Hisano, R. Kitano, and M. M. Nojiri, "Flavor mixing in slepton production at the large hadron collider," *Phys. Rev.* **D65** (2002) 116002.
- [671] H. Baer, C.-h. Chen, F. Paige, and X. Tata, "Trileptons from chargino - neutralino production at the CERN Large Hadron Collider," *Phys. Rev.* **D50** (1994) 4508–4516, arXiv:hep-ph/9404212. doi:10.1103/PhysRevD.50.4508.
- [672] J. R. Ellis, K. A. Olive, and Y. Santoso, "The MSSM parameter space with non-universal Higgs masses," *Phys. Lett.* **B539** (2002) 107–118, arXiv:hep-ph/0204192.
- [673] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, "The hierarchy problem and new dimensions at a millimeter," *Phys. Lett.* **B429** (1998) 263–272, arXiv:hep-ph/9803315.
- [674] J. D. Lykken, E. Poppitz, and S. P. Trivedi, "Branes with GUTs and supersymmetry breaking," *Nucl. Phys.* **B543** (1999) 105–121, arXiv:hep-th/9806080.
- [675] I. Antoniadis, "A Possible new dimension at a few TeV," *Phys. Lett.* **B246** (1990) 377–384. doi:10.1016/0370-2693(90)90617-F.
- [676] J. D. Lykken, "Weak Scale Superstrings," *Phys. Rev.* **D54** (1996) 3693–3697, arXiv:hep-th/9603133. doi:10.1103/PhysRevD.54.R3693.
- [677] G. F. Giudice, R. Rattazzi, and J. D. Wells, "Quantum gravity and extra dimensions at high-energy colliders," *Nucl. Phys.* **B544** (1999) 3–38, arXiv:hep-ph/9811291.
- [678] T. Han, J. D. Lykken, and R.-J. Zhang, "On Kaluza-Klein states from large extra dimensions," *Phys. Rev.* **D59** (1999) 105006, arXiv:hep-ph/9811350. doi:10.1103/PhysRevD.59.105006.
- [679] S. Cullen, M. Perelstein, and M. E. Peskin, "TeV strings and collider probes of large extra dimensions," *Phys. Rev.* **D62** (2000) 055012, arXiv:hep-ph/0001166.
- [680] J. D. Lykken and L. Randall, "The shape of gravity," *JHEP* **06** (2000) 014, arXiv:hep-th/9908076.
- [681] L. Vacavant and I. Hinchliffe, "Signals of models with large extra dimensions in ATLAS," *J. Phys.* **G27** (2001) 1839–1850.
- [682] J. Ruppert, C. Rahmede, and M. Bleicher, "Determination of the fundamental scale of gravity and the number of space-time dimensions from high energetic particle interactions," *Phys. Lett.* **B608** (2005) 240–243, arXiv:hep-ph/0501028.
- [683] E. Dudas and J. Mourad, "String theory predictions for future accelerators," *Nucl. Phys.* **B575** (2000) 3–34, arXiv:hep-th/9911019.
- [684] D. Chialva, R. Iengo, and J. G. Russo, "Cross sections for production of closed superstrings at high energy colliders in brane world models," *Phys. Rev.* **D71** (2005) 106009, arXiv:hep-ph/0503125.
- [685] M. Bando, T. Kugo, T. Noguchi, and K. Yoshioka, "Brane fluctuation and suppression of Kaluza-Klein mode couplings," *Phys. Rev. Lett.* **83** (1999) 3601–3604, arXiv:hep-ph/9906549.



- [686] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, "Phenomenology of the Randall-Sundrum gauge hierarchy model," *Phys. Rev. Lett.* **84** (2000) 2080. doi:10.1103/PhysRevLett.84.2080.
- [687] J. Hewett and M. Spiropulu, "Particle physics probes of extra spacetime dimensions," *Ann. Rev. Nucl. Part. Sci.* **52** (2002) 397–424, arXiv:hep-ph/0205106. doi:10.1146/annurev.nucl.52.050102.090706.
- [688] B. Clerbaux, T. Mahmoud, C. Collard, and P. Miné, "Search  the CMS detector for heavy resonances decaying into an electron pair," *CMS AN ~~NOTE~~: No AN's 2006-045* (2006).
- [689] C. Collard and M.-C. Lemaire, "Search with the CMS Detector for Randall-Sundrum Excitations of Gravitons Decaying Into Electron Pairs," *CMS Note 2004/024* (2004).
- [690] B. Clerbaux, T. Mahmoud, C. Collard, M.-C. Lemaire, and V. Litvin, "TeV electron and photon saturation studies," *CMS Note 2006/004* (2006).
- [691] CDF Collaboration, T. Affolder et al., "Measurement of ds/dM and Forward-Backward Charge Asymmetry for High-mass Drell-Yan e^+e^- Pairs from collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **87** (2001) 131802.
- [692] CTEQ Collaboration, H. L. Lai et al., "Global QCD analysis of parton structure of the nucleon: CTEQ5 parton distributions," *Eur. Phys. J.* **C12** (2000) 375–392, arXiv:hep-ph/9903282. doi:10.1007/s100529900196.
- [693] CTEQ Collaboration, D. Stump et al., "Inclusive Jet Production, Parton Distributions, and the Search for New Physics," *JHEP* **0310** (2003) 046, arXiv:hep-ph/0303013.
- [694] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, "Experimental probes of localized gravity: On and off the wall," *Phys. Rev.* **D63** (2001) 075004, arXiv:hep-ph/0006041.
- [695] K.-m. Cheung and G. Landsberg, "Drell-Yan and diphoton production at hadron colliders and low scale gravity model," *Phys. Rev.* **D62** (2000) 076003, arXiv:hep-ph/9909218.
- [696] R. Mohapatra and J. Pati, "Left-right gauge symmetry and an 'isoconjugate' model of CP violation," *Phys. Rev.* **D11** (1975) 566–571. doi:10.1103/PhysRevD.11.566.
- [697] G. Senjanovic and R. Mohapatra, "Exact Left-Right Symmetry and Spontaneous Violation of Parity," *Phys. Rev.* **D12** (1975) 1502. doi:10.1103/PhysRevD.12.1502.
- [698] R. Mohapatra and G. Senjanovic, "Neutrino mass and spontaneous parity nonconservation," *Phys. Rev. Lett.* **44** (1980) 912. doi:10.1103/PhysRevLett.44.912.
- [699] J. Pati and A. Salam, "Lepton Number as the Fourth Color," *Phys. Rev.* **D10** (1974) 275–289.
- [700] G. Altarelli, B. Mele, and M. Ruiz-Altaba, "Searching for New Heavy Vector Bosons in $p\bar{p}$ Colliders," *Z. Phys.* **C45** (1989) 109. doi:10.1007/BF01556677.

- [701] CDF Collaboration, T. Affolder et al., "Search for quark lepton compositeness and a heavy W' boson using the $e\nu$ channel in p anti- p collisions at $s^{*1/2} = 1.8$ -TeV," *Phys. Rev. Lett.* **87** (2001) 231803, arXiv:hep-ex/0107008.
- [702] K. Gumus, N. Akchurin, S. Esen, and R. Harris, "CMS Sensitivity to Dijet Resonances," *CMS Note* **2006/070** (2006).
- [703] U. Baur, I. Hinchliffe, and D. Zeppenfeld, "Excited Quark Production at Hadron Colliders," *Int. J. Mod. Phys.* **A2** (1987) 1285.
- [704] J. Bagger, C. Schmidt, and S. King, "Axigluon Production in Hadronic Collisions," *Phys. Rev.* **D37** (1988) 1188.
- [705] R. S. Chivukula, A. G. Cohen, and E. H. Simmons, "New Strong Interactions at the Tevatron?," *Phys. Lett.* **B380** (1996) 92–98, arXiv:hep-ph/9603311.
- [706] J. L. Hewett and T. G. Rizzo, "Low-Energy Phenomenology of Superstring Inspired $E(6)$ Models," *Phys. Rept.* **183** (1989) 193.
- [707] K. Lane and S. Mrenna, "The collider phenomenology of technihadrons in the technicolor Straw Man Model," *Phys. Rev.* **D67** (2003) 115011.
doi:10.1103/PhysRevD.67.115011.
- [708] E. Eichten, I. Hinchliffe, K. D. Lane, and C. Quigg, "Super Collider Physics," *Rev. Mod. Phys.* **56** (1984) 579–707.
- [709] CMS Collaboration, M. Pieri, S. Bhattacharya, J. Branson, J. Letts, and V. Litvin, "Generator Level Selection of Multi-Jet and $\gamma +$ Jets Events for $H \rightarrow \gamma\gamma$ Search," *CMS IN* **2005/018** (2005).
- [710] CMS Collaboration, M.-C. Lemaire, H. Newman, and V. Litvin, "Search for Randall-Sundrum K_1 excitations of gravitons decaying into two photons for CMS at LHC," *CMS AN* **NAME: AN's are not generally available 2005-032** (2005).
- [711] CMS Collaboration, B. Clerbaux, T. Mahmoud, C. Collard, M.-C. Lemaire, and V. Litvin, "TeV electron and photon saturation studies," *CMS AN* **NAME: AN's are not generally available 2005-049** (2005).
- [712] D. Acosta et al., "Measurement of the Cross Section for Prompt Diphoton Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV," *Phys. Rev. Lett.* **95** (2005) 022003.
doi:10.1103/PhysRevLett.95.022003.
- [713] R. C. Myers and M. J. Perry, "Black holes in higher dimensional space-times," *Ann. Phys.* **172** (1986) 304.
- [714] S. W. Hawking, "Particle creation by black holes," *Commun. Math. Phys.* **43** (1975) 199–220.
- [715] P. C. Argyres, S. Dimopoulos, and J. March-Russell, "Black holes and sub-millimeter dimensions," *Phys. Lett.* **B441** (1998) 96–104, arXiv:hep-th/9808138.
- [716] C. M. Harris, P. Richardson, and B. R. Webber, "CHARYBDIS: A black hole event generator," *JHEP* **08** (2003) 033, arXiv:hep-ph/0307305.

- [717] K. D. Lane, "Technicolor 2000," arXiv:hep-ph/0007304.
- [718] B. Betev, D. Bourilkov, and S. S. Mavrodiev, "Structure Functions of Pion and Nucleon Determined from High Mass Muon Pair Production," JINR-E2-85-312.
- [719] **Super-Kamiokande** Collaboration, Y. Fukuda et al., "Evidence for oscillation of atmospheric neutrinos," *Phys. Rev. Lett.* **81** (1998) 1562–1567, arXiv:hep-ex/9807003.
- [720] S. Esen and R. Harris, "CMS Sensitivity to Quark Contact Interactions using Dijets," *CMS Note* **2006/071** (2006).
- [721] N. Krasnikov and V. Matveev, "Search for new physics at LHC," *Phys. Usp.* **47** (2004) 643, arXiv:hep-ph/0309200.
- [722] J. C. Pati and A. Salam, "Anomalous lepton-hadron interactions and gauge models," *Phys. Rev.* **D11** (1975) 1137–1154. doi:10.1103/PhysRevD.11.1137.
- [723] A. Datta, M. Guchait, and D. P. Roy, "Prospect of heavy right-handed neutrino search at SSC / LHC energies," *Phys. Rev.* **D47** (1993) 961–966, arXiv:hep-ph/9208228.
- [724] O. Panella, M. Cannoni, C. Carimalo, and Y. N. Srivastava, "Signals of heavy Majorana neutrinos at hadron colliders," *Phys. Rev.* **D65** (2002) 035005, arXiv:hep-ph/0107308.
- [725] A. Ferrari et al., "Sensitivity study for new gauge bosons and right-handed Majorana neutrinos in p p collisions at $s = 14$ TeV," *Phys. Rev.* **D62** (2000) 013001.
- [726] S. Gninenko and N. Krasnikov, "Limits on the magnetic moment of sterile neutrino and two-photon neutrino decay," *Phys. Lett.* **B450** (1999) 165–172, arXiv:hep-ph/9808370.
- [727] S. Gninenko, M. Kirsanov, N. Krasnikov, and V. Matveev, "Prospects of detecting massive isosinglet neutrino at LHC in the CMS detector," arXiv:hep-ph/0301140.
- [728] **Particle Data Group** Collaboration, K. Hagiwara et al., "Review of particle physics," *Phys. Rev.* **D66** (2002) 010001.
- [729] G. Barenboim, J. Bernabeu, J. Prades, and M. Raidal, "Constraints on the W(R) mass and CP-violation in left-right models," *Phys. Rev.* **D55** (1997) 4213–4221, arXiv:hep-ph/9611347.
- [730] **DØ** Collaboration, S. Abachi et al., "Search for Right-Handed W Bosons and Heavy W ν in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **76** (1996) 3271–3276, arXiv:hep-ex/9512007.
- [731] **CTEQ** Collaboration, J. Botts et al., "CTEQ parton distributions and flavor dependence of sea quarks," *Phys. Lett.* **B304** (1993) 159–166, arXiv:hep-ph/9303255. doi:10.1016/0370-2693(93)91416-K.
- [732] **CMS** Collaboration, M. Kirsanov,  "Detection of Heavy Majorana Neutrinos and Right-Handed Bosons," *CMS AN ~~NOTE~~: No AN's 2006-04* (2006).
- [733] E. Lehmann, "A Theory of some Multiple Decision Problems," *The Annals of Mathematical Statistics* **28** 1 (1957).

- [734] R. O'Neill and G. Wetherill, "The Present State of Multiple Comparison Methods," *Journal of the Royal Statistical Society, Series B (Methodological)* **33** 2 (1971).
- [735] J. Hauser, "Search for New Physics in Tails of Distributions," 2006. Private communications, paper in preparation.
- [736] DØ Collaboration, V. M. Abazov et al., "Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using kinematic characteristics of lepton + jets events," *Phys. Lett.* **B626** (2005) 45–54, arXiv:hep-ex/0504043. doi:10.1016/j.physletb.2005.08.104.
- [737] B. Abbott et al., "Search for New Physics in $e\mu X$ Data at DØ Using Sleuth: A Quasi-Model-Independent Search Strategy for New Physics," *Phys. Rev.* **D62** (2000). doi:10.1103/PhysRevD.62.092004.
- [738] B. Abbott et al., "A Quasi-Model-Independent Search for New Physics at Large Transverse Momentum," *Phys. Rev.* **D64** (2001) arXiv:hep-ex/0011067.
- [739] C. Finley et al., "On the evidence for clustering in the arrival directions of AGASA's ultrahigh energy cosmic rays," *Astroparticle Physics* **21** (2004).
- [740] S. Frixione, P. Nason, and B. R. Webber, "Matching NLO QCD and parton showers in heavy flavour production," *JHEP* **08** (2003) 007, arXiv:hep-ph/0305252.
- [741] L. Lonnblad, "Correcting the colour-dipole cascade model with fixed order matrix elements," *JHEP* **05** (2002) 046, arXiv:hep-ph/0112284.
- [742] S. Frixione and B. R. Webber, "Matching NLO QCD computations and parton shower simulations," *JHEP* **06** (2002) 029, arXiv:hep-ph/0204244.
- [743] F. Krauss, "Matrix elements and parton showers in hadronic interactions," *JHEP* **08** (2002) 015, arXiv:hep-ph/0205283.
- [744] S. Mrenna and P. Richardson, "Matching matrix elements and parton showers with HERWIG and PYTHIA," *JHEP* **05** (2004) 040, arXiv:hep-ph/0312274.
- [745] J. Butterworth, S. Butterworth, B. Waugh, W. Stirling, and M. Whalley, "The CEDAR project," arXiv:hep-ph/0412139.
- [746] LEP Collaboration, "A combination of preliminary electroweak measurements and constraints on the standard model," arXiv:hep-ex/0412015.
- [747] D. R. Stump, "A new generation of CTEQ parton distribution functions with uncertainty analysis,". Prepared for 31st International Conference on High Energy Physics (ICHEP 2002), Amsterdam, The Netherlands, 24-31 Jul 2002.
- [748] W. J. Stirling, A. D. Martin, R. G. Roberts, and R. S. Thorne, "MRST parton distributions," *AIP Conf. Proc.* **747** (2005) 16–21.
- [749] M. Botje, "A QCD analysis of HERA and fixed target structure function data," *Eur. Phys. J.* **C14** (2000) 285–297, arXiv:hep-ph/9912439.
- [750] S. I. Alekhin, "Global fit to the charged leptons DIS data: $\alpha(s)$, parton distributions, and high twists," *Phys. Rev.* **D63** (2001) 094022, arXiv:hep-ph/0011002.

- [751] G. Altarelli and G. Parisi, "Asymptotic Freedom in Parton Language," *Nucl. Phys.* **B126** (1977) 298.
- [752] L. Lonnblad, "ARIADNE version 4: A Program for simulation of QCD cascades implementing the color dipole model," *Comput. Phys. Commun.* **71** (1992) 15–31.
- [753] CDF Collaboration, F. Abe et al., "Evidence for color coherence in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev.* **D50** (1994) 5562–5579.
doi:10.1103/PhysRevD.50.5562.
- [754] DØ Collaboration, B. Abbott et al., "Color coherent radiation in multijet events from $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Lett.* **B414** (1997) 419–427.
doi:10.1016/S0370-2693(97)01190-8.
- [755] I. G. Knowles et al., "QCD Event Generators," arXiv:hep-ph/9601212.
- [756] ALEPH Collaboration, D. Buskulic et al., "Properties of hadronic Z decays and test of QCD generators," *Z. Phys.* **C55** (1992) 209–234.
- [757] DELPHI Collaboration, P. Abreu et al., "Tuning and test of fragmentation models based on identified particles and precision event shape data," *Z. Phys.* **C73** (1996) 11–60. doi:10.1007/s002880050295.
- [758] OPAL Collaboration, G. Alexander et al., "A Comparison of b and (u d s) quark jets to gluon jets," *Z. Phys.* **C69** (1996) 543–560.
- [759] B. Andersson, G. Gustafson, and B. Soderberg, "A GENERAL MODEL FOR JET FRAGMENTATION," *Z. Phys.* **C20** (1983) 317.
- [760] ALEPH Collaboration, A. Heister et al., "Study of the fragmentation of b quarks into B mesons at the Z peak," *Phys. Lett.* **B512** (2001) 30–48, arXiv:hep-ex/0106051.
- [761] OPAL Collaboration, G. Abbiendi et al., "Inclusive analysis of the b quark fragmentation function in Z decays at LEP. ((B)), " *Eur. Phys. J.* **C29** (2003) 463–478, arXiv:hep-ex/0210031.
- [762] SLD Collaboration, K. Abe et al., "Measurement of the b-quark fragmentation function in Z^0 decays," *Phys. Rev.* **D65** (2002) 092006.
doi:10.1103/PhysRevD.65.092006.
- [763] M. G. Bowler, "e+ e- Production of Heavy Quarks in the String Model," *Zeit. Phys.* **C11** (1981) 169.
- [764] C. Peterson, D. Schlatter, I. Schmitt, and P. M. Zerwas, "SCALING VIOLATIONS IN INCLUSIVE e+ e- ANNIHILATION SPECTRA," *Phys. Rev.* **D27** (1983) 105.
- [765] V. Kartvelishvili, A. Likhoded, and V. Petrov, "On the fragmentation functions of heavy quarks into hadrons," *Phys. Lett.* **B78** (1978) 615.
- [766] E. Norrbin and T. Sjostrand, "Production and hadronization of heavy quarks," *Eur. Phys. J.* **C17** (2000) 137–161, arXiv:hep-ph/0005110.
- [767] R. Engel and J. Ranft, "Color singlet exchange between jets and the PHOJET Monte Carlo," *Nucl. Phys. Proc. Suppl.* **75A** (1999) 272–274.
doi:10.1016/S0920-5632(99)00263-7.

- [768] CDF Collaboration, T. Affolder et al., "Charged jet evolution and the underlying event in proton anti-proton collisions at 1.8 TeV," *Phys. Rev.* **D65** (2002) 092009. doi:10.1103/PhysRevD.65.092009.
- [769] P. Nason et al., "Bottom production," arXiv:hep-ph/0003142.
- [770] T. Sjostrand and P. Z. Skands, "Transverse-momentum-ordered showers and interleaved multiple interactions," *Eur. Phys. J.* **C39** (2005) 129–154, arXiv:hep-ph/0408302. doi:10.1140/epjc/s2004-02084-y.
- [771] S. Sapeta and K. Golec-Biernat, "Total, elastic and diffractive cross sections at LHC in the Miettinen-Pumplin model," *Phys. Lett.* **B613** (2005) 154–161, arXiv:hep-ph/0502229.
- [772] D. J. Lange, "The EvtGen particle decay simulation package," *Nucl. Instrum. Meth.* **A462** (2001) 152–155.
- [773] P. Bartalini, "Supporting Monte Carlo Generators at the LHC,". Prepared for the 10th International Conference on B Physics at Hadron Machines (BEAUTY 2005), Assisi, Perugia, Italy, 20- 24 Jun 2005. Nuclear Physics B - Proceedings Supplements M. Biasini and S. Erhan editors. Elsevier Publishing.
- [774] Z. Sullivan, "Fully differential W production and decay at next-to-leading order in QCD," *Phys. Rev.* **D66** (2002) 075011, arXiv:hep-ph/0207290.
- [775] M. A. Dobbs et al., "Les Houches guidebook to Monte Carlo generators for hadron collider physics," arXiv:hep-ph/0403045.
- [776] "HIJING Web Site." Located at <http://www-nsdth.lbl.gov/~xnwang/hijing/index.html>.
- [777] T. Gleisberg et al., "Event generator for the LHC," arXiv:hep-ph/0508315.
- [778] E. Boos et al., "Generic user process interface for event generators," arXiv:hep-ph/0109068.
- [779] V. Karimaki et al., "The 's Guide for CMS Physics Generators Interface CMKIN Version 4," *CMS IN : Internal Note 2004-052* (2004).
- [780] I. P. Lokhtin and A. M. Snigirev, "Nuclear geometry of jet quenching," *Eur. Phys. J.* **C16** (2000) 527–536, arXiv:hep-ph/0004176. doi:10.1007/s100520000437.
- [781] I. P. Lokhtin and A. M. Snigirev, "Fast simulation of jet quenching in ultrarelativistic heavy ion collisions," arXiv:hep-ph/0406038. SINP MSU-13/752.
- [782] I. P. Lokhtin and A. M. Snigirev, "Fast simulation of flow effects in central and semi-central heavy ion collisions at LHC," arXiv:hep-ph/0312204.
- [783] CDF Collaboration, R. D. Field, "The underlying event in hard scattering processes," *eConf* **C010630** (2001) P501, arXiv:hep-ph/0201192.
- [784] CompHEP Collaboration, E. Boos, V. Bunichev, M. Dubinin, L. Dudko, V. Edneral, V. Ilyin, A. Kryukov, V. Savrin, A. Semenov, and A. Sherstnev. Web site at <http://theory.sinp.msu.ru/comphep>. See also [47].

- [785] I. P. Lokhtin, L. I. Sarycheva, and A. M. Snigirev, "The method for analysing jet azimuthal anisotropy in ultrarelativistic heavy ion collisions," *Phys. Lett.* **B537** (2002) 261–267, arXiv:hep-ph/0203144. doi:10.1016/S0370-2693(02)01913-5.
- [786] J. H. Holland, "Adaptation in natural and artificial systems". The University of Michigan Press, Ann Arbor, 1975.
- [787] D. E. Goldberg, "Genetic algorithms in search, optimization and machine learning". Addison Wesley, 1989.
- [788] S. Abdullin, "Genetic Algorithm for SUSY Trigger Optimization In CMS Detector At LHC," *NIM A* **502** (2003) 693–695. doi:10.1016/S0168-9002(03)00546-1.
- [789] D. Acosta et al., "CMS Discovery Potential for mSUGRA in Single Muon Events with Jets and Large Missing Transverse Energy in pp Collisions at $\sqrt{s} = 14$ TeV," *CMS Note* **XXXXX**: in preparation (2006).
- [790] D. Acosta et al., "CMS Discovery Potential for mSUGRA in Same Sign Di-muon Events with Jets and Large Missing Transverse Energy in pp collisions at $\sqrt{s} = 14$ TeV," *CMS Note* **XXXXX**: in preparation (2006).
- [791] TOTEM Collaboration, "TOTEM Technical Design Report," *CERN/LHCC* **2004-002** (2004).
- [792] TOTEM Collaboration, "Addendum to the TOTEM-TDR," *CERN/LHCC* **2004-020** (2004).
- [793] V. Avati and K. Österberg, "TOTEM forward measurements: leading proton acceptance," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at <http://www.desy.de/~heralhc/proceedings/wg4avati.pdf>.
- [794] J. Kalliopuska et al., "TOTEM forward measurements: exclusive central diffraction," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at http://www.desy.de/~heralhc/proceedings/wg4pXp_heralhc.pdf.
- [795] "The MAD-X Program, Methodical Accelerator Design." Information available at <http://www.cern.ch/mad>.
- [796] CMS/TOTEM Collaboration, "CMS/TOTEM document on diffractive and forward physics,".
- [797] M. Arneodo et al., "Diffractive Higgs: CMS/TOTEM Level-1 Trigger Studies," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at <http://www.desy.de/~heralhc/proceedings/wg4arneodo.pdf>.
- [798] R. Croft, "**XXXXX**: in preparation". PhD thesis, University of Bristol, 2006.
- [799] F. Oljemark, "First level triggering of diffractively produced low-mass Higgs at The Large Hadron Collider," Master's thesis, University of Helsinki, 2006.
- [800] G. Bruni et al., "Leading proton production in *ep* and *pp*, experiments: how well do high-energy physics Monte Carlos reproduce the data?," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at <http://www.desy.de/~heralhc/proceedings/wg5lps.pdf>.

- [801] F. Ferro, "Diffractive Higgs in CMS/TOTEM: study of L1 trigger conditions from T1 and T2," *TOTEM Note* **04-2005** (2005).
- [802] B. Cox and J. Forshaw, "Herwig for diffractive interactions," *Comput. Phys. Commun.* **144** (2002) 104–110. doi:10.1016/S0010-4655(01)00467-2.
- [803] J. C. Collins, "Light-cone variables, rapidity and all that," arXiv:hep-ph/9705393.
- [804] M. G. Albrow and A. Rostovtsev, "Searching for the Higgs at hadron colliders using the missing mass method," arXiv:hep-ph/0009336.
- [805] M. G. Albrow, "Double pomeron physics at the LHC," *AIP Conf. Proc.* **792** (2005) 509–514, arXiv:hep-ex/0507095.
- [806] J. Branson and E. Trepagnier, "Weighting Bunch Crossings to Calculate Rates in the Presence of Pileup," CMS IN **NAME: No IN's**.
- [807] C. Delaere et al., "HLT Steering Code Prototype," CMS IN **NAME: Internal Note 2003/052** (2003).
- [808] S. Var **NAME** et al., " $2 \times 10^3 \text{ cm}^{-2}\text{s}^{-1}$ Trigger Tables for PTDR, volume 2," CMS Internal Note **NAME: should be Note**.