


# Discovery of the Higgs Boson at the LHC

**Roger Wolf**

17. June 2014

INSTITUTE OF EXPERIMENTAL PARTICLE PHYSICS (IEKP) – PHYSICS FACULTY





BREAKTHROUGH  
of the YEAR  
The **HIGGS**  
**BOSON**

Direct contributions of IEKP:

- Construction and upgrade of silicon track detector.
- Physics with top quarks (→pairwise & single top).
- **Higgs discovery!**

# The Large Hadron Collider

- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec

# The Large Hadron Collider

Energy radiated off per rotation cycle:

$$P = \frac{e^2}{6\pi\epsilon_0 c} |\vec{\beta}|^2 \gamma^4 = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \gamma^4 = \frac{e^4}{6\pi\epsilon_0 \rho^2} \frac{E^2 B^2}{m^4}$$

$$P(p|_{m_p=1 \text{ GeV}}) = 280 \mu\text{W}$$

$$P(e|_{m_e=0.511 \text{ MeV}}) = 450 \text{ kW}$$

- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec



# The Large Hadron Collider


Energy radiated off per rotation cycle:

$$P = \frac{e^2}{6\pi\epsilon_0 c} |\vec{\beta}|^2 \gamma^4 = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \gamma^4 = \frac{e^4}{6\pi\epsilon_0 \rho^2} \frac{E^2 B^2}{m^4}$$


$$P(p|_{m_p=1 \text{ GeV}}) = 280 \mu\text{W}$$

$$P(e|_{m_e=0.511 \text{ MeV}}) = 450 \text{ kW}$$

- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec



- 8.3 T
- 11.8 kA
- 160 cyc



- Energy density  
500 kJ/m
- Tension  
200'000 t/m

# The Large Hadron Collider


Energy radiated off per rotation cycle:

$$P = \frac{e^2}{6\pi\epsilon_0 c} |\vec{\beta}|^2 \gamma^4 = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \gamma^4 = \frac{e^4}{6\pi\epsilon_0 \rho^2} \frac{E^2 B^2}{m^4}$$


$$P(p|_{m_p=1 \text{ GeV}}) = 280 \mu\text{W}$$

$$P(e|_{m_e=0.511 \text{ MeV}}) = 450 \text{ kW}$$


- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec



- 8.3 T
- 11.8 kA
- 160 cyc



- Energy density 500 kJ/m
- Tension 200'000 t/m



# The Large Hadron Collider




Energy radiated off per rotation cycle:

$$P = \frac{e^2}{6\pi\epsilon_0 c} |\vec{\beta}|^2 \gamma^4 = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \gamma^4 = \frac{e^4}{6\pi\epsilon_0 \rho^2} \frac{E^2 B^2}{m^4}$$


$$P(p|_{m_p=1 \text{ GeV}}) = 280 \mu\text{W}$$

$$P(e|_{m_e=0.511 \text{ MeV}}) = 450 \text{ kW}$$



- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec





- 8.3 T
- 11.8 kA
- 160 cyc




- Energy density 500 kJ/m
- Tension 200'000 t/m



# Key demands on Experiments





Vertex  
identification:






# Key demands on Experiments




Vertex identification:



Momentum determination:



$$\vec{p} = e \cdot \vec{r} \times \vec{B}$$

$$\frac{\delta p}{p} = \frac{\delta B}{erB} \oplus \frac{\delta r}{erB}$$




D. Barnum, CERN, February 2009

# Key demands on Experiments



Vertex identification



Momentum determination:

$$\vec{p} = e \cdot \vec{r} \times \vec{B}$$

$$\frac{\delta p}{p} = \frac{\delta B}{e r B} \oplus \frac{\delta r}{e r B}$$


Energy determination:

- Energy resolution
- Stopping power


# The Large Scale Solution (ATLAS)

- Magnet field (solenoid): 2.6 T (inside calorimeter)
- Magnet field (toroid):  $\sim 4$  T (outside calorimeter)
- Tracker: Si/multi-wire chambers
- ECAL/HCAL: LAr (varying granularity)

- Length : 45 m
- Diameter : 22 m
- Weight : 7'000 t



Magnet Field:




Courtesy of CERN/Maximilien Brice

# The Compact Solution (CMS)


- Magnet field: 3.8 T (outside calorimeter)
- Tracker: Si ( $\delta p/p = 0.5\%$  for a 10 GeV track)
- ECAL: PbWO<sub>4</sub> ( $\delta E/E = 1\%$  for a 30 GeV  $e/\gamma$ ,  $X_0 = 28$ )
- HCAL: Sampling (brass scintillator,  $\delta E/E = 10\%$  for a 100 GeV  $\pi^{+/-}$ ,  $\lambda_i = 10$ )

- Length : 21 m
- Diameter : 16 m
- Weight : 12'500 t

Silicon Tracker:



Electromagnetic Calo:




# Worldwide Distribution of Data

- Collaborators:  $\mathcal{O}(3'000)$
- Institutes:  $\mathcal{O}(200)$
- Countries:  $\mathcal{O}(20)$

- Recorded events:  $\mathcal{O}(10^9)$
- Amount of data:  $\mathcal{O}(10 \text{ PB}/a)$

Worldwide Grid:


Data processing in layers:




# Wanted: Higgs Boson (Dead or Alive)

If  $m_H$  is given all properties of the (SM) Higgs boson are known:


Gluon fusion




Vector boson fusion



Associated production




Production (in proton (anti-)proton collisions)




# Wanted: Higgs Boson (Dead or Alive)

If  $m_H$  is given all properties of the (SM) Higgs boson are known:


Gluon fusion




Vector boson fusion




Associated production



Production (in proton (anti-)proton collisions)



Decay



# A Long Road of Theory Developments



$$gg \rightarrow H$$

- NNLO+NNLL( $\alpha_s$ )
- NLO( $\alpha$ )
- Precision 15%

$$qq \rightarrow qqH$$

- NNLO( $\alpha_s$ )
- NLO( $\alpha$ )
- Precision 3%

$$qq \rightarrow VH$$

- NNLO( $\alpha_s$ )
- NLO( $\alpha$ )
- Precision 4%

$$tt \text{ production}$$

- NNLO+NNLL( $\alpha_s$ )
- Precision 4%

$$\text{Single top production}$$

- NNLO( $\alpha_s$ )
- Precision 4%

How this precision was obtained:

$$W + \text{additional jets}$$

- NNLO( $\alpha_s$ )
- Precision 5%

$$Z + \text{additional jets}$$

- NNLO( $\alpha_s$ )
- Precision 5%

$$WW \quad WZ \quad ZZ$$

- NLO( $\alpha_s$ )
- Precision 10%

©matt keikin




# Example: Top Quark Pair Production

$\sigma_{t\bar{t}}$ [pb]	order in $\alpha_s$	uncertainty	
		scale ( $\mu$ )	pdf
158	NLO	+23 -24	
160	NLO	+20 -21	+5 -4
164	NNLO(approx)	+5 -9	+4 -5
163	NNLO(approx)	+7 -5	+9 -9

Kleiss/Stirling '88.

Moch/Uwer '09.

Kidonakis '10.




Scale uncertainty  
(NNLO approx)  $\lesssim 5\%$

Start 10. September 2008:





Incident 19. September 2008:





- Quench in 100 dipoles.
- Set free 6t of He.
- 53 damaged superconducting magnets.

Restart 20. November 2009:



# LHC History (measured in physics measurements)



first MinBias / UE studies, particle multiplicities

first incl. b x-section, 8/nb  $\delta \sim 15\%$

first incl. jet x-section, PF jets 60/nb  $\delta \sim 20-30\%$

first incl. W/Z x-sections, 200/nb  $\delta \sim 4-6\%$ , +11% lumi

first incl.  $J/\psi$  x-section, 100/nb  $\delta \sim 20\%$

first top xsec, 3/pb  $\delta \sim 40\%$

first single top xsec, t-chan., 36/pb  $\delta \sim 36\%$

first  $m_{top}$ , 36/pb  $\Delta \sim 6.5$  GeV

first WW xsec, 36/pb  $\delta \sim 40\%$   
first limit on HWW

first  $q^*$ ,  $Z'$ ,  $W'$  limits, 3-36/pb  $> 1.6, 1.1, 1.4$  TeV

first SUSY limits, 36/pb  $\sim q, \sim g > 500-600$  GeV

first ZZ xsec, 1.1 /fb  $\delta \sim 40\%$

going more differential, e.g.  $Z/W + j,b,c$

first significant limit on  $B_s \rightarrow \mu\mu$ ,  $BR < 1.9 \times 10^{-8}$

first particle discovered by CMS:  $\Xi_b$

BSM searches continue, limits pushed

first  $> 2\sigma$  local excess at low  $m_H$ , 1.1-1.7 /fb


repeating the program at 8 TeV

a new boson is announced, 5 /fb




first spin parity analysis of the boson, 17 /fb

Feb 2014




# Rediscovery of the Top Quark ~2010/2011




- Single top over top quark pair production  $\sim 1/10$ .
- **Typical DGLAP/Regge like** high energy behavior ( $\rightarrow \log(s)$ ).


Top pair production




Single top production



# Rediscovery of the Top Quark ~2010/2011



## CMS Preliminary




# Search for the Higgs Boson 2011-2012




\* for us finding the Higgs it was  
48 years = 1,513,728,000 sec

Explanatory figure (not actual data)



# Most Important Decay Channels


More details in the seminar talks!





# Most Important Decay Channels


More details in the seminar talks!



	Channel	Resolution	S/B
$\kappa_{HVV} = \frac{2m_V^2}{v}$	$H \rightarrow \gamma\gamma$	1-2%	$\mathcal{O}(0.1)$
	$H \rightarrow ZZ$	1-2%	$\mathcal{O}( > 1)$
	$H \rightarrow WW$	20%	$\mathcal{O}(1)$
	$H \rightarrow b\bar{b}$	10%	$\mathcal{O}(0.1)$
	$H \rightarrow \tau\tau$	15%	$\mathcal{O}(0.1)$

# Most Important Decay Channels

More details in the seminar talks!




	Channel	Resolution	S/B
$\kappa_{HVV} = \frac{2m_V^2}{v}$	$H \rightarrow \gamma\gamma$	1-2%	$\mathcal{O}(0.1)$
	$H \rightarrow ZZ$	1-2%	$\mathcal{O}( > 1)$
	$H \rightarrow WW$	20%	$\mathcal{O}(1)$
$\kappa_{Hff} = \frac{m_f}{v}$	$H \rightarrow b\bar{b}$	10%	$\mathcal{O}(0.1)$
	$H \rightarrow \tau\tau$	15%	$\mathcal{O}(0.1)$

# Discovery of a new particle 4<sup>th</sup> July 2012





# Discovery of a new particle 4<sup>th</sup> July 2012


- Scratching magic  $5\sigma$  boundary.
- Discovery driven by  $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ$  (high resolution channels).
- Broad moderate excesses for  $H \rightarrow WW$  and  $H \rightarrow bb$ .
- No signal seen in  $H \rightarrow \tau\tau$ .




# Discovery of a new particle 4<sup>th</sup> July 2012




# Discovery of a new particle 4<sup>th</sup> July 2012



# Discovery of a new particle 4<sup>th</sup> July 2012




# Discovery of a new particle 4<sup>th</sup> July 2012





# Discovery of a new particle 4<sup>th</sup> July 2012



# What Happened Since Then?

- Briefly discuss **each channel and its peculiarities**.
- Go through all five decay channels and discuss **what happened to them since 4<sup>th</sup> July 2012?**
- Make 2 pit-stops:

Status July 2012:

- ICHEP summer conference (Sidney)
- Discovery (with  $\mathcal{L} \approx 10 \text{ fb}^{-1}$  @ 7 TeV & 8 TeV equal share).

# What Happened Since Then?

- Briefly discuss **each channel and its peculiarities**.
- Go through all five decay channels and discuss **what happened to them since 4<sup>th</sup> July 2012?**
- Make 2 pit-stops:

## Status July 2012:

- ICHEP summer conference (Sidney)
- Discovery (with  $\mathcal{L} \approx 10 \text{ fb}^{-1}$  @ 7 TeV & 8 TeV equal share).

## Status March 2013:

- Moriond spring conference (La Thuille)
- Preliminary results based on full dataset (w/  $\mathcal{L} \approx 25 \text{ fb}^{-1}$ ).

# What Happened Since Then?

- Briefly discuss **each channel and its peculiarities**.
- Go through all five decay channels and discuss **what happened to them since 4<sup>th</sup> July 2012?**
- Make 2 pit-stops:

## Status July 2012:

- ICHEP summer conference (Sidney)
- Discovery (with  $\mathcal{L} \approx 10 \text{ fb}^{-1}$  @ 7 TeV & 8 TeV equal share).


## Status March 2013:

- Moriond spring conference (La Thuille)
- Preliminary results based on full dataset (w/  $\mathcal{L} \approx 25 \text{ fb}^{-1}$ ).


## Status Summer 2014:

- Final publications based on full dataset (w/  $\mathcal{L} \approx 25 \text{ fb}^{-1}$ ).
- Final calibrations, alignment, more channels included, more sophisticated analysis methods applied.



# $H \rightarrow \gamma\gamma$ Decay Channel



CMS Experiment at LHC, CERN  
 Data recorded: Mon Sep 26 20:18:07 2011 CEST  
 Run/Event: 177201 / 625786854  
 Lumi section: 450



- **High mass resolution** ( $\mathcal{O}(1-2\%)$ ). Simple reconstruction and event selection.
- **Tiny signal on huge background.**
- Decay via loops:






# $H \rightarrow \gamma\gamma$ Decay Channel

Status **July 2012:**

Status **March 2013:**

Status **Summer 2014:**  
(after complete re-analysis)



$$\mu = 1.6 \pm 0.4$$

$$\sigma = 4.1(\text{obs}) \quad 2.8(\text{exp})$$

@  $m_H \approx 125 \text{ GeV}$



$$\mu = 0.8 \pm 0.2$$

$$\sigma = 3.2(\text{obs}) \quad 4.2(\text{exp})$$

$$\mu = 1.1 \pm 0.2$$


$$\sigma = 5.7(\text{obs}) \quad 5.2(\text{exp})$$

# $H \rightarrow ZZ$ Decay Channel



Summer 2014


- **High mass resolution** ( $\mathcal{O}(1-2\%)$ ). Simple reconstruction and event selection.
- **Obvious signal on small background.**
- Most important search channels:  $4\mu$   $2\mu 2e$   $4e$



March 2013

# $H \rightarrow ZZ$ Decay Channel

Status **July 2012:**




$$\mu = 0.7 \pm_{0.3}^{0.4}$$

$$\sigma = 3.2(\text{obs}) \quad 3.8(\text{exp})$$

@  $m_H \approx 125$  GeV


Status **March 2013:**



$$\mu = 0.9 \pm_{0.2}^{0.3}$$

$$\sigma = 6.7(\text{obs}) \quad 7.2(\text{exp})$$

Status **Summer 2014:**






$$\mu = 0.9 \pm_{0.2}^{0.3}$$

$$\sigma = 6.8(\text{obs}) \quad 6.7(\text{exp})$$




# $H \rightarrow WW$ Decay Channel




- High discovery potential, but bad mass resolution.

$ff$	0-jet	1-jet	2-jet(VBF)
$ff'$	0-jet	1-jet	2-jet(VBF)




# $H \rightarrow WW$ Decay Channel


Status **July 2012:**



Status **March 2013:**



Status **Summer 2014:**



$$\mu = \text{N.A.}$$

$$\sigma = 1.6(\text{obs}) \quad 2.4(\text{exp})$$

@  $m_H \approx 125 \text{ GeV}$

$$\mu = 0.8 \pm 0.2$$

$$\sigma = 4.0(\text{obs}) \quad 5.1(\text{exp})$$





$$\mu = 0.8 \pm 0.2$$

$$\sigma = 4.0(\text{obs}) \quad 5.2(\text{exp})$$

# $H \rightarrow b\bar{b}$ Decay Channel



CMS Experiment at LHC, CERN  
 Data recorded: Mon Jun 27 02:59:42 2011 CEST  
 Run/Event: 167807 / 149404739  
 Lumi section: 134  
 OrbitCrossing: 35103256 / 2259






# $H \rightarrow b\bar{b}$ Decay Channel

Status **July 2012:**

Status **March 2013:**

Status **Summer 2014:**



$$\mu = \text{N.A.}$$

$$\sigma = 0.7(\text{obs}) \quad 1.9(\text{exp})$$

@  $m_H \approx 125$  GeV


$$\mu = 1.0 \pm 0.5$$

$$\sigma = 2.1(\text{obs}) \quad 2.1(\text{exp})$$




$$\mu = 1.0 \pm 0.5$$

$$\sigma = 2.1(\text{obs}) \quad 2.1(\text{exp})$$

# $H \rightarrow \tau\tau$ Decay Channel




- $m_{\tau\tau}$  as main discriminating variable.
- Separation between irreducible  $Z \rightarrow \tau\tau$  background and  $H \rightarrow \tau\tau$  signal.



# $H \rightarrow \tau\tau$ Decay Channel


Status **July 2012:**



$\mu = \text{N.A.}$   
 $\sigma = 0(\text{obs}) \quad 1.4(\text{exp})$

@  $m_H \approx 125 \text{ GeV}$


Status **March 2013:**



$\mu = 1.1 \pm 0.4$   
 $\sigma = 2.9(\text{obs}) \quad 2.6(\text{exp})$

Treating contributions from  $H \rightarrow WW$  as background.

Status **Summer 2014:**






$\mu = 0.8 \pm 0.3$   
 $\sigma = 3.2(\text{obs}) \quad 3.7(\text{exp})$

# $H \rightarrow \tau\tau$ Decay Channel

Status **July 2012:**

Status **March 2013:**

Status **Summer 2014:**



$$\mu = \text{N.A.}$$

$$\sigma = 0(\text{obs}) \quad 1.4(\text{exp})$$

@  $m_H \approx 125$  GeV

$$\mu = 1.1 \pm 0.4$$

$$\sigma = 2.9(\text{obs}) \quad 2.6(\text{exp})$$

Treating contributions from  $H \rightarrow WW$  as background.

$$\mu = 0.8 \pm 0.3$$

$$\sigma = 3.2(\text{obs}) \quad 3.7(\text{exp})$$

# $H \rightarrow \tau\tau$ Decay Channel



Sketch of event categories for 2012, incl  $H \rightarrow \tau\tau$  only.

		0-jet	1-jet		2-jet	
$\mu\tau_h$	$p_{T^{\text{th}}} > 45 \text{ GeV}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$ boosted $p_{T^{\text{th}}} > 100 \text{ GeV}$	loose VBF tag $m_{ij} > 500 \text{ GeV}$ $ \Delta\eta_{ij}  > 3.5$	tight VBF tag (2012 only) $p_{T^{\text{th}}} > 100 \text{ GeV}$ $m_{ij} > 700 \text{ GeV}$ $ \Delta\eta_{ij}  > 4.0$
	baseline	low- $p_{T^{\text{th}}}$	low- $p_{T^{\text{th}}}$			
$e\tau_h$	$p_{T^{\text{th}}} > 45 \text{ GeV}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$ boosted	loose VBF tag	tight VBF tag (2012 only)
	baseline	low- $p_{T^{\text{th}}}$	low- $p_{T^{\text{th}}}$			
$e\mu$	$p_{T^{\text{th}}} > 35 \text{ GeV}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$		loose VBF tag	tight VBF tag (2012 only)
	baseline	low- $p_{T^{\text{th}}}$	low- $p_{T^{\text{th}}}$			
$ee, \mu\mu$	$p_{T^{\text{th}}} > 35 \text{ GeV}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$		2-jet	
	baseline	low- $p_{T^{\text{th}}}$	low- $p_{T^{\text{th}}}$			
$T_h T_h$ (8 TeV only)			boosted	highly boosted	VBF tag	
	baseline					
			$p_{T^{\text{th}}} > 100 \text{ GeV}$	$p_{T^{\text{th}}} > 170 \text{ GeV}$	$p_{T^{\text{th}}} > 100 \text{ GeV}$ $m_{ij} > 500 \text{ GeV}$ $ \Delta\eta_{ij}  > 3.5$	

- Nearly 100 exclusive event categories.
- 6 inclusive decay channels.
- Exclusive decay channels for production in association with  $Z, W$  bosons.
- On 7 TeV and 8 TeV dataset.



# Combination of $H \rightarrow \tau\tau$ & $H \rightarrow b\bar{b}$



$$\mu = 0.8 \pm 0.2$$

$$\sigma = 3.8(\text{obs}) \quad 4.4(\text{exp})$$

@  $m_H \approx 125 \text{ GeV}$

Treating contributions from  $H \rightarrow WW$  as background.

channel	significance		$\mu = \sigma/\sigma_{\text{SM}}$
	expected	observed	
$H \rightarrow \gamma\gamma$	5.2	5.7	$1.1 \pm 0.2$
$H \rightarrow ZZ$	6.7	5.7	$0.9 \pm 0.3$
$H \rightarrow WW$	5.2	4.0	$0.8 \pm 0.2$
$H \rightarrow bb$	2.1	2.1	$1.0 \pm 0.5$
$H \rightarrow \tau\tau$ <sup>(1)</sup>	3.7	3.2	$0.8 \pm 0.3$
$H \rightarrow bb, \tau\tau$ <sup>(1)</sup>	4.4	3.8	$0.8 \pm 0.2$

@  $m_H \approx 125$  GeV

(1) Treating contributions from  $H \rightarrow WW$  as background.

- **Clear evidence in all but one** of the main decay channels.
- **Observation in the high resolution channels** ( $H \rightarrow \gamma\gamma$  &  $H \rightarrow ZZ$ ).
- **Clear evidence for coupling to fermions** ( $H \rightarrow \tau\tau$ ).
- No striking surprises in loops ( $H \rightarrow \gamma\gamma$ ).

# Sneak Preview for Next Week

- We have a **clear discovery of a new particle** at  $m_H = 125$  GeV.
- Next week we will check what are the **properties of this particle**:
  - Exact **mass**?
  - Decay **width**?
  - Compatibility of **couplings** with SM?
  - **Spin and parity**?
- Remaining questions:
  - Is this **A** Higgs bosons?
  - Is this **THE** Higgs bosons?
  - Is there **MORE THAN ONE** Higgs bosons?

- Precision Electroweak Measurements at the Z Resonance.
- Search for the SM Higgs boson at LEP.
- Search for the SM Higgs boson in the di-photon final state.
- Search for the SM Higgs boson in the ZZ final state.
- Search for the SM Higgs boson in the WW final state.
- Search for the SM Higgs boson in the di-tau final state.
- Search for the SM Higgs boson in the final state with two b-quarks.
- Search for the SM Higgs boson in the di-muon final state.
- Search for the SM Higgs boson produced in association with top quarks.
- Search for a Higgs boson decaying into invisible particles.

- Search for neutral MSSM Higgs bosons in the di-tau final state.
- Search for the decay  $H \rightarrow hh$ ,  $A \rightarrow Zh$  in multilepton and photon final states.

**Seminar Dates:** Thursday 03.07. ; Tuesday 08.07. ; Thursday 10.07.

# Backup & Homework Solutions