


Discovery of the Higgs Boson at the LHC

Roger Wolf, Andrew Gilbert
25. June 2015

INSTITUTE OF EXPERIMENTAL PARTICLE PHYSICS (IEKP) – PHYSICS FACULTY





BREAKTHROUGH
of the YEAR
The **HIGGS**
BOSON

Direct contributions of IEKP:

- Construction and upgrade of silicon track detector.
- Physics with top quarks (→pairwise & single top).
- **Higgs discovery!**

The Large Hadron Collider

- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec

The Large Hadron Collider

Energy radiated off per rotation cycle:

$$P = \frac{e^2}{6\pi\epsilon_0 c} |\vec{\beta}|^2 \gamma^4 = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \gamma^4 = \frac{e^4}{6\pi\epsilon_0 \rho^2} \frac{E^2 B^2}{m^4}$$

$$P(p|_{m_p=1 \text{ GeV}}) = 280 \mu\text{W}$$

$$P(e|_{m_e=0.511 \text{ MeV}}) = 450 \text{ kW}$$

- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec



The Large Hadron Collider


Energy radiated off per rotation cycle:

$$P = \frac{e^2}{6\pi\epsilon_0 c} |\vec{\beta}|^2 \gamma^4 = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \gamma^4 = \frac{e^4}{6\pi\epsilon_0 \rho^2} \frac{E^2 B^2}{m^4}$$


$$P(p|_{m_p=1 \text{ GeV}}) = 280 \mu\text{W}$$

$$P(e|_{m_e=0.511 \text{ MeV}}) = 450 \text{ kW}$$

- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec



- 8.3 T
- 11.8 kA
- 160 cyc



- Energy density
500 kJ/m
- Tension
200'000 t/m

The Large Hadron Collider


Energy radiated off per rotation cycle:

$$P = \frac{e^2}{6\pi\epsilon_0 c} |\vec{\beta}|^2 \gamma^4 = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \gamma^4 = \frac{e^4}{6\pi\epsilon_0 \rho^2} \frac{E^2 B^2}{m^4}$$


$$P(p|_{m_p=1 \text{ GeV}}) = 280 \mu\text{W}$$

$$P(e|_{m_e=0.511 \text{ MeV}}) = 450 \text{ kW}$$


- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec



- 8.3 T
- 11.8 kA
- 160 cyc



- Energy density
500 kJ/m
- Tension
200'000 t/m



The Large Hadron Collider




Energy radiated off per rotation cycle:

$$P = \frac{e^2}{6\pi\epsilon_0 c} |\vec{\beta}|^2 \gamma^4 = \frac{e^2 c}{6\pi\epsilon_0 \rho^2} \gamma^4 = \frac{e^4}{6\pi\epsilon_0 \rho^2} \frac{E^2 B^2}{m^4}$$


$$P(p|_{m_p=1 \text{ GeV}}) = 280 \mu\text{W}$$

$$P(e|_{m_e=0.511 \text{ MeV}}) = 450 \text{ kW}$$




- Construction costs: 4.1 billion \$
- Construction time : 14 years
- Circumference : 27 km
- No of dipoles : 1232
- Power : 120 MW
- Luminosity(8TeV) : 8 nb/sec





- 8.3 T
- 11.8 kA
- 160 cyc




- Energy density 500 kJ/m
- Tension 200'000 t/m





Key demands on Experiments




Vertex
identification:



Key demands on Experiments



Vertex identification:





Momentum determination:


$$\vec{p} = e \cdot \vec{r} \times \vec{B}$$

$$\frac{\delta p}{p} = \frac{\delta B}{e r B} \oplus \frac{\delta r}{e r B}$$

Key demands on Experiments



Vertex identification



Momentum determination:

$$\vec{p} = e \cdot \vec{r} \times \vec{B}$$

$$\frac{\delta p}{p} = \frac{\delta B}{e r B} \oplus \frac{\delta r}{e r B}$$


Energy determination:

- Energy resolution
- Stopping power


The Large Scale Solution (ATLAS)

- Magnet field (solenoid): 2.6 T (inside calorimeter)
- Magnet field (toroid): ~ 4 T (outside calorimeter)
- Tracker: Si/multi-wire chambers
- ECAL/HCAL: LAr (varying granularity)

- Length : 45 m
- Diameter : 22 m
- Weight : 7'000 t



Magnet Field:




Courtesy of CERN/Maximilien Brice

The Compact Solution (CMS)


- Magnet field: 3.8 T (outside calorimeter)
- Tracker: Si ($\delta p/p = 0.5\%$ for a 10 GeV track)
- ECAL: PbWO₄ ($\delta E/E = 1\%$ for a 30 GeV e/γ , $X_0 = 28$)
- HCAL: Sampling (brass scintillator, $\delta E/E = 10\%$ for a 100 GeV $\pi^{+/-}$, $\lambda_i = 10$)

- Length : 21 m
- Diameter : 16 m
- Weight : 12'500 t

Silicon Tracker:



Electromagnetic Calo:




Worldwide Distribution of Data

- Collaborators: $\mathcal{O}(3'000)$
- Institutes: $\mathcal{O}(200)$
- Countries: $\mathcal{O}(20)$

- Recorded events: $\mathcal{O}(10^9)$
- Amount of data: $\mathcal{O}(10 \text{ PB}/a)$

Worldwide Grid:


Data processing in layers:




Wanted: Higgs Boson (Dead or Alive)

If m_H is given all properties of the (SM) Higgs boson are known:


Gluon fusion




Vector boson fusion



Associated production




Production (in proton (anti-)proton collisions)




Wanted: Higgs Boson (Dead or Alive)

If m_H is given all properties of the (SM) Higgs boson are known:


Gluon fusion




Vector boson fusion




Associated production



Production (in proton (anti-)proton collisions)



Decay



A Long Road of Theory Developments



$$gg \rightarrow H$$

- NNLO+NNLL(α_s)
- NLO(α)
- Precision 15%

$$qq \rightarrow qqH$$

- NNLO(α_s)
- NLO(α)
- Precision 3%

$$qq \rightarrow VH$$

- NNLO(α_s)
- NLO(α)
- Precision 4%

$$tt \text{ production}$$

- NNLO+NNLL(α_s)
- Precision 4%

$$\text{Single top production}$$

- NNLO(α_s)
- Precision 4%

How this precision was obtained:

$$W + \text{additional jets}$$

- NNLO(α_s)
- Precision 5%

$$Z + \text{additional jets}$$

- NNLO(α_s)
- Precision 5%

$$WW \quad WZ \quad ZZ$$

- NLO(α_s)
- Precision 10%

©matt keikin


Example: Top Quark Pair Production

$\sigma_{t\bar{t}}$ [pb]	order in α_s	uncertainty	
		scale (μ)	pdf
158	NLO	+23 -24	
160	NLO	+20 -21	+5 -4
164	NNLO(approx)	+5 -9	+4 -5
163	NNLO(approx)	+7 -5	+9 -9

Kleiss/Stirling '88.

Moch/Uwer '09.

Kidonakis '10.




Scale uncertainty
(NNLO approx) $\lesssim 5\%$

Start 10. September 2008:





Incident 19. September 2008:





- Quench in 100 dipoles.
- Set free 6t of He.
- 53 damaged superconducting magnets.

Restart 20. November 2009:




LHC History (measured in physics measurements)




- first MinBias / UE studies, particle multiplicities
- first incl. b x-section, 8/nb $\delta \sim 15\%$
- first incl. jet x-section, PF jets 60/nb $\delta \sim 20-30\%$
- first incl. W/Z x-sections, 200/nb $\delta \sim 4-6\%$, +11% lumi
- first incl. J/ψ x-section, 100/nb $\delta \sim 20\%$
- first top xsec, 3/pb $\delta \sim 40\%$
- first single top xsec, t-chan., 36/pb $\delta \sim 36\%$
- first m_{top} , 36/pb $\Delta \sim 6.5$ GeV
- first WW xsec, 36/pb $\delta \sim 40\%$ first limit on HWW
- first q^* , Z', W' limits, 3-36/pb $>1.6, 1.1, 1.4$ TeV
- first SUSY limits, 36/pb $\sim q, \sim g > 500-600$ GeV
- first ZZ xsec, 1.1 /fb $\delta \sim 40\%$
- going more differential, e.g. Z/W + j,b,c
- first significant limit on $B_s \rightarrow \mu\mu$, $\text{BR} < 1.9 \times 10^{-8}$
- BSM searches continue, limits pushed
- first $>2\sigma$ local excess at low m_H , 1.1-1.7 /fb
- repeating the program at 8 TeV
- a new boson is announced, 5 /fb
- first particle discovered by CMS: Ξ_b
- first spin parity analysis of the boson, 17 /fb



Feb 2014




Rediscovery of the Top Quark ~2010/2011




- Single top over top quark pair production $\sim 1/10$.
- **Typical DGLAP/Regge like** high energy behavior ($\rightarrow \log(s)$).


Top pair production




Single top production



Rediscovery of the Top Quark ~2010/2011



CMS Preliminary




Search for the Higgs Boson 2011-2012




* for us finding the Higgs it was
48 years = 1,513,728,000 sec

Explanatory figure (not actual data)




Most Important Decay Channels

More details in the seminar talks!



Most Important Decay Channels


More details in the seminar talks!



	Channel	Resolution	S/B
$\kappa_{HV V} = \frac{2m_V^2}{v}$	$H \rightarrow \gamma\gamma$	1-2%	$\mathcal{O}(0.1)$
	$H \rightarrow ZZ$	1-2%	$\mathcal{O}(> 1)$
	$H \rightarrow WW$	20%	$\mathcal{O}(1)$
	$H \rightarrow b\bar{b}$	10%	$\mathcal{O}(0.1)$
	$H \rightarrow \tau\tau$	15%	$\mathcal{O}(0.1)$


Most Important Decay Channels

More details in the seminar talks!



	Channel	Resolution	S/B
$\kappa_{HV V} = \frac{2m_V^2}{v}$	$H \rightarrow \gamma\gamma$	1-2%	$\mathcal{O}(0.1)$
	$H \rightarrow ZZ$	1-2%	$\mathcal{O}(> 1)$
	$H \rightarrow WW$	20%	$\mathcal{O}(1)$
$\kappa_{H f f} = \frac{m_f}{v}$	$H \rightarrow b\bar{b}$	10%	$\mathcal{O}(0.1)$
	$H \rightarrow \tau\tau$	15%	$\mathcal{O}(0.1)$

- Analysis of full 2011 dataset $\sim 5 \text{ fb}^{-1}$
- Hints start to appear...
- $H \rightarrow \gamma\gamma$:
 - 3.1σ local significance at $m_{\gamma\gamma} = 124 \text{ GeV}$
- This is high, but must be careful!
 - Searching a wide mass range – background fluctuations can appear anywhere: **Look-elsewhere effect**
- “global” significance only 1.8σ
- One of the reasons we demand 5σ for discovery...




Discovery of a new particle 4th July 2012





Discovery of a new particle 4th July 2012


- Scratching magic 5σ boundary.
- Discovery driven by $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ$ (high resolution channels).
- Broad moderate excesses for $H \rightarrow WW$ and $H \rightarrow bb$.
- No signal seen in $H \rightarrow \tau\tau$.




Discovery of a new particle 4th July 2012




Discovery of a new particle 4th July 2012



Discovery of a new particle 4th July 2012



Discovery of a new particle 4th July 2012



What Happened Since Then?

- Briefly discuss **each channel and its peculiarities**.
- Go through all five decay channels and discuss **what happened to them since 4th July 2012?**
- Make 2 pit-stops:

Status July 2012:

- ICHEP summer conference (Sidney)
- Discovery (with $\mathcal{L} \approx 10 \text{ fb}^{-1}$ @ 7 TeV & 8 TeV equal share).

What Happened Since Then?

- Briefly discuss **each channel and its peculiarities**.
- Go through all five decay channels and discuss **what happened to them since 4th July 2012?**
- Make 2 pit-stops:

Status July 2012:

- ICHEP summer conference (Sidney)
- Discovery (with $\mathcal{L} \approx 10 \text{ fb}^{-1}$ @ 7 TeV & 8 TeV equal share).

Status March 2013:

- Moriond spring conference (La Thuille)
- Preliminary results based on full dataset (w/ $\mathcal{L} \approx 25 \text{ fb}^{-1}$).

What Happened Since Then?

- Briefly discuss **each channel and its peculiarities**.
- Go through all five decay channels and discuss **what happened to them since 4th July 2012?**
- Make 2 pit-stops:

Status July 2012:

- ICHEP summer conference (Sidney)
- Discovery (with $\mathcal{L} \approx 10 \text{ fb}^{-1}$ @ 7 TeV & 8 TeV equal share).


Status March 2013:

- Moriond spring conference (La Thuille)
- Preliminary results based on full dataset (w/ $\mathcal{L} \approx 25 \text{ fb}^{-1}$).

Status Summer 2014:




- Final publications based on full dataset (w/ $\mathcal{L} \approx 25 \text{ fb}^{-1}$).
- Final calibrations, alignment, more channels included, more sophisticated analysis methods applied.

$H \rightarrow \gamma\gamma$ Decay Channel



CMS Experiment at LHC, CERN
 Data recorded: Mon Sep 26 20:18:07 2011 CEST
 Run/Event: 177201 / 625786854
 Lumi section: 450

- **High mass resolution** ($\mathcal{O}(1-2\%)$). Simple reconstruction and event selection.
- **Tiny signal on huge background.**
- Decay via loops:






$H \rightarrow \gamma\gamma$ Decay Channel

Status **July 2012:**

Status **March 2013:**

Status **Summer 2014:**
(after complete re-analysis)



$$\mu = 1.6 \pm 0.4$$

$$\sigma = 4.1(\text{obs}) \quad 2.8(\text{exp})$$

$$\mu = 0.8 \pm 0.2$$



$$\sigma = 3.2(\text{obs}) \quad 4.2(\text{exp})$$

$$\mu = 1.1 \pm 0.2$$

$$\sigma = 5.7(\text{obs}) \quad 5.2(\text{exp})$$


@ $m_H \approx 125 \text{ GeV}$

$H \rightarrow ZZ$ Decay Channel



Summer 2014


- High mass resolution ($\mathcal{O}(1-2\%)$). Simple reconstruction and event selection.
- Obvious signal on small background.
- Most important search channels: 4μ $2\mu 2e$ $4e$



March 2013

$H \rightarrow ZZ$ Decay Channel

Status **July 2012:**




$$\mu = 0.7 \pm_{0.3}^{0.4}$$

$$\sigma = 3.2(\text{obs}) \quad 3.8(\text{exp})$$

@ $m_H \approx 125$ GeV


Status **March 2013:**



$$\mu = 0.9 \pm_{0.2}^{0.3}$$

$$\sigma = 6.7(\text{obs}) \quad 7.2(\text{exp})$$




Status **Summer 2014:**



$$\mu = 0.9 \pm_{0.2}^{0.3}$$


$$\sigma = 6.8(\text{obs}) \quad 6.7(\text{exp})$$

$H \rightarrow WW$ Decay Channel




- High discovery potential, but bad mass resolution.

ff	0-jet	1-jet	2-jet(VBF)
ff'	0-jet	1-jet	2-jet(VBF)




$H \rightarrow WW$ Decay Channel


Status **July 2012:**



Status **March 2013:**



Status **Summer 2014:**



$$\mu = \text{N.A.}$$

$$\sigma = 1.6(\text{obs}) \quad 2.4(\text{exp})$$

@ $m_H \approx 125$ GeV

$$\mu = 0.8 \pm 0.2$$

$$\sigma = 4.0(\text{obs}) \quad 5.1(\text{exp})$$





$$\mu = 0.8 \pm 0.2$$

$$\sigma = 4.0(\text{obs}) \quad 5.2(\text{exp})$$

$H \rightarrow bb$ Decay Channel



CMS Experiment at LHC, CERN
 Data recorded: Mon Jun 27 02:59:42 2011 CEST
 Run/Event: 167807 / 149404739
 Lumi section: 134
 OrbitCrossing: 35103256 / 2259






$H \rightarrow b\bar{b}$ Decay Channel

Status **July 2012:**

Status **March 2013:**

Status **Summer 2014:**



$$\mu = \text{N.A.}$$

$$\sigma = 0.7(\text{obs}) \quad 1.9(\text{exp})$$

@ $m_H \approx 125 \text{ GeV}$


$$\mu = 1.0 \pm 0.5$$

$$\sigma = 2.1(\text{obs}) \quad 2.1(\text{exp})$$




$$\mu = 1.0 \pm 0.5$$

$$\sigma = 2.1(\text{obs}) \quad 2.1(\text{exp})$$

$H \rightarrow \tau\tau$ Decay Channel




- $m_{\tau\tau}$ as main discriminating variable.
- Separation between irreducible $Z \rightarrow \tau\tau$ background and $H \rightarrow \tau\tau$ signal.



$H \rightarrow \tau\tau$ Decay Channel


Status **July 2012:**



$\mu = \text{N.A.}$
 $\sigma = 0(\text{obs}) \quad 1.4(\text{exp})$

@ $m_H \approx 125 \text{ GeV}$


Status **March 2013:**



$\mu = 1.1 \pm 0.4$
 $\sigma = 2.9(\text{obs}) \quad 2.6(\text{exp})$

Treating contributions from $H \rightarrow WW$ as background.


Status **Summer 2014:**




$\mu = 0.8 \pm 0.3$
 $\sigma = 3.2(\text{obs}) \quad 3.7(\text{exp})$

$H \rightarrow \tau\tau$ Decay Channel


Status **July 2012:**



Status **March 2013:**



Status **Summer 2014:**



$\mu = \text{N.A.}$
 $\sigma = 0(\text{obs}) \quad 1.4(\text{exp})$

@ $m_H \approx 125 \text{ GeV}$

$\mu = 1.1 \pm 0.4$
 $\sigma = 2.9(\text{obs}) \quad 2.6(\text{exp})$

Treating contributions from

$\mu = 0.8 \pm 0.3$
 $\sigma = 3.2(\text{obs}) \quad 3.7(\text{exp})$

as background.



$H \rightarrow \tau\tau$ Decay Channel

Sketch of event categories for 2012, incl $H \rightarrow \tau\tau$ only.

		0-jet	1-jet		2-jet	
$\mu\tau_h$	$p_{T^{\text{th}}} > 45 \text{ GeV}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$ boosted $p_{T^{\text{th}}} > 100 \text{ GeV}$	loose VBF tag $m_{ij} > 500 \text{ GeV}$ $ \Delta\eta_{ij} > 3.5$	tight VBF tag (2012 only) $p_{T^{\text{th}}} > 100 \text{ GeV}$ $m_{ij} > 700 \text{ GeV}$ $ \Delta\eta_{ij} > 4.0$
	baseline	low- $p_{T^{\text{th}}}$	low- $p_{T^{\text{th}}}$			
$e\tau_h$	$p_{T^{\text{th}}} > 45 \text{ GeV}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$ boosted	loose VBF tag	tight VBF tag (2012 only)
	baseline	low- $p_{T^{\text{th}}}$	low- $p_{T^{\text{th}}}$			
$e\mu$	$p_{T^{\text{th}}} > 35 \text{ GeV}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$		loose VBF tag	tight VBF tag (2012 only)
	baseline	low- $p_{T^{\text{th}}}$	low- $p_{T^{\text{th}}}$			
$ee, \mu\mu$	$p_{T^{\text{th}}} > 35 \text{ GeV}$	high- $p_{T^{\text{th}}}$	high- $p_{T^{\text{th}}}$		2-jet	
	baseline	low- $p_{T^{\text{th}}}$	low- $p_{T^{\text{th}}}$			
$T_h T_h$ (8 TeV only)	baseline		boosted	highly boosted	VBF tag	
			$p_{T^{\text{th}}} > 100 \text{ GeV}$	$p_{T^{\text{th}}} > 170 \text{ GeV}$	$p_{T^{\text{th}}} > 100 \text{ GeV}$ $m_{ij} > 500 \text{ GeV}$ $ \Delta\eta_{ij} > 3.5$	

- Nearly 100 exclusive event categories.
- 6 inclusive decay channels.
- Exclusive decay channels for production in association with Z, W bosons.
- On 7 TeV and 8 TeV dataset.

Combination of $H \rightarrow \tau\tau$ & $H \rightarrow b\bar{b}$



$$\mu = 0.8 \pm 0.2$$





$$\sigma = 3.8(\text{obs}) \quad 4.4(\text{exp})$$

@ $m_H \approx 125 \text{ GeV}$

Treating contributions from $H \rightarrow WW$ as background.


$H \rightarrow \mu\mu$ and $H \rightarrow ee$

- BR ($H \rightarrow \mu\mu$): 2.2×10^{-4}
- BR ($H \rightarrow ee$): 5×10^{-9}
- Looking for a small bump on a falling background
- Set Limits:
 - BR ($H \rightarrow \mu\mu$) < 0.0016
 - BR ($H \rightarrow ee$) < 0.0019
- **Evidence for non-flavour-universality in Higgs to lepton coupling**




ttH Production


- Want to measure top quark Yukawa coupling directly...
 - Indirect measurement from loops
 - m_t (173 GeV) > m_H : no $H \rightarrow tt$ decay
 - **Leaves associated-top production**
- **Small cross section:** 130 fb (ggH is ~ 19 pb)
- Complicated analysis with many channels:
 - **Production** $2 \times t \rightarrow bW^{+/-} \rightarrow (bl^{+/-} \nu \text{ or } bj\bar{j})$
 - **Decay** $H \rightarrow \gamma\gamma, H \rightarrow b\bar{b}, H \rightarrow WW, H \rightarrow ZZ, H \rightarrow \tau\tau$
- Use a **multi-variate approach** to separate signal



$\mu = 2.8 \pm 1.0 @ 125.6 \text{ GeV}$



- $t\bar{t}H(125.6) \times 30$
- $t\bar{t} + lf$
- $t\bar{t} + c\bar{c}$
- $t\bar{t} + b$
- $t\bar{t} + b\bar{b}$
- Single t
- $t\bar{t} + W, Z$
- EWK
- Bkg. Unc.
- ◆ Data



channel	significance		$\mu = \sigma/\sigma_{\text{SM}}$
	expected	observed	
$H \rightarrow \gamma\gamma$	5.2	5.7	1.1 ± 0.2
$H \rightarrow ZZ$	6.7	5.7	0.9 ± 0.3
$H \rightarrow WW$	5.2	4.0	0.8 ± 0.2
$H \rightarrow bb$	2.1	2.1	1.0 ± 0.5
$H \rightarrow \tau\tau$ ⁽¹⁾	3.7	3.2	0.8 ± 0.3
$H \rightarrow bb, \tau\tau$ ⁽¹⁾	4.4	3.8	0.8 ± 0.2

@ $m_H \approx 125$ GeV

(1) Treating contributions from $H \rightarrow WW$ as background.

- **Clear evidence in all but one** of the main decay channels.
- **Observation in the high resolution channels** ($H \rightarrow \gamma\gamma$ & $H \rightarrow ZZ$).
- **Clear evidence for coupling to fermions** ($H \rightarrow \tau\tau$).
- No striking surprises in loops ($H \rightarrow \gamma\gamma$).

Sneak Preview for Next Week

- We have a **clear discovery of a new particle** at $m_H = 125$ GeV.
- Next week we will check what are the **properties of this particle**:
 - Exact **mass**?
 - Decay **width**?
 - Compatibility of **couplings** with SM?
 - **Spin and parity**?
- Remaining questions:
 - Is this **A** Higgs bosons?
 - Is this **THE** Higgs bosons?
 - Is there **MORE THAN ONE** Higgs bosons?

Backup & Homework Solutions